

# A design proposal of the reverse DNA polymerase (3'-5' direction)

Penghui Shi

Gansu Key Lab of Crop Improvement & Germplasm Enhancement, Gansu Provincial Key Laboratory of Aridland Crop Sciences, Gansu Agricultural University, Lanzhou 730070, China  
Email: shiph150@sina.cn

**Abstract:** DNA Polymerase is an enzyme that participates in the process of DNA replication. In general, forward DNA polymerase moves along the template strand in a 3'-5' direction, and the daughter strand is formed in a 5'-3' direction. This paper theoretically analyzes the mechanism of forward DNA polymerase, and gives the design proposal of a new reverse DNA polymerase (3'-5' direction).

**Key words:** forward DNA polymerase, reverse DNA polymerase, design proposal


## Introduction

DNA polymerase is a cellular or viral polymerase enzyme that synthesizes DNA molecules from their nucleotide building blocks. DNA polymerases are essential for DNA replication, and they also play key roles in other processes within cells, including DNA repair, genetic recombination, reverse transcription, and the generation of antibody diversity via the specialized DNA polymerase, terminal deoxynucleotidyl transferase. DNA polymerases are widely used in molecular biology laboratories, notably for the polymerase chain reaction (PCR), DNA sequencing, and molecular cloning. DNA polymerase requires a free 3' OH group for initiation of synthesis, it can synthesize in only one direction by extending the 3' end of the preexisting nucleotide chain. DNA polymerase moves along the template strand in a 3'-5' direction, This results in elongation of the newly forming strand in a 5'-3' direction. The phosphodiester bond is formed in an SN2 reaction in which the hydroxyl group at the 3' end of the primer strand attacks the  $\alpha$ -phosphoryl group of the incoming nucleoside triphosphate. The leaving group for the reaction is pyrophosphate, which arises from the release of the  $\beta$ - and  $\gamma$ -phosphates of the nucleotide substrate. Hydrolysis of pyrophosphate is the driving force for DNA synthesis [1-3].


Generally, natural DNA polymerase (forward DNA polymerase) can add free nucleotides only to the 3' end of the template strand. Hence, elongation of the newly forming strand in a 5'-3' direction. Would we be able to design a new reverse DNA polymerase (3'-5' direction)? This enzyme can synthesize new DNA strand in a 3'-5' direction. We believe that it possibly plays a vital role in modern molecular biological technology.

The known forward DNA polymerases have highly conserved structure, which means that their overall catalytic subunits vary very little from species to species, independent of their domain structures. The shape can be described as resembling a right hand with thumb, finger and palm domains (Figure 1). The palm domain appears to function in catalyzing the transfer of phosphoryl groups in the phosphoryl transfer reaction. It is composed of a  $\beta$  sheet and contains the primary elements of the catalytic site. In particular, this region of forward DNA polymerase binds two divalent metal ions that alter the chemical environment around the correctly base-paired dNTP and


the 3'OH of the primer (Figure 2). One metal ion reduces the affinity of the 3'OH for its hydrogen. This generates a 3'O<sup>-</sup> that is primed for the nucleophilic attack of the  $\alpha$ -phosphate of the incoming dNTP. The second metal ion coordinates the negative charges of the  $\beta$ - and  $\gamma$ -phosphates of the dNTP and stabilizes the pyrophosphate produced by joining the primer and the incoming nucleotide. Besides, the palm domain also monitors the accuracy of base-pairing for the most recently added nucleotides. The finger domain functions to bind the nucleotide triphosphate with the template base. Several residues located within the fingers bind to the incoming dNTP. More importantly, once a correct base pair is formed between the incoming dNTP and the template, the finger domain moves to enclose the dNTP (Figure 3). This closed form of the polymerase hand stimulates catalysis by moving the incoming nucleotide in close contact with the catalytic metal ions. Compared with the fingers and the palm, the thumb domain is not intimately involved in catalysis. Instead, the thumb interacts with the DNA that has been most recently synthesized. On the one hand, it maintains the correct position of the primer and the active site. On the other hand, the thumb helps to maintain a strong association between the DNA polymerase and its substrate. This association contributes to the ability of the DNA polymerase to add many dNTPs each time it binds a primer:template junction. Moreover, the thumb domain plays a potential role in the processivity, translocation and positioning of the DNA. Forward DNA polymerase uses a single active site to catalyze the addition of any of the four deoxynucleoside triphosphates. It accomplishes this catalytic flexibility by exploiting the nearly identical geometry of the A:T and G:C base pairs. The forward DNA polymerase monitors the ability of the incoming nucleotide to form an A:T or G:C base pair rather than detecting the exact nucleotide that enters the active site. Only when a correct base pair is formed are the 3'OH of the primer and the  $\alpha$ -phosphate of the incoming nucleoside triphosphate in the optimum position for catalysis to occur. Incorrect base-pairing leads to dramatically lower rates of nucleotide addition due to a catalytically unfavorable alignment of these substrates [4-8].



**Figure 1 Structure of the Klenow fragment with duplex DNA bound at the Polymerase active site.**



**Figure 2 Illustration of the active site of the forward DNA polymerase.** The two metal ions are held in place by interactions with two highly conserved Aspartate residues. Metal ion A primarily interacts with the 3'OH resulting in reduced association between the H and the O. This leaves a nucleophilic 3'O<sup>-</sup>. Metal ion B interacts with the triphosphates of the incoming dNTP to neutralize their negative charge. At the end of reaction, the pyrophosphate product is stabilized through similar interactions with metal ion B [10,11].




**Figure 3 The changes in forward DNA polymerase structure after the incoming nucleotide base-pairs correctly to the template DNA.** The primary change is a 40° rotation of one of the helices in the finger domain called the O-helix. In the open conformation this helix is distant from the incoming nucleotide. When the polymerase is in the closed conformation, this helix moves and makes several important interactions with the incoming dNTP. A tyrosine makes stacking interactions with the base of the dNTP and two charged residues associate with the triphosphate. The combination of these interactions positions the dNTP for catalysis mediated by the two metal ions bound to the forward DNA polymerase [10,11].

## Design proposal and discussion


On the basis of the mentioned analysis, A new reverse DNA polymerase (3'-5' direction) design proposal is proposed. This may be achieved by reforming the structure and substrate of forward DNA polymerase, the proposal includes 3 aspects as follows: (1) The reform of substrate: We uses 2'-deoxynucleoside-3'-phosphate as the reverse DNA polymerase substrate (Figure 4). (2) The reform of primer: New primers (reverse primer) are designed and synthesized by means of the new substrate, and they should have an exposed 5'OH adjacent to the single-stranded region of the template (Figure 5). (3) The reform of enzyme: The two metal ions are held in place by interactions with two highly conserved Aspartate residues to form the activity center of forward DNA polymerase. The changing in structures of substrates and primers leads to dramatically lower rates of nucleotide addition due to a catalytically unfavorable alignment of these substrates. Base on stereochemical basic principle, in order to maintain the chemical environment around the correctly base-paired dNTP and the 5'OH of the primer. we replace two aspartic acid with glutamic acid and Alpha amino malonic acid (or glutamic acid). Only in this way, the 5'OH of the primer and the  $\alpha$ -phosphate of the incoming nucleoside triphosphate in the optimum position for catalysis to occur (Figure 6).

DNA polymerase uses a single active site to catalyze the addition of any of the four deoxynucleoside triphosphates, which shows it has lower substrate specificity. Its catalytic activity largely depend on two divalent metal ions of polymerase active center, and the extensional direction of the newly forming strand rely on the free 3' OH group for initiation of synthesis, and 2'-deoxynucleoside -5'-phosphate and 2'-deoxynucleoside -3'-phosphate show a very similar spatial structure. We believe that the design proposal should be reasonable.





**Figure 4** Illustration of the reform of substrate

(a)-(b) The structure formulas of 2'-deoxynucleoside -5'-phosphate and 2'-deoxynucleoside-3'-phosphate. Unlike the general DNA substrates, new nucleoside triphosphates have three phosphoryl groups which are attached via the 3' hydroxyl of the 2'-deoxyribose, and they must have an exposed 5'OH. (c)-(d) Chemical stereoscopic structure of 2'-deoxynucleoside -5'-phosphate and 2'-deoxynucleoside -3'-phosphate. They show a very similar spatial structure.



**Figure 5 Illustration of the reform of primer**

(a) The structure formulas of forward primer. (b) The structure formulas of reverse primer. Forward primer must have a free 3'OH, and new DNA synthesis extends the 3' end of the primer. In contrast, reverse primer must have a free 5'OH, and new DNA synthesis extends the 5' end of the primer.



### **Figure 6 Illustration of the reform of the DNA polymerase active center**

The changing in structures of substrates and primers leads to dramatically lower rates of nucleotide addition due to a catalytically unfavorable alignment of these substrates. Compared with 3'OH of forward primer, the position of 5'OH of reverse primer may have been altered (size of about '-CH<sub>2</sub>-' group). A similar shift is also playing out in the phosphate group of the incoming nucleotide. In order to maintain the reverse DNA polymerase activity, it is necessary to correct the position of these metal ions. And on this basis we replace two aspartic acid with glutamic acid and Alpha amino malonic acid (or glutamic acid) within DNA polymerase active center.

### **Conclusions**

In this paper, we propose a preliminary design scheme of the reverse DNA polymerase (3'-5' direction). It can synthesize new DNA strand in a 3'-5' direction. If successful, this invention will be useful in molecular biology research. Our future target is to implement this design proposal, we believe that some details may need to be revised, but the basic idea for such a design is reasonable. Therefore, more studies will be necessary to prove its feasibility and efficiency.

### **References**

- [1] Yamtich J, Sweasy JB (2010). DNA polymerase family X: function, structure, and cellular roles. *Biochim. Biophys. Acta* **1804** (5): 1136-50.
- [2] Sutton MD, Walker GC (2001). Managing DNA polymerases: coordinating DNA replication, DNA repair, and DNA recombination. *Proc. Natl. Acad. Sci. U.S.A.* **98** (15): 8342-9.
- [3] Steitz TA (1999). DNA polymerases: structural diversity and common mechanisms. *J. Biol. Chem.* **274** (25): 17395-8.
- [4] Brautigam C.A. and Steitz T.A (1998). Structural and functional insights provided by crystal structures of DNA polymerases. *Curr. Opin. Struct. Biol.* **8**: 54-63.
- [5] Jager J. and Pata J.D (1999). Getting a grip: Polymerases and their substrate complexes. *Curr. Opin. Struct. Biol.* **9**: 21-28.
- [6] Chung DW, Zhang JA, Tan CK, Davie EW, So AG, Downey KM (1991). Primary structure of the catalytic subunit of human DNA polymerase delta and chromosomal location of the gene. *Proc. Natl. Acad. Sci. U.S.A.* **88** (24): 11197-201
- [7] Doublié S. and Ellenberger T (1998). The mechanism of action of T7 DNA polymerase. *Curr. Opin. Struct. Biol.* **8**: 704-712.
- [8] Steitz T.A (1998). A mechanism for all polymerases. *Nature* **391**: 231-232.
- [9] Beese L.S, Derbyshire. V, Steitz T.A (1993). Structure of dna polymerase I klenow fragment bound to duplex dna. *Science*.
- [10] Doublié S, Tabor S, Long A.M, Richardson C.C. and Ellenberger T (1998). *Nature* 391: 251.
- [11] James D. Watson, Tania A. Baker, Stephen P. Bell, Alexander Gann, Michael Levine and Richard Losick. 2004. *Molecular Biology of the Gene*, 5nd edition. Pearson Education, Inc, publishing as Benjamin Cummings. USA.