#### Characteristics of conservations laws of chiral-type systems

A.V. Balandin

Department of Mathematics and Mechanics N.I.Lobatchevsky Nizhny Novgorod State University 23 Gagarin ave., 603950 Nizhny Novgorod, Russia e-mail: balandin@mm.unn.ru

#### Abstract

In this note a new way to construct characteristics of conservations laws of integrable chiral-type systems is proposed. Some examples of such characteristics are considered.

### **MSC**(2010): 35Q51,37K10,37K05

**Key words**: chiral-type systems, Lax representation, characteristics of conservation laws, Killing fields.

### 1 Introduction

Chiral-type systems (see, for example, [2]) are systems of partial differential equations of the form

$$\Delta^{\alpha} \equiv U^{\alpha}_{xy} + G^{\alpha}_{\beta\gamma} U^{\beta}_{x} U^{\gamma}_{y} + Q^{\alpha} = 0.$$
 (1)

Here, Greek indices  $\alpha$ ,  $\beta$ ,  $\gamma$  range from 1 to n, and subscripts denote partial derivatives with respect to the independent variables x and y. The coefficients  $G^{\alpha}_{\beta\gamma}, Q^{\alpha}$  are assumed to be smooth functions of variables  $U^1, U^2, ..., U^n$ . The summation rule over the repeated indices is also assumed.

If system (1) is a system of Euler-Lagrange equations, then it is called a nonlinear generalized sigma model.

Following [10], recall that the characteristic of a conservation law  $L = (L_1, L_2)$  of system (1) is a set of functions  $R = \{R_\alpha\}$  such that

$$Div \ L = D_x L_1 + D_y L_2 = R_\alpha \Delta^\alpha,$$

where  $\Delta^{\alpha}$  denotes the l.h.s. of Eq. (1).

By integrable systems, we mean systems admitting a Lax representation, i.e., systems for which there exist a matrix Lie algebra  $\mathfrak{g}$  and the matrix  $\mathfrak{g}$ -valued functions  $\widetilde{A}, \widetilde{B}, S_{\alpha}$  such that the following identity is fulfilled,

$$D_y \tilde{A} - D_x \tilde{B} + [\tilde{A}, \tilde{B}] = S_\alpha \Delta^\alpha.$$
<sup>(2)</sup>

**Remark 1.** Note that the set of such matrices  $S_{\alpha}$  in the general case was investigated by M.Marvan [8], and named a characteristic element of a Lax representation.

Since system (1) does not contain  $U_{xx}^{\alpha}, U_{yy}^{\alpha}$ , the Lax matrices  $\widetilde{A}, \widetilde{B}$  satisfy the conditions  $\widetilde{A} = \widetilde{A}(U_x^{\alpha}, U^{\beta}), \ \widetilde{B} = \widetilde{B}(U_y^{\alpha}, U^{\beta}).$ 

Choose these matrices in the form

$$\widetilde{A} = A_{\alpha}U_x^{\alpha} + M, \quad \widetilde{B} = B_{\alpha}U_y^{\alpha} + N, \tag{3}$$

where  $A_{\alpha}, B_{\alpha}, M, N$  are smooth functions of the variables  $U^1, U^2, ..., U^n$  taking values in a matrix Lie algebra  $\mathfrak{g}$ .

**Remark 2**. In all cases known to the author, the Lax matrices for systems (1) are of form (3) (see, for example, [6]).

Substituting (3) into (2) and collecting the terms at  $U_{xy}^{\alpha}, U_{y}^{\alpha}, U_{x}^{\alpha}$ , we find that

$$S_{\alpha} = A_{\alpha} - B_{\alpha},\tag{4}$$

$$M_{,\alpha} = [B_{\alpha}, M], \tag{5}$$

$$N_{,\alpha} = [A_{\alpha}, N]. \tag{6}$$

Here and further, partial derivatives are denoted by comma, i.e.,  $P_{,\alpha} = \frac{\partial P}{\partial U^{\alpha}}$ .

**Remark 3**. Note that form (3) of the Lax representation and the form of the chiral-type system (1) require Eqs. (5),(6).

**Remark 4.** Note that Lax matrices (3) do not contain a spectral parameter. However, if  $M \neq 0$  or  $N \neq 0$  it can be introduced automatically by the transformation  $x \to \lambda x, y \to \frac{1}{\lambda}y$ . Now, we obtain the following Lax representation with the spectral parameter

$$\widetilde{A} = A_{\alpha}U_x^{\alpha} + \lambda M, \quad \widetilde{B} = B_{\alpha}U_y^{\alpha} + \frac{1}{\lambda}N.$$
(7)

Let  $f: \underbrace{\mathfrak{g} \times \mathfrak{g} \times ...\mathfrak{g}}_{p} \mapsto C$  be a symmetric p-linear ad-invariant form on  $\mathfrak{g}$ , i.e., for all  $x_1, x_2, ..., x_p, y \in \mathfrak{g}$  the following identity holds:

$$f([y, x_1], x_2, ..., x_p) + f(x_1, [y, x_2], x_3, ..., x_p) + ... + f(x_1, x_2, ..., [y, x_p]) = 0.$$

For p = 2 one can take as f the Killing metric of the Lie algebra  $\mathfrak{g}$ .

**Remark 5**. Let G be a matrix Lie group with the Lie algebra  $\mathfrak{g}$ . Consider a gauge transformation

$$\widetilde{A} \to T_x T^{-1} + T \widetilde{A} T^{-1}, \ \widetilde{B} \to T_y T^{-1} + T \widetilde{B} T^{-1}$$

generated by a matrix  $T \in G$ . Under such transformation, the functions  $S_{\alpha}, M, N$  are transformed according to the rule:

$$S_{\alpha} \to TS_{\alpha}T^{-1}, M \to TMT^{-1}, N \to TNT^{-1},$$

and the tensor fields obtained by substituting functions  $S_{\alpha}$ , M, N into adinvariant forms on  $\mathfrak{g}$  are invariant. Thus, it seems reasonable to assume that these tensor fields carry important information about the integrable PDE system.

The purpose of this paper is to prove the following main theorem that states that the tensor fields  $R_{\alpha}$  and  $\tilde{R}_{\alpha}$  defined by the expressions

$$R_{\alpha} = f(S_{\alpha}, \underbrace{M, \dots, M}_{p-1})), \tag{8}$$

$$\widetilde{R}_{\alpha} = f(S_{\alpha}, \underbrace{N, \dots, N}_{p-1})) \tag{9}$$

are characteristics of conservation laws of system (1).

### 2 Main theorem and examples

Denote the  $\alpha$ -th Euler operator by

$$E_{\alpha} = \sum_{J} (-D)_{J} (\frac{\partial}{\partial U_{J}^{\alpha}}),$$

where the sum extends over all multi-indices  $J = (j_1, j_2)$ .

For the sequel, we need the following technical lemma 1.

**Lemma 1** Let matrices  $\widetilde{A}$  and  $\widetilde{B}$  be of form (3) where  $A_{\alpha}, B_{\alpha}, M, N$  are smooth functions of the variables  $U^1, U^2, ..., U^n$  that take values in a matrix Lie algebra  $\mathfrak{g}$  and satisfy (5),(6). Assume that f is a symmetric p-linear ad-invariant form on  $\mathfrak{g}$ .

Then the following identities hold:

$$E_{\alpha}(f(D_y\widetilde{A} - D_x\widetilde{B} + [\widetilde{A}, \widetilde{B}], \underbrace{M, \dots, M}_{p-1})) = 0,$$
(10)

$$E_{\alpha}(f(D_y\widetilde{A} - D_x\widetilde{B} + [\widetilde{A}, \widetilde{B}], \underbrace{N, \dots, N}_{p-1})) = 0.$$
(11)

We will prove this lemma in the appendix.

**Theorem 1** Let the chiral-type system (1) admit the Lax representation of the form (2),(3) in a matrix Lie algebra  $\mathfrak{g}$ . Then for each ad-invariant symmetric p-form f of the Lie algebra  $\mathfrak{g}$ , the sets of functions (8),(9) are characteristics of conservation laws of the system (1).

### Proof.

We will consider the case of Eq. (8). The case of Eq. (9) can be proved in a similar way. One can easily verify that coefficients at the first-order derivatives  $U_x^{\alpha}, U_y^{\beta}$  in l.h.s. of Eq. (2) vanish whenever Eqs. (5),(6) are fulfilled.

Substituting (2) in the polynomial f as the first argument, and M as the remaining arguments, we obtain the relations

$$f(D_y\widetilde{A} - D_x\widetilde{B} + [\widetilde{A}, \widetilde{B}], M, ..., M) = R_\alpha [U_{xy}^\alpha + G_{\beta\gamma}^\alpha U_x^\beta U_y^\gamma + Q^\alpha].$$

The use of the lemma 1 completes the proof.

Next, we discuss invariant properties of characteristics of conservation laws.

One can readily verify that under arbitrary non-degenerate transformations of the variables  $U^1, U^2, ..., U^n$  the functions  $G^{\alpha}_{\beta\gamma}$  in (1) are transformed as coefficients of an affine connection. Therefore, we assume that  $G^{\alpha}_{\beta\gamma}$  are Christoffel symbols of an affine connection in the local coordinate system  $U^1, U^2, ..., U^n$  of a space  $V^n$ . We say that this connection is associated to system (1), and denote covariant derivatives w.r.t. this connection by  $\nabla_{\alpha}$ .

Unless otherwise indicated, we consider characteristics of conservation laws of the form  $R_{\alpha} = R_{\alpha}(U^1, U^2, ..., U^n)$  only. **Theorem 2** A set  $R_{\alpha}$  is the characteristic of a conservation law of system (1) iff the following conditions are satisfied:

1)  $R_{\alpha}$  is a Killing covector field, i.e.,

$$\nabla_{(\alpha} R_{\beta)} = 0; \tag{12}$$

2)

$$R_{\alpha}Q^{\alpha} = const; \tag{13}$$

3) the form  $\nabla_{\alpha}R_{\beta}dU^{\alpha} \wedge dU^{\beta}$  is closed, i.e.,

$$d(\nabla_{\alpha}R_{\beta}dU^{\alpha}\wedge dU^{\beta}) = 0.$$
(14)

### Proof.

As it is well known (see for example [10]), a function  $R_{\alpha}\Delta^{\alpha}$  is a divergence iff  $E_{\beta}(R_{\alpha}\Delta^{\alpha}) = 0$ .

One can readily verify that

$$E_{\alpha}(R_{\beta}\Delta^{\beta}) = K_{\alpha\beta}U_{xy}^{\beta} + \widehat{L}_{\alpha\beta\gamma}U_{x}^{\beta}U_{y}^{\gamma} + L_{\alpha\gamma}$$

where

$$K_{\alpha\beta} = \nabla_{(\alpha}R_{\beta)}, \ L_{\alpha} = (R_{\delta}Q^{\delta})_{,\alpha},$$
$$\widehat{L}_{\alpha\beta\gamma} = R_{\alpha,\beta\gamma} - (R_{\delta}G^{\delta}_{\alpha\gamma})_{,\beta} - (R_{\delta}G^{\delta}_{\beta\alpha})_{,\gamma} + (R_{\delta}G^{\delta}_{\beta\gamma})_{,\alpha}.$$
(15)

Now, one can see that conditions (12) and (13) are fulfilled. Taking into account condition (12), Eq. (15) results in

$$\widehat{L}_{\alpha\beta\gamma} = (\nabla_{\gamma}R_{\alpha})_{\beta} + (\nabla_{\beta}R_{\gamma})_{\alpha} + (\nabla_{\alpha}R_{\beta})_{\gamma}.$$

This completes the proof.

**Corollary 1** Let the chiral-type system (1) admit the Lax representation of the form (2),(3) in a compact semisimple Lie algebra  $\mathfrak{g}$ , where:

1) not all of the coefficients  $Q^{\alpha}$  vanish;

2) the number n of components of the system (1) and the dimension of the Lie algebra  $\mathfrak{g}$  satisfy the condition:  $n \leq \dim \mathfrak{g} \leq n+1$ ;

3) the matrices  $S_{\alpha}$  are linearly independent.

Then the system (1) admits at least one non-trivial conservation law whose characteristic  $R_{\alpha}$  depends on  $U^1, U^2, ..., U^n$  and satisfies the conditions (12),(14), as well as  $R_{\alpha}Q^{\alpha} = 0$ .

### Proof.

First, using the Lax representation (2), we find that  $2S_{\alpha}Q^{\alpha} = [M, N]$ . In view of linearly independence of  $S_{\alpha}$  we obtain that M and N are linearly independent too. Let the 2-form f be the Killing metric on the Lie algebra  $\mathfrak{g}$ . Taking into account non-degeneracy and positivity of the form f, as well as and considerations of the dimensionality, we conclude that at least one of the fields  $R_{\alpha} = f(S_{\alpha}, M), \tilde{R}_{\alpha} = f(S_{\alpha}, N)$  is non-vanishing. Now, the use of theorem 2 and the observation that  $R_{\alpha}Q^{\alpha} = f(Q^{\alpha}S_{\alpha}, M) = f(\frac{[M,N]}{2}, M) = 0$ complete the proof.

**Remark 6.** In general it seems hard to say how many nonzero characteristics can be constructed by the proposed method. In the following example 1, to each semisimple Lie algebra of rank r, we assign a system for which at least r linearly independent characteristics can be constructed using theorem 1.

**Example 1**: Systems associated to semisimple Lie groups [6],[1]. Let  $U^1, U^2, ... U^n$  be local coordinates on a semisimple Lie group G of rank  $r, \mathfrak{g}$  be Lie algebra of G, and  $C^{\alpha}_{\beta\gamma}$  the structure constants of the group G corresponding to a basis of left-invariant differential forms  $\Theta^{\alpha} = T^{\alpha}_{\beta} dU^{\beta}$ . Thus, the following equations are fulfilled

$$d\Theta^{\alpha} = C^{\alpha}_{\beta\gamma}\Theta^{\gamma} \wedge \Theta^{\beta},$$
  

$$T^{\alpha}_{[\mu,\nu]} = C^{\alpha}_{\beta\gamma}T^{\beta}_{\mu}T^{\gamma}_{\nu}.$$
(16)

Assume that the matrices  $\widetilde{A} = (\widetilde{A}^{\alpha}_{\beta}), \widetilde{B} = (\widetilde{B}^{\alpha}_{\beta})$  are of the form

$$\widetilde{A}^{\alpha}_{\beta} = 2C^{\alpha}_{\beta\gamma}(T^{\gamma}_{\lambda}U^{\lambda}_{x} + M^{\gamma}), \quad \widetilde{B}^{\alpha}_{\beta} = 2C^{\alpha}_{\beta\gamma}N^{\gamma},$$

where  $M^{\alpha} = const$  and  $N^{\alpha}$  satisfy the equations

$$N^{\alpha}_{,\gamma} = 2C^{\alpha}_{\beta\delta}T^{\delta}_{\gamma}N^{\beta}.$$
 (17)

Using Eqs. (16), one can find that the system (17) is completely integrable, and its solutions are determined by the set of n constants  $N_0 = (n_1, n_2, ..., n_n)$ .

Now, one can verify that these matrices form the Lax representation of chiral-type system (1), where

$$G^{\sigma}_{\mu\nu} = \tilde{T}^{\sigma}_{\lambda} T^{\lambda}_{\mu,\nu}, \ Q^{\sigma} = 2\tilde{T}^{\sigma}_{\lambda} C^{\lambda}_{\mu\nu} N^{\nu} M^{\mu},$$

and  $\tilde{T}$  denotes the inverse matrix of T. Thus, such integrable systems are determined by two elements  $M, N_0$  of the algebra  $\mathfrak{g}$ . Note that characteristic elements  $S_{\alpha}$  of the Lax representation for these systems are the matrices  $S_{\lambda} = ||2C^{\alpha}_{\beta\gamma}T^{\gamma}_{\lambda}||$ . These matrices  $S_{\lambda}$  form a basis of the matrix algebra  $\mathfrak{g}$ .

Let  $I(\mathfrak{g})$  be the ring of *ad*-invariant polynomials on  $\mathfrak{g}$ . The known theorem (see, for example, [3]) states that the ring  $I(\mathfrak{g})$  is generated by precisely r algebraically independent homogeneous polynomials. Denote the symmetric *ad*-invariant forms on  $\mathfrak{g}$  obtained by the polarization of the generating polynomials by  $f_1, f_2, ..., f_r$ , of degrees  $p_1, p_2, ..., p_r$ , respectively.

Fix  $M \in \mathfrak{g}$  and consider the linear functional  $f_i$  on  $\mathfrak{g}$  defined by the equality

$$f_i(X) = f_i(X, \underbrace{M, M, \dots, M}_{p_i-1}), X \in \mathfrak{g}.$$

Then, by the result of Kostant [5] (see, also, [12]), the functionals  $\tilde{f}_1, \tilde{f}_2, ..., \tilde{f}_r$  are linearly independent iff M is a regular element of the algebra  $\mathfrak{g}$ .

Now, turning back to the chiral-type system, assume that M is a regular element of the algebra  $\mathfrak{g}$ .

Note that the characteristic elements  $S_{\alpha}$  of the Lax representation for these systems are the matrices  $S_{\lambda} = ||2C_{\beta\gamma}^{\alpha}T_{\lambda}^{\gamma}||$ . Such matrices  $S_{\lambda}$  form a basis of the matrix algebra  $\mathfrak{g}$ . Thus, the characteristics of conservation laws of the form  $F_i = (F_{i\alpha}) = (f_i(S_{\alpha}, \underbrace{M, M, ..., M}_{p_i-1})), (i = \overline{1, r})$  are linearly independent.

**Remark 7**. Note that the characteristics constructed using theorem 1 are characteristics of order zero, i.e., they do not depend on the derivatives, and are of the form  $F_{\alpha} = F_{\alpha}(U^1, U^2, ..., U^n)$ .

Let a chiral-type system be a system obtained from a Lagrangian of the form

$$L = h_{\alpha\beta}(U^{1}, ..., U^{n})U_{x}^{\alpha}U_{y}^{\beta} + Q(U^{1}, ..., U^{n}), \ det(h_{\alpha\beta}) \neq 0.$$

For such systems characteristics obtained using theorem 1 correspond to symmetries of the form

$$V = v^{\alpha}(U^1, U^2, ..., U^n) \frac{\partial}{\partial U^{\alpha}}.$$
(18)

Thus, the number of different characteristics, in the case of variational systems, is bounded by the number of symmetries of the form (18).

Suppose that a variational system does not admit symmetries of the form (18); then all of the characteristics  $F_{\alpha}$ ,  $\tilde{F}_{\alpha}$  vanish. For example, nonlinear Klein-Gordon equation,  $U_{xy} = f(U)$ , does not have symmetries of the form  $V = v(U)\frac{\partial}{\partial U}$ . Thus, all characteristics constructed in the proposed way will automatically vanish.

**Remark 8.** If  $\mathfrak{g}$  is  $\mathfrak{sl}(2)$  or  $\mathfrak{so}(3)$ , then one can construct first order characteristics of conservation laws similarly to theorem 1. By a direct computation analogous to the proof of theorem 1, one can verify that functions

$$Y_{\alpha} = f([S_{\alpha}, M], [S_{\beta}, M], M, ..., M)U_{x}^{\beta},$$
  
$$\widetilde{Y}_{\alpha} = f([S_{\alpha}, N], [S_{\beta}, N], M, ..., M)U_{y}^{\beta},$$
(19)

where f is a symmetric ad-invariant form on  $\mathfrak{g}$ , are the first order characteristics of conservation laws.

This construction, and theorem 1, are illustrated in examples 2,3,4 below. Example 2. Consider the Pohlmeier-Lund-Regge system [7],

$$\Delta^{1} = U_{xy}^{1} + \frac{1}{\sin U^{2}} (U_{x}^{1} U_{y}^{2} + U_{y}^{1} U_{x}^{2}) = 0,$$
  
$$\Delta^{2} = U_{xy}^{2} - \frac{\sin U^{2}}{(1 + \cos U^{2})^{2}} U_{x}^{1} U_{y}^{1} - p \sin U^{2} = 0,$$

where p is an arbitrary constant. It will be convenient to write the Lax representation of PLR system in form (2),(3), where

$$\begin{split} \widetilde{A} &= \begin{pmatrix} 0 & \lambda p - \frac{\cos U^2 U_x^1}{2 \cos^2 \frac{U^2}{2}} & -tg \frac{U^2}{2} U_x^1 \\ -(p\lambda - \frac{\cos U^2 U_x^1}{2 \cos^2 \frac{U^2}{2}}) & 0 & U_x^2 \\ tg \frac{U^2}{2} U_x^1 & -U_x^2 & 0 \end{pmatrix}, \\ \widetilde{B} &= \begin{pmatrix} 0 & -(\frac{\cos U^2}{\lambda} + \frac{U_y^1}{2 \cos^2 \frac{U^2}{2}}) & -\frac{\sin U^2}{\lambda} \\ \frac{\cos U^2}{\lambda} + \frac{U_y^1}{2 \cos^2 \frac{U^2}{2}} & 0 & 0 \\ -\frac{\sin U^2}{\lambda} & 0 & 0 \end{pmatrix}. \end{split}$$

Choose the following basis B of the Lie algebra  $\mathfrak{so}(\mathfrak{z})$ 

$$\overrightarrow{e}_{1} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \ \overrightarrow{e}_{2} = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \ \overrightarrow{e}_{3} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix};$$

then we have

$$D_{y}\widetilde{A} - D_{x}\widetilde{B} + [\widetilde{A}, \widetilde{B}] = S_{\alpha}\Delta^{\alpha} = \begin{pmatrix} 0 & tg^{2}\frac{U^{2}}{2} & -tg\frac{U^{2}}{2} \\ -tg^{2}\frac{U^{2}}{2} & 0 & 0 \\ tg\frac{U^{2}}{2} & 0 & 0 \end{pmatrix} \Delta^{1} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix} \Delta^{2}.$$

Thus, w. r. t. the basis B,

$$S_1 = (0, tg\frac{U^2}{2}, tg^2\frac{U^2}{2}), \ S_2 = (1, 0, 0),$$
$$M = (0, 0, p), \ N = (0, \sin U^2, -\cos U^2).$$

Assume that the 2-form f is the Killing metric of the Lie algebra  $\mathfrak{so}(3)$  which, w. r. t. the basis B, is  $\delta_{ij}$  up to a constant factor. Next, we obtain

$$R = (f(S_1, M), f(S_2, M)) = (ptg^2 \frac{U^2}{2}, 0),$$
$$\widetilde{R} = (f(S_1, N), f(S_2, N)) = (tg^2 \frac{U^2}{2}, 0).$$

Indeed, one can verify that the 1-form  $\phi = tg^2 \frac{U^2}{2} (U_x^1 dx - U_y^1 dy)$  is the conservation law of PLR system with the characteristic  $\widetilde{R} = \frac{1}{p}R$ .

Let us construct a characteristic of the first order. Using expressions (19) and assuming that f is the Killing form, we obtain

$$Y_1 = p^2 U_x^1 \tan^2 \frac{U^2}{2}, \ Y_2 = p^2 U_x^2, \ \widetilde{Y}_1 = U_y^1 \tan^2 \frac{U^2}{2}, \ \widetilde{Y}_2 = U_y^2.$$

One can verify that Y and  $\tilde{Y}$  are the characteristics of the following conservation laws:  $p^2[(U_x^1)^2 \tan^2 \frac{U^2}{2} + (U_x^2)^2]dx - 2p^2 \cos U^2 dy$  and  $[(U_x^1)^2 \tan^2 \frac{U^2}{2} + (U_y^2)^2]dy - 2\cos U^2 dx$ , respectively.

**Remark 9.** It can be proved that PLR-system admits only one conservation law whose characteristic depends on  $U^1, U^2$ . It would be interesting to investigate in which cases of integrable chiral-type systems all of the characteristics depending on  $U^1, U^2, ..., U^n$  can be found by using expression (8),(9).

**Example 3**. Consider the 3-component system

$$U_{xy}^{1} + U_{x}^{3}U_{y}^{1}ctgU^{3} - \frac{1}{\sin U^{3}}U_{y}^{3}U_{x}^{2} + = 0,$$
  
$$U_{xy}^{2} + U_{y}^{3}U_{x}^{2}ctgU^{3} - \frac{1}{\sin U^{3}}U_{x}^{3}U_{y}^{1} = 0,$$
  
$$U_{xy}^{3} + U_{y}^{1}U_{x}^{2}\sin U^{3} - p\sin U^{3} = 0,$$

where p is an arbitrary constant. This system admits the Lax representation of the form (2),(3), where [1]:

$$\widetilde{A} = \begin{pmatrix} 0 & i\lambda M^3 & -i\lambda M^2 \\ -\lambda M^3 & 0 & i\lambda M^1 \\ i\lambda M^2 & -i\lambda M^1 & 0 \end{pmatrix},$$
  
$$\widetilde{B} = \begin{pmatrix} 0 & \frac{i}{\lambda} - (\cos U^3 U_y^1 + U_y^2) & -b_{31} \\ -\frac{i}{\lambda} + (\cos U^3 U_y^1 + U_y^2) & 0 & b_{23} \\ b_{31} & -b_{23} & 0 \end{pmatrix},$$
  
$$M^1 = p \sin U^3 \sin U^2, \ M^2 = -p \sin U^3 \cos U^2, \ M^3 = p \cos U^3,$$
  
$$b_{31} = \sin U^3 \cos U^2 U_y^1 - \sin U^2 U_y^3, \ b_{23} = -\cos U^2 U_y^3 - \sin U^2 \sin U^3 U_y^1$$

Consider the same basis B and the same 2-form f as in example 1. Then, we find  $S_1 = (\sin U^2 \sin U^3, -\cos U^2 \sin U^3, \cos U^3)$ ,  $S_2 = (0, 0, 1)$ ,  $S_3 = (\cos U^2, \sin U^2, 0)$ , and  $R = \{f(S_\alpha, M)\} = \{p, p \cos U^3, 0\}, \tilde{R} = \{f(S_\alpha, M)\} = \{\cos U^3, 1, 0\}$ . Now, one can see that the set R, up to factor p, is the characteristic of the conservation law  $(\cos U^3 U_x^2 + U_x^1) dx$ , and the set  $\tilde{R}$  is the characteristic of the conservation law  $(\cos U^3 U_y^1 + U_y^2) dy$ .

Again, using (19) and assuming f to be the Killing form, one can find  $Y = (0, U_x^2 \sin^2 U^3, U_x^3), \tilde{Y} = (U_y^1 \sin^2 U^3, -U_x^2, 0)$  which are the first-order characteristics of the conservation laws  $\frac{dx}{2}[(U_x^3)^2 + (U_x^2 \sin U^3)^2] - p \cos U^3 dy$  and  $\frac{dy}{2}[(U_y^3)^2 + (U_y^1 \sin U^3)^2] - p \cos U^3 dx$ .

**Example 4**. The Lax representation for the Sine-Gordon equation is of the form:

$$\widetilde{A} = \begin{pmatrix} i\lambda & i\frac{U_x}{2} \\ i\frac{U_x}{2} & -i\lambda \end{pmatrix}, \widetilde{B} = \frac{1}{4i\lambda} \begin{pmatrix} \cos U & -i\sin U \\ i\sin U & -\cos U \end{pmatrix},$$

then  $S = \begin{pmatrix} 0 & \frac{i}{2} \\ \frac{i}{2} & 0 \end{pmatrix}$ ,  $M = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ . Now, using Eq. (19), one can obtain a characteristic of the fist order  $Y = U_x$ . Similarly, one gets  $\widetilde{Y} = U_y$ . One can see that characteristics  $Y, \widetilde{Y}$  correspond to the conservation laws  $\frac{1}{2}U_x^2dx - \cos Udy$ ,  $\frac{1}{2}U_y^2dy - \cos Udx$ , respectively.

**Remark 10.** Using (7), one can rewrite (8),(9), up to a constant factor, in the following way

$$R_{\alpha} = f(S_{\alpha}, \frac{\partial \widetilde{A}}{\partial \lambda}, \frac{\partial \widetilde{A}}{\partial \lambda}, ..., \frac{\partial \widetilde{A}}{\partial \lambda}), \quad \widetilde{R}_{\alpha} = f(S_{\alpha}, \frac{\partial \widetilde{B}}{\partial \lambda}, \frac{\partial \widetilde{B}}{\partial \lambda}, ..., \frac{\partial \widetilde{B}}{\partial \lambda})$$

It turns out that these expressions for characteristics are still valid not just for chiral-type systems, but in some cases of evolution equations. This is illustrated in examples 5-8.

In the examples below, we consider  $\mathfrak{sl}(2)$ -valued Lax representation, assume that f is the Killing metric on  $\mathfrak{sl}(2)$ , and that  $S_{\alpha}$  is a characteristic element of the Lax representation.

**Example 5**: Korteweg de Vries equation. Write the Lax representation of KdV in the form [9]:

$$D_t \widetilde{A} - D_x \widetilde{B} + [\widetilde{A}, \widetilde{B}] = i(U_t - 6UU_x + U_{xxx}) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

where

$$\widetilde{A} = i\lambda \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + i \begin{pmatrix} 0 & U \\ 1 & 0 \end{pmatrix},$$
$$\widetilde{B} = -4\lambda^2 \widetilde{A} - 2i\lambda \begin{pmatrix} -U & -iU_x \\ 0 & U \end{pmatrix} + \begin{pmatrix} U_x & iU_{xx} + 2iU^2 \\ 2iU & -U_x \end{pmatrix}$$

Then, we find  $S = i \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ ,  $f(S, \frac{\partial \tilde{A}}{\partial \lambda}) = 0$ ,  $f(S, \frac{\partial \tilde{B}}{\partial \lambda}) = 8i\lambda$  and  $f(S, \frac{\partial \tilde{B}}{\partial \lambda})(U_t - 6UU_x + U_{xxx}) = D_t(U) + D_x(-3U_x^2 + U_{xx}).$ 

**Example 6**: Zakharov-Shabat, or Ablowitz-Kaup-Newell-Segur system has the form

$$\triangle^1 = U_t - U_{xx} - 2U^2 V = 0, \ \triangle^2 = V_t + V_{xx} + 2UV^2 = 0$$

ZS-AKNS system has the Lax representation of the form [11]:

$$\widetilde{A} = \begin{pmatrix} \lambda & -V \\ U & -\lambda \end{pmatrix}, \widetilde{B} = -2\lambda \widetilde{A} - \begin{pmatrix} -UV & V_x \\ U_x & UV \end{pmatrix},$$
$$D_t \widetilde{A} - D_x \widetilde{B} + [\widetilde{A}, \widetilde{B}] = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \triangle^1 + \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix} \triangle^2.$$

Then, we obtain

$$f(S_{\alpha}, \frac{\partial \widetilde{A}}{\partial \lambda}) = 0$$

and

$$f(S_1, \frac{\partial \widetilde{B}}{\partial \lambda}) = -2V, \ f(S_2, \frac{\partial \widetilde{B}}{\partial \lambda}) = -2U.$$

Now, the following identity holds:

$$-2V\Delta^1 - 2U\Delta^2 = -D_t(2UV) - D_x(UV_x - VU_x).$$

**Example 7**: AKNS  $2 \times 2$  system of the third order [4]. Let us consider the system

$$\Delta_1 = -U_t - \frac{1}{4}\alpha_0 U_{xxx} - \frac{1}{2}\alpha_1 U_{xx} + \frac{3}{2}\alpha_0 U_x UV - \alpha_2 U_x + \alpha_1 U^2 V - 2\alpha_3 U = 0,$$
  
$$\Delta_2 = -V_t - \frac{1}{4}\alpha_0 V_{xxx} + \frac{1}{2}\alpha_1 V_{xx} + \frac{3}{2}\alpha_0 V_x UV - \alpha_2 V_x + \alpha_1 UV^2 + 2\alpha_3 V = 0,$$

where  $\alpha_i$   $(i = \overline{0,3})$  are arbitrary constants. This system admits the Lax matrices of the form

$$\widetilde{A} = \begin{pmatrix} \lambda & U \\ V & -\lambda \end{pmatrix}, \ \widetilde{B} = \begin{pmatrix} \widetilde{b}_{11} & \widetilde{b}_{12} \\ \widetilde{b}_{21} & -\widetilde{b}_{11} \end{pmatrix}.$$

Here

$$\widetilde{b}_{11} = \alpha_0 \lambda^3 + \alpha_1 \lambda^2 + \lambda (\alpha_2 - \frac{1}{2}UV) + \frac{1}{4}(UV_x - VU_x) - \frac{1}{2}\alpha_1 UV + \alpha_3,$$
  

$$\widetilde{b}_{12} = \alpha_1 U\lambda^2 + \lambda (\frac{1}{2}\alpha_0 U_x + \alpha_1 U) + \frac{1}{4}\alpha_0 (U_{xx} - 2U^2V) + \frac{1}{2}\alpha_1 U_x + \alpha_2 U,$$
  

$$\widetilde{b}_{21} = \alpha_1 V\lambda^2 + \lambda (-\frac{1}{2}\alpha_0 V_x + \alpha_1 V) + \frac{1}{4}\alpha_0 (V_{xx} - 2UV) - \frac{1}{2}\alpha_1 V_x + \alpha_2 V.$$

Then, we obtain  $f(S_{\alpha}, \frac{\partial \widetilde{A}}{\partial \lambda}) = 0$ , and the equality  $f(S_{\alpha}, \frac{\partial \widetilde{B}}{\partial \lambda})\Delta^{\alpha} = \alpha_0\lambda(V\Delta^1 + U\Delta^2) + (\alpha_1V - \frac{1}{2}\alpha_0V_x)\Delta^1 + (\alpha_1U + \frac{1}{2}\alpha_0U_x)\Delta^2$ . Now, one can verify that

$$(\alpha_0 U, \alpha_0 V); (\alpha_1 V - \frac{1}{2}\alpha_0 V_x, \ \alpha_1 U + \frac{1}{2}\alpha_0 U_x)$$

are the characteristics of conservation laws.

**Example 8**: AKNS  $2 \times 2$  system of the forth order [4]. Here we consider the system

$$\Delta_1 = -U_t - U_{xxxx} + 8UVU_{xx} + 6U_x^2V + 4UU_xV_x + 2U^2V_{xx} - 6U^3V^2 = 0,$$
  
$$\Delta_2 = -V_t + V_{xxxx} - 8UVV_{xx} - 6V_x^2U - 4VU_xV_x - 2V^2U_{xx} + 6U^3V^2 = 0.$$

The Lax matrices for this system are:

$$\widetilde{A} = \begin{pmatrix} \lambda & U \\ V & -\lambda \end{pmatrix}, \ \widetilde{B} = \begin{pmatrix} \widetilde{b}_{11} & \widetilde{b}_{12} \\ \widetilde{b}_{21} & -\widetilde{b}_{11} \end{pmatrix},$$

where

$$\begin{split} \widetilde{b}_{11} &= -8\lambda^4 + 4UV\lambda^2 + 2\lambda(VU_x - UV_x) + UV_{xx} + VU_{xx} - U_xV_x - 3U^2V^2, \\ \widetilde{b}_{12} &= -8U\lambda^3 - 4\lambda^2U_x - 2\lambda U_{xx} + 4U^2V\lambda - U_{xxx} + 6UU_xV, \\ \widetilde{b}_{21} &= -8V\lambda^3 + 4\lambda^2V_x - 2\lambda V_{xx} + 4UV^2\lambda + V_{xxx} - 6VV_xU. \end{split}$$

In this case again  $f(S_{\alpha}, \frac{\partial \tilde{A}}{\partial \lambda}) = 0$ . Collecting terms at different powers of  $\lambda$  in the expressions  $f(S_{\alpha}, \frac{\partial \tilde{B}}{\partial \lambda})$ , similarly to example 7, we find the following three characteristics of conservation laws

$$(V, U); (V_x, U_x); (-2V_{xx} + 4UV^2, -2U_{xx} + 4U^2V).$$

**Example 9.** The following evolution equation (which is nothing but MKdV up to a transformation of t),

$$\Delta = -U_t + t(U_{xxx} + 6U^2U_x) = 0,$$

admits the Lax matrices [4]

$$\widetilde{A} = \begin{pmatrix} \lambda & U \\ -U & -\lambda \end{pmatrix}, \quad \widetilde{B} = \begin{pmatrix} \widetilde{b}_{11} & \widetilde{b}_{12} \\ \widetilde{b}_{21} & -\widetilde{b}_{11} \end{pmatrix},$$

where

$$\tilde{b}_{11} = -4\lambda^3 t - 2\lambda t U^2, \ \tilde{b}_{12} = -4\lambda^2 t U - 2\lambda t U_x - t(U_{xx} + 2U^3),$$
  
 $\tilde{b}_{21} = 4\lambda^2 t U - 2\lambda t U_x + t(U_{xx} + 2U^3).$ 

One can find that  $f(S, \frac{\partial \tilde{B}}{\partial \lambda}) = 16tU\lambda$ . But in this case the equality  $E_U(tU\Delta) = 0$  fails. This example shows that formula  $f(S, \frac{\partial \tilde{B}}{\partial \lambda})$  does not always give characteristics of conservation laws.

# 3 Conclusion

In this paper we prove that expressions (8),(9) are characteristics of conservations laws for chiral-type systems admitting a Lax representation. According to the examples shown above we can propose the following conjecture. Assume that an evolutionary AKNS system, whose coefficients do not depend on t, admits a  $\mathfrak{g}$ -valued Lax representation. Note that by construction of [4], coefficients of such systems do not depend on x. Then the expressions

$$R_{\alpha} = f(S_{\alpha}, \frac{\partial \widetilde{A}}{\partial \lambda}), \ \widetilde{R}_{\alpha} = f(S_{\alpha}, \frac{\partial \widetilde{B}}{\partial \lambda}),$$

where f is the Killing metric on the Lie algebra  $\mathfrak{g}$ , and  $S_{\alpha}$  is the characteristic element of the Lax representation, form the characteristic of a conservation laws of the system under consideration.

## 4 Acknowledgment

The author is grateful to E.V. Ferapontov for attention to this work and to Y.V. Tuzov, M.I. Kuznetsov, O.N. Kashcheeva, and E.M. Makarov for discussions. The author is thankful to the referees for useful remarks.

This work was supported by the project 1014 of Russian Ministry of Education and Science.

# 5 Appendix

Here, we give proof of lemma 1.

Let us first prove Eq. (10). The case of Eq. (11) can be proved in a similar way. Denote by  $R_{1\alpha} = E_{\alpha}(f(D_y\widetilde{A}, \underbrace{M, ..., M}_{p-1})), R_{2\alpha} = E_{\alpha}(D_x\widetilde{B}, \underbrace{M, ..., M}_{p-1})), R_{3\alpha} = E_{\alpha}(f([\widetilde{A}, \widetilde{B}]), \underbrace{M, ..., M}_{p-1}))$ . It is readily verified that collecting the terms at  $U_{xy}^{\beta}, U_x^{\beta}U_y^{\gamma}, U_x^{\beta}$  in  $R_{i\alpha}, i = 1, 2, 3$ , taking into account Eqs.(5), (6), and adinvariancy of the form f, we obtain the relations

$$R_{i\alpha} = Z_{i\alpha\beta}U^{\beta}_{xy} + T_{i\alpha\beta\gamma}U^{\beta}_{x}U^{\gamma}_{y} + W_{i\alpha\beta}U^{\beta}_{x},$$

where

$$Z_{1\alpha\beta} = 2(p-1)f(A_{(\alpha}, M_{,\beta}), M, ..., M),$$

$$Z_{2\alpha\beta} = 2(p-1)f(B_{(\alpha}, M_{,\beta}), M, ..., M)$$

$$= 2(p-1)f(B_{(\alpha}, [B_{,\beta}), M], M, ..., M) = 0,$$
(20)

and

$$Z_{3\alpha\beta} = -2(p-1)f([A_{(\alpha}, B_{\beta)}], M, ..., M),$$
(21)

$$T_{1\alpha\beta\gamma} = (p-1)\{2f(A_{[\alpha,\beta]}, M_{,\gamma}, M, ..., M) + f(A_{\beta,\gamma}, M_{,\alpha}, M, ..., M) + f(A_{\alpha}, M_{,\gamma\beta}, M, ..., M) + (p-2)f(A_{\alpha}, M_{,\gamma}, M_{,\beta}, \underbrace{M, ..., M}_{p-3})\}, \quad (22)$$

$$T_{2\alpha\beta\gamma} = (p-1)\{2f(B_{[\alpha,\gamma]}, M_{,\beta}, M, ..., M) + f(B_{\gamma,\beta}, M_{,\alpha}, M, ..., M) + f(B_{\alpha}, M_{,\gamma\beta}, M, ..., M) + (p-2)f(B_{\alpha}, M_{,\gamma}, M_{,\beta}, \underbrace{M_{,\alpha}, M_{,\alpha}, M_{,\alpha}}_{p-3})\}, \quad (23)$$

$$T_{3\alpha\beta\gamma} = f([A_{\beta}, B_{\gamma}]_{,\alpha}, M, ..., M) - f([A_{\alpha}, B_{\gamma}]_{,\beta}, M, ..., M) -f([A_{\beta}, B_{\alpha}]_{,\gamma}, M, ..., M) + (p-1)\{f([A_{\beta}, B_{\gamma}], M_{,\alpha}M, ..., M) -f([A_{\alpha}, B_{\gamma}], M_{,\beta}, M, ..., M) - f([A_{\beta}, B_{\alpha}], M_{,\gamma}, M, ..., M)\},$$
(24)

$$W_{1\alpha\beta} = 0, \ W_{2\alpha\beta} = (p-1)(f(N_{,\beta}, M_{,\alpha}, M, ..., M)) - f(N_{,\alpha}, M_{,\beta}, M, ..., M)),$$
(25)

$$W_{3\alpha\beta} = f([A_{\beta}, N]_{,\alpha}, M, ..., M) - f([A_{\alpha}, N]_{,\beta}, M, ..., M) + (p-1)(f([A_{\beta}, N], M_{,\alpha}, M, ..., M)) - f([A_{\alpha}, N], M_{,\beta}, M, ..., M)).$$
(26)

We claim that all of the coefficients of the expression  $R_{1\alpha} - R_{2\alpha} + R_{3\alpha}$  vanish.

In view of relations (20),(21), and (5), we obtain the condition

$$Z_{1\alpha\beta} - Z_{2\alpha\beta} + Z_{3\alpha\beta} = 0$$

Further, using Eqs. (22)-(24) and the identities of the form  $f([A_{\beta}, B_{\gamma}]_{,\alpha}, M, ..., M) = f([A_{\beta}, B_{\gamma,\alpha}], M, ..., M) + (p-1)f(A_{\beta,\alpha}, M_{\gamma}, ..., M),$ we obtain the equations:

$$T_{1\alpha\beta\gamma} - T_{2\alpha\beta\gamma} + T_{3\alpha\beta\gamma} = (p-1)\{\underbrace{f(A_{\alpha}, M_{,\gamma\beta}, M, ..., M)}_{p-3} + (p-2)f(A_{\alpha}, M_{,\gamma}, M_{,\beta}, \underbrace{M, ..., M}_{p-3}) - 2f(B_{[\alpha,\gamma]}, M_{,\beta}, M, ..., M) - \underbrace{f(B_{\gamma,\beta}, M_{,\alpha}, M, ..., M) - f(B_{\alpha}, M_{,\gamma\beta}, M, ..., M)}_{-(p-2)f(B_{\alpha}, M_{,\gamma}, M_{,\beta}, \underbrace{M, ..., M}_{p-3}) + f([A_{\beta}, B_{\gamma}], M_{,\alpha}M, ..., M) - \underbrace{f([A_{\alpha}, B_{\gamma}], M_{,\beta}, M, ..., M)}_{p-3} - f([A_{\beta}, B_{\alpha}], M_{,\gamma}, M, ..., M)\} + f([A_{\beta}, B_{\gamma,\alpha}], M, ..., M) - \underbrace{f([A_{\alpha}, B_{\gamma,\beta}], M, ..., M)}_{-(p-2)f(B_{\alpha}, M_{,\gamma}, M, ..., M)} - f([A_{\beta}, B_{\alpha}], M_{,\gamma}, M, ..., M)\}$$

Denote the sum of the underlined terms by  $D_{\alpha\beta\gamma}$ . We claim that  $D_{\alpha\beta\gamma} = 0$ . Indeed, using the identities

$$f([A_{\alpha}, B_{\gamma,\beta}], M, ..., M) = (p-1)f(A_{\alpha}, [B_{\gamma,\beta}, M], M, ..., M),$$

$$f([A_{\alpha}, B_{\gamma}], M_{\beta}, M, ..., M) = f(A_{\alpha}, [B_{\gamma}, M_{\beta}], M, ..., M) + (p-2)f(A_{\alpha}, M_{\gamma}, M_{\beta}, \underbrace{M, ..., M}_{p-3}),$$
(27)

rewrite  $D_{\alpha\beta\gamma}$  in the form

$$D_{\alpha\beta\gamma} = (p-1)[f(A_{\alpha}, M_{,\gamma\beta}, M, ..., M) + (p-2)f(A_{\alpha}, M_{,\gamma}, M_{,\beta}, \underbrace{M, ..., M}_{p-3}) - f(A_{\alpha}, [B_{\gamma}, M_{\beta}], M, ..., M) - (p-2)f(A_{\alpha}, M_{\gamma}, M_{\beta}, \underbrace{M, ..., M}_{p-3}) - f(A_{\alpha}, [B_{\gamma,\beta}, M], M, ..., M)].$$

Thus we conclude that  $D_{\alpha\beta\gamma} = 0$ .

Define  $D_{1\alpha\beta\gamma}$  as follows:

$$D_{1\alpha\beta\gamma} = (p-1)[f([A_{\beta}, B_{\gamma}], M_{,\alpha}M, ..., M) - f([A_{\beta}, B_{\alpha}], M_{,\gamma}, M, ..., M)] + f([A_{\beta}, B_{\gamma,\alpha}], M, ..., M) - f([A_{\beta}, B_{\alpha,\gamma}], M, ..., M).$$

Using Eq. (27), one can verify that  $D_{1\alpha\beta\gamma}$  results in

$$D_{1\alpha\beta\gamma} = (p-1)f(A_{\beta}, M_{\gamma\alpha} - M_{\alpha\gamma}, M, ..., M),$$

i.e.,  $D_{1\alpha\beta\gamma} = 0$ .

Denote by  $D_{2\alpha\beta\gamma} = T_{1\alpha\beta\gamma} - T_{2\alpha\beta\gamma} + T_{3\alpha\beta\gamma}$ . Now taking into account the previous computations,  $D_{2\alpha\beta\gamma}$  yields that

$$D_{2\alpha\beta\gamma} = -(p-1)\{f(B_{\alpha,\gamma}, M_{,\beta}, M, ..., M) - f(B_{\gamma,\alpha}, M_{,\beta}, M, ..., M) + f(B_{\gamma,\beta}, M_{,\alpha}, M, ..., M) + f(B_{\alpha}, M_{,\beta\gamma}, M, ..., M) + (p-2)f(B_{\alpha}, M_{,\gamma}, M_{,\beta}, \underbrace{M, ..., M}_{p-3})\}.$$

Further, using the following identities:

$$f(B_{\alpha}, M_{,\beta\gamma}, M, ..., M) = f(B_{\alpha}, [B_{\beta,\gamma}, M], M, ..., M) +f(B_{\alpha}, [B_{\gamma}, M_{,\beta}], M, ..., M) = -f(M_{,\alpha}, B_{\beta,\gamma}, M, ..., M) -f([B_{\beta}, B_{\alpha}], M_{,\gamma}, M, ..., M) - (p-2)f(B_{\alpha}, M_{,\gamma}, M_{,\beta}, M, ..., M),$$

 $D_{2\alpha\beta\gamma}$  results in

$$D_{2\alpha\beta\gamma} = -(p-1)\{2f(B_{[\gamma,\beta]}, M_{,\alpha}, M, ..., M) + 2f(B_{[\alpha,\gamma]}, M_{,\beta}, M, ..., M) - f([B_{\beta}, B_{\alpha}], M_{,\gamma}, M, ..., M).$$

Taking into account the identities:

$$2[B_{[\alpha,\beta]}, M] = -[B_{\alpha}, M_{\beta}] + [B_{\beta}, M_{\alpha}] = -[[B_{\alpha}, B_{\beta}], M],$$

$$2f(B_{[\gamma,\beta]}, M_{,\alpha}, M, ..., M) = 2f(B_{[\gamma,\beta]}, [B_{\alpha}, M], M, ..., M) = -2f([B_{[\gamma,\beta]}, M], B_{\alpha}, M, ..., M) = f([[B_{\gamma}, B_{\beta}], M]], B_{\alpha}, M, ..., M) = -f([B_{\gamma}, B_{\beta}], M_{\alpha}, M, ..., M),$$

transform  $D_{2\alpha\beta\gamma}$  to the form:

$$D_{2\alpha\beta\gamma} = (p-1)\{f([B_{\gamma}, B_{\beta}], M_{,\alpha}, M, ..., M) + f([B_{\alpha}, B_{\gamma}], M_{,\beta}, M, ..., M) + f([B_{\beta}, B_{\alpha}], M_{,\gamma}, M, ..., M)\}.$$

In view of relations

$$f([B_{\alpha}[B_{\gamma}, B_{\beta}]], M, ..., M) = -(p-1)f([B_{\gamma}, B_{\beta}], M_{\alpha}, M, ..., M)$$

and Jacobi's identity,  $D_{2\alpha\beta\gamma}$  vanishes.

Finally, denote by

$$\Delta_{1\alpha\beta} = W_{1\alpha\beta} - W_{2\alpha\beta} + W_{3\alpha\beta}.$$
 (28)

Substituting Eq. (25) and (26) in (28), we obtain the relations

$$\Delta_{1\alpha\beta} = f([A_{\beta}, N]_{,\alpha}, M, ..., M) - f([A_{\alpha}, N]_{,\beta}, M, ..., M).$$

This equation results in

$$\Delta_{1\alpha\beta} = f(N_{,\alpha\beta} - N_{,\beta\alpha}, M, ..., M) = 0.$$

# References

- [1] Balandin A.V., Pakhareva O.N., Potemin G.V.: Lax representation of the chiral-type field equations, Phys. Lett. A. 23 168 -176 (2001).
- [2] Demskoi D.K., Meshkov A.G.: Zero-curvature representation for a chiral-type three-field system, Inverse Problems. 19 563–571 (2003)

- [3] Goto M., Grosshans F.D.: Semisimple Lie algebras. Lecture Notes in Pure and Applied Mathematics. Vol 38, (1978)
- [4] Gu C., Hu H., Zhou Z.: Darboux Transformations in Integrable Systems. Theory and their Applications to Geometry, Springer (2005)
- [5] Kostant, B.: Lie group representations on polynomial rings, Amer. J. Math. 85 327-404 (1963)
- [6] Leznov A.N., Savel'ev M.V.: Group-Theoretical Methods for Integration of Nonlinear Dynamical Systems, Birkhauser, (1992)
- [7] Lund F., Regge T.: Unified approach to strings and vortices with soliton solutions, Phys.Rew. D. 14 1524–1535 (1976)
- [8] Marvan M.: On zero-curvature representations of partial differential equations. in: Kowalski O., Krupka D. (eds.) 5th International Conference on Differential Geometry and Its Applications. pp. 103-122, Opava, Czech Republic (1993), (http://www.emis.de/proceedings/5ICDGA).
- [9] Novokshenov V.Y.: Vvedenie v teoriju solitonov, IKI, Izhevsk (2002) (in Russian)
- [10] Olver P.: Applications of Lie Groups to Differential Equations, 2nd ed. Springer, New York (1993)
- [11] Shabat A.B.(ed): Encyclopedia of integrable systems, version 0043, L.D. Landau Institute for Theoretical Physics, Moscow (2010)
- [12] Varadarajan V.S.: On the ring of invariant polynomials on a semisimple Lie algebra, Amer. J. Math. 90 308-317 (1968)