

TITLE: Sex-specific recombination rates and allele frequencies
affect the invasion of sexually antagonistic variation on autosomes

ARTICLE TYPE: Research article

RUNNING HEAD: Sex-specific recombination rates and allele frequencies

AUTHORS: Minyoung J. Wyman^{1,2}
Mark C. Wyman^{3,4}

ADDRESS: ¹minyong.yi@mail.utoronto.ca
²Department of Ecology & Evolutionary Biology
University of Toronto
25 Willcocks Street
Toronto, ON M5S 3B2
CANADA

³Department of Astronomy & Astrophysics

⁴Kavli Institute for Cosmological Physics, Enrico Fermi Institute
University of Chicago
Chicago, IL 60637
USA

KEYWORDS: sexual dimorphism, heterochiasmy, sexual conflict, sexual antagonism

1 Abstract

The introduction and persistence of novel sexually antagonistic alleles can depend upon factors that differ between males and females. Understanding the conditions for invasion in a two-locus model can elucidate these processes. For instance, selection can act differently upon the sexes, or sex-linkage can facilitate the invasion of genetic variation with opposing fitness effects between the sexes. Two factors that deserve further attention are recombination rates and allele frequencies – both of which can vary substantially between the sexes. We find that sex-specific recombination rates in a two-locus diploid model can affect the invasion outcome of sexually antagonistic alleles and that the sex-averaged recombination rate is not necessarily sufficient to predict invasion. We confirm that the range of permissible recombination rates is smaller in the sex benefitting from invasion and larger in the sex harmed by invasion. However, within the invasion space, male recombination rate can be greater than, equal to, or less than female recombination rate in order for a male-benefit, female-detriment allele to invade (and similarly for a female-benefit, male-detriment allele). We further show that a novel, sexually antagonistic allele that is also associated with a lowered recombination rate can invade more easily when present in the double heterozygote genotype. Finally, we find that sexual dimorphism in resident allele frequencies can impact the invasion of new sexually antagonistic alleles at a second locus. Our results suggest that accounting for sex-specific recombination rates and allele frequencies can determine the difference between invasion and non-invasion of novel sexually antagonistic alleles in a two-locus model.

2 Introduction

Selection can act differently in males versus females (Arnqvist and Rowe 2005), resulting in antagonism over the expression of shared traits. Fitness itself can be under sexual conflict so that reproductive success has a negative genetic correlation between the sexes (e.g., Chippindale et al. 2001; Foerster et al. 2007). Alternatively, individual traits may also be under sexual conflict because trait expression in one sex has opposing fitness effects in the other sex. Among sexually dimorphic traits, $\sim 17\%$ of selection estimates were sexually antagonistic (Cox and Calsbeek 2009). Sexual conflict can operate at even finer scale levels: the expression of $\sim 8\%$ of genes is beneficial for one sex but detrimental in the other in *Drosophila melanogaster* (Innocenti and Morrow 2010). Understanding sexual antagonism requires experimental (e.g., Long et al. 2012) as well as theoretical approaches. In a one-locus model, Rice (1984) suggested that sex-linkage can facilitate the initial spread of sexually antagonistic alleles. His model demonstrated that recessivity and hemizyosity (possessing one major sex chromosome) can shield novel antagonistic alleles from selection in the non-benefitting sex. However, Fry (2010) has shown that differences in sex-specific dominance may affect the invasion advantage of sex-linkage.

Recent theoretical work on the invasion of sexually antagonistic alleles has focused on two-locus models (Connallon and Clark 2010; Patten et al. 2010; Úbeda et al. 2011) which introduce realistic factors such as epistasis, linkage disequilibrium, and recombination. These factors are known to alter the invasion outcomes for novel sexually antagonistic alleles. For instance, Connallon and Clark (2010) employed a model whereby sexual antagonism can be introduced through epistatic interactions between two different loci. When epistasis is present, the recombination rate difference between males and females affects the invasion of sexually antagonistic allelic combinations. In a related model, Patten et al. (2010) showed that a second locus can both increase the opportunities for polymorphic equilibria relative to a one-locus model and preserve allelic combinations that increase fitness variation through linkage disequilibrium.

Differences in sex-specific recombination rates are often dramatic (e.g., Mank 2009; Lenormand and Dutheil 2005; Brandvain and Coop 2012), suggesting the need for further study into its implications. Up to 75% of recombining species demonstrate >5% overall rate difference between the sexes (Burt et al. 1991; Lenormand 2003). Recombination rates on a local scale can also vary among individuals (Coop et al. 2008; Baudat et al. 2010; Fledel-Alon et al. 2011) and between the sexes (e.g., Hansson et al. 2005; Berset-Brändli et al. 2008; Kong et al. 2010). In fact, recombination rates have very little intersexual covariance (Coop et al. 2008), so that tracking the rates in males and females separately may be necessary. It is important to understand what, if any, implications sexual dimorphism in recombination might have on the introduction of sexually antagonistic phenotypes. For instance, it has been suggested that lower recombination can maintain combinations of genes beneficial to males because the genes have successfully undergone sexual selection (Trivers 1988). Although sex-specific selection at the diploid stage is unlikely to facilitate the evolution of heterochiasmy (Lenormand 2003), once heterochiasmy is established, it may subsequently affect the formation of male- or female-benefit regions in the genome. Indeed, lower sex-specific recombination makes it easier for alleles benefitting that same sex to invade on sex chromosomes (Connallon and Clark 2010).

Here we expand upon previous models and investigate the effect of heterochiasmy in three ways. First, we study the effect of sexually dimorphic recombination rates on the invasion of sexually antagonistic alleles. Specifically, we examine whether dimorphism in recombination can hinder or facilitate the spread of an autosomal male-benefit, female-detriment allele (and similarly for female-benefit, male-detriment alleles). Second, to capture the reality of sex-specific and individual variation in recombination rates, we study how sex- and genotype-specific recombination rates affect invasion outcomes. Finally, we analyze the impact of sexually dimorphic allele frequencies at one locus for invasion outcome at a second locus. Considered together, these results suggest that novel sexually antagonistic variation can potentially spread more easily due to pre-existing sexual

dimorphism in recombination and allele frequencies.

3 Model

We constructed an autosomal two-locus diploid model. The two-locus diploid model has been studied extensively in other contexts (e.g., Otto 1991; Curtsinger and Heisler 1988; Albert and Otto 2005; Patten et al. 2010). Much of our notation is adopted from Connallon and Clark (2010) to facilitate comparison. The A and B loci both have two alleles so that there are 10 unique male and 10 unique female genotypes and associated fitnesses (see Table 1). Recombination between the A and B locus occurs at a rate r_m in males and r_f in females. Discrete time recursion equations for the AB genotypes are reproduced in the Appendix B (and in a `Mathematica` notebook provided in the online supplementary materials). The genotype frequencies in the next generation depend upon the maternal and paternal allele frequencies at the A and B loci, as well as upon recombination and linkage disequilibrium between the two loci.

First, we analyze recombination rates that are non-zero (i.e., no achiasmy) and different in each sex for the polymorphic equilibrium case (e.g., A_1 is not fixed upon the invasion of B_2). We first work with a general model to derive our expressions, and then plug in a specific fitness parameterization to investigate the effects of dominance and epistasis on our results (parameterization in Appendix A). Because we only specialize at the last step, any fitness parameterization may be used, and our general results are not contingent upon them. Although we are using a two-locus diploid model in the context of sexual antagonism, our model is not to be conflated with models of interlocus sexual conflict (i.e., different loci in each sex interact to affect the expression of a shared trait). Second, we treat recombination rate as a novel sexually antagonistic trait in our model and study its effect on invasion. Lastly, we study the model's behavior on the invasion of new alleles when resident allele frequencies are different between the sexes.

4 Results and Discussion

4.1 Sex-averaged versus sex-specific recombination rates

Upon first consideration, it may appear that only the sex-averaged recombination rate can matter for invasion (e.g., Hedrick 2007). After all, recombination shuffles loci in males and females, so that any particular allelic combination has approximately the same likelihood of appearing in both sons and daughters. These combinations experience sex-specific selection, but then get re-shuffled in males and females of the second generation. As re-shuffling is inevitable, it may seem that the effects of sex-specific recombination are averaged out between the sexes over time. However, a few lines of evidence suggest that this view is too simplistic. First, sexual antagonism can preserve stable linkage disequilibrium in the face of recombination so that sex-specific patterns of variation persist as polymorphic equilibria (Patten et al. 2010; Úbeda et al. 2011). Second, when sex-specific recombination rates are explicitly modeled, their particular values appear to affect the conditions for invasion (Connallon and Clark 2010). Finally, whether a novel allele spreads or not depends only upon the present conditions, not the long-term conditions. Selection precedes fertilization, so that recombination occurs in a subset of individuals different from the total set initially preset at birth. Selection may occur at the adult stage, but may also occur at the gamete stage via fertility selection. Thus, the interplay of sexually antagonistic fitness and sex-specific recombination rates may lead to small but potentially consequential sex differences in the frequency of novel alleles, affecting their invasion.

We can study these dynamics by conducting a local stability analysis of the recursion equations governing a two-locus model with sex-specific recombination (Appendix B). We calculated the eigenvalues of the Jacobian matrix for the recursion equations; whenever the dominant eigenvalue describing an invasive allele is greater than one, the population is subject to invasion by that novel allele. We conducted this analysis under the condition that the first A locus is polymorphic for A_1 (frequency q) and A_2 (frequency p)

and the second B locus is fixed for B_1 upon the introduction of B_2 . The polymorphic case may be more interesting under the assumption that a novel mutation may appear on a background that is genetically variable throughout a population (NB: The eigenvalue for the equilibrium whereby A_1 is fixed may be recovered by setting $p = 0$ and $q = 1$).

The dominant eigenvalue (λ_{CC}) for the invasion of a novel B_2 allele when the male and female frequencies are equal ($p_m = p_f$) at the A locus is given by equation 6a in Connallon and Clark (2010):

$$\lambda_{CC} = \frac{1}{2\bar{w}_f} (f_{32}p + f_{22C}(1-p)(1-r_f)) + \frac{1}{2\bar{w}_m} (m_{32}p + m_{22C}(1-p)(1-r_m)). \quad (1)$$

As Connallon and Clark (2010), we have used f_{type} and m_{type} to designate female and male fitnesses, respectively (Table 1). For instance, genotype $A_2B_1A_2B_2$ has the fitness type 32. In addition, the 22 types are either 22C or 22R, because 22 is heterozygous at both loci, but may occur as coupling or repulsion (Table 1). \bar{w}_f and \bar{w}_m represent the mean female and male fitnesses, respectively.

λ_{CC} can be re-written by expressing fitness values as a deviation from 1. For instance, for the $A_2B_1A_2B_2$ genotype the female fitness can be re-expressed as $f_{32} = 1 - u \delta f_{32}$, where δf_{32} is the deviation of f_{32} from 1. The u term is a counting parameter for doing a series expansion when u is small. Terms with no u (i.e., u^0) are larger than terms with u^1 , and terms with u^1 are larger than terms with u^2 , etc. This expansion is made assuming weak selection; we also assume that mutation is weak. Finally, we further assume that recombination rates are small and also multiply them by u . Equation 1 can now be re-written as an expansion in u :

$$\begin{aligned} \lambda_{CC} \simeq & \frac{1}{2} \left(\frac{\bar{w}_f + \bar{w}_m}{\bar{w}_f \bar{w}_m} \right) + \\ & - \frac{u}{2\bar{w}_f \bar{w}_m} (\bar{w}_f ((1-p)\delta m_{22C} + p\delta m_{32}) + \bar{w}_m ((1-p)\delta f_{22C} + p\delta f_{32}) + (1-p)(\bar{w}_m r_f + \bar{w}_f r_m)) \\ & + \frac{u^2}{2\bar{w}_f \bar{w}_m} (1-p) (\delta f_{22C} \bar{w}_m r_f + \delta m_{22C} \bar{w}_f r_m) + \mathcal{O}(u^3). \end{aligned} \quad (2)$$

This approximation has a few nice features that aid our understanding. The first line contains just the mean fitnesses and is approximately 1. The second line contains terms that have one power of u ; the third line contains even smaller pieces (where \mathcal{O} indicates the order in u). Taking $\bar{w}_f = \bar{w}_m = 1$ simplifies things further:

$$\begin{aligned} \lambda_{CC} \simeq & 1 - \frac{u}{2} [(1-p)\delta m_{22C} + p\delta m_{32} + (1-p)\delta f_{22C} + p\delta f_{32} + (1-p)(r_f + r_m)] \\ & + \frac{u^2}{2} (1-p) [\delta f_{22C} r_f + \delta m_{22C} r_m] + \mathcal{O}(u^3). \end{aligned} \quad (3)$$

We observe that the main recombination effect is indeed the sex-averaged recombination rate (the last term in the first line of equation 3), as is typically assumed (Hedrick 2007). However, when males and females differ in fitness for the genotype $A_1B_1A_2B_2$ – i.e., when $\delta f_{22} \neq \delta m_{22}$ (as is expected when B_2 is sexually antagonistic) – the sex-averaged rate is adjusted by the second line (i.e., $\mathcal{O}(u^2)$). The strength of the second line is proportional to $1-p$ and affects genotypes bearing B_2 . As the strength of the sexual antagonism increases, the effect of heterochiasmy also increases.

In general, the $\mathcal{O}(u^2)$ terms will be smaller than the $\mathcal{O}(u^1)$ terms. However this assumption is not met in two cases. Most obviously, the approximation is not valid under strong selection or strong sexual antagonism (i.e., genotype fitnesses are not close to 1) so that the approximation $\mathcal{O}(u^2) \ll \mathcal{O}(u^1)$ breaks down. A more subtle scenario (that does not violate the approximation) is when the terms on the first line almost cancel out, allowing order 1 terms to be order 2:

$$\frac{u}{2} [(1-p)\delta m_{22C} + p\delta m_{32} + (1-p)\delta f_{22C} + p\delta f_{32} + (1-p)(r_f + r_m)] \simeq u^2. \quad (4)$$

For example, for the stable polymorphic equilibrium $p = 0.25$ [i.e., polymorphic equilibrium is determined by $p = (W_{AA} - W_{Aa}) / (2W_{Aa} - W_{AA} - W_{aa})$, in the notation of Otto and Day (2007)], the δm terms are all slightly negative (male-benefit) and the δf terms are all slightly positive (female-detriment). As a result, the magnitude of $\delta m_{22} \simeq$ the magnitude

of δf_{22} , and the magnitude of $\delta m_{32} \simeq$ magnitude of δf_{32} , but with the overall sum of the four terms having a slightly negative effect. In other words, the A_2 and B_2 alleles are only slightly male-benefit and slightly female-detriment. Such alleles may be important to study under the assumption that new mutations may have minor phenotypic effects in relation to the current phenotype, as we assume under an infinitesimal model. Then the sum within the brackets of equation 4 may be very small, of $\mathcal{O}(u^1)$. Since r_m and r_f are of $\mathcal{O}(u^1)$, the “first” order term is the same or smaller magnitude as the second order term. In such a case, the second order term makes a substantial difference to the outcome of invasion and must be included for consistency; so, the sex-averaged rates and the relative rates may contribute equally when equation 4 holds. Even though this special case may only persist for a short time in a population at large, it can affect the invasion of new alleles during that time. (NB: We emphasize that when $\mathcal{O}(u) \simeq \mathcal{O}(u^2)$ due to an accidental cancellation, the approximation is still valid because we still expect $\mathcal{O}(u^3) \ll \mathcal{O}(u^2)$). We have demonstrated qualitatively how equation 4 may be relevant. Below we also present an example using a particular fitness model to highlight both the qualitative and quantitative implications of equation 4 to invasion.

4.2 Implications of sex-specific recombination

The analysis above shows that the sex-specific recombination rates matter for invasion, in addition to the sex-averaged rate. A specific fitness model (see Appendix A) can be used for further investigation. We determine whether B_2 can invade on a polymorphic A background, given that B_2 decreases fitness in both males and females, but also interacts epistatically with A_2 to increase only male fitness and not alter female fitness (i.e., the A_2B_2 combination is sexually antagonistic overall). The fitness model assumes linear dominance (dominance coefficients h and g for the A and B loci, respectively). The selection coefficients are s and t (for A and B loci, respectively); positive values indicate a fitness detriment and negative values indicate a fitness benefit. Epistasis (ϵ) is additive;

positive values indicate a fitness detriment and negative values indicate a fitness benefit.

For the A locus we assumed equal dominance and selection coefficients between the sexes: $h_f = -0.1$, $h_m = -0.1$, $s_f = 0.1$, and $s_m = 0.1$. For the B locus we also assumed equal dominance and selection coefficients: $g_f = 0.5$, $g_m = 0.5$, $t_f = 0.05$, and $t_m = 0.05$. For these set of parameters, the B_2 allele is detrimental to both males and females. However, we assumed epistasis to be $\epsilon_m = -0.25$ and $\epsilon_f = 0$. As a result, when A_2 and B_2 are together, the allele combination confers an overall benefit to males but not females (since $t_m = t_f$ but $\epsilon_m \neq \epsilon_f$). It would be easy enough to make the alleles themselves sexually antagonistic by setting selection to have opposite signs between the sexes (i.e., $s_m, t_m > 0$ and $s_f, t_f < 0$). However, doing so at the A locus would violate the assumption of equal allele frequencies in males and females (see more below). Meanwhile, making the B locus sexually antagonistic independently of the A locus would reduce the dynamics to a one-locus model.

In Fig. 1a demonstrates a few features of the model. The solid line is the critical invasionary eigenvalue of one; invasion occurs to the left, and not to the right, of this line. First, we observe that the range of recombination rates required for the new B_2 allele to invade due to its epistatic interaction with A_2 , differs between male and females. Males have a smaller range of recombination rates conducive to invasion than females. So, overall, $r_m < r_f$. Connallon and Clark (2010) also show this result but present it in terms of the wait times for successful co-invasion of epistatically beneficial alleles. An intuitive explanation for this theoretical result rests in the fact that keeping A_2 and B_2 together in males requires suppressing recombination to some degree. By contrast, because B_2 does not benefit females with an A_2 background, a wider range of recombination values that can potentially break up the detrimental combinations benefits females. In support of this interpretation, when B_2 hurts females ($t_m < 0$ and $t_f > 0$), as in the case of sexual antagonism, this range of female recombination rates increases.

A second, and key, observation is that within the overall invasion space (left of the




Figure 1: Sex-specific recombination rates. See text for parameter selection. a) The dashed line represents the sex-averaged recombination dependent term; it connects the same recombination value on the r_f and r_m axes. The solid line represents λ_{CC} which provides the correction to the sex-averaged estimate. Because the individual recombination rates matter, the solid line does not connect the same values on both axes. Both lines represent the critical eigenvalue of one. Invasion occurs to the right of the each line; invasion does not occur to the left. In order for recombination to facilitate the invasion of the second male-benefit, female-detriment allele, the range of male recombination rates has to be smaller than the range of female recombination rates. However, within the invasion space, $r_m = r_f$, $r_m > r_f$ and $r_m < r_f$ can all permit invasion. b) In region II, the recombination rates ($r_m = 0.08$, $r_f = 0.012$) allow invasion of B_2 even though the sex-averaged rate would not predict invasion. In region I the recombination rates ($r_m = 0.01$, $r_f = 0.079$) do not allow invasion of B_2 , even though the sex-averaged rate would predict invasion.

solid line), $r_m = r_f$, $r_m > r_f$, and $r_m < r_f$ all yield valid conditions for the successful invasion of B_2 . This may explain why we might not expect to see a consistent fine-scale correlation between sex-specific recombination rates and the concentration of male- or female-benefit alleles – even though, the range of invasionary recombination rates differs between the sexes. So while the male recombination rate should be lower than the female recombination rate overall in regions enriched with genes beneficial to males, in particular cases they may not be. Conversely, the female recombination rate may not necessarily be lower in regions of female-benefit genes. Positive correlations between the concentration of

sex-specific beneficial alleles and lower recombination rates may only be apparent in the most extreme circumstances: e.g., when male and female recombination values are near the invasion border and when males do not recombine (i.e., along the x-axis of Fig. 1a). As Connallon and Clark (2010) have shown, the lack of male recombination and the fact that the X-chromosome experiences selection more often in females than in males may explain empirically why the concentration of male-biased genes is lower on the X chromosome of a male non-recombining species such as *Drosophila*. This must be the case if male-bias can be roughly equated with male-benefit, as the current evidence suggests (Pröschel et al. 2006; Andolfatto et al. 2010; Innocenti and Morrow 2010; Wyman et al. 2010).

Finally, Fig. 1a demonstrates that the correction (solid line) to the sex-averaged (dotted line) recombination rate makes the greatest impact near the border of invasion. This borderline region is where the $\mathcal{O}(u)$ effects and the $\mathcal{O}(u^2)$ effects compete with each other to determine invasion. Both lines indicate the slope of the critical invasionary eigenvalue. However, in region II, the sex-averaged rate will incorrectly predict the lack of invasion for this given set of parameters; conversely, in region I, the sex-averaged rate will incorrectly predict successful invasion. The numerical results support this mismatch between the two critical eigenvalues (Fig. 1b). Reliably predicting the persistence or disappearance of novel sexually antagonistic alleles based upon the sex-averaged recombination rate will require that the rate is well within or beyond this borderline.

Many factors lead to variation in the rate of recombination among individuals – such as age, genetic background, and environmental stress (see Tedman-Aucoin and Agrawal 2011 for review). Sources of variation relevant to this study include re-mating rate (Priest et al. 2007) and male-genotype (Stevison 2011). Sexually antagonistic allele combinations may invade more easily in populations in which males can lower their own recombination rates, or can induce lower average recombination rates in females, relative to the critical invasionary values. Conversely, male-benefit, female-detriment alleles may invade less easily in populations where males do not decrease (or rather, increase) their

own recombination rate or their partner's.

4.3 Individual variation in sex-specific recombination rates

The analyses above support that the sex-averaged recombination rate is not the only quantity of interest. The sex-specific rates can matter through sexually antagonistic effects on fitness from epistasis. However, because recombination itself is a phenotype, it is also important to consider its variation among individuals (Coop et al. 2008; Baudat et al. 2010; Fledel-Alon et al. 2011) as well as between the sexes. In fact, variation in recombination rate is known to affect variation in fitness; mothers with higher average recombination rate have more children (Kong et al. 2004; Fledel-Alon et al. 2011).

To capture these biological realities, our model can be modified so that each sex by genotype combination has a distinct recombination rate, e.g., an adult $A_1B_1A_1B_1$ male has a recombination rate, r_{11}^m (adopting the notation in Table 1). Thus, A and B affect both fitness and recombination rate. In the new eigenvalues of the modified model, recombination only affects terms that carry the double heterozygote genotype ($22 = A_1A_2B_1B_2$; see Table 1):

$$\lambda \supset f_{22C}(1-p)(1-r_{22C}^f) + m_{22C}(1-p)(1-r_{22C}^m) \quad (5)$$

or

$$\lambda \supset f_{22R}p(1-r_{22R}^f) + m_{22R}p(1-r_{22R}^m). \quad (6)$$

This result comports nicely with intuition. Recombination in the double heterozygotes matters because recombination destroys A_2B_2 (from the $22C$ genotype) or creates A_2B_2 (from the $22R$ genotype). Keeping track of these processes will determine the size of the respective eigenvalues. The other genotypes with the B_2 allele are either extremely rare (i.e., B_2B_2) or have no effective recombination (i.e., recombination occurs but does not change the haplotypes).

More generally, equations 5 and 6 have the form:

$$\lambda \supset \text{female fitness} + \text{male fitness} - (\text{female fitness} \times r_f) - (\text{male fitness} \times r_m) \quad (7)$$

If fitness is always a positive number, and the new allele is male-benefit and female-detriment, the fourth term is the largest negative term. As a result, invasion becomes more likely (i.e., $\lambda > 1$) as r_m becomes smaller. In particular, r_{22C}^m or r_{22R}^m must be smaller, and not any other genotype-specific male recombination rate (e.g., r_{11}^m , r_{12}^m , r_{21}^m , etc.). Furthermore, in order for a sexually antagonistic allele to invade, the recombination rate must be low in the sex benefitting from the new allele, and in particular, recombination must be low in the double heterozygote genotype, which carries one copy of the new allele. Interestingly, it is also true that r_{22C}^f or r_{22R}^f , and not any other genotype-specific female recombination rate, should be smaller for a male-benefit allele to invade.

4.4 Sexually dimorphic allele frequencies

In the previous sections, we allowed for a polymorphic equilibrium at the A locus, but made the assumption that allele frequencies in males and females were equivalent, $p_{female} = p_{male}$. While assuming equal allele frequencies in the sexes simplifies the math and is a fair approximation, as a precise statement it may have limited biological relevance. After all, when an allele is sexually antagonistic, allele frequencies should be slightly higher in the helped sex and slightly lower in the harmed sex. Although male and female offspring are equally likely to inherit their alleles from their fathers and mothers, sexually antagonistic selection and fertility selection can alter allele frequencies prior to mating and fertilization (e.g., Patten et al. 2010; Úbeda et al. 2011).

To assess and estimate the importance of this mismatch between male and female allele frequencies, we used the recursion equations and found polymorphic equilibria for $p_{male} \neq p_{female}$. The equilibrium allele frequencies can be re-written in terms of two

numbers p_f and d_m :

$$\begin{aligned} x_1 &= 1 - p_f & y_1 &= 1 - p_f - d_m \\ x_2 &= p_f & y_2 &= p_f + d_m \end{aligned} \tag{8}$$

where $x_1 = 1 - p_f$ is the proportion of the ova carrying the A_1 allele, x_2 the proportion carrying A_2 . y_1 is the proportion of sperm carrying A_1 , y_2 carrying A_2 . d_m is the difference in allele frequency between males and females, $d_m = p_m - p_f$. Plugging equations 8 into the recursion formulas, using $f = 1 - u \delta f$, and expanding out the mean fitnesses \bar{w}_m and \bar{w}_f and keeping only u^1 terms, we found the following approximate equilibrium solutions:

$$p_f = \frac{\delta f_{11} - \delta f_{21} + \delta m_{11} - \delta m_{21}}{\delta f_{31} - 2\delta f_{21} + \delta f_{11} + \delta m_{31} - 2\delta m_{21} + \delta m_{11}} \tag{9}$$

$$\begin{aligned} d_m &= 2u [\delta f_{31} - \delta f_{21} + \delta m_{31} - \delta m_{21})(\delta f_{21} - \delta f_{11} + \delta m_{21} - \delta m_{11}) \times \\ &\quad \delta f_{11}(\delta m_{31} - \delta m_{21}) + \delta f_{31}(\delta m_{21} - \delta m_{11}) + \delta f_{21}(-\delta m_{31} + \delta m_{11})] \times \\ &\quad [\delta f_{31} - 2\delta f_{21} + \delta f_{11} + \delta m_{31} - 2\delta m_{21} + \delta m_{11}]^{-3}. \end{aligned} \tag{10}$$

Although the full expression (to higher order in u) is necessary to find accurate equilibria (see **Mathematica** notebook in online supplement of the journal version of this article for details), this first order approximation has a few instructive properties. p_f appears with no factors of u , which makes sense as allele frequencies can be large relative to recombination when A is polymorphic. However, d_m appears as a multiple of u^1 (four factors of δf or δm in the numerator and three in the denominator of d_m). Thus, the difference between p_m and p_f has the same size effect on the eigenvalue as the sex-specific fitnesses do. The assumption $p_m = p_f$, in addition to its limited biological appeal, is not necessarily consistent.

We can calculate how much the eigenvalue is changed relative to λ_{CC} when $d_m \neq 0$

by determining the value of $\Delta\lambda$, defined as $\Delta\lambda = \lambda - \lambda_{CC}$ (where λ is the full eigenvalue). We can re-write d_m as $d_m u$ to make its size obvious in the counting parameter u . $\Delta\lambda$ is a long expression found by solving for the characteristic polynomial; the most important piece is new first order contribution to the eigenvalue from d_m :

$$\Delta\lambda = \frac{1}{2}d_m u + \mathcal{O}(u^2). \quad (11)$$

Invasion is substantially impacted by including sexually dimorphic allele frequencies, as represented by d_m . Moving onto the next smaller set of terms, we find additional $\mathcal{O}(u^2)$ contributions to the eigenvalue that affect d_m . Since our main focus is on sexually dimorphic recombination rates, we show only terms with r_m or r_f :

$$\Delta\lambda \supset \frac{1}{4}d_m u^2 (-p_f r_f - 2r_m + p_f r_m) + \mathcal{O}(u^3). \quad (12)$$

Hence, whenever the sex-specific recombination rates are important (i.e., borderline of Fig. 1) we should also keep track of the new parts of the eigenvalue that are proportional to d_m .

To interpret d_m , we can again plug in our fitness model (see Appendix A):

$$p_f = -\frac{h_f s_f + h_m s_m}{s_f - 2h_f s_f + s_m - 2h_m s_m} \quad (13)$$

$$d_m = -\frac{2(h_f - h_m)s_f s_m((h_f - 1)s_f + (h_m - 1)s_m)(h_f s_f + h_m s_m)}{(s_f - 2h_f s_f + s_m - 2h_m s_m)^3}. \quad (14)$$

The equation for d_m shows that when male and female dominances are equal, $d_m = 0$, so that male and female allele frequencies are identical. Thus, sexual dimorphism in dominance can affect the invasion probabilities, in addition to the effect from sexual dimorphism in recombination rates.

In Fig. 2 we numerically confirm that d_m has the potential to impact whether a sexually antagonistic allele B_2 can invade in the population. For the same averaged male




Figure 2: Differences in allele frequencies of A_1 between the sexes. Invasion of the B_2 allele (males, upper black points; females, lower black points) can occur in a population which has sexually dimorphic frequencies of A_1 , with $d_m = 0.005$ ($h_m = -0.9$, $h_f = -0.1$, $s_m = 0.15$, $s_f = 0.05$, $g_m = 0.1$, $g_f = 0.2$, $t_m = 0.05$, $t_f = 0.2$, $\epsilon_m = -0.1$, $\epsilon_f = 0$). However, invasion of B_2 does not occur (both sexes, grey points) when A_1 frequency is monomorphic, $d_m \simeq 0$ ($h_m = h_f = -0.69$, $s_m = 0.1$, $s_f = 0.1$, $g_m = 0.1$, $g_f = 0.2$, $t_m = 0.05$, $t_f = 0.2$, $\epsilon_m = -0.1$, $\epsilon_f = 0$). This difference in invasion outcome exists even though the sex-averaged allele frequencies were initially identical and even though recombination was the same in both populations ($r_m = r_f = 0.006$).

and female allele frequencies at locus A , B_2 can eventually invade in one population (black points: $d_m = 0.005$) but not in the other population (grey points: $d_m = 0$). This occurs despite the fact that both populations have the same recombination parameters ($r_m = r_f = 0.006$) and the same sex-averaged p . We note that invasion depends on the size and sign of d_m (larger d_m has a greater impact). That is, when $d_m > 0$ and a new allele is male-benefit, invasion is easier. On the other hand, parameter selection that results in

$d_m < 0$ will hinder the invasion of a male-benefit sexually antagonistic combination (or favor female-benefit alleles). In summary, sex differences in allele frequencies can have definite effects on the possibility of invasion.

4.5 Limitations of the model

As with any model, the results are subject to limitations of the assumptions. Two factors may mitigate the applicability of these results. First, the role of sexually dimorphic recombination was only investigated for weak selection. When the strength of selection increases in either sex, the approximation (equation 3) breaks down, and the sex-averaged recombination rate determines invasion outcome over the sex-specific recombination rates. Second, since populations are finite, it is important to assess the effect of the modified eigenvalues for realistic conditions. When the eigenvalue is near one (i.e., sex-specific recombination rates make a difference to invasion), we must worry about genetic drift. Drift dominates when $N(\lambda - 1) \ll 1$, where N is the population size and $\lambda - 1$ is the deviation of the eigenvalue for invasion from one. $1/(\lambda - 1)$ measures the time to invasion for a novel allele. Hence, in a case when the fitnesses and recombination rates are on the order of 1%, $(\lambda - 1) \sim 10^{-4}$. Drift becomes less important when $N > 10^4$, a modest population size.

5 Conclusions

Recombination is important for adaptation because it shuffles loci and places favorable alleles together. However, recombination also hinders adaptation by breaking up the very same favorable combinations. The model studied here demonstrates that the advantages and drawbacks of recombination occur alongside each other, albeit in the different sexes. In fact, sex-specific recombination affects invasion in a manner distinct from simply the sex-averaged recombination rate, especially at the border of invasion. Low recombination permits the helped sex to keep favorable sexually antagonistic combinations

together. Low recombination also permits stable linkage disequilibria to maintain sexually antagonistic variation (Patten et al. 2010; Úbeda et al. 2011). Both effects occur because low sex-specific recombination increases allele frequencies in the benefitting sex over the harmed sex without the differences washing out over time.

However, within the invasion space, we find that male recombination can be greater than, less than, or equal to female recombination and still allow a male-benefit allele combination to invade. In other words, successful invasion does not always require that the benefitting sex have a lower recombination rate than the harmed sex. Empirically, the model suggests that low male recombination rates may not always be correlated to a high concentration of male-benefit genes on chromosomes.

When recombination is treated as the phenotype of interest and allowed to vary in a genotype-specific manner, only the recombination rate of the double heterozygotes matters for invasion outcome. The double heterozygotes are the only genotype in which recombination will make a difference to increasing or decreasing the frequency of the new sexually antagonistic allele. This is a sex-specific effect so that a male-benefit, female-detriment allele that simultaneously increases male fitness and decreases male recombination rate invade more easily than those alleles that do only one or the other.

We also find that sexual dimorphism in allele frequencies affects invasion outcomes. As sexually antagonistic alleles benefit only one sex, allele frequencies may be higher in the helped sex (Patten et al. 2010; Úbeda et al. 2011). Here, sex differences in allele frequencies at one locus impact invasion of novel alleles at a different locus (Fig. 2). The magnitude of this effect can be as important as that of sex-specific recombination. These results suggest that maintaining sexually dimorphic allele frequencies at one locus would be a way to both facilitate or hinder the spread of sexually antagonistic alleles at a second locus.

In sum, there is a delicate interplay among sex-specific selection, sex-specific recombination rates, and sex-specific allele frequencies. The impact of sexual dimorphism in recombination rates and allele frequencies became more apparent by doing the

calculations than by relying upon intuition alone. Two-locus models of sexual antagonism have a great deal of complexity unavailable to one-locus models that might explain the presence of widespread sexual antagonism in nature.

6 Acknowledgments

We thank L. Rowe, A.F. Agrawal, J.C. Perry, and 2 anonymous referees for comments. M.J.W. was supported by the University of Toronto Connaught Scholarship and Doctoral Completion Award and by NSERC grants to L. Rowe. M.C.W. was supported by U.S. Dept. of Energy contract DE-FG02-90ER-40560 and by the Kavli Institute for Cosmological Physics at the University of Chicago through grant NSF PHY-1125897 and an endowment from the Kavli Foundation and its founder Fred Kavli.

References

- Albert, A. Y. K. and S. P. Otto. 2005. Sexual selection can resolve sex-linked sexual antagonism. *Science* 310:119–121.
- Andolfatto, P., K. Wong, and D. Bachtrog. 2010. Effective population size and the efficacy of selection on the X chromosomes of two closely related *Drosophila species*. *Genome Biology and Evolution* 3:114–128.
- Arnqvist, G. and L. Rowe. 2005. *Sexual conflict*. Princeton University Press, Princeton.
- Baudat, F., J. Buard, C. Grey, A. Fledel-Alon, C. Ober, M. Przeworski, G. Coop, and B. de Massy. 2010. PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice. *Science* 327:836–840.
- Berset-Brändli, L., J. Jaquiéry, T. Broquet, Y. Ulrich, and N. Perrin. 2008. Extreme heterochiasmy and nascent sex chromosomes in European tree frogs. *Proceedings of the Royal Society B-Biological Sciences* 275:1577–1585.
- Brandvain, Y. and G. Coop. 2012. Scrambling eggs: meiotic drive and the evolution of female recombination rates. *Genetics* 190:709–723.
- Burt, A., G. Bell, and P. Harvey. 1991. Sex differences in recombination. *Journal of Evolutionary Biology* 4:259–277.
- Chippindale, A. K., J. R. Gibson, and W. R. Rice. 2001. Negative genetic correlation for fitness between sexes reveals ontogenetic conflict in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 98:1671–1675.
- Connallon, T. and A. G. Clark. 2010. Sex-linkage, sex-specific selection, and the role of recombination in the evolution of sexually dimorphic gene expression. *Evolution* 64:3417–3442.

- Coop, G., X. Wen, C. Ober, J. K. Pritchard, and M. Przeworski. 2008. High-resolution mapping of crossovers reveals extensive variation in fine-scale recombination patterns among humans. *Science* 319:1395–1398.
- Cox, R. M. and R. Calsbeek. 2009. Sexually antagonistic selection, sexual dimorphism, and the resolution of intralocus sexual conflict. *American Naturalist* 173:176–187.
- Curtsinger, J. W. and I. L. Heisler. 1988. A diploid sexy son model. *American Naturalist* 132:437–453.
- Fledel-Alon, A., E. M. Leffler, Y. Guan, M. Stephens, G. Coop, and M. Przeworski. 2011. Variation in human recombination rates and its genetic determinants. *Plos One* 6:e20321.
- Foerster, K., T. Coulson, B. C. Sheldon, J. M. Pemberton, T. H. Clutton-Brock, and L. E. B. Kruuk. 2007. Sexually antagonistic genetic variation for fitness in red deer. *Nature* 447:1107–1110.
- Fry, J. D. 2010. The genomic location of sexually antagonistic variation: some cautionary comments. *Evolution* 64:1510–1516.
- Hansson, B., M. Akesson, J. Slate, and J. Pemberton. 2005. Linkage mapping reveals sex-dimorphic map distances in a passerine bird. *Proceedings of the Royal Society B-Biological Sciences* 272:2289–2298.
- Hedrick, P. W. 2007. Sex: Differences in mutation, recombination, selection, gene flow, and genetic drift. *Evolution* 61:2750–2771.
- Innocenti, P. and E. H. Morrow. 2010. The sexually antagonistic genes of *Drosophila melanogaster*. *PLoS Biology* 8:e1000335.
- Kong, A., J. Barnard, D. F. Gudbjartsson, G. Thorleifsson, G. Jonsdottir, S. Sigurdardottir, B. Richardsson, J. Jonsdottir, T. Thorgeirsson, M. L. Frigge, N. E.

- Lamb, S. Sherman, J. R. Gulcher, and K. Stefansson. 2004. Recombination rate and reproductive success in humans. *Nature Genetics* 36:1203–1206.
- Kong, A., G. Thorleifsson, D. F. Gudbjartsson, G. Masson, A. Sigurdsson, A. Jonasdottir, G. B. Walters, A. Jonasdottir, A. Gylfason, K. T. Kristinsson, S. A. Gudjonsson, M. L. Frigge, A. Helgason, U. Thorsteinsdottir, and K. Stefansson. 2010. Fine-scale recombination rate differences between sexes, populations and individuals. *Nature* 467:1099–1103.
- Lenormand, T. 2003. The evolution of sex dimorphism in recombination. *Genetics* 163:811–822.
- Lenormand, T. and J. Dutheil. 2005. Recombination difference between sexes: A role for haploid selection. *PLoS Biology* 3:e63.
- Long, T. A. F., A. F. Agrawal, and L. Rowe. 2012. The effect of sexual selection on offspring fitness depends on the nature of genetic variation. *Current Biology* 22:204–208.
- Mank, J. E. 2009. The evolution of heterochiasmy: the role of sexual selection and sperm competition in determining sex-specific recombination rates in eutherian mammals. *Genetical Research* 91:355–363.
- Otto, S. and T. Day. 2007. *A biologist's guide to mathematical modeling in ecology and evolution*. Princeton University Press, Princeton.
- Otto, S. P. 1991. On evolution under sexual and viability selection: a two-locus diploid model. *Evolution* 45:1443–1457.
- Patten, M., D. Haig, and F. Úbeda. 2010. Fitness variation due to sexual antagonism and linkage disequilibrium. *Evolution* 64:3638–3642.
- Priest, N. K., D. A. Roach, and L. F. Galloway. 2007. Mating-induced recombination in fruit flies. *Evolution* 61:160–167.

- Pröschel, M., Z. Zhang, and J. Parsch. 2006. Widespread adaptive evolution of *Drosophila* genes with sex-biased expression. *Genetics* 174:893–900.
- Rice, W. R. 1984. Sex-chromosomes and the evolution of sexual dimorphism. *Evolution* 38:735–742.
- Stevison, L. S. 2011. Male-mediated effects on female meiotic recombination. *Evolution* 66:905–911.
- Tedman-Aucoin, K. and A. F. Agrawal. 2011. The effect of deleterious mutations and age on recombination in *Drosophila melanogaster*. *Evolution* 66:575–585.
- Trivers, R., 1988. Sex differences in rates of recombination and sexual selection. Pp. 270–286, *in* R. Michod and B. Levin, eds. *The evolution of sex*. Sinauer Associates, Sunderland, MA.
- Úbeda, F., D. Haig, and M. M. Patten. 2011. Stable linkage disequilibrium owing to sexual antagonism. *Proceedings of the Royal Society B - Biological Sciences* 278:855–862.
- Wyman, M. J., A. F. Agrawal, and L. Rowe. 2010. Condition-dependence of the sexually dimorphic transcriptome in *Drosophila melanogaster*. *Evolution* 64:1836–1848.

A Model of selection and dominance

The m and f prefix or subscript indicate male and female, respectively. The h and s terms describe dominance and selection at the A locus. The g and t terms describe dominance and selection at the B locus. Epistasis, ϵ , is additive; having two copies of B_2 has twice the effect of having one copy. This model assumes *cis*-epistasis since haploid gametes are being modeled; recombination occurs between the A and B locus within a chromosome.

$$\begin{array}{ll}
 fA_1B_1A_1B_1 = 1 & mA_1B_1A_1B_1 = 1 \\
 fA_2B_1A_1B_1 = 1 - h_f s_f & mA_2B_1A_1B_1 = 1 - h_m s_m \\
 fA_1B_2A_1B_1 = 1 - g_f t_f & mA_1B_2A_1B_1 = 1 - g_m t_m \\
 fA_2B_2A_1B_1 = 1 - h_f s_f - g_f t_f - (1/2)\epsilon_f & mA_2B_2A_1B_1 = 1 - h_m s_m - g_m t_m - (1/2)\epsilon_m \\
 fA_2B_1A_2B_1 = 1 - s_f & mA_2B_1A_2B_1 = 1 - s_m \\
 fA_1B_2A_2B_1 = 1 - h_f s_f - g_f t_f & mA_1B_2A_2B_1 = 1 - h_m s_m - g_m t_m \\
 fA_2B_2A_2B_1 = 1 - s_f - g_f t_f - (1/2)\epsilon_f & mA_2B_2A_2B_1 = 1 - s_m - g_m t_m - (1/2)\epsilon_m \\
 fA_1B_2A_1B_2 = 1 - t_f & mA_1B_2A_1B_2 = 1 - t_m \\
 fA_2B_2A_1B_2 = 1 - h_f s_f - t_f - (1/2)\epsilon_f & mA_2B_2A_1B_2 = 1 - h_m s_m - t_m - (1/2)\epsilon_m \\
 fA_2B_2A_2B_2 = 1 - s_f - t_f - \epsilon_f & mA_2B_2A_2B_2 = 1 - s_m - t_m - \epsilon_m
 \end{array}$$

B Recursion equations and stability analysis

We write the frequency of haplotype $A_i B_j$ in generation n abstractly as $x^{ij}(n)$ (for eggs) and $y^{ij}(n)$ (for sperm). The recursion equations for these frequencies are given by the

general equations

$$x^{ij}(n+1) = \frac{1}{2\bar{w}_f} \left(x^{ij}(n) \sum_{\ell m} f_{ij\ell m} y^{\ell m}(n) + y^{ij}(n) \sum_{\ell m} f_{\ell mij} x^{\ell m}(n) \right) - (-1)^{i+j} r_f LD_f \quad (15)$$

$$y^{ij}(n+1) = \frac{1}{2\bar{w}_m} \left(x^{ij}(n) \sum_{\ell m} m_{ij\ell m} y^{\ell m}(n) + y^{ij}(n) \sum_{\ell m} m_{\ell mij} x^{\ell m}(n) \right) - (-1)^{i+j} r_m LD_m \quad (16)$$

where $\{i, j, \ell, m\} \in \{1, 2\}$, $f_{ij\ell m}(m_{ij\ell m})$ is the fitness of a female (male) with a maternal $A_i B_j$ and paternal $A_\ell B_m$ genotype, $r_f(r_m)$ is the recombination rate of females (males), $\bar{w}_f(\bar{w}_m)$ is the mean female (male) fitness, and $LD_f(LD_m)$ is the female (male) linkage disequilibrium rate. Mean fitness is defined as

$$\bar{w}_f = \sum_{ijkl=1}^2 x^{ij} y^{\ell m} f_{ij\ell m} \quad (17)$$

$$\bar{w}_m = \sum_{ijkl=1}^2 x^{ij} y^{\ell m} m_{ij\ell m} \quad (18)$$

We define the sex specific linkage disequilibrium factors by

$$LD_f = \frac{1}{2\bar{w}_f} \sum_{ij\ell m}^{i \neq \ell, k \neq m} (-1)^{i+j} x^{ij} y^{\ell m} f_{ij\ell m} \quad (19)$$

$$LD_m = \frac{1}{2\bar{w}_m} \sum_{ij\ell m}^{i \neq \ell, k \neq m} (-1)^{i+j} x^{ij} y^{\ell m} m_{ij\ell m} \quad (20)$$

If we put the haplotype frequencies in a vector:

$$X(n) = \{x_{11}(n), y_{11}(n), x_{21}(n), y_{21}(n), x_{12}(n), y_{12}(n), x_{22}(n), y_{22}(n)\}, \quad (21)$$

then the i, j entry of the Jacobian stability matrix J associated with these recursion equations is given by

$$J_{ij} = \frac{\partial X_i(n+1)}{\partial X_j(n)}. \quad (22)$$

The eigenvalues associated with the lower right 4×4 sub-matrix of \mathbf{J} will be the ones relevant to the invasion of the B_2 allele, the one relevant to our study. These are reproduced in the main text. We derive them by first finding the characteristic in full generality, and then solve it order by order in an expansion in a small parameter. In our calculations, we do not need to assume weak selection, but always assume weak recombination and a small intersexual fitness difference. That is, we assume all recombination rates are $\ll 1$, and that the difference between the male and female equilibrium allele frequencies is also small – i.e., $|p_m - p_f| \ll 1$. In the main text of the paper, we have also assumed weak selection when we write most of our equations. These calculations, including equations without the assumption of weak selection, are all reproduced in the associated `Mathematica` notebook (see journal version’s online supplementary materials).

Table 1: Shorthand for genotype fitnesses.

Diploid genotype	Female fitness	Male fitness
$A_1B_1 A_1B_1$	f_{11}	m_{11}
$A_1B_1 A_2B_1$	f_{21}	m_{21}
$A_1B_1 A_1B_2$	f_{12}	m_{12}
$A_1B_1 A_2B_2$	f_{22C}	m_{22C}
$A_1B_2 A_2B_1$	f_{22R}	m_{22R}
$A_2B_1 A_2B_1$	f_{31}	m_{31}
$A_1B_2 A_1B_2$	f_{13}	m_{13}
$A_2B_1 A_2B_2$	f_{32}	m_{32}
$A_2B_2 A_1B_2$	f_{23}	m_{23}
$A_2B_2 A_2B_2$	f_{33}	m_{33}

C Online Supplementary Material: eigenvalues to higher order in r_f and r_m

By performing a perturbative, order-by-order solution of the characteristic polynomial of the recursion equations, we were able to find an expression for the eigenvalue for invasion of the B_2 allele with the A locus polymorphic to fourth order in r , under the simplifying assumption that $p_m = p_f$. This expression is given by:

$$\begin{aligned}
\lambda = & \frac{1}{2w_f} (f_{32}p + f_{22C}q(1 - r_f)) + \frac{1}{2w_m} (m_{32}p + m_{22C}q(1 - r_m)) \tag{23} \\
& + \frac{pq(f_{22C}w_m r_f + m_{22C}w_f r_m)(f_{22R}w_m r_f + m_{22R}w_f r_m)}{2w_m w_f (m_{32}w_f p - m_{22R}w_f p + m_{22C}w_f q - m_{12}w_f q + f_{32}w_m p - f_{12}w_m q + f_{22C}w_m q - m_{22R}w_m p)} \\
& + \frac{pq(f_{22C}w_m r_f + m_{22C}w_f r_m)(f_{22R}w_m r_f + m_{22R}w_f r_m)((f_{22R}p - f_{22C}q)w_m r_f + (m_{22R}p - m_{22C}q)w_f r_m)}{2w_m w_f (m_{32}w_f p - m_{22R}w_f p + m_{22C}w_f q - m_{12}w_f q + f_{32}w_m p - f_{12}w_m q + f_{22C}w_m q - m_{22R}w_m p)^2} \\
& + \frac{pq(f_{22C}w_m r_f + m_{22C}w_f r_m)(f_{22R}w_m r_f + m_{22R}w_f r_m)\mathcal{A}}{2w_m w_f (m_{32}w_f p - m_{22R}w_f p + m_{22C}w_f q - m_{12}w_f q + f_{32}w_m p - f_{12}w_m q + f_{22C}w_m q - m_{22R}w_m p)^3}
\end{aligned}$$

where we have rewritten part of the fourth numerator as

$$\begin{aligned}\mathcal{A} &= w_m^2 (f_{22C}^2 q^2 + 3f_{22C} f_{22Rp} q + f_{22Rp}^2) r_f^2 \\ &\quad + w_m w_f (f_{22Rp} (3m_{22C} q + 2m_{22Rp}) + f_{22C} q (2m_{22C} q + 3m_{22Rp})) r_f r_m \\ &\quad + w_f^2 (m_{22C}^2 q^2 + 3m_{22C} m_{22Rp} q + m_{22Rp}^2) r_m^2\end{aligned}$$

This extremely complex expression is curious because the denominators for the second, third, and fourth order terms are small numbers, which is not what one would expect. If we write $q = (1 - p)$, set the mean fitnesses to one for simplicity, and expand the fitnesses as $f = 1 - \delta f$, we have

$$\begin{aligned}\text{denominator} &\propto \delta m_{22C} - \delta m_{12} + \delta f_{22C} - \delta f_{12} \\ &\quad + p(\delta m_{32} - \delta m_{22R} - \delta m_{22C} + \delta m_{12} + \delta f_{32} - \delta f_{22R} - \delta f_{22C} + \delta f_{12}).\end{aligned}\quad (24)$$

All of the δf and δm terms are small numbers, so their sum (equation 24) is also small. Thus, even if the numerators of the each term in the fourth order expansion of equation 23 are small, each term is still relatively large. This means that the subsequent higher order corrections to the eigenvalue adjust the eigenvalue by significantly more than one would have expected. Normally, higher order corrections have a smaller impact. This effect will have a greater impact for larger values of r (since this was an expansion in r). In other words, when sex-specific recombination rates are near the borderline of invasion ($\lambda_{CC} \simeq 1$), adding higher order corrections to λ_{CC} will affect the slope and location of the line demarcating invasion.