Goal oriented adaptivity in the IRGNM for parameter identification in PDEs II: all-at once formulations

B. Kaltenbacher A. Kirchner B. Vexler

September 26, 2018

Abstract

In this paper we investigate adaptive discretization of the iteratively regularized Gauss-Newton method IRGNM. All-at-once formulations considering the PDE and the measurement equation simultaneously allow to avoid (approximate) solution of a potentially nonlinear PDE in each Newton step as compared to the reduced form [22]. We analyze a least squares and a generalized Gauss-Newton formulation and in both cases prove convergence and convergence rates with a posteriori choice of the regularization parameters in each Newton step and of the stopping index under certain accuracy requirements on four quantities of interest. Estimation of the error in these quantities by means of a weighted dual residual method is discussed, which leads to an algorithm for adaptive mesh refinement. Numerical experiments with an implementation of this algorithm show the numerical efficiency of this approach, which especially for strongly nonlinear PDEs outperforms the nonlinear Tikhonov regularization considered in [21].

1 Introduction

We consider the problem of identifying a parameter q in a PDE

$$A(q,u) = f \tag{1}$$

from measurements of the state u

$$C(u) = g, \qquad (2)$$

where $q \in Q$, $u \in V$, $g \in G$, Q, V, G are Hilbert spaces and $A: Q \times V \to W^*$ with W^* denoting the dual space of some Hilbert space W and $C: V \to G$ differential and observation operators, respectively. Among many others, for example the classical model problem of identifying the diffusion coefficient q in the linear elliptic PDE

$$-\nabla(q\nabla u) = f$$
 in Ω

from measurements of u in Ω can be cast in this form with $Q \subseteq L^{\infty}(\Omega)$, $V, W \subseteq H^{1}(\Omega)$, $G = L^{2}(\Omega)$, $A(q, u) = -\nabla(q\nabla u)$ and C the embedding of $H^{1}(\Omega)$ into $L^{2}(\Omega)$.

The usual approach for tackling such inverse problems is to reduce them to an operator equation

$$F(q) = g, (3)$$

where $F = C \circ S$ is the composition of the parameter-to-solution map for (1)

with the measurement operator C. The forward operator F will then be a nonlinear operator between Q and G with typically unbounded inverse, so that recovery of q is an ill-posed problem. Since the given data g^{δ} are noisy with some noise level δ

$$||g - g^{\delta}|| \le \delta,\tag{5}$$

regularization is needed.

We will here as in [22] consider the paradigm of the Iteratively Regularized Gauss-Newton Method (IRGNM) cf., e.g., [3, 4, 7, 18, 20, 23] and its adaptive discetization. However, instead of reducing to (3), we will simulteously consider the measurement equation and the PDE:

$$C(u) = g \text{ in } G \tag{6}$$

$$A(q,u) = f \text{ in } W^* \tag{7}$$

as a system of operator equations for (q, u), which we will abbreviate by

$$\mathbf{F}(q,u) = \mathbf{g},\tag{8}$$

where

$$\mathbf{F} \colon Q \times V \to G \times W^* \,, \quad \mathbf{F}(q, u) = \begin{pmatrix} C(u) \\ A(q, u) \end{pmatrix} \,, \qquad \text{and} \qquad \mathbf{g} = \begin{pmatrix} g \\ f \end{pmatrix} \in G \times W^* \,. \tag{9}$$

The noisy data for this all-at-once formulation is denoted by

$$\mathbf{g}^{\delta} = \begin{pmatrix} g^{\delta} \\ f \end{pmatrix} \in G \times W^*$$
.

This will allow us to avoid a major drawback of the method in [22], namely the necessity of solving the possibly nonlinear PDE (to a certain precision) in each Newton step in order to evaluate F(q) = C(S(q)). Another key difference to the paper [22] is that here the *u* part of the previous iterate will not be subject to new discretization in the current iteration but keep its (usually coarser, hence cheaper) discretization from the previous step.

Therewith, we will arrive at iterations of the form

$$\begin{aligned} (q^{k}, u^{k}) \\ &= \arg\min_{q, u} \quad \varrho \|A_{q}'(q^{k-1}, u^{k-1})(q - q^{k-1}) + A_{u}'(q^{k-1}, u^{k-1})(u - u^{k-1}) + A(q^{k-1}, u^{k-1}) - f\|_{W^{*}}^{r} \\ &+ \|C(u^{k-1}) + C'(u^{k-1})(u - u^{k-1}) - g^{\delta}\|_{G}^{2} + \alpha_{k} \|q - q_{0}\|_{Q}^{2} . \end{aligned}$$

$$(10)$$

with $\rho > 0, r \in \{1, 2\}.$

For r = 2, this yields a least squares formulation, see Section 2. In case r = 1 and ρ sufficiently large, by exactness of the norm with exponent one as a penalty, this leads to a Generalized Gauss-Newton type [8] form of the IRGNM

$$\begin{aligned} (q^{k}, u^{k}) \\ &= \arg\min_{q, u} \|C(u^{k-1}) + C'(u^{k-1})(u - u^{k-1}) - g^{\delta}\|_{G}^{2} + \alpha_{k} \|q - q_{0}\|_{Q}^{2} \\ &\text{s.t. } A'_{q}(q^{k-1}, u^{k-1})(q - q^{k-1}) + A'_{u}(q^{k-1}, u^{k-1})(u - u^{k-1}) + A(q^{k-1}, u^{k-1}) = f \text{ in } \mathbb{M}^{1}. \end{aligned}$$

see Section 3.

Remark 1. Although q^k, u^k obviously depend on δ , i.e. $q^k = q^{k,\delta}, u^k = u^{k,\delta}$, we omit the superscript δ for better readability.

All-at-once formulations have also been considered, e.g., in [1, 2, 9, 10], however, our approach focuses on adaptive discretization using a posteriori error estimators. Additionally it differs from the previous ones in the following sense: In [9, 10] a Levenberg-Marquardt approach is considered, whereas we work with an iterative regularized Gauss-Newton approach which allows us to also prove convegergence rates (which is an involved task in a Levenberg-Marquardt setting, that has been resolved only relatively recently, [16]). Moreover we use a different regularization parameter choice in each Newton step than [9, 10]. The papers [1, 2] put more emphasis on computational aspects and applications than we do here.

For both cases r = 1, r = 2 in (10) we will investigate convergence and convergence rates in the continuous and adaptively discretized setting with discrepancy type choice of α_k (which in most of what follows will be replaced by $\frac{1}{\beta_k}$) and the overal stopping index k_* . The discretization errors with respect to certain quantities of interest will serve as refinement criteria during the Gauss-Newton iteration, where at the same time, we control the size of the regularization parameter. In order to estimate this discretization error we use goal-oriented error estimators (cf. [5, 6]). For the least squares case we will (for the sake of completeness but not in the main steam of this paper) also provide a result on convergence with a priori parameter choice in the continuous setting, see the appendix. In Section 4, we will provide numerical results and in Section 5 some conclusions.

Throughout this paper, we will make the following assumptions:

Assumption 1. There exists a solution $(q^{\dagger}, u^{\dagger}) \in \mathcal{B}_{\rho}(q_0, u_0) \subset \mathcal{D}(A) \cap (Q \times \mathcal{D}(C)) \subseteq Q \times V$ to (8), where (q_0, u_0) is some initial guess and ρ (not to be confused with the penalty parameter ρ in (10)) is the radius of the neighborhood in which local convergence of the Newton type iterations under consideration will be shown.

Assumption 2. The PDE (1) and especially also its linearization at (q, u) is uniquely and stably solvable.

Assumption 3. The norms in G, Q, as well as the operator C and the semilinear form $a: Q \times V \times W \to \mathbb{R}$ defined by the relation $a(q, u)(v) = \langle A(q, u), v \rangle_{W^*, W}$ (where $\langle ., . \rangle_{W^*, W}$ denotes the duality pairing between W^* and W) are assumed to be evaluated exactly.

2 A least squares formulation

Direct application of the IRGNM to (6), (7), i.e., to the all-at-once system (8) yields the iteration

$$\begin{pmatrix} q^{k} \\ u^{k} \end{pmatrix} = \begin{pmatrix} q^{k-1} \\ u^{k-1} \end{pmatrix} - \begin{pmatrix} \mathbf{F}'(q^{k-1}, u^{k-1})^{*} \mathbf{F}'(q^{k-1}, u^{k-1}) + \begin{pmatrix} \alpha_{k} \operatorname{id} & 0 \\ 0 & \mu_{k} \operatorname{id} \end{pmatrix} \end{pmatrix}^{-1} \\ \cdot \begin{pmatrix} \mathbf{F}'(q^{k-1}, u^{k-1})^{*} (\mathbf{F}(q^{k-1}, u^{k-1}) - \mathbf{g}^{\delta}) + \begin{pmatrix} \alpha_{k}(q^{k-1} - q_{0}) \\ \mu_{k}(u^{k-1} - u_{0}) \end{pmatrix} \end{pmatrix}$$
(12)

with regularization parameters α_k , μ_k for the q and u part of the iterates, respectively.

We will first of all show that Assumption 2 allows us to set the regularization parameter μ_k for the *u* part to zero. For this purpose, we introduce the abbreviations

$$K \colon V \to W^*, \quad K \coloneqq A'_u(q, u) \quad \text{and} \quad L \colon Q \to W^*, \quad L \coloneqq A'_q(q, u)$$
(13)

with Hilbert space adjoints $K^* \colon W^* \to V$ and $L^* \colon W^* \to Q$, i.e.,

$$(Lq, w^*)_{W^*} = (q, L^* w^*)_Q \qquad \forall q \in Q, w^* \in W^*, (Kv, w^*)_{W^*} = (v, K^* w^*)_V \qquad \forall v \in V, w^* \in W^*,$$
(14)

where $(., .)_{W^*}$ and $(., .)_V$ denote the inner products in W^* and V.

In the same way we define the Hilbert space adjoint $C'(u)^* \colon G \to V$ for $C'(u) \colon V \to G$, i.e.,

$$(C'(u)(\delta u), \varphi)_G = (\delta u, C'(u)^* \varphi)_V,$$

where $(.,.)_G$ denotes the inner product in G.

We denote the derivate of **F** at a pair (q, u) by **T**, i.e.,

$$\mathbf{T} \colon Q \times V \to G \times W^*, \quad \mathbf{T} = \mathbf{F}'(q, u) = \begin{pmatrix} 0 & C'(u) \\ A'_q(q, u) & A'_u(q, u) \end{pmatrix} = \begin{pmatrix} 0 & C'(u) \\ L & K \end{pmatrix}$$
(15)

and define the norm

$$\left\| \begin{pmatrix} q \\ u \end{pmatrix} \right\|_{Q \times V}^2 \coloneqq \left\| q \right\|_Q^2 + \left\| u \right\|_V^2 \quad \text{and the operator norm} \quad \left\| T \right\|_{Q \times V} \coloneqq \sup_{x \in Q \times V, x \neq 0} \frac{\left\| Tx \right\|_{Q \times V}}{\left\| x \right\|_{Q \times V}}.$$
(16)

for some $x \in Q \times V$ and some operator $T \colon Q \times V \to Q \times V$.

Further we define

$$\mathbf{Y}_{\alpha,\mu} \coloneqq \left(\mathbf{T}^* \mathbf{T} + \begin{pmatrix} \alpha \, \mathrm{id} & 0 \\ 0 & \mu \, \mathrm{id} \end{pmatrix} \right)$$
(17)

for $\alpha > 0, \ \mu \ge 0$.

Lemma 1. Under Assumption 2

(i) for any $\alpha > 0$, $\mu \ge 0$ the inverse $\mathbf{Y}_{\alpha,\mu}^{-1}$ of $\mathbf{Y}_{\alpha,\mu}$ exists

$$\left\|\mathbf{Y}_{\alpha,\mu}^{-1}\mathbf{T}^{*}\mathbf{T}\right\|_{Q\times V} \leq 1 + \max\{\alpha,\mu\} \left\|\mathbf{Y}_{\alpha,\mu}^{-1}\right\|_{Q\times V},$$

(iii)

(ii)

$$\left\|\mathbf{Y}_{\alpha,\mu}^{-1}\right\|_{Q\times V} \le c_T \left(\frac{1}{\alpha} + 1\right) \tag{18}$$

for all $\alpha \in (0,1]$, $\mu \geq 0$ and some $c_T > 0$ independent of α, μ , where the bound c_T in (18) is independent of q and u, if the operators K, K^{-1} and L, are bounded uniformly in (q, u).

Proof. (i): With the abbreviations

$$P = L^*L + \alpha \operatorname{id}$$
 and $M = C'(u)^*C'(u) + K^*K + \mu \operatorname{id}$

we have

$$\mathbf{Y}_{\alpha,\mu} = \left(\begin{array}{cc} P & L^*K \\ K^*L & M \end{array}\right) \,.$$

Since Assumption 3 implies that K is invertible, M^{-1} exists, such that we can define some kind of Schur complement

$$N := P - L^* K M^{-1} K^* L = L^* L + \alpha \operatorname{id} - L^* K M^{-1} K^* L.$$

We will now show that N is also invertible. Using the fact that

$$\begin{split} \|M^{-1/2}K^*\|_{W^* \to V}^2 &= \|KM^{-1/2}\|_{V \to W^*}^2 \\ &= \sup_{v \in V, v \neq 0} \frac{\|KM^{-1/2}v\|_{W^*}^2}{\|v\|_V^2} \\ &= \sup_{v \in V, v \neq 0} \frac{\|Kv\|_{W^*}^2}{\|M^{1/2}v\|_V^2} \\ &= \sup_{v \in V, v \neq 0} \frac{\|Kv\|_{W^*}^2}{\|C'(u)(v)\|_G^2 + \|Kv\|_{W^*}^2 + \mu \|v\|_V^2} \leq 1 \,. \end{split}$$

for any $q\in Q$ we get

$$(Nq,q)_Q = (L^*Lq + \alpha q - L^*KM^{-1}K^*Lq, q)_Q$$

$$\geq \|Lq\|_{W^*}^2 + \alpha \|q\|_Q^2 - \|M^{-1/2}K^*\|_{W^* \to V}^2 \|Lq\|_{W^*}^2$$

$$\geq \alpha \|q\|_Q^2,$$

which implies the existence of N^{-1} , since M and therewith also N is self-adjoint. For

$$O_{\alpha\mu} \coloneqq \begin{pmatrix} N^{-1} & -N^{-1}L^*KM^{-1} \\ -M^{-1}K^*LN^{-1} & M^{-1} + M^{-1}K^*LN^{-1}L^*KM^{-1} \end{pmatrix}$$
(19)

there holds

$$O_{\alpha\mu} \left(\begin{array}{cc} P & L^*K \\ K^*L & M \end{array} \right) = \left(\begin{array}{cc} A & B \\ C & D \end{array} \right)$$

with

$$\begin{split} &A \coloneqq N^{-1} \left(P - L^* K M^{-1} K^* L \right) = \mathrm{id} \\ &B \coloneqq N^{-1} L^* K + -N^{-1} L^* K M^{-1} M = 0 \\ &C \coloneqq -M^{-1} K^* L N^{-1} P + \left(M^{-1} + M^{-1} K^* L N^{-1} L^* K M^{-1} \right) K^* L \\ &= -M^{-1} K^* L \left[N^{-1} \left(P - L^* K M^{-1} K^* L \right) - \mathrm{id} \right] = 0 \\ &D \coloneqq -M^{-1} K^* L N^{-1} L^* K + \left(M^{-1} + M^{-1} K^* L N^{-1} L^* K M^{-1} \right) M = \mathrm{id} \,, \end{split}$$

we have

$$O_{\alpha\mu} = \mathbf{Y}_{\alpha,\mu}^{-1} \,. \tag{20}$$

(ii):

$$\begin{split} \left\| \mathbf{Y}_{\alpha,\mu}^{-1} \mathbf{T}^* \mathbf{T} \right\|_{Q \times V} &= \left\| \operatorname{id} - \mathbf{Y}_{\alpha,\mu}^{-1} \begin{pmatrix} \alpha \operatorname{id} & 0 \\ 0 & \mu \operatorname{id} \end{pmatrix} \right\|_{Q \times V} \\ &\leq 1 + \max\{\alpha,\mu\} \left\| \mathbf{Y}_{\alpha,\mu}^{-1} \right\|_{Q \times V} \end{split}$$

(iii): For any $v \in V$ we get

$$(Mv,v)_{V} = \|C'(u)(v)\|_{G}^{2} + \|Kv\|_{W^{*}}^{2} + \mu\|v\|_{V}^{2} \ge \|Kv\|_{W^{*}}^{2} \ge \frac{1}{\|K^{-1}\|_{W^{*} \to V}^{2}} \|v\|_{V}^{2},$$

hence we have

$$\|N^{-1}\|_{Q \to Q} \le \frac{1}{\alpha} , \quad \|M^{-1}\|_{V \to V} \le \|K^{-1}\|_{W^* \to V}^2 .$$
(21)

For $O_{\alpha\mu}$ (cf. (19)) this yields

$$\begin{split} \|O_{\alpha\mu}\|_{Q\times V}^{2} &\leq \sup_{(q,u)\in Q\times V, (q,u)\neq 0} \frac{\left\|N^{-1}q + N^{-1}L^{*}KM^{-1}u\right\|_{Q}^{2}}{\|q\|_{Q}^{2} + \|u\|_{V}^{2}} \\ &+ \sup_{(q,u)\in Q\times V, (q,u)\neq 0} \frac{\left\|-M^{-1}KLN^{-1}q + \left(M^{-1} + M^{-1}K^{*}LN^{-1}L^{*}KM^{-1}\right)u\right\|_{V}^{2}}{\|q\|_{Q}^{2} + \|u\|_{V}^{2}} \\ &\leq 2\left(\left\|N^{-1}\right\|_{Q\rightarrow Q}^{2} + \left\|N^{-1}L^{*}KM^{-1}\right\|_{V\rightarrow Q}^{2} \\ &+ \left\|M^{-1}K^{*}LN^{-1}\right\|_{Q\rightarrow V}^{2} + \left\|M^{-1} + M^{-1}K^{*}LN^{-1}L^{*}KM^{-1}\right\|_{V\rightarrow V}^{2}\right) \\ &\leq 2\left\|N^{-1}\right\|_{Q\rightarrow Q}^{2}\left(1 + \left\|L\right\|_{Q\rightarrow W^{*}}^{2}\left\|K\right\|_{V\rightarrow W^{*}}^{2}\left\|M^{-1}\right\|_{V\rightarrow V}^{2}\right)^{2} + 2\left\|M^{-1}\right\|_{V\rightarrow V}^{2} \\ &\leq \frac{2}{\alpha^{2}}(1 + \left\|L\right\|_{Q\rightarrow W^{*}}^{2}\left\|K\right\|_{V\rightarrow W^{*}}^{2}\left\|K^{-1}\right\|_{W^{*}\rightarrow V}^{4}\right)^{2} + 2\left\|K^{-1}\right\|_{W^{*}\rightarrow V}^{4} \end{split}$$

Motivated by Lemma 1, and setting $\alpha_k = \frac{1}{\beta_k}$ we define a regularized iteration by

$$\begin{pmatrix} q^{k} \\ u^{k} \end{pmatrix} = \begin{pmatrix} q^{k-1} \\ u^{k-1} \end{pmatrix} - \left(\mathbf{F}'(q^{k-1}, u^{k-1})^{*} \mathbf{F}'(q^{k-1}, u^{k-1}) + \frac{1}{\beta_{k}} \begin{pmatrix} \operatorname{id} & 0 \\ 0 & 0 \end{pmatrix} \right)^{-1} \cdot \left(\mathbf{F}'(q^{k-1}, u^{k-1})^{*} (\mathbf{F}(q^{k-1}, u^{k-1}) - \mathbf{g}^{\delta}) + \frac{1}{\beta_{k}} \begin{pmatrix} q^{k-1} - q_{0} \\ 0 \end{pmatrix} \right)$$
(22)

or equivalently $\begin{pmatrix} q^k \\ u^k \end{pmatrix}$ as solution to the unconstrained minimization problem

$$\min_{(q,u)\in Q\times V} \mathcal{T}_{\beta_k}(q,u) \coloneqq \|L_{k-1}(q-q^{k-1}) + K_{k-1}(u-u^{k-1}) + A(q^{k-1}, u^{k-1}) - f\|_{W^*}^2
+ \|C(u^{k-1}) + C'(u^{k-1})(u-u^{k-1}) - g^{\delta}\|_G^2 + \frac{1}{\beta_k}\|q-q_0\|_Q^2$$
(23)

with the abbreviations

$$L_{k-1} = A'_{q}(q^{k-1}, u^{k-1}) \quad \text{and} \quad K_{k-1} = A'_{u}(q^{k-1}, u^{k-1}), \quad (24)$$

where we have set the regularization parameter for the component u to zero, which is justified by (i) in Lemma 1.

The optimality conditions of first order for (23) read

$$\begin{aligned} 0 &= (\mathcal{T}_{\beta_k})'_q(q,u)(\delta q) \\ &= 2(L_{k-1}(q-q^{k-1}) + K_{k-1}(u-u^{k-1}) + A(q^{k-1},u^{k-1}) - f, L_{k-1}(\delta q))_{W^*} + 2(q-q_0,\delta q)_Q \\ 0 &= (\mathcal{T}_{\beta_k})'_u(q,u)(\delta u) \\ &= 2(L_{k-1}(q-q^{k-1}) + K_{k-1}(u-u^{k-1}) + A(q^{k-1},u^{k-1}) - f, K_{k-1}(\delta u))_{W^*} \\ &+ 2(C(u^{k-1}) + C'(u^{k-1})(u-u^{k-1}) - g^{\delta}, C'(u^{k-1})(\delta u))_G \end{aligned}$$

We refer to the Appendix for a convergence and convergence rates results for (22) with a priori choice of the regularization parameters and in a continuous setting.

Here we are rather interested in a posteriori parameter choice rules and adaptive discretization. So in each step k we will replace the infinite dimensional spaces Q, V, W in (22) by finite dimensional ones $Q_h, V_h, W_h = Q_{h_k}, V_{h_k}, W_{h_k}$

$$\begin{pmatrix} q^{k} \\ u^{k} \end{pmatrix} = \arg\min_{q \in Q_{h}, u \in V_{h}} \|L_{k-1}(q - q_{\text{old}}) + K_{k-1}(u - u_{\text{old}}) + A(q_{\text{old}}, u_{\text{old}}) - f\|_{W_{h}^{*}}^{2} + \|C(u_{\text{old}}) + C'(u_{\text{old}})(u - u_{\text{old}}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta_{k}}\|q - q_{0}\|_{Q}^{2}.$$

$$(25)$$

where $(q_{\text{old}}, u_{\text{old}}) = (q^{k-1}, u^{k-1}) = (q^{k-1}_{h_{k-1}}, u^{k-1}_{h_{k-1}})$ is the previous iterate, which itself is discretized by the use of spaces $Q_{h_{k-1}}, V_{h_{k-1}}, W_{h_{k-1}}$. The discretization h_k may be different in each Newton step (typically it will get finer for increasing k), but we suppress dependence of h on k in our notation in most of what follows.

To still obtain convergence of these discretized iterates, it is essential to control the discretization error in certain quantities, which are defined, analogously to [22], via the functionals

$$I_1: Q \times V \times Q \times V \times \mathbb{R}, \to \mathbb{R}$$
$$I_2: Q \times V \times Q \times V, \to \mathbb{R}$$
$$I_3: Q \times V, \to \mathbb{R}$$
$$I_4: Q \times V, \to \mathbb{R}$$

where we insert the previous and current iterates $(q_{old}, u_{old}), (q, u)$, respectively:

$$I_{1}(q_{\text{old}}, u_{\text{old}}, q, u, \beta) = \left\| \mathbf{F}'(q_{\text{old}}, u_{\text{old}}) \begin{pmatrix} q - q_{\text{old}} \\ u - u_{\text{old}} \end{pmatrix} + \mathbf{F}(q_{\text{old}}, u_{\text{old}}) - \mathbf{g}^{\delta} \right\|_{G \times W^{*}}^{2} + \frac{1}{\beta} \|q - q_{0}\|_{Q}^{2}$$

$$= \left\| A'_{q}(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}}) + A'_{u}(q_{\text{old}}, u_{\text{old}})(u - u_{\text{old}}) + A(q_{\text{old}}, u_{\text{old}}) - f \right\|_{W^{*}}^{2}$$

$$+ \left\| C'(u)(u - u_{\text{old}}) + C(u_{\text{old}}) - g^{\delta} \right\|_{G}^{2} + \frac{1}{\beta} \|q - q_{0}\|_{Q}^{2}$$

$$I_{2}(q_{\text{old}}, u_{\text{old}}, q, u) = \left\| \mathbf{F}'(q_{\text{old}}, u_{\text{old}}) \begin{pmatrix} q - q_{\text{old}} \\ u - u_{\text{old}} \end{pmatrix} + \mathbf{F}(q_{\text{old}}, u_{\text{old}}) - \mathbf{g}^{\delta} \right\|_{G \times W^{*}}^{2}$$

$$= \left\| A'_{q}(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}}) + A'_{u}(q_{\text{old}}, u_{\text{old}}) - \mathbf{g}^{\delta} \right\|_{G \times W^{*}}^{2}$$

$$I_{3}(q_{\text{old}}, u_{\text{old}}) = \left\| \mathbf{F}(q_{\text{old}}, u_{\text{old}}) - \mathbf{g}^{\delta} \right\|_{G \times W^{*}}^{2}$$

$$= \left\| A(q_{\text{old}}, u_{\text{old}}) - g^{\delta} \right\|_{G \times W^{*}}^{2}$$

$$= \left\| A(q_{\text{old}}, u_{\text{old}}) - g^{\delta} \right\|_{G \times W^{*}}^{2}$$

$$I_{4}(q, u) = \left\| \mathbf{F}(q, u) - \mathbf{g}^{\delta} \right\|_{G \times W^{*}}^{2}$$

$$= \left\| A(q, u) - f \right\|_{W^{*}}^{2} + \left\| C(u) - g^{\delta} \right\|_{G}^{2}.$$
(26)

and quantities of interest

$$I_{1}^{k} = I_{1}(q_{\text{old}}^{k}, u_{\text{old}}^{k}, q^{k}, u^{k}, \beta_{k})$$

$$I_{2}^{k} = I_{2}(q_{\text{old}}^{k}, u_{\text{old}}^{k}, q^{k}, u^{k})$$

$$I_{3}^{k} = I_{3}(q_{\text{old}}^{k}, u_{\text{old}}^{k})$$

$$I_{4}^{k} = I_{4}(q^{k}, u^{k}).$$
(27)

Their discrete analogs are correspondingly defined by

$$I_{1,h}: Q \times V \times Q \times V \times \mathbb{R}, \to \mathbb{R}$$

$$I_{2,h}: Q \times V \times Q \times V, \to \mathbb{R}$$

$$I_{3,h}: Q \times V, \to \mathbb{R}$$

$$I_{3,h}: Q \times V, \to \mathbb{R}$$

$$I_{4,h}: Q \times V, \to \mathbb{R}$$

$$I_{4,h}: Q \times V, \to \mathbb{R}$$

$$I_{1,h}(q_{\text{old}}, u_{\text{old}}, q, u, \beta) = \left\| \mathbf{F}'(q_{\text{old}}, u_{\text{old}}) \begin{pmatrix} q - q_{\text{old}} \\ u - u_{\text{old}} \end{pmatrix} + \mathbf{F}(q_{\text{old}}, u_{\text{old}}) - \mathbf{g}^{\delta} \right\|_{G \times W_{h}^{*}}^{2} + \frac{1}{\beta} \|q - q_{0}\|_{Q}^{2}$$

$$= \|A'_{q}(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}}) + A'_{u}(q_{\text{old}}, u_{\text{old}})(u - u_{\text{old}}) + A(q_{\text{old}}, u_{\text{old}}) - f \|_{W_{h}^{*}}^{2}$$

$$+ \|C'(u)(u - u_{\text{old}}) + C(u_{\text{old}}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta} \|q - q_{0}\|_{Q}^{2}$$

$$I_{2,h}(q_{\text{old}}, u_{\text{old}}, q, u) = \left\|\mathbf{F}'(q_{\text{old}}, u_{\text{old}}) \begin{pmatrix} q - q_{\text{old}} \\ u - u_{\text{old}} \end{pmatrix} + \mathbf{F}(q_{\text{old}}, u_{\text{old}}) - \mathbf{g}^{\delta} \right\|_{G \times W_{h}^{*}}^{2}$$

$$= \|A'_{q}(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}}) + A'_{u}(q_{\text{old}}, u_{\text{old}}) - \mathbf{g}^{\delta} \|_{G \times W_{h}^{*}}^{2}$$

$$I_{3,h}(q_{\text{old}}, u_{\text{old}}) = \left\|\mathbf{F}(q_{\text{old}}, u_{\text{old}}) - \mathbf{g}^{\delta} \right\|_{G \times W_{h}^{*}}^{2}$$

$$= \|A(q_{\text{old}}, u_{\text{old}}) - f \|_{W_{h}^{*}}^{2} + \|C(u_{\text{old}}) - g^{\delta} \|_{G}^{2}$$

$$I_{4,h}(q, u) = \left\|\mathbf{F}(q, u) - \mathbf{g}^{\delta} \right\|_{G \times W_{h}^{*}}^{2}$$

$$= \|A(q, u) - f \|_{W_{h}^{*}}^{2} + \|C(u) - g^{\delta} \|_{G}^{2}$$
(28)

and

$$I_{1,h}^{k} = I_{1,h}(q_{\text{old}}^{k}, u_{\text{old}}^{k}, q_{h}^{k}, u_{h}^{k}, \beta_{k})$$

$$I_{2,h}^{k} = I_{2,h}(q_{\text{old}}^{k}, u_{\text{old}}^{k}, q_{h}^{k}, u_{h}^{k},)$$

$$I_{3,h}^{k} = I_{3,h}(q_{\text{old}}^{k}, u_{\text{old}}^{k})$$

$$I_{4,h}^{k} = I_{4,h}(q_{h}^{k}, u_{h}^{k}).$$
(29)

At the end of each iteration step we set

$$q_{\text{old}}^{k+1} = q_h^k \quad \text{and} \quad u_{\text{old}}^{k+1} = u_h^k \,. \tag{30}$$

Remark 2. Note that here neither q_{old} nor u_{old} are subject to new adaptive discretization in the current step, but they are taken as fixed quantities from the previous step. This is different from [22], where u_{old} also depends on the current discretization.

For (27) and (29) we assume that the norms in G and Q are evaluated exactly cf. Assumption 3.

In our convergence proofs we will compare the quantities of interest $I_{i,h}^k$ with those I_i^k that would be obtained with exact computation on the infinite dimensional spaces, starting from the same $(q_{\text{old}}, u_{\text{old}}) = (q_{\text{old}_{h_{k-1}}}, u_{\text{old}_{h_{k-1}}})$ as the one underlying $I_{i,h}^k$. Thus, in our analysis besides the actually computed sequence $(q_h^k, u_h^k) = (q_{h_k}^k, u_{h_k}^k)$ there appears an auxiliary sequence (q^k, u^k) , see Figure 1.

We assume the knowledge about bounds η^k_i on the error in the quantities of interest due to discretization

$$|I_{i,h}^{k} - I_{i}^{k}| \le \eta_{i}^{k}, \quad i \in \{1, 2, 3, 4\}$$
(31)

(, which can, at least partly, be computed by goal oriented error estimators, see e.g., [5, 6, 14, 21] and Section2.1) and to refine adaptively according to these bounds. On the other hand, we will now impose conditions on such upper bounds for the discretization error that enable to prove convergence and convergence rates results, see Assumption 7 below.

Additionally, we will make some assumptions on the forward operator



Figure 1: Sequence of discretized iterates and auxiliary sequence of continuous iterates for the all-at-once formulation of IRGNM

Assumption 4. Let the reduced forward operator \mathbf{F} be continuous and weakly sequentially closed, *i.e.*

$$(q_n \to q \land u_n \to u \land C(u_n) \to g \land A(q_n, u_n) \to f)$$

$$\Rightarrow (u \in \mathcal{D}(C) \land (q, u) \in \mathcal{D}(A) \land C(u) = g \land A(q, u) = f)$$

for all sequences $((q_n, u_n))_{n \in \mathbb{N}} \subseteq Q \times V$.

We also transfer the usual tangential cone condition to the all-at once setting from this section, which yields

Assumption 5. Let

 $\begin{aligned} \|C(u) - C(\bar{u}) - C'(u)(u - \bar{u})\|_{G} + \|A(q, u) - A(\bar{q}, \bar{u}) - A'_{q}(q, u)(q - \bar{q}) - A'_{u}(q, u)(u - \bar{u})\|_{W^{*}} \\ &\leq c_{tc} \left(\|C(u) - C(\bar{u})\|_{G} + \|A(q, u) - A(\bar{q}, \bar{u})\|_{W^{*}} \right) \end{aligned}$

hold for all $(q, u), (\bar{q}, \bar{u}) \in \mathcal{B}_{\rho}(q_0, u_0) \subset (Q \times V)$ and some $0 < c_{tc} < 1$.

The choice of the regularization parameter β_k will be done a posteriori according to an inexact Newton /discrepancy principle, which with the quantities introduced above reads as

$$\tilde{\underline{\theta}}I_{3,h}^k \le I_{2,h}^k \le \overline{\theta}I_{3,h}^k \,. \tag{32}$$

A discrepancy type principle will also be used for the choice of the overall stopping index

$$k_* = \min\{k \in \mathbb{N} : I_{3,h}^k \le \tau^2 \delta^2\}.$$
(33)

The parameters used there have to satisfy the following assumption.

Assumption 6. Let τ and $\underline{\tilde{\theta}}$ be chosen sufficiently large and $\underline{\tilde{\theta}}$ sufficiently small (see (32),(33)), such that

$$2\left(c_{tc}^{2} + \frac{(1+c_{tc})^{2}}{\tau^{2}}\right) < \tilde{\underline{\theta}} \quad and \quad \frac{2\overline{\theta} + 4c_{tc}^{2}}{1-4c_{tc}^{2}} < 1.$$

$$(34)$$

Therewith, we can also formulate our conditions on precision in the quantities of interest:

Assumption 7. Let for the discretization error with respect to the quantities of interest estimate (31) hold, where η_1^k , η_2^k , η_3^k , η_4^k are selected such that

$$\eta_1^k + 4c_{tc}^2 \eta_3^k \le \left(\frac{\tilde{\theta}}{\ell} - 2\left(2c_{tc}^2 + \frac{(1+2c_{tc})^2}{\tau^2}\right)\right) I_{3,h}^k \tag{35}$$

$$\eta_3^k \le c_1 I_{3,h}^k \quad and \quad \eta_2^k \to 0, \quad \eta_3^k \to 0, \quad \eta_4^k \to 0 \quad as \quad k \to \infty$$

$$(36)$$

$$I_{3,h}^{k} \le (1+c_3)I_{4,h}^{k-1} + r^k \quad and \quad (1+c_3)\frac{2\overline{\theta} + 4c_{tc}^2}{1 - 4c_{tc}^2} \le c_2 < 1 \tag{37}$$

for some constants $c_1, c_2, c_3 > 0$, and a sequence $r^k \to 0$ as $k \to \infty$ (where the second condition in (37) is possible due to the right inequality in (34)).

Exactly along the lines of the proofs of Theorems 1 and 2 in in [22], replacing F there by **F** according to (9), we therewith obtain convergence and convergence rates results:

Theorem 1. Let the Assumptions 1, 2, 3, 4 and 5 with c_{tc} sufficiently small be satisfied and let Assumption 6 hold. For the quantities of interest (27) and (29), let, further, the estimate (31) hold with η_i satisfying Assumption 7.

Then with β_k , $h = h_k$ fulfilling (32), k_* selected according to (33), and $(q_{h_k}^k, u_{h_k}^k)$ defined by (25) there holds

$$\|q_{h_k}^k - q_0\|_Q^2 + \|u_{h_k}^k - u_0\|_V^2 \le \|q^{\dagger} - q_0\|_Q^2 + \|u^{\dagger} - u_0\|_V^2 \qquad \forall 0 \le k \le k_* \,; \tag{38}$$

(ii) k_* is finite;

(iii) $(q_h^{k_*}, u_h^{k_*}) = (q_{h_{k_*}(\delta)}^{k_*}, u_{h_{k_*}(\delta)}^{k_*})$ converges (weakly) subsequentially to a solution of (8) as $\delta \to 0$ in the sense that it has a weakly convergent subsequence and each weakly convergent subsequence converges strongly to a solution of (8). If the solution $(q^{\dagger}, u^{\dagger})$ to (8) is unique, then $(q_h^{k_*}, u_h^{k_*})$ converges strongly to $(q^{\dagger}, u^{\dagger})$ as $\delta \to 0$.

For proving rates, as usual (cf. e.g. [4, 11, 18, 23]) source conditions are assumed

Assumption 8. Let

$$(q^{\dagger} - q_0, u^{\dagger} - u_0) \in \mathcal{R}\left(\kappa\left(\mathbf{F}'(q^{\dagger}, u^{\dagger})^*\mathbf{F}'(q^{\dagger}, u^{\dagger})
ight)\right)$$

(cf. Assumption 1) hold with some $\kappa \colon \mathbb{R}^+ \to \mathbb{R}^+$ such that κ^2 is strictly monotonically increasing on $(0, \|\mathbf{F}(q^{\dagger}, u^{\dagger})\|_{Q \times V}^2]$, ϕ defined by $\phi^{-1}(\lambda) = \kappa^2(\lambda)$ is convex and ψ defined by $\psi(\lambda) = \kappa(\lambda)\sqrt{\lambda}$ is strictly monotonically increasing on $(0, \|\mathbf{F}(q^{\dagger}, u^{\dagger})\|_{Q \times V}^2]$. Here, for some selfadjoint nonnegative operator A, the operator function $\kappa(A)$ is defined via functional calculus based on the spectral theorem (cf. e.g. [11]).

Theorem 2. Let the conditions of Theorem 1 and additionally the source condition Assumption 8 be fulfiled.

Then there exists a $\overline{\delta} > 0$ and a constant $\overline{C} > 0$ independent of δ such that for all $\delta \in (0, \overline{\delta}]$ the convergence rates

$$\|q_{\text{old}}^{k_*} - q^{\dagger}\|_Q^2 + \|u_{\text{old}}^{k_*} - u^{\dagger}\|_V^2 = \mathcal{O}\left(\frac{\delta^2}{\psi^{-1}(\bar{C}\delta)}\right).$$
(39)

are obtained.

Remark 3. We compare the source conditions for the reduced formulation

$$q^{\dagger} - q_0 \in \mathcal{R}\left(\kappa\left(F'(q^{\dagger})^*F'(q^{\dagger})\right)\right)$$
(40)

with Assumption 8 for the all-at-once formulation, e.g. in the case $\kappa(\lambda) = \sqrt{\lambda}$. Namely, in that case (40) reads: There exists $\overline{g} \in G$ such that

$$q^{\dagger} - q_0 = F'(q^{\dagger})^* \overline{g} = S'(q^{\dagger})^* C'(S(q^{\dagger}))^* \overline{g}.$$
(41)

On the other hand, Assumption 8 with the same κ reads: There exists $\tilde{\mathbf{g}} = \begin{pmatrix} \tilde{g} \\ \tilde{f} \end{pmatrix} \in G \times W^*$ such that

$$(q^{\dagger} - q_0, u^{\dagger} - u_0) = \mathbf{F}'(q^{\dagger}, u^{\dagger})^* \tilde{\mathbf{g}} = \begin{pmatrix} 0 & A'_q(q^{\dagger}, u^{\dagger})^* \\ C'(u^{\dagger})^* & A'_u(q^{\dagger}, u^{\dagger})^* \end{pmatrix} \tilde{\mathbf{g}},$$

which is equivalent to

$$\begin{array}{rcl} q^{\dagger} - q_{0} &=& A_{q}^{\prime}(q^{\dagger}, u^{\dagger})^{*}\tilde{f} \\ u^{\dagger} - u_{0} &=& C^{\prime}(u^{\dagger})^{*}\tilde{g} + A_{u}^{\prime}(q^{\dagger}, u^{\dagger})^{*}\tilde{f} \end{array}$$

and by elimination of \tilde{f} and use of the identities $u^{\dagger} = S(q^{\dagger})$ and $S'(q^{\dagger}) = -A'_u(q^{\dagger}, u^{\dagger})^{-1}A'_q(q^{\dagger}, u^{\dagger})$ we get

$$q^{\dagger} - q_0 = S'(q^{\dagger})^* \left(C'(S(q^{\dagger}))^* \tilde{g} + u_0 - u^{\dagger} \right),$$

which, setting $\overline{g} = \tilde{g} + C'(u^{\dagger})^{-*}(u_0 - u^{\dagger})$ becomes (41), provided $u_0 - u^{\dagger} \in \mathcal{R}(C'(u^{\dagger})^*)$.

2.1 Computation of the error estimators

Theoretically the error estimators for this subsection can be computed similarly to those from [22]. The fact that we consider an unconstrained optimization problem should make things easier, but we get another problem in return: For estimating I_1 and I_2 we would have to estimate terms like

$$||E(q, u)||_{W^*} - ||E(q_h, u_h)||_{W^*_u}$$

for some operator $E: Q \times V \to W^*$, which would be quite an effort to do via goal oriented error estimators. For this reason, the presented least squares formulation will not be implemented and we will not go into more detail concerning the error estimators for this section.

3 A Generalized Gauss-Newton formulation

A drawback of the unconstrained formulation (23) is the necessity of computing the W^* -norm of the (linearized) residual and especially of computing error estimators for this quantity of interest. Besides, a rescaling of the state equation (7) changes the solution of the optimization problem. Moreover, depending on the given inverse problem and its application, in some cases, it does not make sense to only minimize the residual of the linearized state equation, instead of setting it to zero.

A formulation that is much better tractable is obtained by defining $(q^k, u^k) = (q^{k,\delta}, u^{k,\delta})$ as a solution to the PDE constrained minimization problem

$$\min_{(q,u)\in Q\times V} \mathcal{T}_{\beta_k}(q,u) \coloneqq \|C(u^{k-1}) + C'(u^{k-1})(u - u^{k-1}) - g^{\delta}\|_G^2 + \frac{1}{\beta_k} \left(\|q - q_0\|_Q^2 + \|u - u_0\|_V^2\right)$$
(42)

s.t.
$$L_{k-1}(q-q^{k-1}) + K_{k-1}(u-u^{k-1}) + A(q^{k-1}, u^{k-1}) = f$$
 in W^* (43)

(see also [9], [10]) with the abbreviations (24).

We consider the Lagrangian L: $Q \times V \times W \to {\rm I\!R}$

$$\mathcal{L}(q, u, z) \coloneqq \mathcal{T}_{\beta_k}(q, u) + \langle f - A(q^{k-1}, u^{k-1}) - L_{k-1}(q - q^{k-1}) - K_{k-1}(u - u^{k-1}), z \rangle_{W^*, W}$$

and formulate the optimality conditions of first order for (42):

$$\mathbf{L}_{z}'(q,u,z)(\delta z) = \langle f - A(q^{k-1}, u^{k-1}) - L_{k-1}(q - q^{k-1}) - K_{k-1}(u - u^{k-1}), \delta z \rangle_{W^{*},W} = 0,$$
(44)

$$L'_{u}(q, u, z)(\delta u) = 2(C(u^{k-1}) + C'(u^{k-1})(u - u^{k-1}) - g^{\delta}, C'(u^{k-1})(\delta u))_{G} + \frac{2}{\beta_{k}}(u - u_{0}, \delta u)_{V} - \langle K_{k-1}\delta u, z \rangle_{W^{*}, W} = 0, \qquad (45)$$

$$\mathbf{L}_{q}'(q,u,z)(\delta q) = \frac{2}{\beta_{k}}(q-q_{0},\delta q)_{Q} - \langle L_{k-1}\delta q,z \rangle_{W^{*},W} = 0$$

$$\tag{46}$$

for all $\delta q \in Q$, $\delta u \in V$, $\delta z \in W$.

We assume boundedness of the operators $A(q, u), L_{k-1}, K_{k-1}^*, K_{k-1}^{-1}, C(u)$ and C'(u) in the following sense.

Assumption 9. There holds

 $\sup_{(q,u)\in\mathcal{B}_{\rho}(q_{0},u_{0})}\|A(q,u)\|_{W^{*}}+\|A_{q}'(q,u)\|_{Q\to W^{*}}+\|A_{u}'(q,u)^{*}\|_{W^{*}\to V}\|A_{u}'(q,u)^{-1}\|_{W^{*}\to V}<\infty$

and

$$\sup_{u\in\mathcal{B}_{\rho}(u_0)}\{\|C(u)\|_G+\|C'(u)\|_{V\to G}\}<\infty.$$

The following lemma about boundedness of the adjoint variable will serve as tool for unformly bounding the penalty parameter ρ .

Lemma 2. Under Assumption 9 and provided $(q^{k-1}, u^{k-1}) \in \mathcal{B}_{\rho}(q_0, u_0)$, for a stationary point $(q^k, u^k, z^k) \in Q \times V \times W$ of L (cf. (44) - (46)) there holds the estimate

$$||z^{k}||_{W} \le c_{\text{adj}} \left(||q^{k-1} - q_{0}||_{Q} + ||u^{k-1} - u_{0}||_{V} + 1 \right),$$
(47)

with a constant c_{adj} independent of k.

Proof. To formulate the optimality system (44)-(46) in a matrix-vector form, we introduce another dual variable $p \in W^*$ defined by

$$p = J_{W^*} z \in W^* \tag{48}$$

via the map J_{W^*} , which maps $z \in W$ to the Riesz representation $J_{W^*}z \in W^*$ of the linear functional $W^* \to \mathbb{R}, w^* \mapsto w^*(p)$, such that

$$\mathcal{L}(q,u,z) = \mathcal{T}_{\beta_k}(q,u) + (f - A(q^{k-1}, u^{k-1}) - L_{k-1}(q - q^{k-1}) - K_{k-1}(u - u^{k-1},), p)_{W^*}.$$

Using the abbreviations (24) and

$$C_{k-1} \coloneqq C'(u^{k-1}), \quad r_{k-1}^f \coloneqq A(q^{k-1}, u^{k-1}) - f, \quad r_{k-1}^g \coloneqq C(u^{k-1}) - g^\delta$$
(49)

the optimality system (44)-(46) can be written as

$$\begin{split} \mathbf{L}'_{z}(q,u,z)(\delta z) &= \langle f - A(q^{k-1},u^{k-1}) - L_{k-1}(q-q^{k-1}) - K_{k-1}(u-u^{k-1}), \delta z \rangle_{W^{*},W} = 0 \,, \\ \mathbf{L}'_{u}(q,u,z)(\delta u) &= 2(r^{g}_{k-1} + C_{k-1}(u-u^{k-1}), C_{k-1}\delta u)_{G} + \frac{2}{\beta_{k}}(u-u_{0},\delta u)_{V} - (K\delta u,p)_{W^{*}} \\ &= (2C^{*}_{k-1}[r^{g}_{k-1} + C_{k-1}(u-u^{k-1})] + \frac{2}{\beta_{k}}(u-u_{0}) - K^{*}_{k-1}p, \delta u)_{V} = 0 \,, \\ \mathbf{L}'_{q}(q,u,z)(\delta q) &= \frac{2}{\beta_{k}}(q-q_{0},\delta q)_{Q} - (L_{k-1}\delta q,p)_{W^{*}} = (\frac{2}{\beta_{k}}(q-q_{0}) - L^{*}_{k-1}p,\delta q)_{Q} = 0 \end{split}$$

for all $\delta q \in Q$, $\delta u \in V$ and $\delta z \in W$, or equivalently as

$$q^{k} = q_{0} + \frac{\beta_{k}}{2} L_{k-1}^{*} p^{k}$$
$$u^{k} = \left[\frac{2}{\beta_{k}} \operatorname{id} + 2C_{k-1}^{*} C_{k-1}\right]^{-1} \left(2C_{k-1}^{*} \left(C_{k-1}(u^{k-1}) - r_{k-1}^{g}\right) + \frac{2}{\beta_{k}}u_{0} + K_{k-1}^{*} p^{k}\right)$$
$$u^{k} = K_{k-1}^{-1} \left(L_{k-1}q^{k-1} + K_{k-1}u^{k-1} - r_{k-1}^{f} - L_{k-1}q^{k}\right).$$

Eliminating q^k and u^k this yields

0

$$\left[\frac{2}{\beta_k} \operatorname{id} + 2C_{k-1}^* C_{k-1} \right]^{-1} \left(2C_{k-1}^* \left(C_{k-1}(u^{k-1}) - r_{k-1}^g \right) + \frac{2}{\beta_k} u_0 + K_{k-1}^* p^k \right)$$

= $K_{k-1}^{-1} \left(L_{k-1} q^{k-1} + K_{k-1} u^{k-1} - r_{k-1}^f - L_{k-1} \left(q_0 + \frac{\beta_k}{2} L_{k-1}^* p^k \right) \right) ,$

which we reformulate as

$$-\frac{\beta_k}{2}K_{k-1}^{-1}L_{k-1}L_{k-1}^*p^k - \left[\frac{2}{\beta_k}\operatorname{id} + 2C_{k-1}^*C_{k-1}\right]^{-1}K_{k-1}^*p^k$$
$$= \left[\frac{2}{\beta_k}\operatorname{id} + 2C_{k-1}^*C_{k-1}\right]^{-1}\left(2C_{k-1}^*\left(C_{k-1}(u^{k-1}) - r_{k-1}^g\right) + \frac{2}{\beta_k}u_0\right)$$
$$-K_{k-1}^{-1}\left(L_{k-1}(q^{k-1} - q_0) + K_{k-1}u^{k-1} - r_{k-1}^f\right)$$

and finally

$$-\left[\frac{1}{\beta_{k}}\operatorname{id}+C_{k-1}^{*}C_{k-1}\right]\beta_{k}K_{k-1}^{-1}L_{k-1}L_{k-1}^{*}p^{k}-K_{k-1}^{*}p^{k}$$
$$=2C_{k-1}^{*}\left(C_{k-1}(u^{k-1})-r_{k-1}^{g}\right)+\frac{2}{\beta_{k}}u_{0}$$
$$-2\left[\frac{1}{\beta_{k}}\operatorname{id}+C_{k-1}^{*}C_{k-1}\right]K_{k-1}^{-1}\left(L_{k-1}(q^{k-1}-q_{0})+K_{k-1}u^{k-1}-r_{k-1}^{f}\right).$$

With

$$C_{\beta} \coloneqq \left(\frac{1}{\beta_k} \operatorname{id} + C_{k-1}^* C_{k-1}\right)^{1/2}$$

this is equivalent to

$$-\beta_k C_\beta^2 K_{k-1}^{-1} L_{k-1} L_{k-1}^* p^k - K_{k-1}^* p^k$$

= $2C_{k-1}^* \left(C_{k-1}(u^{k-1}) - r_{k-1}^g \right) + \frac{2}{\beta_k} u_0$
 $- 2C_\beta^2 K_{k-1}^{-1} \left(L_{k-1}(q^{k-1} - q_0) + K_{k-1}u^{k-1} - r_{k-1}^f \right) ,$

which upon premultiplication with C_{β}^{-1} becomes

$$\begin{split} &-\left(\beta_{k}C_{\beta}K_{k-1}^{-1}L_{k-1}(K_{k-1}^{-1}L_{k-1})^{*}C_{\beta}+\mathrm{id}\right)C_{\beta}^{-1}K_{k-1}^{*}p^{k} \\ &=-2C_{\beta}^{-1}C_{k-1}^{*}r_{k-1}^{g}+\frac{2}{\beta_{k}}C_{\beta}^{-1}(u_{0}-u^{k-1})+2C_{\beta}K_{k-1}^{-1}\left(L_{k-1}(q_{0}-q^{k-1})+r_{k-1}^{f}\right) \\ &+\left[2C_{\beta}^{-1}C_{k-1}^{*}C_{k-1}-2C_{\beta}+\frac{2}{\beta_{k}}C_{\beta}^{-1}\right]u^{k-1} \\ &=-2C_{\beta}^{-1}C_{k-1}^{*}r_{k-1}^{g}+\frac{2}{\beta_{k}}C_{\beta}^{-1}(u_{0}-u^{k-1})+2C_{\beta}K_{k-1}^{-1}\left(L_{k-1}(q_{0}-q^{k-1})+r_{k-1}^{f}\right) \\ &+\left[2C_{\beta}^{-1}\left(\frac{1}{\beta_{k}}\mathrm{id}+C_{k-1}^{*}C_{k-1}\right)-2C_{\beta}\right]u^{k-1} \\ &=-2C_{\beta}^{-1}C_{k-1}^{*}r_{k-1}^{g}+\frac{2}{\beta_{k}}C_{\beta}^{-1}(u_{0}-u^{k-1})+2C_{\beta}K_{k-1}^{-1}\left(L_{k-1}(q_{0}-q^{k-1})+r_{k-1}^{f}\right). \end{split}$$

Since $\beta_k C_\beta K_{k-1}^{-1} L_{k-1} (K_{k-1}^{-1} L_{k-1})^* C_\beta = \beta_k (C_\beta K_{k-1}^{-1} L_{k-1}) (C_\beta K_{k-1}^{-1} L_{k-1})^*$ is positive semidefinite, we can conclude

$$\left\| C_{\beta}^{-1} K_{k-1}^{*} p^{k} \right\|_{V} \\ \leq \left\| -2C_{\beta}^{-1} C_{k-1}^{*} r_{k-1}^{g} + \frac{2}{\beta_{k}} C_{\beta}^{-1} (u_{0} - u^{k-1}) + 2C_{\beta} K_{k-1}^{-1} \left(L_{k-1} (q_{0} - q^{k-1}) + r_{k-1}^{f} \right) \right\|_{V},$$

and with the estimates

$$\|C_{\beta}^{-1}C_{k-1}^{*}\|_{G \to V} \le 1, \quad \|C_{\beta}^{-1}\|_{V} \le \beta^{\frac{1}{2}}, \qquad \text{and} \quad \|C_{\beta}\|_{V} \le \left(\frac{1}{\beta_{k}} + \|C_{k-1}\|_{V \to G}^{2}\right)^{\frac{1}{2}}$$

we have

$$\begin{split} \left\| K_{k-1}^* p^k \right\|_V &\leq \left\| C_\beta C_\beta^{-1} K_{k-1}^* p^k \right\|_V \\ &\leq \left(\frac{1}{\beta_k} + \| C_{k-1} \|_{V \to G}^2 \right)^{\frac{1}{2}} \left\| C_\beta^{-1} K_{k-1}^* p^k \right\|_V \\ &\leq 2 \left(\frac{1}{\beta_k} + \| C_{k-1} \|_{V \to G}^2 \right)^{\frac{1}{2}} \left\{ \| r_{k-1}^g \| + \frac{1}{\sqrt{\beta_k}} \| u_0 - u^{k-1} \|_V \\ &+ \left(\frac{1}{\beta_k} + \| C_{k-1} \|_{V \to G}^2 \right)^{\frac{1}{2}} \| K_{k-1}^{-1} \left(L_{k-1} (q_0 - q^{k-1}) + r_{k-1}^f \| \right) \right\}, \end{split}$$

which by Assumption 9 and $(q^{k-1}, u^{k-1}) \in \mathcal{B}_{\rho}(q_0, u_0)$ yields (47).

We will prove inductively that the iterates indeed remain in $\mathcal{B}_{\rho}(q_0, u_0)$, see estimate (63) below. Thus, due to Lemma 2, which remains valid in the discretized setting (52), we get uniform boundedness of the dual variables by some sufficiently large ϱ , namely

$$\varrho \ge c_{\rm adj} \left(\|q^{\dagger} - q_0\|_Q + \|u^{\dagger} - u_0\|_V + 1 \right) \,. \tag{50}$$

Hence we can use exactness of the norm with exponent one as a penalty (cf., e.g., Theorem 5.11 in [13]), which implies that a solution (q^k, u^k) of (42), (43) coincides with the unique solution of the unconstrained minimization problem

$$\min_{(q,u)\in Q\times V} \varrho \|A'_{q}(q^{k-1}, u^{k-1})(q-q^{k-1}) + A'_{u}(q^{k-1}, u^{k-1})(u-u^{k-1}) + A(q^{k-1}, u^{k-1}) - f\|_{W^{*}} \\
+ \|C(u^{k-1}) + C'(u^{k-1})(u-u^{k-1}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta_{k}}(\|q-q_{0}\|_{Q}^{2} + \|u-u_{0}\|_{V}^{2}),$$
(51)

for ρ larger than the norm of the dual variable. The formulation (51) of (42), (43) will be used in the convergence proofs only. For a practical implementation we will directly discretize (42), (43).

The discrete version of (42), (43) reads

$$\min_{(q,u)\in Q_{h_k}\times V_{h_k}} \|C(u_{\text{old}}) + C'(u_{\text{old}})(u - u_{\text{old}}) - g^{\delta}\|_G^2
+ \frac{1}{\beta_k} \left(\|q - q_0\|_Q^2 + \|u - u_0\|_{V_{h_k}}^2 \right)$$
(52)

s.t.
$$L_{k-1}(q - q_{\text{old}}) + K_{k-1}(u - u_{\text{old}}) + A(q_{\text{old}}, u_{\text{old}}) = f$$
 in $W_{h_k}^*$, (53)

where $(q_{\text{old}}, u_{\text{old}}) = (q^{k-1}, u^{k-1}) = (q^{k-1}_{h_{k-1}}, u^{k-1}_{h_{k-1}})$ is the previous iterate and we assume again that the norms in G and W as well as A and C are evaluated exactly (cf. Assumption 3).

With ρ chosen sufficiently large such that (50) holds, we define the quantities of interest as follows

$$I_{1}: V \times Q \times V \times \mathbb{R} \to \mathbb{R}, \quad (u_{\text{old}}, q, u, \beta) \mapsto \left\| C'(u_{\text{old}})(u - u_{\text{old}}) + C(u_{\text{old}}) - g^{\delta} \right\|_{G}^{2} \\ + \frac{1}{\beta} \left(\left\| q - q_{0} \right\|_{Q}^{2} + \left\| u - u_{0} \right\|_{V}^{2} \right) \\ I_{2}: V \times V \to \mathbb{R}, \quad (u_{\text{old}}, u) \mapsto \left\| C'(u_{\text{old}})(u - u_{\text{old}}) + C(u_{\text{old}}) - g^{\delta} \right\|_{G}^{2} \\ I_{3}: Q \times V \to \mathbb{R}, \quad (q_{\text{old}}, u_{\text{old}}) \mapsto \left\| C(u_{\text{old}}) - g^{\delta} \right\|_{G}^{2} + \varrho \left\| A(q_{\text{old}}, u_{\text{old}}) - f \right\|_{W^{*}} \\ I_{4}: Q \times V \to \mathbb{R}, \quad (q, u) \mapsto \left\| C(u) - g^{\delta} \right\|_{G}^{2} + \varrho \left\| A(q, u) - f \right\|_{W^{*}}$$

$$(54)$$

(cf. (26)) and

$$I_{1}^{k} = I_{1}(u_{\text{old}}^{k}, q^{k}, u^{k}, \beta_{k})$$

$$I_{2}^{k} = I_{2}(u_{\text{old}}^{k}, u^{k})$$

$$I_{3}^{k} = I_{3}(q_{\text{old}}^{k}, u_{\text{old}}^{k})$$

$$I_{4}^{k} = I_{4}(q^{k}, u^{k}),$$
(55)

(cf. (27)), where q_{old}^k , u_{old}^k are fixed from the previous step and q^k , u^k are coupled by the linearized state equation (43) (or the third line of (??) respectively) for $q^{k-1} = q_{\text{old}}^k$ and $u^{k-1} = u_{\text{old}}^k$.

Consistently, the discrete counterparts to (54) and (55) are

$$I_{1,h}: V \times Q \times V \times \mathbb{R} \to \mathbb{R}, \quad (u_{\text{old}}, q, u, \beta) \mapsto \left\| C'(u_{\text{old}})(u - u_{\text{old}}) + C(u_{\text{old}}) - g^{\delta} \right\|_{G}^{2} \\ + \frac{1}{\beta} \left(\left\| q - q_{0} \right\|_{Q}^{2} + \left\| u - u_{0} \right\|_{V_{h_{k}}}^{2} \right) \\ I_{2,h}: V \times V \to \mathbb{R}, \qquad (u_{\text{old}}, u) \mapsto I_{2}(u_{\text{old}}, u) \\ I_{3,h}: Q \times V \to \mathbb{R}, \qquad (q_{\text{old}}, u_{\text{old}}) \mapsto \left\| C(u_{\text{old}}) - g^{\delta} \right\|_{G}^{2} + \varrho \left\| A(q_{\text{old}}, u_{\text{old}}) - f \right\|_{W_{h_{k}}^{*}} \\ I_{4,h}: Q \times V \to \mathbb{R}, \qquad (q, u) \mapsto \left\| C(u) - g^{\delta} \right\|_{G}^{2} + \varrho \left\| A(q, u) - f \right\|_{W_{h_{k}}^{*}}$$
(56)

and

$$I_{1,h}^{k} = I_{1,h}(u_{\text{old}}^{k}, q_{h_{k}}^{k}, u_{h_{k}}^{k}, \beta_{k})$$

$$I_{2,h}^{k} = I_{2,h}(u_{\text{old}}^{k}, u_{h_{k}}^{k})$$

$$I_{3,h}^{k} = I_{3,h}(q_{\text{old}}^{k}, u_{\text{old}}^{k})$$

$$I_{4,h}^{k} = I_{4,h}(q_{h_{k}}^{k}, u_{h_{k}}^{k})$$
(57)

(cf. (29)), where q_{old}^k , u_{old}^k are fixed from the previous step, since (like in (30)) we set $q_{\text{old}}^{k+1} = q_{h_k}^k$ and $u_{\text{old}}^{k+1} = u_{h_k}^k$ at the end of each iteration step.

Remark 4. Here, as compared to (27), we have removed the W^* -norms in the definition of I_1^k and I_2^k .

The W^* -norm still appears in I_3^k , but only in connection with the old iterative $(q_{\text{old}}^k, u_{\text{old}}^k)$, such that the only source of error in I_3^k is the evaluation of the W^* norm. That means that with respect to Section 2, we have replaced the problematic expression

$$||E(q, u)||_{W^*} - ||E(q_h, u_h)||_{W^*_h}$$

(cf. Section 2.1) by an expression of the form

$$\|E(q_{\text{old}}, u_{\text{old}})\|_{W^*} - \|E(q_{\text{old}}, u_{\text{old}})\|_{W^*_t} , \qquad (58)$$

For the very typical case $W = V = H_0^1(\Omega)$ (see Section 4), we can indeed estimate such an error using goal oriented error estimators:

Let $v \in V$, $v_h \in V_h$ solve the equations

$$(\nabla v, \nabla \varphi)_{L^{2}(\Omega)} = \langle E(q_{\text{old}}, u_{\text{old}}), \varphi \rangle_{V^{*}, V} \quad \forall \varphi \in V ,$$

$$(\nabla v_{h}, \nabla \varphi)_{L^{2}(\Omega)} = \langle E(q_{\text{old}}, u_{\text{old}}), \varphi \rangle_{V^{*}, V} \quad \forall \varphi \in V_{h} ,$$

where $(.,.)_{L^2(\Omega)}$ denotes the scalar product in $L^2(\Omega)$ and $\langle .,. \rangle_{V^*,V}$ denotes the duality pairing between V^* and V. Then there holds

$$\|E(q_{\text{old}}, u_{\text{old}})\|_{V^*} = \|\nabla v\|_{L^2(\Omega)}$$
 and $\|E(q_{\text{old}}, u_{\text{old}})\|_{V^*_h} = \|\nabla v_h\|_{L^2(\Omega)}$.

We define the functional

$$\Psi(v) \coloneqq \left\| \nabla v \right\|_{L^2(\Omega)}$$

and the Lagrangian

$$L(v,w) \coloneqq \Psi(v) + \langle E(q_{\text{old}}, u_{\text{old}}), w \rangle_{V^*, V} - (\nabla v, \nabla w)_{L^2(\Omega)}$$

Let (v, w) and (v_h, w_h) be continuous and discrete stationary points of L, i.e.

$$L'_{v}(v,w)(\varphi) = (\|\nabla v\|_{L^{2}(\Omega)})^{-1} (\nabla v, \nabla \varphi)_{L^{2}(\Omega)} - (\nabla \varphi, \nabla w)_{L^{2}(\Omega)} = 0 \qquad \forall \varphi \in V,$$
(59)
$$L'_{w}(v,w)(\varphi) = \langle E(q_{\text{old}}, u_{\text{old}}), \varphi \rangle_{V^{*},V} - (\nabla v, \nabla \varphi)_{L^{2}(\Omega)} = 0 \qquad \forall \varphi \in V,$$

$$L'_{v}(v_{h}, w_{h})(\varphi) = (\|\nabla v_{h}\|_{L^{2}(\Omega)})^{-1} (\nabla v_{h}, \nabla \varphi)_{L^{2}(\Omega)} - (\nabla \varphi, \nabla w_{h})_{L^{2}(\Omega)} = 0 \quad \forall \varphi \in V_{h},$$
(60)

$$L'_{w}(v_{h}, w_{h})(\varphi) = \langle E(q_{\text{old}}, u_{\text{old}}), \varphi \rangle_{V^{*}, V} - (\nabla v_{h}, \nabla \varphi)_{L^{2}(\Omega)} = 0 \qquad \forall \varphi \in V_{h}.$$

Then (by (59) and (60)) we have

$$w = (\|\nabla v\|_{L^{2}(\Omega)})^{-1}v \quad and \quad w_{h} = (\|\nabla v_{h}\|_{L^{2}(\Omega)})^{-1}v_{h}.$$
(61)

For the error (58) then holds

 $||E(q_{\text{old}}, u_{\text{old}})||_{V^*} - ||E(q_{\text{old}}, u_{\text{old}})||_{V_h^*}$

$$\begin{split} &= \Psi(v) - \Psi(v_h) \\ &= \frac{1}{2}L'(v_h, w_h)(v - \tilde{v_h}, w - \tilde{w_h}) + R \\ &= \frac{1}{2}(\|\nabla v_h\|_{L^2(\Omega)})^{-1}(\nabla v_h, \nabla (v - \tilde{v_h}))_{L^2(\Omega)} - \frac{1}{2}(\nabla (v - \tilde{v_h}), \nabla w_h)_{L^2(\Omega)} \\ &\quad + \frac{1}{2}\langle E(q_{\text{old}}, u_{\text{old}}), w - \tilde{w_h}\rangle_{V^*, V} - \frac{1}{2}(\nabla v_h, \nabla (w - \tilde{w_h}))_{L^2(\Omega)} \\ &= \frac{1}{2}(\|\nabla v_h\|_{L^2(\Omega)})^{-1}(\nabla v_h, \nabla (v - \tilde{v_h}))_{L^2(\Omega)} - \frac{1}{2}(\|\nabla v_h\|_{L^2(\Omega)})^{-1}(\nabla (v - \tilde{v_h}), \nabla v_h)_{L^2(\Omega)} \\ &\quad + \frac{1}{2}\langle E(q_{\text{old}}, u_{\text{old}}), w - \tilde{w_h}\rangle_{V^*, V} - \frac{1}{2}(\nabla v_h, \nabla (w - \tilde{w_h}))_{L^2(\Omega)} \\ &= \frac{1}{2}\langle E(q_{\text{old}}, u_{\text{old}}), w - \tilde{w_h}\rangle_{V^*, V} - \frac{1}{2}(\nabla v_h, \nabla (w - \tilde{w_h}))_{L^2(\Omega)} \end{split}$$

for arbitrary $\tilde{v}_h, \tilde{w}_h \in V_h$, where R is a third order remainder term (see e.g. [5, 6], Section 3.1). Please note that due to the relation (61) no additional system of equations has to be solved in order to obtain the additional variable w_h .

Another way to deal with the discretization error in I_3^k is the following: Tracking the upcoming convergence proof (cf. Theorem 3) the reader should realize that the discretization for $I_{3,h}^k$ does not have to be the same as for $I_{1,h}^k$, $I_{2,h}^k$, such that $I_{3,h}^k$ could be evaluated on a very fine separate mesh, such that η_3^k could be neglected. This alternative is of course, more costly, but since everything else is still done on the adaptively refined (coarser) mesh, the proposed method could still lead to an efficient algorithm.

The W^{*}-norm also appears in I_4^k , and unfortunately, in combination with the current qand u, which are subject to discretization, such that in principle we face the same situation as in the least squares formulation from Section 2 (cf. Subsection 2.1). Since, however, η_4^k only appears in connection with the very weak assumption $\eta_4^k \to 0$ as $k \to \infty$ (cf. (36)), as in [22], we save ourselves the computational effort of computing an error estimator for I_4^k .

Like in Section 2 we need the weak sequential closedness of \mathbf{F} , i.e. Assumption 4 and the following tangential cone condition (cf. Assumption 5).

Assumption 10. There exist $0 < c_{tc} < 1$ and $\rho > 0$ such that

$$\|C(u) - C(\bar{u}) - C'(u)(u - \bar{u})\|_{G} \leq c_{tc} \|C(u) - C(\bar{u})\|_{G} \|A(q, u) - A(\bar{q}, \bar{u}) - A'_{q}(q, u)(q - \bar{q}) - A'_{u}(q, u)(u - \bar{u})\|_{W^{*}} \leq 4c_{tc}^{2} \|A(q, u) - A(\bar{q}, \bar{u})\|_{W^{*}},$$

holds for all $(q, u), (\overline{q}, \overline{u}) \in \mathcal{B}_{\rho}(q_0, u_0) \subset Q \times V$ (cf. Assumption 1).

By means of Lemma 2 and the Assumptions 4, 10 we can now formulate a convergence result like in Theorems 1 and 3 in [22] and Theorem 1 here for (42). This can be done similarly to the proof of Theorem 3 in [22], replacing F there by **F** according to (9) and setting

$$S\left(\begin{pmatrix} y_C\\ y_A \end{pmatrix}, \begin{pmatrix} \tilde{y}_C\\ \tilde{y}_A \end{pmatrix}\right) = \|y_C - \tilde{y}_C\|_G^2 + \varrho \|y_A - \tilde{y}_A\|_{W^*},$$

$$\mathcal{R}\left(\begin{pmatrix} q\\ u \end{pmatrix}\right) = \|q - q_0\|_Q^2 + \|u - u_0\|_V^2,$$

$$c_S = 2,$$
(62)

there. For clarity of exposition we provide the full convergence proof (Theorem 3) without making use of the equivalence to (51) here. Only for the convergence rates result Theorem 4 we refer to Theorem 4 in [22] with (62) and the equivalence to (51). So in the proof of Theorem 3 we will not use minimality wrt (51) but only wrt the original formulation (42), (43) (actually we are using KKT points instead of minimizers, but this make no real difference due to convexity of the problem).

Theorem 3. Let the Assumptions 1, 2, 3, 4 and 10 with c_{tc} sufficiently small be satisfied and let Assumption 6 hold. For the quantities of interest (55) and (57), let, further, the estimate (31) hold with η_i satisfying Assumption 7.

Then with β_k , $h = h_k$ fulfilling (32), k_* selected according to (33), and $(q_{h_k}^k, u_{h_k}^k)$ defined as the primal part of a KKT point of (52), (53) there holds

$$\|q_{h_k}^k - q_0\|^2 + \|u_{h_k}^k - u_0\|^2 \le \|q^{\dagger} - q_0\|^2 + \|u^{\dagger} - u_0\|^2 \qquad \forall 0 \le k \le k_* \,; \tag{63}$$

(ii) k_* is finite;

(iii) $(q_h^{k_*}, u_h^{k_*}) = (q_{h_{k_*}(\delta)}^{k_*(\delta),\delta}, u_{h_{k_*}(\delta)}^{k_*(\delta),\delta})$ converges (weakly) subsequentially to a solution of (8) as $\delta \to 0$ in the sense that it has a weakly convergent subsequence and each weakly convergent subsequence converges strongly to a solution of (8). If the solution $(q^{\dagger}, u^{\dagger})$ to (8) is unique, then $(q_h^{k_*}, u_h^{k_*})$ converges strongly to $(q^{\dagger}, u^{\dagger})$ as $\delta \to 0$.

We mention in passing that this is a new result also in the continuous case $\eta_i^k = 0$.

Proof. (i): We will prove (63) by induction. The base case k = 0 is trivial. To carry out the induction step, we assume that

$$\|q_{h_{k-1}}^{k-1} - q_0\|_Q^2 + \|u_{h_{k-1}}^{k-1} - u_0\|_V^2 \le \|q^{\dagger} - q_0\|_Q^2 + \|u^{\dagger} - u_0\|_V^2 \qquad \forall 1 \le k \le k_*$$
(64)

holds. We consider a continuous step emerging from discrete $q_{\text{old}}^k = q_{h_{k-1}}^{k-1}$, $u_{\text{old}}^k = u_{h_{k-1}}^{k-1}$ (cf. Figure 1), i.e. let (q^k, u^k) be a solution to (42) for $q^{k-1} = q_{\text{old}}^k = q_{h_{k-1}}^{k-1}$. Then the KKT conditions $L'(q^k, u^k, z^k) = 0$ (cf. (44)-(46)) imply

$$0 = (C_{h,k-1}(u^{k} - u_{h_{k-1}}^{k-1}) + r_{k-1}^{g}, C_{h,k-1}\delta u)_{G} + \frac{1}{\beta_{k}} \left[(q^{k} - q_{0}, \delta q)_{Q} + (u^{k} - u_{0}, \delta u)_{V} \right] \\ + \frac{1}{2} (L_{h,k-1}\delta q + K_{h,k-1}\delta u, p^{k})_{W^{*}}$$

for all $\delta q \in Q$ and $\delta u \in V$, where we have used the same abbreviations as in (24) and (49), as well as p^k defined by (48).

Setting $\delta q = q^k - q^{\dagger}$, $\delta u = u^k - u^{\dagger}$, this yields

$$0 = \|C_{h,k-1}(u^{k} - u_{h_{k-1}}^{k-1}) + r_{k-1}^{g}\|_{G}^{2}$$

- $(C_{h,k-1}(u^{k} - u_{h_{k-1}}^{k-1}) + r_{k-1}^{g}, C_{h,k-1}(u^{\dagger} - u_{h_{k-1}}^{k-1}) + r_{k-1}^{g})_{G}$
+ $\frac{1}{\beta_{k}}\|q^{k} - q_{0}\|_{Q}^{2} - \frac{1}{\beta_{k}}(q^{k} - q_{0}, q^{\dagger} - q_{0})_{Q} + \frac{1}{\beta_{k}}\|u^{k} - u_{0}\|^{2} - \frac{1}{\beta_{k}}(u^{k} - u_{0}, u^{\dagger} - u_{0})_{V}$
- $\frac{1}{2}(L_{h,k-1}(q^{\dagger} - q_{h_{k-1}}^{k-1}) + K_{h,k-1}(u^{\dagger} - u_{h_{k-1}}^{k-1}) + r_{k-1}^{f}, p^{k})_{W^{*}},$

where we have used the fact that (q^k, u^k) satisfies the linearized state equation (43), i.e.

$$L_{h,k-1}(q^{k}-q_{h_{k-1}}^{k-1})K_{h,k-1}(u^{k}-u_{h_{k-1}}^{k-1})+r_{k-1}^{f}=0.$$

Hence by Cauchy-Schwarz and the fact that $ab \leq \frac{1}{2}a^2 + \frac{1}{2}b^2$ for all $a,b \in {\rm I\!R}$

$$\begin{split} I_{1}^{k} &\leq \|C_{h,k-1}(u^{k}-u_{h_{k-1}}^{k-1}) + r_{k-1}^{g}\|_{G}\|C_{h,k-1}(u^{\dagger}-u_{h_{k-1}}^{k-1}) + r_{k-1}^{g}\|_{G} \\ &+ \frac{1}{\beta_{k}}\|q^{k}-q_{0}\|_{Q}\|q^{\dagger}-q_{0}\|_{Q} + \frac{1}{\beta_{k}}\|u^{k}-u_{0}\|_{V}\|u^{\dagger}-u_{0}\|_{V} \\ &+ \frac{1}{2}\left\|L_{h,k-1}(q^{\dagger}-q_{h_{k-1}}^{k-1}) + K_{h,k-1}(u^{\dagger}-u_{h_{k-1}}^{k-1}) + r_{k-1}^{f}\right\|_{W^{*}}\|p^{k}\|_{W^{*}} \\ &\leq \frac{1}{2}\|C_{h,k-1}(u^{k}-u_{h_{k-1}}^{k-1}) + r_{k-1}^{g}\|_{G}^{2} + \frac{1}{2}\|C_{h,k-1}(u^{\dagger}-u_{h_{k-1}}^{k-1}) + r_{k-1}^{g}\|_{G}^{2} \\ &+ \frac{1}{2\beta_{k}}\|q^{k}-q_{0}\|_{Q}^{2} + \frac{1}{2\beta_{k}}\|q^{\dagger}-q_{0}\|_{Q}^{2} + \frac{1}{2\beta_{k}}\|u^{k}-u_{0}\|_{V}^{2} + \frac{1}{2\beta_{k}}\|u^{\dagger}-u_{0}\|_{V}^{2} \\ &+ \frac{1}{2}\left\|L_{h,k-1}(q^{\dagger}-q_{h_{k-1}}^{k-1}) + K_{h,k-1}(u^{\dagger}-u_{h_{k-1}}^{k-1}) + r_{k-1}^{f}\right\|_{W^{*}}\|p^{k}\|_{W^{*}} \\ &= \frac{1}{2}I_{1}^{k} + \frac{1}{2}\|C_{h,k-1}(u^{\dagger}-u_{h_{k-1}}^{k-1}) + r_{k-1}^{g}\|_{G}^{2} \\ &+ \frac{1}{2\beta_{k}}\|q^{\dagger}-q_{0}\|_{Q}^{2} + \frac{1}{2\beta_{k}}\|u^{\dagger}-u_{0}\|_{V}^{2} \\ &+ \frac{1}{2}\left\|L_{h,k-1}(q^{\dagger}-q_{h_{k-1}}^{k-1}) + K_{h,k-1}(u^{\dagger}-u_{h_{k-1}}^{k-1}) + r_{k-1}^{f}\right\|_{W^{*}}\|p^{k}\|_{W^{*}} , \end{split}$$

(i)

which dividing by 2 and applying Lemma 2 with (50), and (64) leads to

$$\begin{split} I_{1}^{k} &\leq \|C_{h,k-1}(u^{\dagger} - u_{h_{k-1}}^{k-1}) + r_{k-1}^{g}\|_{G}^{2} + \frac{1}{\beta_{k}}\|q^{\dagger} - q_{0}\|_{Q}^{2} + \frac{1}{\beta_{k}}\|u^{\dagger} - u_{0}\|_{V}^{2} \\ &+ \left\|L_{h,k-1}(q^{\dagger} - q_{h_{k-1}}^{k-1}) + K_{h,k-1}(u^{\dagger} - u_{h_{k-1}}^{k-1}) + r_{k-1}^{f}\right\|_{W^{*}}\|p^{k}\|_{W^{*}} \\ &\leq \|C'(u_{h_{k-1}}^{k-1})(u^{\dagger} - u_{h_{k-1}}^{k-1}) + C(u_{h_{k-1}}^{k-1}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta_{k}}\|q^{\dagger} - q_{0}\|_{Q}^{2} + \frac{1}{\beta_{k}}\|u^{\dagger} - u_{0}\|_{V}^{2} \\ &+ \varrho \left\|A'_{q}(q_{h_{k-1}}^{k-1})(q^{\dagger} - q_{h_{k-1}}^{k-1}) + A'_{u}(u_{h_{k-1}}^{k-1})(u^{\dagger} - u_{h_{k-1}}^{k-1}) + A(q_{h_{k-1}}^{k-1}, u_{h_{k-1}}^{k-1}) - A(q^{\dagger}, u^{\dagger})\right\|_{W^{*}} \end{split}$$
(65)

for all $k < k_*$.

The rest of the proof basically follows the lines of the proof of Theorem 3 in [22] with the choice (62), but for convenience of the reader we will follow through the proof anyway. Using the fact that $(a + b)^2 \leq 2a^2 + 2b^2$ for all $a, b \in \mathbb{R}$ and Assumption 10 from (65) we get

$$\begin{split} I_{1}^{k} &\leq 2c_{tc}^{2} \|C(u_{h_{k-1}}^{k-1}) - C(u^{\dagger})\|_{G}^{2} + 2\delta^{2} + \frac{1}{\beta_{k}} \|q^{\dagger} - q_{0}\|_{Q}^{2} + \frac{1}{\beta_{k}} \|u^{\dagger} - u_{0}\|_{V}^{2} \\ &+ 4c_{tc}^{2} \varrho \left\|A(q_{h_{k-1}}^{k-1}, u_{h_{k-1}}^{k-1}) - A(q^{\dagger}, u^{\dagger})\right\|_{W^{*}} \\ &\leq 4c_{tc}^{2} \|C(u_{h_{k-1}}^{k-1}) - g^{\delta}\|_{G}^{2} + 2(1 + 2c_{tc}^{2})\delta^{2} + \frac{1}{\beta_{k}} \|q^{\dagger} - q_{0}\|_{Q}^{2} + \frac{1}{\beta_{k}} \|u^{\dagger} - u_{0}\|_{V}^{2} \\ &+ 4c_{tc}^{2} \varrho \left\|A(q_{h_{k-1}}^{k-1}, u_{h_{k-1}}^{k-1}) - f\right\|_{W^{*}} \\ &\leq 4c_{tc}^{2} I_{3}^{k} + \frac{2(1 + 2c_{tc}^{2})}{\tau^{2}} I_{3,h}^{k} + \frac{1}{\beta_{k}} \|q^{\dagger} - q_{0}\|_{Q}^{2} + \frac{1}{\beta_{k}} \|u^{\dagger} - u_{0}\|_{V}^{2} \\ &\leq 2 \left(2c_{tc}^{2} + \frac{1 + 2c_{tc}^{2}}{\tau^{2}}\right) I_{3,h}^{k} + 4c_{tc}^{2} \eta_{3}^{k} + \frac{1}{\beta_{k}} \left(\|q^{\dagger} - q_{0}\|_{Q}^{2} + \|u^{\dagger} - u_{0}\|_{V}^{2}\right) \end{split}$$

for all $k < k_*$. This together with (32) and the fact that $I_{1,h}^k = I_{2,h}^k + \frac{1}{\beta_k} \left(\|q_h^k - q_0\|_Q^2 + \|u_h^k - u_0\|_V^2 \right)$ yields

$$\begin{split} &\tilde{\underline{\theta}}I_{3,h}^{k} + \frac{1}{\beta_{k}}(\|q_{h_{k}}^{k} - q_{0}\|^{2} + \|u_{h_{k}}^{k} - u_{0}\|^{2}) \\ &\leq I_{2,h}^{k} + \frac{1}{\beta_{k}}(\|q_{h_{k}}^{k} - q_{0}\|^{2} + \|u_{h_{k}}^{k} - u_{0}\|^{2}) \\ &\leq I_{1}^{k} + \eta_{1}^{k} \\ &\leq 2\left(2c_{tc}^{2} + \frac{(1 + 2c_{tc})^{2}}{\tau^{2}}\right)I_{3,h}^{k} + \frac{1}{\beta_{k}}(\|q^{\dagger} - q_{0}\|^{2} + \|u^{\dagger} - u_{0}\|^{2}) + \eta_{1}^{k} + 4c_{tc}^{2}\eta_{3}^{k} \,. \end{split}$$

hence by (35) we get (63).

(ii): By the triangle inequality as well as (32), Assumption 10 and the fact that (q^k, u^k) satisfies the linearized state equation (43), we have

$$\begin{split} I_4^k &= \|C(u^k) - g^{\delta}\|_G^2 + \varrho \|A(q^k, u^k) - f\|_{W^*} \\ &\leq 2\|C'(u_{\text{old}}^k)(u^k - u_{\text{old}}^k) + C(u_{\text{old}}^k) - g^{\delta}\|_G^2 + 2\|C'(u_{\text{old}}^k)(u^k - u_{\text{old}}^k) + C(u_{\text{old}}^k) - C(u^k)\|_G^2 \\ &+ \varrho \|A'_q(q_{\text{old}}^k, u_{\text{old}}^k)(q^k - q_{\text{old}}^k) + A'_u(q_{\text{old}}^k, u_{\text{old}}^k)(u^k - u_{\text{old}}^k) + A(q_{\text{old}}^k, u_{\text{old}}^k) - f\|_{W^*} \\ &+ \varrho \|A'_q(q_{\text{old}}^k, u_{\text{old}}^k)(q^k - q_{\text{old}}^k) + A'_u(q_{\text{old}}^k, u_{\text{old}}^k)(u^k - u_{\text{old}}^k) + A(q_{\text{old}}^k, u_{\text{old}}^k) - A(q^k, u^k)\|_{W^*} \\ &\leq 2I_2^k + 2c_{tc}^2 \|C(u^k) - C(u_{\text{old}}^k)\|_G^2 + 4c_{tc}^2 \varrho \|A(q_{\text{old}}^k, u_{\text{old}}^k) - A(q^k, u^k)\|_{W^*} \\ &\leq 2I_2^k + 4c_{tc}^2 \left(\|C(u^k) - g^{\delta}\|_G^2 + \|C(u_{\text{old}}^k) - g^{\delta}\|_G^2\right) \\ &+ 4c_{tc}^2 \varrho \left(\|A(q_{\text{old}}^k, u_{\text{old}}^k) - f\|_{W^*} + \|A(q^k, u^k) - f\|_{W^*}\right) \\ &\leq 2 \left(\tilde{\bar{\theta}}I_{3,h}^k + \eta_2^k\right) + 4c_{tc}^2 (I_4^k + I_3^k), \end{split}$$

which implies

$$I_4^k \le \frac{1}{1 - 4c_{tc}^2} \left(2\tilde{\bar{\theta}} I_{3,h}^k + 2\eta_2^k + 4c_{tc}^2 I_3^k \right)$$

From this, using (31) and (37) we can deduce exactly as in the proof of Theorem 3 (ii) in [22] that

$$I_{4,h}^{k} \le c_{2}^{k} I_{4,h}^{0} + \sum_{j=0}^{k-1} c_{2}^{j} a^{k-j} .$$
(66)

with

$$a^{i} \coloneqq \frac{1}{1 - 4c_{tc}^{2}} \left((2\tilde{\bar{\theta}} + 4c_{tc}^{2})r^{i} + 2\eta_{2}^{i} + 4c_{tc}^{2}\eta_{3}^{i} \right) + \eta_{4}^{i} \quad \forall i \in \{1, 2, \dots, k\}.$$
(67)

So since the right hand side of (66) tends to zero as $k \to \infty$, $I_{4,h}^k$ and therewith $I_{3,h}^k$ (cf. (37)) eventually has to fall below $\tau^2 \delta^2$ for some finite index k.

(iii): With (5), (31), (36) and the definition of k_* , we have

$$\|C(u_{\text{old}}^{k_*}) - g\|_G^2 + \varrho \|A(q_{\text{old}}^{k_*}, u_{\text{old}}^{k_*}) - f\|_{W^*} \le 2I_3^{k_*} + 2\delta^2$$

$$\le 2\left(I_{3,h}^{k_*} + \eta_3^k + \delta^2\right)$$

$$\le 2\left((1 + c_1)I_{3,h}^{k_*} + \delta^2\right)$$

$$\le 2\delta^2\left((1 + c_1)\tau^2 + 1\right) \to 0$$
(68)

as $\delta \to 0$. Thus, due to (ii) (63) $(q_{\text{old}}^{k_*}, u_{\text{old}}^{k_*}) = (q_{h_{k_*-1}}^{\delta,k_*-1}, u_{h_{k_*-1}}^{\delta,k_*-1})$ has a weakly convergent subsequence $\left((q_{\text{old}}^{k_*(\delta_l)}, u_{\text{old}}^{k_*(\delta_l)})\right)_{l \in \mathbb{N}}$ and with Assumption 4 and (68) the limit of every weakly convergent subsequence is a solution to (8). Strong convergence of any weakly convergent subsequence again follows by a standard argument like in [22] using (63).

Corollary 1. The sequence $(z^k)_{k \in \mathbb{N}, k \leq k_*}$ is bounded, i.e.

$$\bar{\varrho} = \sup_{k \le k_*} \|z^k\|_W \le c_{\text{adj}}(\|q^{\dagger} - q_0\|^2 + \|u^{\dagger} - u_0\|^2 + 1)$$

Proof. The assertion follows directly from Theorem 3 (i) and Lemma 2.

The convergences rates from Theorem 4 in [22] also hold for the all-at-once formulation (8), due to equivalence with (51) which we formulate in the following theorem.

Instead of source conditions we use variational source conditions (cf., e.g., [12, 19, 20, 26]) due to the nonquadratic penalty term in (51).

Assumption 11. Let

$$\begin{aligned} |(q^{\dagger} - q_{0}, q - q^{\dagger})_{Q} + (u^{\dagger} - u_{0}, u - u^{\dagger})_{V}| \\ &\leq c \sqrt{\|q - q^{\dagger}\|_{Q}^{2} + \|u - u^{\dagger}\|_{V}^{2}} \kappa \left(\frac{\|C(u) - C(u^{\dagger})\|_{G}^{2} + \varrho\|A(q, u) - A(q^{\dagger}, u^{\dagger})\|_{W^{*}}}{\|q - q^{\dagger}\|_{Q}^{2} + \|u - u^{\dagger}\|_{V}^{2}} \right) (69) \\ &\qquad (q, u) \in \mathcal{D}(A), \ u \in \mathcal{D}(C), \end{aligned}$$

with ϱ sufficiently large (cf. (50)) and independent from q, u, hold with some $\kappa \colon \mathbb{R}^+ \to \mathbb{R}^+$ such that κ^2 is strictly monotonically increasing on $(0, \|\mathbf{F}(q^{\dagger}, u^{\dagger})\|_{Q \times V}^2]$, ϕ defined by $\phi^{-1}(\lambda) = \kappa^2(\lambda)$ is convex and ψ defined by $\psi(\lambda) = \kappa(\lambda)\sqrt{\lambda}$ is strictly monotonically increasing on $(0, \|\mathbf{F}(q^{\dagger}, u^{\dagger})\|_{Q \times V}^2]$.

Theorem 4. Let the conditions of Theorem 3 and additionally the variational inequality Assumption 11 be fulfilled.

Then there exists a $\overline{\delta} > 0$ and a constant $\overline{C} > 0$ independent of δ such that for all $\delta \in (0, \overline{\delta}]$ the convergence rates

$$\|q_{\text{old}}^{k_*} - q^{\dagger}\|_Q^2 + \|u_{\text{old}}^{k_*} - u^{\dagger}\|_V^2 = \mathcal{O}\left(\frac{\delta^2}{\psi^{-1}(\bar{C}\delta)}\right),$$
(71)

with $q_{\text{old}}^{k_*} = q_{\beta_{k_*-1},h_{k_*-1}}^{\delta,k_*-1}$, $u_{\text{old}}^{k_*} = u_{\beta_{k_*-1},h_{k_*-1}}^{\delta,k_*-1}$ are obtained.

Proof. With (62) the rate follows directly from Theorem 4 in [22] due to Theorem 3 (especially (63)) and (68). \Box

Remark 5. In fact, no regularization of the u part would be needed for proving just stability of the single Gauss-Newton steps, since by Assumption 2 the terms $\varrho \|A'_q(q^{k-1}, u^{k-1})(q-q^{k-1}) + A'_u(q^{k-1}, u^{k-1})(u-u^{k-1}) + A(q^{k-1}, u^{k-1}) - f\|_{W^*}$ and $\frac{1}{\beta_k} \|q-q_0\|_Q^2$ in (51) as regularization term together ensure weak compactness of the level sets of the Tikhonov functional (cf. Item 6 in Assumption 2 in [22]). However, we require even uniform boundedness of $u_{h_k}^k$ in order to uniformly bound the dual variable and come up with a penalty parameter ϱ that is independent of k, cf. the discrete version of Lemma 2. Using the equality constraint (43) (for $q^{k-1} = q_{h_{k-1}}^{k-1}$) together with the tangential cone condition Assumption 10 for $c_{tc}^2 < \frac{1}{8}$ would only enable to bound $A(q^k, u^k) - f$:

$$\begin{split} \|A(q^{k}, u^{k}) - f\|_{W^{*}} \\ &= \|L_{h,k-1}(q^{k} - q_{h_{k-1}}^{k-1}) + K_{h,k-1}(u^{k} - u_{h_{k-1}}^{k-1}) + A(q_{h_{k-1}}^{k-1}, u_{h_{k-1}}^{k-1}) - A(q^{k}, u^{k})\|_{W^{*}} \\ &\leq 4c_{tc}^{2} \|A(q_{h_{k-1}}^{k-1}, u_{h_{k-1}}^{k-1}) - A(q^{k}, u^{k})\|_{W^{*}} \\ &\leq 4c_{tc}^{2} \|A(q_{h_{k-1}}^{k-1}, u_{h_{k-1}}^{k-1}) - f\|_{W^{*}} + 4c_{tc}^{2} \|A(q^{k}, u^{k}) - f\|_{W^{*}} \,, \end{split}$$

such that

$$\|A(q^k, u^k) - f\|_{W^*} \le \frac{4c_{tc}^2}{1 - 4c_{tc}^2} \|A(q_{h_{k-1}}^{k-1}, u_{h_{k-1}}^{k-1}) - f\|_{W^*}.$$

However, without error estimators on the difference between $||A(q^k, u^k) - f||_{W^*}$ and its discretized version, this does not give a recursion

$$||A(q_{h_k}^k, u_{h_k}^k) - f||_{W^*} \le c ||A(q_{h_{k-1}}^{k-1}, u_{h_{k-1}}^{k-1}) - f||_{W^*}$$

(from which, by uniform boundedness of $q_{h_k}^k$ and Assumption 9 we could conclude uniform boundedness of $u_{h_k}^k$).

Thus, in order to obtain uniform boundedness of $u_{h_k}^k$ we introduce the term $\frac{1}{\beta_k} ||u - u_0||_V^2$ here for theoretical purposes. For our practical computations we will assume that the error by discretization between $||A(q^k, u^k) - f||_{W^*}$ and $||A(q_{h_k}^k, u_{h_k}^k) - f||_{W^*}$ is small enough so that the mentioned gap in this argument for uniform boundedness of $u_{h_k}^k$ can be neglected and the part $\frac{1}{\beta_k} ||u - u_0||_V^2$ of the regularization term is omitted.

3.1 Computation of the error estimators

Since – different to $[22] - u_{old}$ ist not subject to new discretization in the *k*th step here, the computation of the error estimators is easier and can be done exactly as in [14] and [21]. Thus we omit the arguments q_{old} and u_{old} in the quantities of interest in this subsection and we also omit the iteration index k and the explicit dependence on β .

Error estimator for I_1 : We consider

$$I_1(q, u) = \left\| C'(u_{\text{old}})(u - u_{\text{old}}) + C(u_{\text{old}}) - g^{\delta} \right\|_G^2 + \alpha (\|q - q_0\|_G^2 + \|u - u_0\|_V^2)$$

and the Lagrange functional

$$\mathcal{L}(q, u, z) \coloneqq I_1(q, u) + h(z) - B(q, u)(z), \qquad (72)$$

with $h \in W^*$ and $B(q, u) \in W^*$ defined as

 $h \coloneqq f - A(q_{\text{old}}, u_{\text{old}}) - A'_q(q_{\text{old}}, u_{\text{old}})(q_{\text{old}}) - A'_u(q_{\text{old}}, u_{\text{old}})(u_{\text{old}})$

and

$$B(q, u) \coloneqq A'_q(q_{\text{old}}, u_{\text{old}})(q) + A'_u(q_{\text{old}}, u_{\text{old}})(u) \,.$$

There holds a similar result to Proposition 1 in [22](see also [14]), which allows to estimate the difference $I_1(q, u) - I_1(q_h, u_h)$ by computing a discrete stationary point $x_h = (q_h, u_h, z_h) \in$

 $X_h = Q_h \times V_h \times W_h$ of L. This is done by solving the equations

$$z_h \in W_h: \qquad A'_u(q_{\text{old}}, u_{\text{old}})(du)(z_h) = I'_{1,u}(q_h, u_h)(du) \qquad \qquad \forall du \in V_h \tag{73}$$

$$u_h \in V_h: \qquad A'_q(q_{\text{old}}, u_{\text{old}})(q_h)(dz) + A'_u(q_{\text{old}}, u_{\text{old}})(u_h)(dz) = h(dz) \qquad \forall dz \in W_h \tag{74}$$

$$q_h \in Q_h: \qquad I'_q(q_h, u_h)(dq) = A'_q(q_{\text{old}}, u_{\text{old}})(dq)(z_h) \qquad \qquad \forall dq \in Q_h \qquad (75)$$

Then the error estimator η_1 for I_1 can be computed as

$$I_1 - I_{1,h} = I_1(q, u) - I_2(q_h, u_h) \approx \frac{1}{2} \mathbf{L}'(x_h)(\pi_h x_h - x_h) = \eta_1$$
(76)

(cf. [22, 21, 14]).

Remark 6. Please note that the equations (73)-(75) are solved anyway in the process of solving the optimization problem (42), (43).

Error estimator for I_2 : The computation of the error estimator for I_2 can be done similarly to the computation of η_2 in [22] (or η^I in [14]) by means of the Lagrange functional L. We consider

$$I_2(u) \coloneqq \left\| C'(u_{\text{old}})(u - u_{\text{old}}) + C(u_{\text{old}}) - g^{\delta} \right\|_G^2$$

and compute a discrete stationary point $y_h := (x_h, x_h^1) \in X_h \times X_h$ of the auxiliary Lagrange functional

$$\mathcal{M}(y) \coloneqq I_2(u) + \mathcal{L}'(x)(x_1)$$

by solving the equations

$$\begin{aligned}
x_h \in X_h: & \mathcal{L}'(x_h)(dx_1) = 0 & \forall dx_1 \in X_h \\
x_h^1 \in X_h: & \mathcal{L}''(x_h)(x_h^1, dx) = -I'_2(u_h)(du) & \forall dx \in X_h & (77)
\end{aligned}$$

(78)

(with dx = (dq, du, dz)). Then we compute the error estimator for I_2 by

$$\eta_2 \coloneqq \frac{1}{2} \mathcal{M}'(y_h)(\pi_h y_h - y_h) \approx I_2(u) - I_2(u_h) = I_2 - I_{2,h}$$

Remark 7. To avoid the computation of second order information in (77) we would like to refer to [6], where (77) is replaced by an approximate equation of first order.

Error estimator for I_3 : In Remark 4, we already mentioned that the W^* norm in I_3 can be evaluated on a separate very fine mesh, so that we will neglect the difference between $||A(q_{\text{old}}, u_{\text{old}}) - f||_{W^*}$ and $||A(q_{\text{old}}, u_{\text{old}}) - f||_{W^*_h}$. This implies that we do not need to compute the error estimator η_3 , since $I_3 = I_{3,h}$, so that (37), and the first part of (36) is trivially fullfilled.

Error estimator for I_4 : We also mentioned in Remark 4 that we will not compute η_4 , as the error $|I_4 - I_{4,h}|$ needs to be controlled only through the very weak assumption $\eta_4^k \to 0$ as $k \to \infty$ (cf. (36)), which in practice we will simply make sure by altogether decreasing the mesh size in the course of the iteration.

3.1.1 Algorithm

Since we only know about the existence of an upper bound $\overline{\varrho}$ of $||z^k||_W$ (cf. Corollary 1), but not its value, we choose ϱ (cf. (50)) heuristically, i.e. in each iteration step we set $\varrho = \varrho_k = \max\{\varrho_{k-1}, ||z_{h_k}^k||_{W_{h_k}}\}$ for the discrete counterpart $z_{h_k}^k$ of z^k .

Remark 8. Theoretically one should use $\rho = \|z_{h_k}^k\|_{W_H}$ on a very fine discretization H in order to get a better approximation to $\|z^k\|_W$. However, since we only need the correct order of magnitude and not the exact value, we just use the current mesh h_k .

In view of Remark 5 we omit the part $\frac{1}{\beta_k} ||u - u_0||_V^2$ of the regularization term. Also, as motivated in Section 3.1, we assume $\eta_3^k = 0$ for all k, such that we neither compute η_3 nor η_4 .

Thus we only check for the condition

$$\eta_1^k \le \left(\frac{\tilde{\theta}}{2} - 2\left(2c_{tc}^2 + \frac{(1+2c_{tc})^2}{\tau^2}\right)\right) I_{3,h}^k \tag{79}$$

on η_1^k in Assumption 7.

For simplicity, we evaluate $I_{3,h}^k$ on the current mesh instead of a very fine mesh as explained in Remark 4.

For computing β_k , $h = h_k$ fulfilling (32), we can resort to the Algorithm from [14], which also contains refinement with respect to the quantity of interest $I_{2,h}^k$ and repeated solution of

$$\min_{(q,v)\in Q_h\times V_h} \|C'(u_{\text{old},h}^k)(v) + C(u_{\text{old},h}^k) - g^{\delta}\|_G^2 + \frac{1}{\beta_k} \|q - q_0\|_Q^2$$
(80)

 $A'_{u}(q^{k}_{\mathrm{old},h}, u^{k}_{\mathrm{old},h})(v)(\varphi) + A'_{a}(q^{k}_{\mathrm{old},h}, u^{k}_{\mathrm{old},h})(q - q^{k}_{\mathrm{old},h})(\varphi) + A(q^{k}_{\mathrm{old},h}, u^{k}_{\mathrm{old},h})(\varphi) - f(\varphi) = 0$ $\forall \varphi \in W_h$ s.t. for $(q, v) \in Q_h \times V_h$.

The presented Generalized Gauss-Newton formulation can be implemented according to the following Algorithm 1.

Algorithm 1. Generalized Gauss-Newton Method

- 1: Choose τ , τ_{β} , $\tilde{\tau}_{\beta}$, $\underline{\tilde{\theta}}$, $\overline{\tilde{\theta}}$ such that $0 < \underline{\tilde{\theta}} \leq \overline{\tilde{\theta}} < 1$ and Assumption 6 holds. $\overline{\theta} = (\underline{\tilde{\theta}} + \overline{\tilde{\theta}})/2$ and $\max\{1, \tilde{\tau}_{\beta}\} < \tau_{\beta} \leq \tau$, and choose c_1 , c_2 and c_3 , such that the second part of (37) is fulfilled.
- 2: Choose a discretization $h = h_0$ and starting value $q_h^0 (= q_{h_0}^0)$ (not necessarily coinciding with q_0 in the regularization term) and set $q_{\text{old}}^0 = q_h^0$. 3: Choose starting value $u_h^0 (= u_{h_0}^0)$ (e.g. by solving the PDE $A(q_{\text{old}}^0, u_{\text{old}}^0) = f$) and set
- $u_{\text{old}}^0 = u_h^0.$
- 4: Compute the adjoint state $z_h^0(=z_{h_0}^0)$ (see (45)), evaluate $\left\|z_h^0\right\|_{W_1}$, set $\varrho_0 = \|z_h^0\|_{W_h}$. and $\begin{array}{ll} \mbox{evaluate } I^0_{3,h} \ (cf. \ (29)). \\ 5: \ Set \ k = 0 \ and \ h = h^1_0 = h_0. \\ 6: \ \mbox{while } I^k_{3,h} > \tau^2 \delta^2 \ \ do \\ 7: \ \ Set \ h = h^1_k. \end{array}$

Solve the optimization problem (80) Set $h_{\perp}^2 = h_{\perp}^1$ and $\delta_{\perp}^2 = \tilde{\theta} I_{\perp h}^k$. 8:

9: Set
$$h_k^2 = h_k^1$$
 and $\delta_\beta^2 = \theta I_{3,h}^k$

10: if
$$I_{2,h}^k > \left(au_{eta}^2 + rac{ au_{eta}^2}{2}
ight) \delta_{eta}^2$$
 then

- With q_{old}^k , u_{old}^k fixed, apply the Algorithm from [14] (with quantity of interest I_2^k 11: and noise level $\delta_{\beta}^2 = \tilde{\theta} I_{3,h}^k$ starting with the current mesh $h(=h_k^1)$ to obtain a regularization parameter β_k and a possibly different discretization h_k^2 such that (32) holds; Therwith, also the corresponding $v_h^k = v_{h_k^2}^k$, $q_h^k = q_{h_k^2}^k$ according to (80) are computed.
- Set $h = h_k^2$. 12:
- Evaluate the error estimator η_1^k (cf. (27),(29)). 13:
- Set $h_k^3 = h_k^2$. 14:
- while (79) is violated do 15:
- Refine grid with respect to η_1^k such that we obtain a finer discretization h_k^3 . 16:
- Solve the the optimization problem (80) and evaluate η_1^k . 17:
- Set $h = h_k^3$. 18:
- 19:
- Set $h = n_k$. Set $q_{\text{old}}^{k+1} = q_h^k$, $u_{\text{old}}^{k+1} = u_{\text{old}}^k + v_h^k$. Compute the adjoint state $z_h^{k+1} (= z_{h_k}^{k+1})$ (see (45)), evaluate $||z_h^{k+1}||_{W_h}$, set $\varrho_k = \max\{\varrho_{k-1}, ||z_h^{k+1}||_{W_h}\}$. and evaluate $I_{3,h}^{k+1}$. Set $h_{k+1}^1 = h_k^3$ (i.e. use the current mesh as a starting mesh for the next iteration). 20:
- 21:
- Set k = k + 1. 22:

Remark 9. In practice, we replace the "while"-loop on lines 15-19 of Algorithm 1 and in the Algorithm from [14] which only serve as refinement loops by an "if"-condition in order to prevent over-refinement. Since we want to either refine or make a Gauss Newton step, lines 15-19 are replaced by

Algorithm 2.

1: if (79) is violated then 2: Refine grid with respect to η_1^k such that we obtain a finer discretization h_k^3 3: $h = h_k^3$. 4: else 5: Set $q_{old}^{k+1} = q_h^k$, $u_{old}^{k+1} = u_{old}^k + v_h^k$. The trust way of the larger is the same as in Algorithm 1 from [22] but have

The structure of the loops is the same as in Algorithm 1 from [22], but here, we only have to solve linear PDEs (i.e. Step 6 in Algorithm 4 in [22] is replaced by "Solve linear PDE"), which justifies the drawback of one additional loop in comparison to [21] (see also Algorithm 5 in [22]. This motivates the implementation and assumes the gain of computation time for strongly nonlinear problems, which will be considered in terms of numerical tests in Section 4.

4 Numerical Results

For illustrating the performance of the proposed method according to Algorithm 1, we apply it to the example PDE

$$\begin{cases} -\Delta u + \zeta u^3 = q & \text{in } \Omega \\ u = 0 & \text{on } \partial \Omega \end{cases}$$

where we aim to identify the parameter $q \in Q = L^2(\Omega)$ from noisy measurements $g^{\delta} \in G$ of the state $u \in H_0^1(\Omega)$ in $\Omega = (0,1)^2 \subset \mathbb{R}^2$, where $\zeta > 0$ is a given constant. As for the measurements we consider two cases:

- (i) via point functionals in n_m uniformly distributed points ξ_i , $i = 1, 2, ..., n_m$ and perturbed by uniformly distributed random noise of some percentage p > 0. Then the observation space is chosen as $G = \mathbb{R}^{n_m}$ and the observation operator is defined by $(C(v))_i = v(\xi_i)$ for $i = 1, ..., n_m$.
- (ii) via L^2 -projection. Then $G = L^2(\Omega)$, C = id, and

$$g^{\delta} = g + \delta \frac{r}{\|r\|_{L^{2}(\Omega)}} = g + p \left\|g\right\|_{L^{2}(\Omega)} \frac{r}{\|r\|_{L^{2}(\Omega)}}$$

where r denotes some uniformly distributed random noise and p the percentage of perturbation. The exact state u^{\dagger} is simulated on a very fine mesh with 1050625 nodes and equally sized quadratic cells, and we denote the corresponding finite element space by V_{h_L} . In order to evaluate $||C(u) - g^{\delta}||_{L^2(\Omega)} = ||u - g^{\delta}||_{L^2(\Omega)}$ on coarser meshes and the corresponding finite element spaces V_{h_l} with $l = 0, 1, \ldots, L$ during the optimization algorithm, g^{δ} has to be transferred from V_{h_L} to the current grid V_{h_l} . As usual in the finite element context, this is done by the L^2 -projection as the restriction operator.

We consider configurations with three different exact sources q^{\dagger} :

(a) A Gaussian distribution

$$q^{\dagger} = \frac{c}{2\pi\sigma^2} \exp\left(-\frac{1}{2}\left(\left(\frac{sx-\mu}{\sigma}\right)^2 + \left(\frac{sy-\mu}{\sigma}\right)^2\right)\right)$$

with c = 10, $\mu = 0.5$, $\sigma = 0.1$, and s = 2.

(b) Two Gaussian distributions added up to one distribution

$$q^{\mathsf{T}} = q_1 + q_2,$$

where

$$q_{1} = \frac{c_{1}}{2\pi\sigma^{2}} \exp\left(-\frac{1}{2}\left(\left(\frac{s_{1}x-\mu}{\sigma}\right)^{2} + \left(\frac{s_{1}y-\mu}{\sigma}\right)^{2}\right)\right),$$
$$q_{2} = \frac{c_{2}}{2\pi\sigma^{2}} \exp\left(-\frac{1}{2}\left(\left(\frac{s_{2}x-\mu}{\sigma}\right)^{2} + \left(\frac{s_{2}y-\mu}{\sigma}\right)^{2}\right)\right)$$

with $\sigma = 0.1$, $\mu = 0.5$, $s_1 = 2$, $s_2 = 0.8$, $c_1 = 1$, and $c_2 = 1$.

(c) The step function

$$q^{\dagger} = \begin{cases} 0 & \text{for } x \ge \frac{1}{2} \\ 1 & \text{for } x < \frac{1}{2} \end{cases}.$$

The concrete choice of the parameters for the numerical tests is as follows: $c_{tc} = 10^{-7}$, $\tilde{\theta} = 0.4999$, $\tilde{\theta} = 0.2$, $\tau = 5$, $\tau_{\beta} = 1.66$, $\tilde{\tau}_{\beta} = 1$, $(c_2 = 0.9999, c_3 = 0.0001)$. The coarsest (starting mesh) consists of 25 nodes and 16 equally sized squares, the initial values for the control and the state are $q_0 = 0$ and $u_0 = 0$ and we start with a regularization parameter $\beta = 10$.

Considering the numerical tests, we are mainly interested in saving computation time compared to the Algorithm from [21], where the inexact Newton method for the determination of the regularization parameter β is applied directly to the nonlinear problem, instead of the linearized subproblems (80). That is why besides the numerical results for the Generalized Gauss-Newton (GGN) method presented in section 3, we also present the results from the "Nonlinear Tikhonov" (NT) Algorithm from [21].

The choice of the parameters for (NT) is the following: $\tilde{\tau} = 0.1, \underline{\tau} = 3.1, \tau = 4, \overline{\tau} = 5, c_{tc} = 10^{-7}, c_1 = 0.9, c_2 = 0.4$. This setting implies that both algorithms (NT) and (GGN) are stopped, if the concerning quantities of interest fall below the same bound ($\overline{\tau}^2 \delta^2$ for (NT) and $\tau^2 \delta^2$ for (GGN)).



Figure 2: FLTR: exact control q^{\dagger} , reconstructed control by NT, reconstructed control by GGN for example (a) (i) with $\zeta = 100, 1\%$ noise



Figure 3: FLTR: exact state u^{\dagger} , reconstructed state by NT, reconstructed state by GGN for example (a) (i) with $\zeta = 100, 1\%$ noise

The figures 2 and 3 show the exact source distribution q^{\dagger} and the corresponding simulated state u^{\dagger} , as well as the reconstructions of the control and the state obtained by Algorithm 1 (GGN), as well as the ones obtained by the algorithm from [21] (NT) for the example (a)(i) with $\zeta = 100$ and 1% noise. In Figure 4 we see the very fine mesh for simulating the data, the adaptively refined mesh obtained by (NT), and the adaptively refined mesh obtained by (GGN).

In table 1 we present the respective results for different choices of ζ (first column). In the second and fifth column one can see the relative control error $\frac{\left\|q_{h}^{*}-q^{\dagger}\right\|_{Q}^{2}}{\left\|q^{\dagger}\right\|_{Q}}$, in the third and sixth column is the number of nodes in the adaptively refined mesh, and in the forth and seventh column one can see the regularization parameter obtained by (GGN) and (NT) respectively.



Figure 4: FLTR: adaptively refined mesh by NT , adaptively refined mesh by GGN for example (a) (i) with $\zeta = 100, 1\%$ noise

Table 1: Algorithm 1 (GGN) versus the algorithm from [21] (NT) for Example (a)(i) for different choices of ζ with 1% noise. CTR: Computation time reduction using (GGN) in comparison to (NT)

ζ	NT			GGN			CTR
	error	β	# nodes	error	β	# nodes	
1	0.418	2985	2499	0.412	4600	3873	-65%
10	0.417	3194	2473	0.411	4918	3965	-59%
100	0.408	5014	6653	0.417	6773	9813	39%
500	0.418	9421	11851	0.404	13756	821	97%
1000	0.439	11486	44391	0.426	16355	793	99%

The eighth column shows the gain of computation time using (GGN) instead of (NT). The higher the factor ζ is, the more computation time we save with (GGN). This is probably due to the higher number of iterations needed for "more nonlinear" problems. Already for the choice $\zeta = 100$ replacing the nonlinear PDEs by linear ones (getting an additional loop in return cf. subsection 3.1.1) seems to pay off. In that case (GGN) refines more than (NT) (see Figure 4), but it is still faster (see table 1). For higher $\zeta = 500$ and $\zeta = 1000$ (GGN) is even much faster than (NT), because in addition to the cheaper linear PDEs, it also refines less. At the same time, the relative control error is about the same as with (NT).

In table 2 the reader can see the results for the same example with $\zeta = 100$ for different noise levels using (GGN). The numerical results confirm what we would expect: the larger the noise, the larger the error, the stronger the regularization, the coarser the discretization.

Taking a look at Figure 4 the reader can track the behavior of Algorithm 1 (GGN) for the considered example (a)(i) with $\zeta = 100$ and 1% noise. The algorithm goes from right to left in Figure 4, where the quantities of interest I_2 and I_3 (or rather their discrete counterparts $I_{3,h}$ and $I_{2,h}$) are rather large. The noise level for the inner iteration $\tilde{\theta}I_{3,h}$ is about 0.52 in the beginning. For this noise level the stopping criterion for the β -algorithm (step 10,11 in Algorithm 1) is already fulfilled, such that only one Gauss-Newton step is made without refining or updating β . This decreases the noise level $\tilde{\theta}I_{3,h}$ to about 0.33. Then the β -algorithm comes into play, with one refinement step, two β -steps and again one refinement step, which in total reduces I_2 from 0.90 to 0.33, with which the β -algorithm terminates. The subsequent run of the β -algorithm consists only of three β -enlargement steps and finally after 7 Gauss-Newton iterations, both quantites of interest I_2 and I_3 fulfill the required smallness conditions such that the whole Gauss-Newton Algorithm terminates.

Due to the observation above concerning the nonlinearity of the PDE, we restrict our considerations to the case $\zeta = 1000$ for the rest of this section. The figures 6, 7, and 8 again show the results for example (a) with 1% noise, but for the case (ii), i.e. via L^2 -projection.

(GGN) yields a regularization parameter $\beta = 4400806$, a discretization with 1125 nodes and a relative control error of 0.268. (NT) leads to a much larger error of 1.472, a finer discretization with 1405 nodes and a much larger regularization parameter $\beta = 60875207$. Although (GGN) refines only a little less than (NT), (GGN) is much faster than (NT), namely 81%. Compared to the point measurement evaluation, the L^2 -projection causes smoother



Figure 5: Behavior of (GGN) for example (a)(i) with $\zeta = 100$ and 1% noise



Figure 6: FLTR: exact control q^{\dagger} , reconstructed control by NT, reconstructed control by GGN for example (a) (ii) with $\zeta = 1000, 1\%$ noise



Figure 7: FLTR: exact state u^{\dagger} , reconstructed state by NT, reconstructed state by GGN for example (a) (ii) with $\zeta = 1000, 1\%$ noise

Table 2: Example (a)(i) for different noise levels with $\zeta = 100$

noise	error	β	# nodes
0.5%	0.385	12260	9867
1%	0.417	6773	9813
2%	0.553	1687	413
4%	0.700	599	137
8%	0.937	42	137



Figure 8: FLTR: adaptively refined mesh by NT , adaptively refined mesh by GGN for example (a) (ii) with $\zeta = 1000, 1\%$ noise

solutions, which seem to reconstruct the exact data better, but at the same time this is probably the less realistic case with respect to real applications.

In the figures 9 and 10, we can see the results using (GGN) and (NT) for a different source, namely example (b) with point measurements (i) and again $\zeta = 1000$ and 1% noise. Since we are interested in idenfying the parameter q, we take a pass on presenting the reconstructed states and only show the reconstructed controls, as well as the adaptively refined meshes.



Figure 9: FLTR: exact control q^{\dagger} , reconstructed control by NT, reconstructed control by GGN for example (b) (i) with $\zeta = 1000, 1\%$ noise

(GGN) stops with a regularization parameter $\beta = 1616$, a mesh with 6697 nodes, and a reconstruction yielding a relative error of 0.247, whereas (NT) terminates with $\beta = 539$, 10063 nodes and a larger error of 0.366. Due to the much coarser discretization obtained by (GGN), it is not surprising, that we save about 26% of computation time in this case.

The corresponding results for the source (c) are shown in Figure 11 and Figure 12.

Using (GGN) we obtain a regularization parameter $\beta = 379$, a discretization with 9565 nodes and a relative control error of 0.433, while (NT) yields $\beta = 24$, 5367 nodes and an error of 0.615. Also for this configuration (GGN) is faster than (NT), if only 9.4%. To put this in perspective, we would like to mention that the step function (c) is a very challenging example, since the initial guess $q_0 = 0$ and the source q^{\dagger} have different values on the boundary. Moreover, for piecewise constant functions total variation regularization is known to yield much better results than L^2 regularization.



Figure 10: FLTR: exact (very fine) mesh, adaptively refined mesh by NT , adaptively refined mesh by GGN for example (b) (i) with $\zeta=1000,\,1\%$ noise



Figure 11: FLTR: exact control q^{\dagger} , reconstructed control by NT, reconstructed control by GGN for example (c) (i) with $\zeta = 1000, 1\%$ noise



Figure 12: FLTR: exact (very fine) mesh, adaptively refined mesh by NT , adaptively refined mesh by GGN for example (c) (i) with $\zeta=1000,\,1\%$ noise

5 Conclusions and Remarks

In this paper we consider all-at-once formulations of the iteratively regularized Gauss-Newton method and their adaptive discretizations using a posteriori error estimators. This allows us to consider only the linearized PDE (instead of the full potentially nonlinear one) as a constraint in each Newton step, which safes computational effort. Alternatively, in a least squares approach, the measurement equation and the PDE are treated simultaneously via unconstrained minimization of the squared residual. In both cases we show convergence and convergence rates which we carry over to the discretized setting by controlling precision only in four real valued quantities per Newton step. The choices of the regularization parameters in each Newton step and of the overall stopping index are done a posteriori, via a discrepancy type principle. From the numerical tests we have seen, that the presented method yields reasonable reconstructions and can even lead to a large reduction of computation time compared to similar non-iterative methods.

6 Acknowledgments

The authors would like to thank the Federal Ministry of Education and Research (BMBF) for financial support within the grant 05M2013 "ExtremSimOpt: Modeling, Simulation and Optimization of Fluids in Extreme Conditions", as well as the German Science Foundation (DFG) for their support within the grant KA 1778/5-1 and VE 368/2-1 "Adaptive Discretization Methods for the Regularization of Inverse Problems".

References

- U. ASCHER AND E. HABER, A multigrid method for distributed parameter estimation problems, ETNA 15, (2003), 1–12.
- G. BIROS AND O. GHATTAS, Parallel Lagrange-Newton-Krylov-Schur Methods for PDE-Constrained Optimization. Part I: The Krylov-Schur Solver, SIAM Journal on Scientific Computing, 27 (2005) 687–713
- [3] A. B. BAKUSHINSKII, The problem of the convergence of the iteratively regularized Gauss-Newton method, Comput. Math. Math. Phys. 32 (1992), 1353–1359.
- [4] A.B. BAKUSHINSKY AND M. KOKURIN: Iterative Methods for Approximate solution of Inverse Problems. Kluwer Academic Publishers, Dordrecht, 2004.
- [5] R. BECKER, R. RANNACHER, An Optimal Control Approach to a-Posteriori Error Estimation Acta Numerica 2011 ed. A Iseries, (Cambridge University Press), 1–102.
- [6] R. BECKER, B. VEXLER, A posteriori error estimation for finite element discretizations of parameter identification problems Numer. Math. 96 (2004), 435–59
- [7] B. BLASCHKE(-KALTENBACHER), A. NEUBAUER, O. SCHERZER, On convergence rates for the iteratively regularized Gauβ-Newton method, IMA J. Numer. Anal. 17 (1997), 421–436.
- [8] H.-G. BOCK, Randwertproblemmethoden zur Parameteridentifizierung in Systemen nichtlinearer Differentialgleichungen, Bonner Mathematische Schriften 183, Bonn, 1987.
- M. BURGER AND W. MÜHLHUBER: Iterative regularization of parameter identification problems by sequential quadratic programming methods, Inverse Problems 18 (2002) 943–969.
- [10] M. BURGER AND W. MÜHLHUBER: Numerical approximation of an SQP-type method for parameter identification, SIAM J. Numer. Anal. 40 (2002), 1775–1797.
- [11] H.W. ENGL, M. HANKE, A. NEUBAUER, Regularization of Inverse Problems, Kluwer, Dordrecht, 1996.
- [12] J. FLEMMING: Theory and examples of variational regularisation with non-metric fitting functionals, Journal of Inverse and Ill-Posed Problems, 18(6), 2010.
- [13] C. GEIGER, C. KANZOW, Theorie und Numerik restringierter Optimierungsaufgaben, Springer, New York, 2002.

- [14] A. Griesbaum, B. Kaltenbacher, B. Vexler: Efficient computation of the Tikhonov regularization parameter by goal-oriented adaptive discretization, Inverse Problems 24 (2008),
- [15] M. HANKE, A regularization Levenberg-Marquardt scheme, with applications to inverse groundwater filtration problems, Inverse Problems 13 (1997), 79–95.
- [16] M. HANKE, The regularizing Levenberg-Marquardt scheme is of optimal order, Journal of Integral Equations and Applications 22(2), (2010).
- [17] , M. HANKE AND A. NEUBAUER AND O. SCHERZER: A convergence analysis of the Landweber iteration for nonlinear ill-posed problems, *Numer. Math.* 72 (1995), p. 21-37.
- [18] T. HOHAGE, Iterative Methods in Inverse Obstacle Scattering: Regularization Theory of Linear and Nonlinear Exponentially Ill-Posed Problems, PhD thesis, University of Linz, 1999.
- [19] T. HOHAGE AND F. WERNER: Iteratively regularized Newton methods with general data misfit functionals and applications to Poisson data, Numerische Mathematik 123 (2013), 745-779.
- [20] B. KALTENBACHER, B. HOFMANN: Convergence rates for the Iteratively Regularized Gauss-Newton method in Banach spaces, Inverse Problems 26 (2010) 035007.
- [21] B. KALTENBACHER, A. KIRCHNER, B. VEXLER: Adaptive discretizations for the choice of a Tikhonov regularization parameter in nonlinear inverse problems, Inverse Problems 27 (2011) 125008.
- [22] B. KALTENBACHER, A. KIRCHNER, S. VELJOVIĆ: Goal oriented adaptivity in the IRGNM for parameter identification in PDEs I: reduced formulation, submitted.
- [23] B. KALTENBACHER, A. NEUBAUER, O. SCHERZER: Iterative Regularization Methods for Nonlinear Ill-Posed Problems. Walter de Gruyter, Berlin – New York, 2008.
- [24] B. KALTENBACHER, F. SCHÖPFER, AND T. SCHUSTER: Convergence of some iterative methods for the regularization of nonlinear ill-posed problems in Banach spaces. *Inverse Problems* 25 (2009), 065003 (19pp). DOI:10.1088/0266-5611/25/6/065003.
- [25] M.T. NAIR AND E. SCHOCK AND U. TAUTENHAHN: Morozov's discrepancy principle under general source conditions. Anal. Anw. 22 (2003), p. 199-214.
- [26] C. PÖSCHL. Tikhonov Regularization with General Residual Term. PhD thesis, Universität Innsbruck, October 2008.
- [27] A. RIEDER, On the regularization of nonlinear ill-posed problems via inexact Newton iterations, Inverse Problems 15 (1999), 309–327.
- [28] O. SCHERZER: Convergence criteria of iterative methods based on Landweber iteration for nonlinear problems, J. Math. Anal. Appl. 194 (1995), p. 911-933.

Appendix

For proving convergence rates of the iterates according to (22), we consider source conditions of the form

$$\exists (s,v) \in Q \times V \text{ s.t. } (q^{\dagger} - q_0, u^{\dagger} - u_0) = \kappa(\mathbf{F}'(q^{\dagger}, u^{\dagger})^* \mathbf{F}'(q^{\dagger}, u^{\dagger}))(s,v).$$

$$\tag{81}$$

with $\kappa = \kappa_{\nu}$ or $\kappa = \kappa_{p}$ as in the following Lemma. The case $\kappa = \kappa_{\nu}$ with $\nu = 0$ corresponds to the pure convergence case without rates. Using the interpolation inequality and Lemma 3.13 in [18] we immediately get the following result, that is crucial for convergence and convergence rates.

Lemma 3. Under Assumption 2 we have for $\alpha > 0$ $\mu \in [0, \alpha]$, as well as any $\nu \in [0, 1]$ and any p > 0

$$\alpha \left\| \left(\mathbf{T}^* \mathbf{T} + \begin{pmatrix} \alpha I & 0 \\ 0 & \mu I \end{pmatrix} \right)^{-1} \kappa_{\nu} (\mathbf{T}^* \mathbf{T}) \right\| \le C_{\nu} \alpha^{\nu}$$
(82)

$$\alpha \left\| \left(\mathbf{T}^* \mathbf{T} + \begin{pmatrix} \alpha I & 0 \\ 0 & \mu I \end{pmatrix} \right)^{-1} \kappa_p \left(\frac{1}{e \|\mathbf{T}\|^2} \mathbf{T}^* \mathbf{T} \right) \right\| \le C_p f_p(\alpha)$$
(83)

where $\kappa_{\nu}(\lambda) = \lambda^{\nu}, \ \kappa_{p}(\lambda) = \ln(\frac{1}{\lambda})^{-p}, \ \lambda \in (0, 1/e].$

Therewith, the following convergence and convergence rates result with a priori chosen sequence α_k and stopping index k_* follow directly along the lines of the proofs of Theorem 2.4 in [7] and Theorem 4.7 in [18], see also Theorem 4.12 in [23]:

Theorem 5. Let β_k be a positive sequence decreasing monotonically to zero and satisfying

$$\sup_{k\in\mathbb{N}}\frac{\beta_{k+1}}{\beta_k}<\infty\,,$$

and let $k_* = k_*(\delta)$ be chosen according to

$$k_* \to \infty$$
 and $\eta \ge \delta \beta_{k_*}^{\frac{1}{2}} \to 0$ as $\delta \to 0$

and

$$\eta \beta_{k_*}^{-\nu - \frac{1}{2}} \le \delta < \eta \beta_k^{-\nu - \frac{1}{2}} , \qquad 0 \le k < k_* ,$$

in case of (81) with $\kappa(\lambda) = \lambda^{\nu}, \nu \in (0, 1]$ or

$$\frac{\eta}{\beta_{k_*}} \le \delta < \frac{\eta}{\beta_k} \,, \qquad 0 \le k < k_* \;,$$

in case of (81) with $\kappa(\lambda) = \ln(\frac{1}{\lambda})^{-p}$, p > 0 respectively.

1. If (81) holds with $\kappa(\lambda) = \lambda^{\nu}, \ \nu \in [0, \frac{1}{2}]$ or $\kappa(\lambda) = \ln(\frac{1}{\lambda})^{-p}, \ p > 0$, we assume that

$$\mathbf{F}'(\tilde{q}, \tilde{u}) = R((\tilde{q}, \tilde{u}), (q, u))\mathbf{F}'(q, u) + Q((\tilde{q}, \tilde{u}), (q, u)) \\
\|I - R((\tilde{q}, \tilde{u}), (q, u))\| \leq c_R \\
\|Q((\tilde{q}, \tilde{u}), (q, u))\| \leq c_Q \|\mathbf{F}'(q^{\dagger}, u^{\dagger})((\tilde{q}, \tilde{u}) - (q, u))\|$$

for all $(q, u), (\tilde{q}, \tilde{u}) \in \mathcal{B}_{\rho}(q_0, u_0)$ and that $||(q^{\dagger}, u^{\dagger}) - (q_0, u_0)||$, ||(s, v)||, η , ρ , c_R are sufficiently small.

2. If (81) holds with $\kappa(\lambda) = \lambda^{\nu}$, $1/2 \leq \nu \leq 1$, we assume that

 $\|\mathbf{F}'((\tilde{q}, \tilde{u})) - \mathbf{F}'(q, u)\| \le L \|(\tilde{q}, \tilde{u}) - (q, u)\|$

for all $(q, u), (\tilde{q}, \tilde{u}) \in \mathcal{B}_{\rho}(q_0, u_0)$ and $||(q^{\dagger}, u^{\dagger}) - (q_0, u_0)||$, ||(s, v)||, η , ρ are sufficiently small.

Then for $(q^{k_*,\delta}, u^{k_*,\delta})$ defined by (22) (i.e., (23)), we obtain convergence $(q^{k_*,\delta}, u^{k_*,\delta}) \rightarrow (q^{\dagger}, u^{\dagger})$ as $\delta \rightarrow 0$ and convergence rates

$$\|q_h^{k_*} - q^{\dagger}\|^2 + \|u_h^{k_*} - u^{\dagger}\|^2 \le \frac{\bar{C}^2 \delta^2}{\Theta^{-1}\left(\frac{\bar{C}}{2\|(s,v)\|}\delta\right)} = 4\|(s,v)\|^2 \kappa^2 (\Theta^{-1}(\frac{\bar{C}}{2\|(s,v)\|}\delta))$$
(84)

where $\Theta(\lambda) \coloneqq \kappa(\lambda) \sqrt{\lambda}$.