The automorphism group of the \mathbb{Z}_2 -orbifold of the Barnes-Wall lattice vertex operator algebra of central charge 32

Hiroki Shimakura*

Research Center for Pure and Applied Mathematics, Graduate School of Information Sciences, Tohoku University, Aramaki aza Aoba 6-3-09, Aoba-ku Sendai-city, 980-8579, Japan e-mail: shimakura@m.tohoku.ac.jp

2010 Mathematics Subject Classification. Primary 17B69; Secondary 20B25.

Abstract

In this article, we prove that the full automorphism group of the \mathbb{Z}_2 -orbifold of the Barnes-Wall lattice vertex operator algebra of central charge 32 has the shape $2^{27} \cdot E_6(2)$. In order to identify the group structure, we introduce a graph structure on the Griess algebra and show that it is a rank 3 graph associated to $E_6(2)$.

Introduction

The automorphism group of a vertex operator algebra (VOA) relates to many areas in mathematics. For example, the automorphism group of the moonshine VOA is the Monster simple group, and this realization plays important roles in the moonshine phenomena, an amazing relation between group theory, VOA theory and number theory ([Bo92]). Hence it has been an important problem to determine the automorphism group of a VOA.

One of the standard methods of determining the automorphism group Aut V of a VOA V is to restrict its action to the non-trivial lowest weight subspace. If the weight 1 subspace V_1 is non-zero, then Aut V can be studied by using the Lie algebra structure on V_1 . If the weight 1 subspace is zero, then the weight 2 subspace V_2 forms a commutative (non-associative) algebra, called the Griess algebra. Indeed, the automorphism groups of some VOAs were calculated by using the Griess algebras ([DG98, DG05, FLM88, Gr98, MM00, LSY07]). However, in general, it is difficult to determine the automorphism group of the Griess algebra.

The Barnes-Wall lattices BW_{2^n} of rank 2^n are even lattices in \mathbb{R}^{2^n} ([BW59]). They have remarkable automorphism groups of shape $2^{1+2n}_+ \Omega^+(2n,2)$ if $n \ge 4$, and play important roles in lattice theory, such as sphere packings (cf. [CS99]). Since there are many analogues between lattice theory and VOA theory, it is believed that VOAs with large symmetry can be obtained from Barnes-Wall lattices; indeed, the VOAs $V^+_{\sqrt{2}BW_8}$ and $V^+_{BW_{16}}$ are such examples ([Sh04, Sh06, Hö08]). The next example would be the holomorphic VOA $\tilde{V}_{BW_{32}}$ of central charge 32 obtained as the \mathbb{Z}_2 -orbifold of the lattice VOA $V_{BW_{32}}$; in [Mi04, Mi], an infinite series of holomorphic framed VOAs with finite automorphism groups are constructed, and $\tilde{V}_{BW_{32}}$ turns out to be the VOA next to the moonshine VOA.

^{*}The author was partially supported by JSPS KAKENHI Grant Numbers 18740001, 20549004, 23540013.

The automorphism group of the lattice VOA V_L associated to even lattice L was determined in [DN99] as $N(V_L) \cdot O(\hat{L})$, where $N(V_L) = \langle \exp(a_0) | a \in (V_L)_1 \rangle$ and $O(\hat{L})$ is the automorphism group of the central extension \hat{L} of L by $\mathbb{Z}/2\mathbb{Z}$. On the other hand, the automorphism group of the \mathbb{Z}_2 -orbifold \tilde{V}_L has not been well studied except for the moonshine VOA (cf. [FLM88]).

The main theorem of this article is the following:

Theorem 2.30. Let $\tilde{V}_{BW_{32}}$ be the \mathbb{Z}_2 -orbifold of the lattice VOA associated to the Barnes-Wall lattice BW₃₂ of rank 32. Then its automorphism group has the shape $2^{27} \cdot E_6(2)$.

Note that this theorem was conjectured by Griess in a workshop held on July 2004 at Edinburgh.

Let us provide two related topics, VOAs of class S^n and trace functions.

A VOA is said to be of class S^n if its trivial component with respect to the full automorphism group coincides with the subVOA generated by the conformal element up to weight n ([Ma01]). This is a sufficient condition that all homogeneous subspaces are conformal n-designs, introduced in [Hö08]; the notion of conformal designs based on VOAs is a natural analogue of the notion of block and spherical designs based on binary codes and lattices, respectively. Although the VOAs of class S^6 with $V_1 \neq 0$ are classified in [Hö08], few candidates of VOAs of class S^6 with $V_1 = 0$ are known ([Ma01, Hö08]); $\tilde{V}_{BW_{32}}$ is one of them ([Hö08, Example 3.2]).

For a VOA $V = \bigoplus_{i=0}^{\infty} V_i$ and $g \in \operatorname{Aut} V$, a trace function is given by $T_g(q) = \sum_{i=0}^{\infty} \operatorname{Tr}(g_{|V_i})q^{i-c/24}$. For example, trace functions for the moonshine VOA provide a correspondence between the conjugacy classes of the Monster simple group and certain modular functions ([Bo92]). Hence, it would be an interesting problem to study similar phenomena on trace functions for $\tilde{V}_{BW_{32}}$ and its automorphism group of shape $2^{27} \cdot E_6(2)$; notice that $T_1(q)$ is just the graded dimension and $T_1(q) = \sqrt[3]{j}(j-992)$, where $\sqrt[3]{j} = q^{-1/3}(1+248q+4124q^2+\dots)$ (cf. [Hö95]).

Let G be the automorphism group of $\tilde{V}_{BW_{32}}$. The proof of the main theorem is divided into three steps:

- (i) Construct an elementary abelian 2-subgroup A of G of order 2^{27} ;
- (ii) Prove that $N_G(A)/A$ is isomorphic to $E_6(2)$, where $N_G(A)$ is the normalizer of A in G;
- (iii) Prove that A is normal in G.

Let us explain these steps in more detail.

(i) Set $V = \tilde{V}_{BW_{32}}$ and $U = V_{BW_{16}}^+$. By the description of BW_{32} as an overlattice of $BW_{16} \oplus BW_{16}$ in [Gr05], V contains $U^{\otimes 2}$ as a subVOA such that as $U^{\otimes 2}$ -modules,

$$V \cong \bigoplus_{[M] \in S} M^{\otimes 2},$$

where S is the set of all isomorphism classes of irreducible U-modules. Since S forms an elementary abelian 2-group of order 2^{10} under the fusion rules ([AD04, ADL05, Sh04]), V is a simple current extension of $U^{\otimes 2}$ graded by S. Hence the dual S^* of S acts on V as an automorphism group. By [Sh04, Theorem 3.3] the restriction group homomorphism

$$\rho: N_G(S^*) \to \{g \in \operatorname{Aut} U^{\otimes 2} \mid \{g \circ (M^{\otimes 2}) \mid M \in S\} = \{M^{\otimes 2} \mid M \in S\}\}$$

is surjective and its kernel is identical to S^* , where $N_G(S^*)$ is the normalizer of S^* in G; the image of ρ is described by $\operatorname{Aut} U^{\otimes 2} \cong \operatorname{Aut} U \wr \mathbb{Z}_2$ ([Sh11]) and the description of the action of Aut U on S in [Sh04]. Let us consider the subgroup A of G generated by S^* and $\rho^{-1}(\langle \Delta(O_2(\operatorname{Aut} U)), \tau_U \rangle)$, where $\Delta(O_2(\operatorname{Aut} U))$ is the subgroup $\{(h, h) \mid h \in O_2(\operatorname{Aut} U)\}$ of Aut $U \wr \mathbb{Z}_2$ and $\tau_U \in \operatorname{Aut} U^{\otimes 2}$ sends $x \otimes y$ to $y \otimes x$; one can see that A is abelian, of order 2^{27} and normal in $N_G(S^*)$. We view A as a subgroup of $C_G(z)$, where z is the involution in G associated to the \mathbb{Z}_2 -grading $V = V_{\mathrm{BW}_{32}}^+ \oplus V_{\mathrm{BW}_{32}}^{T,+}$. By using the explicit description of $C_G(z)$ in terms of Aut $V_{\mathrm{BW}_{32}}^+$ ([FLM88]), we prove that A is normal in $C_G(z)$ and that A is elementary abelian.

(ii) By the arguments in (i), $N_G(A)$ contains both $C_G(z)$ and $N_G(S^*)$. In order to identify the structure of $N_G(A)$, we consider its action on the graph Γ defined as follows: the vertex set $V\Gamma$ is the set of eigenspaces of A in the orthogonal complement B of the conformal element in V_2 , and two distinct vertices B_1 and B_2 are adjacent if there exist $x_i \in B_i$ such that $x_1 \cdot x_2 \neq 0$, where \cdot is the product in V_2 ; $N_G(A)/A$ acts on this graph as an automorphism group. By the action of $N_G(S^*)$ and $C_G(z)$ on B, B is an irreducible module for $N_G(A)$. Hence $N_G(A)/A$ acts transitively on $V\Gamma$. For the irreducible character $\mu \in A^*$ specified in (2.5), we show that the dimension of the eigenspace B_{μ} with character μ is 1 and its stabilizer in $N_G(A)/A$ is identical to $N_G(S^*)/A$. Moreover, we determine the vertices adjacent to B_{μ} . Hence, the cardinality of $V\Gamma$ is equal to dim B(= 139503), the valency is 4590, and the stabilizer of a vertex in $N_G(A)/A$ is isomorphic to $N_G(S^*)/A$. In addition, we also prove in Proposition 2.20 that the action of $N_G(A)/A$ on $V\Gamma$ is rank 3. According to the classification of primitive permutation groups of rank 3 ([Li87] and references therein), $N_G(A)/A$ must be $E_6(2)$.

(iii) Since V is finitely generated, G is algebraic ([DG02]). By an argument similar to [Ti84], G is discrete, and hence G is finite. Using finite group theory, we prove that A is normal in G (Proposition 2.29); remark that the classification of finite simple groups is used to prove the following: there are no finite simple groups K with an involution t such that $C_K(t) \cong 2^{1+32} \cdot 2^{10} \cdot \Omega^+(10, 2)$.

The organization of this article is as follows: In §1, we recall and give some definitions and facts necessary in this article, including a review of $V_{BW_{16}}^+$ in §1.5. In §2.1 and §2.2, we discuss the structure of two subgroups $C_G(z)$ and $N_G(S^*)$ of G and their actions on B, respectively. In §2.3, we construct an elementary abelian 2-subgroup A of order 2^{27} , which is a normal subgroup of both $C_G(z)$ and $N_G(S^*)$. In §2.4, we introduce a graph Γ , and in §2.5, we study its structure and the action of $N_G(A)/A$ on it. In §2.6, we prove that the action of $N_G(A)/A$ on Γ is rank 3, and in §2.7 we review the classification of permutation groups of rank 3 and identify $N_G(A)/A$. In §2.8, we prove that A is normal in G.

Notations

1_V	the vacuum vector of a VOA V .
2^n	an elementary abelian 2-group of order 2^n .
2^{1+2m}_{+}	an extraspecial 2-group of order 2^{1+2m} of plus type.
A	the elementary abelian 2-subgroup of $\operatorname{Aut} V$ defined in (2.4).
A^*	the set of all irreducible characters of A .
$\operatorname{Aut} X$	the automorphism group of X , where X is a lattice, a graph, or a VOA.
BW_{2^n}	the Barnes-Wall lattice of rank 2^n .
B	the orthogonal complement of the conformal element ω_V in the Griess algebra V_2 .
B_{χ}	the eigenspace of A in B with character χ .
C	the subgroup of Aut U which acts trivially on S, and in fact, $C = O_2(\operatorname{Aut} U)$.
$C_H(K)$	the centralizer of K in a group H .
$\Delta(H)$	the subgroup $\{(h,h) \in H \times H \mid h \in H\}$ of $H \wr \mathbb{Z}_2$.
G	the automorphism group of $V = \tilde{V}_{BW_{22}}$ in Section 2.

Γ	the graph associated to B and A (see Definition 2.12).
η	the canonical group homomorphism from $O(\hat{L})$ to Aut $L/\langle -1 \rangle$ defined in (1.6).
L^*	the dual lattice of L .
L(n)	the subset $\{a \in L \mid \langle a, a \rangle = n\}$ of a lattice L.
[M]	the isomorphism class of a module M for a VOA.
$N_H(K)$	the normalizer of a subgroup K in a group H .
$O_2(H)$	the maximal normal 2-subgroup of a group H .
$O(\hat{L})$	the automorphism group of the central extension \hat{L} of L by $\mathbb{Z}/2\mathbb{Z}$
ρ	the restriction group homomorphism from $N_G(S^*)$ to Aut $U^{\otimes 2}$ defined in (2.2).
S = R(U)	the set of all isomorphism classes of irreducible U -modules for the VOA U .
S^*	the subgroup of $\operatorname{Aut} V$ induced from the dual group of S .
θ	an involution in $O(\hat{L})$ whose image under the map $O(\hat{L}) \to \operatorname{Aut} L$ is -1
ω_V	the conformal element of a VOA V .
U, V	$U = V_{BW_{16}}^+$ and $V = \tilde{V}_{BW_{32}}$ in Section 2.
$V\Gamma$	the set of all vertices in a graph Γ .
V_L^+	the subVOA of the lattice VOA V_L fixed by θ .
\tilde{V}_L^-	the \mathbb{Z}_2 -orbifold of the holomorphic lattice VOA V_L associated to θ .
ν	the canonical group homomorphism from $C_{\text{Aut }\tilde{V}_L}(z)$ to $\text{Aut }V_L^+$ defined in (1.8).
μ	the irreducible character of A defined in (2.5).
X.Y	a group extension with normal subgroup X and quotient Y .
z	the automorphism of \tilde{V}_L which acts by 1 and -1 on V_L^+ and $V_L^{T,+}$, respectively.
\mathbb{Z}_2	a cyclic group of order 2.
Z(H)	the center of a group H .
$\Omega^{+}(10,2)$	the simple normal subgroup of the orthogonal group $O^+(10,2)$ of degree 10 over \mathbb{F}_2 .

1 Preliminaries

In this section, we recall or give some definitions and facts necessary in this article.

1.1 Barnes-Wall lattices

In this subsection, we review the Barnes-Wall lattices of rank 16 and 32 (cf. [BW59, CS99, Gr05]).

Let us start by recalling an explicit construction of the Barnes-Wall lattice BW₁₆ of rank 16. Let $\alpha_1, \alpha_2, \ldots, \alpha_{16}$ be an orthogonal basis of \mathbb{R}^{16} of (squared) norm 2. Let RM(1, 4) denote the first order Reed-Muller code of length 16, that is, RM(1, 4) is a 5-dimensional subspace of \mathbb{F}_2^{16} spanned by (1^{16}) , $((10)^8)$, $((1100)^4)$, $(1^40^41^40^4)$, and (1^80^8) . For an element $c = (c_i) \in \mathbb{F}_2^{16}$, set $\alpha_c = \sum_{i=1}^{16} c_i \alpha_i$, where we view c_i as an element in $\{0, 1\}$. Then

$$BW_{16} \cong \sum_{1 \le i,j \le 16} \mathbb{Z}(\alpha_i + \alpha_j) + \sum_{c \in RM(1,4)} \mathbb{Z}\frac{\alpha_c}{2}.$$
 (1.1)

Notice that $BW_{16}^*/BW_{16} \cong 2^8$ and $Aut BW_{16} \cong 2^{1+8}_+$. $\Omega^+(8,2)$ (cf. [CS99, §10 in Chapter 4]).

Let BW₃₂ denote the Barnes-Wall lattice of rank 32. It is well known that BW₃₂ is even and unimodular, it is generated by vectors of norm 4 and Aut BW₃₂ is of shape 2^{1+10}_{+} . $\Omega^{+}(10,2)$ (cf. [Gr05] and references therein). In [Gr05, Definition 6.2], BW₃₂ was constructed from BW₁₆ as follows:

$$BW_{32} \cong BW_{16} \oplus BW_{16} + \{(v, v) \in \mathbb{R}^{32} \mid v \in BW_{16}^*\},$$
(1.2)

where BW_{16}^* is the dual lattice of BW_{16} . Under the isomorphism (1.2), the sublattice $BW_{32}[1]$ of BW_{32} was defined in [Gr05, Definition 6.2 and Lemma 7.6] by

$$BW_{32}[1] \cong 2BW_{16}^* \oplus 2BW_{16}^* + \{(v, v) \in \mathbb{R}^{32} \mid v \in BW_{16}\}.$$
(1.3)

It follows from $\sqrt{2}BW_{16}^* \cong BW_{16}$ that $BW_{32}[1] \cong \sqrt{2}BW_{32}$. Hence $BW_{32}/BW_{32}[1] \cong 2^{16}$. By [Gr05, Corollary 6.5], Aut BW_{32} acts on $BW_{32}[1]$ as Aut $BW_{32}[1]$. By the same procedure, we obtain the sublattice $(BW_{32}[1])[1]$ of $BW_{32}[1]$; clearly $(BW_{32}[1])[1] = 2BW_{32}$.

Lemma 1.1. (cf. [Gr05, Corollary 6.5 and Lemmas 7.6 and 7.13]) The automorphism group of BW₃₂ preserves BW₃₂[1], and the subgroup of Aut BW₃₂ which acts trivially on BW₃₂[1]/2BW₃₂ is $O_2(\text{Aut BW}_{32}) \cong 2^{1+10}_+$. Moreover, BW₃₂/BW₃₂[1] and BW₃₂[1]/2BW₃₂ are irreducible modules over \mathbb{F}_2 for Aut BW₃₂.

The following lemma can be proved by the explicit construction of irreducible modules for extraspecial 2-groups of plus type (cf. [FLM88, §5.5]).

Lemma 1.2. Let $H \cong 2^{1+2m}_+$ and let W be a 2^m -dimensional irreducible H-module over \mathbb{C} on which the central element of H acts by -1. Then $W \otimes_{\mathbb{C}} W$ is a 2^{2m} -dimensional irreducible H * H-module by the natural action $(g, h) : v \otimes w \mapsto g(v) \otimes h(w)$, where * is the central product and $H * H \cong 2^{1+4m}_+$.

Later, we will apply this lemma to the 2⁴-dimensional irreducible module $\mathbb{C} \otimes_{\mathbb{Z}} BW_{16}$ for $O_2(Aut BW_{16}) \cong 2^{1+8}_+$.

1.2 Vertex operator algebras and modules

In this subsection, we recall the notion of vertex operator algebras (VOAs) and modules from [Bo86, FLM88, FHL93]. Throughout this article, the ground field is \mathbb{C} if not otherwise stated.

A vertex operator algebra (VOA) $(V, Y, \mathbf{1}_V, \omega_V)$ is a $\mathbb{Z}_{\geq 0}$ -graded vector space $V = \bigoplus_{m \in \mathbb{Z}_{\geq 0}} V_m$ equipped with a linear map

$$Y(a, z) = \sum_{i \in \mathbb{Z}} a_{(i)} z^{-i-1} \in (\text{End } V)[[z, z^{-1}]], \quad a \in V,$$

the vacuum vector $\mathbf{1}_V$ and the conformal element ω_V satisfying a number of conditions ([Bo86, FLM88]). We often denote it by V or (V, Y) simply.

Two VOAs $(V, Y, \mathbf{1}_V, \omega_V)$ and $(V', Y', \mathbf{1}_{V'}, \omega_{V'})$ are said to be *isomorphic* if there exists a linear isomorphism g from V to V' satisfying

$$g\omega_V = \omega_{V'}$$
 and $gY(v, z) = Y'(gv, z)g$ for $\forall v \in V$.

When V = V', such an isomorphism is called an *automorphism*. The group of all automorphisms of V is called the *automorphism group* of V and is denoted by Aut V.

A vertex operator subalgebra (or a subVOA) is a graded subspace of V which has a structure of a VOA such that the operations and its grading agree with the restriction of those of V and that they share the vacuum vector. When they also share the conformal element ω_V , we will call it a full subVOA.

An (ordinary) module (M, Y_M) for a VOA V is a C-graded vector space $M = \bigoplus_{m \in \mathbb{C}} M_m$ equipped with a linear map

$$Y_M(a, z) = \sum_{i \in \mathbb{Z}} a_{(i)} z^{-i-1} \in (\text{End } M)[[z, z^{-1}]], \quad a \in V$$

satisfying a number of conditions ([FHL93]). We often denote it by M and its isomorphism class by [M]. The *weight* of a homogeneous vector $v \in M_k$ is k. If M is irreducible then there exists the *lowest weight* $h \in \mathbb{C}$ of M such that $M = \bigoplus_{m \in \mathbb{Z}_{>0}} M_{h+m}$ and $M_h \neq 0$.

A VOA is said to be *simple* if V is irreducible as a V-module. A VOA is said to be *rational* if any (admissible) module is completely reducible. A rational VOA is said to be *holomorphic* if itself is the only irreducible module up to isomorphism. A VOA is said to be of CFT type if $V_0 = \mathbb{C}\mathbf{1}_V$, and is said to be C_2 -cofinite if dim $V/\text{Span}_{\mathbb{C}}\{u_{(-2)}v \mid u, v \in V\} < \infty$. Let R(V) denote the set of all isomorphism classes of irreducible V-modules.

Let M_a and M_b be modules for a simple, rational and C_2 -cofinite VOA V of CFT type. Then, up to isomorphism, there exists a module $M_a \boxtimes_V M_b$ for V satisfying some conditions, called the *fusion product* (cf. [HL95]). Note that $M_a \boxtimes_V M_b$ is uniquely determined by the isomorphism classes of M_a and M_b , up to isomorphism, and that its V-module structure is described by the fusion rules (cf. [FHL93]).

Let $M = (M, Y_M)$ be a module for a VOA V. For $g \in \operatorname{Aut} V$, let $g \circ M = (M, Y_{g \circ M})$ denote the V-module defined by $Y_{g \circ M}(v, z) = Y_M(g^{-1}v, z)$. Note that if M is irreducible then so is $g \circ M$ and that if $M_a \cong M_b$ then $g \circ M_a \cong g \circ M_b$. Hence Aut V acts on R(V). Since this action preserves the fusion rules (cf. [Sh11, Lemma 1.3]), it also does the fusion product.

Let $V = \bigoplus_{i=0}^{\infty} V_i$ be a VOA such that $V_0 = \mathbb{C} \mathbf{1}_V$ and $V_1 = 0$. Consider the first non-trivial subspace V_2 and set

$$a \cdot b = a_{(1)}b, \quad (a,b)\mathbf{1}_V = a_{(3)}b, \quad \text{for } a, b \in V_2.$$

Then \cdot is a commutative (non-associative) product and (,) is an invariant bilinear form. The space V_2 equipped with them is called the *Griess algebra* of V. Since Aut V fixes ω_V and preserves (,), it also acts on $\omega_V^{\perp} = \{v \in V_2 \mid (v, \omega_V) = 0\}.$

1.3 Simple currents and simple current extensions

In this subsection, we recall the notion of simple currents and simple current extensions.

Let V(0) be a simple, rational and C_2 -cofinite VOA of CFT type. An irreducible V(0)module M_a is called a *simple current* if for any irreducible V(0)-module M_b , there exists a unique irreducible V(0)-module M_c satisfying $M_a \boxtimes_{V(0)} M_b \cong M_c$, up to isomorphism. Let $\{V(\alpha) \mid \alpha \in D\}$ be a set of inequivalent irreducible V(0)-modules indexed by an abelian group D. A simple VOA $V = \bigoplus_{\alpha \in D} V(\alpha)$ is called a *simple current extension* of V(0) if it carries a D-grading and every $V(\alpha)$ is a simple current. Note that $V(\alpha) \boxtimes_{V(0)} V(\beta) \cong V(\alpha + \beta)$.

Proposition 1.3. ([DM04, Proposition 5.3]) Let V(0) be a simple, rational and C_2 -cofinite VOA of CFT type and let $V = \bigoplus_{\alpha \in D} V(\alpha)$ and $\tilde{V} = \bigoplus_{\alpha \in D} \tilde{V}(\alpha)$ be simple current extensions of $V(0) = \tilde{V}(0)$. If $V(\alpha) \cong \tilde{V}(\alpha)$ as V(0)-modules for all $\alpha \in D$, then V and \tilde{V} are isomorphic.

By the *D*-grading of a simple current extension $V = \bigoplus_{\alpha \in D} V(\alpha)$ of V(0), the group D^* consisting of irreducible characters of *D* acts on *V* as an automorphism group; $\chi \in D^*$ acts on $V(\alpha)$ by the scalar multiplication $\chi(\alpha)$ for each $\alpha \in D$. The normalizer of D^* in Aut *V* is describe as follows:

Proposition 1.4. ([Sh04, Theorem 3.3]) The restriction homomorphism

$$N_{\operatorname{Aut} V}(D^*) \to \left\{ g \in \operatorname{Aut} V(0) \mid \{ g \circ [V(\alpha)] \mid \alpha \in D \} = \{ [V(\alpha)] \mid \alpha \in D \} \right\}$$
(1.4)

is surjective, and its kernel is D^* .

Lemma 1.5. Let V be a simple current extension of V(0). Then $N_{\operatorname{Aut} V}(D^*) = \{g \in \operatorname{Aut} V \mid g(V(0)) = V(0)\}.$

Proof. Since V(0) is the subVOA of V fixed by D^* , $N_{\operatorname{Aut} V}(D^*)$ preserves V(0). Let $g \in \operatorname{Aut} V$ with g(V(0)) = V(0). Since V is a simple current extension of V(0), the multiplicity of any irreducible V(0)-submodule of V is 1. Hence g permutes irreducible V(0)-submodules of V, and thus $g \in N_{\operatorname{Aut} V}(D^*)$.

1.4 Lattice VOA and the \mathbb{Z}_2 -orbifold

Let L be an even lattice of rank n and let V_L denote the lattice VOA associated with L ([Bo86, FLM88]). Let \hat{L} be the central extension of L by $\{\pm 1\} \cong \mathbb{Z}/2\mathbb{Z}$ with commutator relation $[e^{\alpha}, e^{\beta}] = (-1)^{\langle \alpha, \beta \rangle}$. The automorphism group $O(\hat{L})$ of \hat{L} is described as follows ([FLM88, Proposition 5.4.1])

$$1 \to \operatorname{Hom}_{\mathbb{Z}}(L, \mathbb{Z}_2) \to O(\hat{L}) \xrightarrow{\gamma} \operatorname{Aut} L \to 1;$$
(1.5)

notice that $O(\hat{L})$ acts on V_L as an automorphism group ([FLM88, DN99]). We often call $g \in O(\hat{L})$ a lift of $h \in \operatorname{Aut} L$ if $\gamma(g) = h$. Let $\theta \in \gamma^{-1}(-1)$, a lift of $-1 \in \operatorname{Aut} L$. Let V_L^+ denote the subspace of V_L fixed by θ ; it is a simple VOA of CFT type of central charge n; notice that the VOA structure of V_L^+ is independent of the choice of θ up to isomorphism ([DGH98, Corollary D.7]). Clearly, $O(\hat{L})/\langle \theta \rangle \subset \operatorname{Aut} V_L^+$ and

$$1 \to \operatorname{Hom}_{\mathbb{Z}}(L, \mathbb{Z}_2) \to O(\hat{L})/\langle \theta \rangle \xrightarrow{\eta} \operatorname{Aut} L/\langle -1 \rangle \to 1$$
(1.6)

is exact. Indeed $\operatorname{Hom}_{\mathbb{Z}}(L, \mathbb{Z}_2)$ is a subgroup of $\operatorname{Aut} V_L$ which commute with θ . The explicit action of $\operatorname{Hom}_{\mathbb{Z}}(L, \mathbb{Z}_2)$ on V_L shows the following lemma (cf. [FLM88, §10.4]):

Lemma 1.6. Let $f \in \text{Hom}_{\mathbb{Z}}(L, \mathbb{Z}_2) \subset \text{Aut} V_L$. Then the following hold:

- (1) f acts trivially on the subspace $\{h(-1)\mathbf{1}_V \mid h \in \mathbb{C} \otimes_{\mathbb{Z}} L\} \cong \mathbb{C} \otimes_{\mathbb{Z}} L$ of $(V_L)_1$;
- (2) The subspace of V_L^+ fixed by f is V_N^+ , where $N = \{v \in L \mid f(v) = 0\}$.

We often identify $\operatorname{Hom}_{\mathbb{Z}}(L, \mathbb{Z}_2)$ with $L^*/2L^*$ as follows:

$$L^*/2L^* \to \operatorname{Hom}_{\mathbb{Z}}(L, \mathbb{Z}_2), \quad v + 2L^* \mapsto \langle v, \cdot \rangle,$$
(1.7)

where \langle , \rangle is the inner product on $\mathbb{R} \otimes_{\mathbb{Z}} L$.

Assume that L is unimodular, that is, the dual lattice L^* of L is identical to L. Then V_L is holomorphic ([Do93]). By [DLM00, Theorem 1.2], V_L possesses a unique irreducible θ -twisted module V_L^T , up to isomorphism. It follows from the explicit construction of an irreducible θ twisted module in [FLM88, Chapter 9] that the weights of V_L^T are half-integers. Let $V_L^{T,+}$ be the subspace of V_L^T with integral weights; it is an irreducible module for V_L^+ . Set

$$\tilde{V}_L = V_L^+ \oplus V_L^{T,+}.$$

It is well known that \tilde{V}_L has a unique holomorphic VOA structure by extending its V_L^+ -module structure (cf. [FLM88, DGM96]). Note that the central charge of \tilde{V}_L is n. The VOA \tilde{V}_L is often called the \mathbb{Z}_2 -orbifold¹ of V_L associated to θ . It is known that \tilde{V}_L is simple and of CFT type and that it is a simple current extension of V_L^+ graded by \mathbb{Z}_2 (cf. [ADL05]). Note that if $n \geq 24$ and the minimum norm of L is greater than or equal to 4, then $(\tilde{V}_L)_1 = 0$.

¹In some literature, V_L^+ is called the \mathbb{Z}_2 -orbifold of V_L associated to θ .

Let z denote the automorphism of \tilde{V}_L which acts by 1 and -1 on V_L^+ and $V_L^{T,+}$ respectively. Since V is a simple current extension of V_L^+ , we have $z \in \operatorname{Aut} \tilde{V}_L$. By Proposition 1.4 we obtain the exact sequence

$$1 \to \langle z \rangle \to C_{\operatorname{Aut}\tilde{V}_L}(z) \xrightarrow{\nu} \operatorname{Aut} V_L^+ \to 1.$$
 (1.8)

Notice that Aut $V_L^+ = O(\hat{L})/\langle\theta\rangle$ ([Sh04, Theorem 4.1]) and $\nu^{-1}(\operatorname{Hom}_{\mathbb{Z}}(L,\mathbb{Z}_2)) \cong 2^{1+n}_+$ (cf. [FLM88, §10.4]) for even unimodular lattice L of rank n.

Assume that $L(2) = \emptyset$ and $L = \langle L(4) \rangle_{\mathbb{Z}}$; by the same argument as in [FLM88, Proposition 12.2.5], we can show that V_L^+ is generated by the weight 2 subspaces as a VOA. We also assume that the rank of L is 24 or 32; the lowest weight of $V_L^{T,+}$ is 2. The irreducibility of $V_L^{T,+}$ shows the following:

Lemma 1.7. Let L be an even unimodular lattice of rank 24 or 32. Assume $L(2) = \emptyset$ and $L = \langle L(4) \rangle_{\mathbb{Z}}$. Then the VOA \tilde{V}_L is generated by the weight 2 subspace $(\tilde{V}_L)_2$ as a VOA.

1.5 VOA associated to the Barnes-Wall lattice of rank 16

In this subsection, we review properties of the VOA $U = V_{BW_{16}}^+$.

By (1.1) BW₁₆ has an orthogonal basis of norm 4; U is a framed VOA. By [DGH98], U is rational and C_2 -cofinite. Since BW₁₆ is generated by norm 4 vectors, U is also generated by the weight 2 subspace of U as a VOA. It follows from BW₁₆(2) = \emptyset that $U_1 = 0$.

Let S denote the set R(U) of all isomorphism classes of irreducible U-modules. By the determination of $R(V_L^+)$ in [AD04], we have $|S| = 2^{10}$. Moreover, by the fusion rules of V_L^+ ([ADL05]), any irreducible U-module is a simple current, and S becomes an elementary abelian 2-group of order 2^{10} under the fusion product. We view S as a 10-dimensional vector space over \mathbb{F}_2 .

Let $q_U: S \to \mathbb{F}_2$ be the map defined by setting $q_U([M]) = 1$ if M is $(1/2 + \mathbb{Z})$ -graded, and $q_U([M]) = 0$ if M is \mathbb{Z} -graded; notice that for $[M] \in S$, the lowest weight of M is 0, 1, 3/2 if [M] is zero, non-zero singular, non-singular, respectively. It was shown in [Sh04, Theorem 3.8] that q_U is an Aut U-invariant non-singular quadratic form on S of plus type; there exists a natural group homomorphism from Aut U to the orthogonal group $O(S, q_U)$; its kernel C is an elementary abelian 2-group of order 2^{16} , and its image is the simple subgroup $\Omega^+(10, 2)$ of $O(S, q_U)$ of index 2 ([Sh04, Theorem 4.8]); Aut $U \cong 2^{16} \cdot \Omega^+(10, 2)$ and $C = O_2(Aut U)$. Since C acts trivially on S and $O(\widehat{BW_{16}})/\langle\theta\rangle$ is the stabilizer of $[V_L^-]$ ([Sh04, Propositions 2.8 and 3.10]), C is a subgroup of $O(\widehat{BW_{16}})/\langle\theta\rangle$. By [Sh04, §4.3], the image $\eta(C)$ and kernel of $\eta_{|C}$ in (1.6) are described as follows:

$$C \cap \text{Ker } \eta = \{ f \in \text{Hom}_{\mathbb{Z}}(BW_{16}, \mathbb{Z}_2) \mid f(2BW_{16}^*) = 0 \} \cong 2^8, \tag{1.9}$$

$$\eta(C) = O_2(\operatorname{Aut} BW_{16})/\langle -1 \rangle \cong 2^8.$$
 (1.10)

By the same argument as in [Sh11, Proposition 4.16], we have $\operatorname{Aut} U^{\otimes 2} \cong \operatorname{Aut} U \wr \mathbb{Z}_2$.

Let us consider the action of Aut U on ω_U^{\perp} . The orthogonal complement ω_U^{\perp} of the conformal element ω_U in U_2 is a direct sum of $S^2(\mathbb{C} \otimes_{\mathbb{Z}} BW_{16}) \cap \omega_U^{\perp}$ and the subspace $\sum_{\alpha \in BW_{16}(4)} \mathbb{C}(e^{\alpha} + \theta(e^{\alpha}))$ of the group algebra $\mathbb{C}[BW_{16}]$ as a vector space, where $S^2(\cdot)$ is the symmetric tensor and $BW_{16}(4) = \{v \in BW_{16} \mid \langle v, v \rangle = 4\}$. By

$$\dim S^2(\mathbb{C} \otimes_{\mathbb{Z}} BW_{16}) \cap \omega_U^{\perp} = 16 + \binom{16}{2} - 1 = 135, \ \dim \sum_{\alpha \in BW_{16}(4)} \mathbb{C}(e^{\alpha} + \theta(e^{\alpha})) = \frac{|BW_{16}(4)|}{2} = 2160,$$

we have dim $\omega_U^{\perp} = 2295$. By a similar argument as in [FLM88, Remark 10.4.13], these two subspaces are irreducible for $O(\widehat{BW}_{16})/\langle\theta\rangle$. By [FLM88, Chapter 11] (cf. [Sh04, §2.3]) U has an extra automorphism not preserving them. Hence ω_U^{\perp} is irreducible for Aut U.

Let $(\omega_U^{\perp})^C = \{v \in \omega_U^{\perp} \mid g(v) = v \text{ for all } g \in C\}$. Since C is normal in Aut U, $(\omega_U^{\perp})^C$ is a submodule of ω_U^{\perp} for Aut U. Since the action of C on ω_U^{\perp} is not trivial on ω_U^{\perp} and ω_U^{\perp} is irreducible for Aut U, we have $(\omega_U^{\perp})^C = 0$, that is, the subspace of U_2 fixed by C is $\mathbb{C}\omega_U$.

Ising vectors, vectors in the weight 2 subspace of a VOA generating a simple Virasoro VOA L(1/2,0) of central charge 1/2 as a subVOA, are classified for V_L^+ with $L(2) = \emptyset$ in [Sh12] as follows: any Ising vector has a form $\omega^{\pm}(\alpha)$ or $\omega(E,\varphi)$, where $\alpha \in L(4)$, E is a sublattice of L isomorphic to $\sqrt{2}E_8$ and $\varphi \in \text{Hom}_{\mathbb{Z}}(E,\mathbb{Z}/2\mathbb{Z})$. For an Ising vector e, the automorphism τ_e of a VOA, called the τ -involution, is introduced in [Mi96].

Let I be the set of all Ising vectors in U. Let us show that the normal subgroup $N = \langle \tau_e | e \in I \rangle$ is identical to $O_2(\operatorname{Aut} U)$. Let I_1 (resp. I_2) be the set of all Ising vectors in U of form $\omega^{\pm}(\alpha)$ (resp. $\omega(E, \varphi)$); by the classification in [Sh12] $I = I_1 \cup I_2$. By Aut $U \cong 2^{16} \Omega^+(10, 2)$ and $N \neq \{1\}$, N must be $O_2(\operatorname{Aut} U)$ or Aut U. One can easily see that if $e = \omega^{\pm}(\alpha)$, then $\tau_e \in C \cap \operatorname{Ker} \eta \subset O_2(\operatorname{Aut} U)$ (cf. (1.9)). Hence it suffices to show that Aut U is transitive on I.

The transitivity of $O(BW_{16})/\langle\theta\rangle$ on I_1 follows from that of Aut BW₁₆ on the set of norm 4 vectors in BW₁₆. Let us consider the action of $O(\widehat{BW_{16}})/\langle\theta\rangle$ on I_2 . Let E be a sublattice of BW₁₆ isomorphic to $\sqrt{2}E_8$ and $\varphi \in \operatorname{Hom}_{\mathbb{Z}}(E,\mathbb{Z}_2)$. Since a basis of E is contained in one of BW₁₆, there exists $\tilde{\varphi} \in \operatorname{Hom}_{\mathbb{Z}}(BW_{16},\mathbb{Z}_2)$ such that its restriction to E is identical to φ ; clearly $\tilde{\varphi}(\omega(E,0)) = \omega(E,\varphi)$. By [Gr05, Theorem 12.1], Aut BW₁₆ also acts transitively on the set of sublattices of BW₁₆ isomorphic to $\sqrt{2}E_8$. Thus $O(\widehat{BW_{16}})/\langle\theta\rangle$ acts transitively on I_2 . In addition, U has an extra automorphism σ ([FLM88, Chapter 11]) such that $\sigma(I_1) \cap I_2 \neq \emptyset$. Thus Aut U acts transitively on I.

The following proposition is a summary of this subsection:

Proposition 1.8. (1) The VOA U is simple, rational, C_2 -cofinite, and of CFT type;

- (2) $U_1 = 0$, and U is generated by U_2 as a VOA;
- (3) (S, q_U) is a non-singular 10-dimensional quadratic space of plus type over \mathbb{F}_2 . Moreover, for $[M] \in S$, the lowest weight of M is 0, 1, and 3/2 if [M] is zero, non-zero singular, and non-singular, respectively;
- (4) Aut $U \cong 2^{16} \Omega^+(10,2)$. Moreover, $C = O_2(\text{Aut } U)$ satisfies (1.9) and (1.10);
- (5) C is the subgroup of Aut U acting on S trivially, and S is irreducible for Aut U/C;
- (6) Aut $U^{\otimes 2} \cong$ Aut $U \wr \mathbb{Z}_2$;
- (7) dim $\omega_U^{\perp} = 2295$ and ω_U^{\perp} is irreducible for Aut U;
- (8) $U_2^C = \{ v \in U_2 \mid g(v) = v \text{ for all } g \in O_2(\operatorname{Aut} U) \} = \mathbb{C}\omega_U;$
- (9) C is identical to the subgroup generated by all τ -involutions.

2 Automorphism group of the VOA $\tilde{V}_{BW_{32}}$

Throughout this section, we denote $\tilde{V}_{BW_{32}}$, $V^+_{BW_{16}}$ and Aut $\tilde{V}_{BW_{32}}$ by V, U and G, respectively. Let B denote the orthogonal complement of ω_V in V_2 .

2.1 Centralizer of the involution associated to the \mathbb{Z}_2 -grading

Let z denote the automorphism of V acting by 1 and -1 on $V_{BW_{32}}^+$ and $V_{BW_{32}}^{T,+}$ respectively. In this subsection, we study the centralizer of z in G, and its action on B.

Recall from §1.4 that $\operatorname{Aut} V_{\mathrm{BW}_{32}}^+ = O(\widehat{\mathrm{BW}_{32}})/\langle\theta\rangle$ and $\nu^{-1}(\operatorname{Hom}_{\mathbb{Z}}(\mathrm{BW}_{32}, \mathbb{Z}_2)) \cong 2^{1+32}_+$. By $\operatorname{Aut} \mathrm{BW}_{32} \cong 2^{1+10}_+ . \Omega^+(10, 2), (1.6)$ and (1.8), we obtain the following proposition.

Proposition 2.1. The centralizer $C_G(z)$ in G of z has the shape $2^{1+32}_+.(2^{10}.\Omega^+(10,2)).$

Note 2.2. In [Mi], the shape of $C_G(z)$ was determined by a different method.

By the similar arguments as in [FLM88, Remark 10.4.13], we obtain the following lemma.

Lemma 2.3. Under the action of $C_G(z)$, B breaks into the three irreducible submodules

$$B(1) := S^2(\mathbb{C} \otimes_{\mathbb{Z}} BW_{32}) \cap B, \qquad B(2) := \sum_{\alpha \in BW_{32}(4)} \mathbb{C}(e^\alpha + \theta(e^\alpha)), \qquad B(3) := T$$

of dimensions 527, 73440, 65536, respectively.

2.2 Normalizer of the elementary abelian 2-subgroup of order 2^{10}

Let $U^{\otimes 2}$ be the full subVOA of V based on the embedding of BW₁₆ \oplus BW₁₆ into BW₃₂ in (1.2), where $U \cong V^+_{\text{BW}_{16}}$. By (1.2) and [Sh07, Proposition 4.2], as $U^{\otimes 2}$ -modules,

$$V \cong \bigoplus_{[M] \in S} M^{\otimes 2}, \tag{2.1}$$

where S = R(U) is the set of all isomorphism classes of irreducible U-modules. Since any element in S is a simple current, V is a simple current extension of $U^{\otimes 2}$ graded by $S \cong 2^{10}$. Let S^* denote the dual of the group S. Then S^* acts on V as an automorphism group. In this subsection, we study the structure of the normalizer of S^* in G and its action on B.

Set $C = \{g \in \operatorname{Aut} U \mid g \circ U \cong U \forall U \in S\}$; by Proposition 1.8 (4) and (5), $C = O_2(\operatorname{Aut} U) \cong 2^{16}$. Let σ_U denote the automorphism of $U^{\otimes 2}$ which sends $x \otimes y$ to $y \otimes x$. By Proposition 1.8 (5) and (6), the stabilizer of $\{W \otimes W \mid W \in S\}$ in $\operatorname{Aut} U^{\otimes 2}$ is $(C \wr \langle \sigma_U \rangle) \Delta(\operatorname{Aut} U)$, where $\Delta(\operatorname{Aut} U) = \{(h, h) \mid h \in \operatorname{Aut} U\}$ is the subgroup of $\operatorname{Aut} U \times \operatorname{Aut} U$ commuting with σ_U . Applying Proposition 1.4 to (2.1), we obtain the following exact sequences:

$$1 \to S^* \to N_G(S^*) \xrightarrow{\rho} (C \wr \langle \sigma_U \rangle) \Delta(\operatorname{Aut} U) \to 1,$$
(2.2)

$$1 \to S^* \to C_G(S^*) \xrightarrow{\rho} (C \wr \langle \sigma_U \rangle) \to 1,$$
(2.3)

where ρ is the restriction homomorphism from $N_G(S^*)$ to Aut $U^{\otimes 2}$. Clearly $C\wr\langle\sigma_U\rangle\cap\Delta(\operatorname{Aut} U) = \Delta(C)$.

Proposition 2.4. The normalizer $N_G(S^*)$ and the centralizer $C_G(S^*)$ of S^* in G have the shapes $2^{10} \cdot (2^{16} \wr \mathbb{Z}_2) \cdot \Omega^+(10,2)$ and $2^{10} \cdot (2^{16} \wr \mathbb{Z}_2)$, respectively.

Let us consider the action of $N_G(S^*)$ on B. By Lemma 1.5, $N_G(S^*)$ preserves $(U^{\otimes 2})_2$.

Lemma 2.5. Under the action of $N_G(S^*)$, the subspace $(U^{\otimes 2})_2 \cap B$ breaks into the two irreducible submodules:

$$B(4) := \mathbb{C}(\omega_U \otimes \mathbf{1}_U - \mathbf{1}_U \otimes \omega_U), \qquad B(5) := (\omega_U^{\perp} \otimes \mathbf{1}_U) \oplus (\mathbf{1}_U \otimes \omega_U^{\perp})$$

of dimensions 1 and 4590, respectively.

Proof. Since $N_G(S^*)$ preserves $U^{\otimes 2}$ and Aut $U^{\otimes 2}$ does B(4) by Proposition 1.8 (6), B(4) is an (irreducible) submodule for $N_G(S^*)$. Since B(5) is the orthogonal complement of B(4) in $(U^{\otimes 2})_2 \cap B$, it is also a submodule for $N_G(S^*)$. Hence it suffices to show that B(5) is irreducible. By Proposition 1.8 (7), B(5) is the direct sum of two irreducible modules $\{v \otimes \mathbf{1}_U + \mathbf{1}_U \otimes v \mid v \in \omega_U^{\perp}\}$ and $\{v \otimes \mathbf{1}_U - \mathbf{1}_U \otimes v \mid v \in \omega_U^{\perp}\}$ for $\rho^{-1}(\Delta(\operatorname{Aut} U))$. Considering the action of $\rho^{-1}(C \times 1)$, we have the irreducibility of B(5) for $N_G(S^*)$ by Proposition 1.8 (8).

Proposition 2.6. As a $\langle C_G(z), N_G(S^*) \rangle$ -module, B is irreducible.

Proof. Let B(i) $(1 \le i \le 5)$ be the subspaces of B given in Lemmas 2.3 and 2.5. It suffices to show that B(1), B(2) and B(3) are not preserved by $N_G(S^*)$.

By $(U^{\otimes 2})_2 \cap B \subset (V^+_{BW_{32}})_2 \cap B = B(1) \oplus B(2)$ and the dimension of B(i), we have $B(5) \cap B(1) \neq 0$ and $B(5) \cap B(2) \neq 0$. Hence B(1) and B(2) are not preserved by $N_G(S^*)$ since B(5) is irreducible.

Let M be an irreducible U-module of twisted type with $q_U(M) = 0$; $(M \otimes M)_2 \cap B(3) \neq 0$. By the transitivity of Aut $U/C \cong \Omega^+(10, 2)$ on the set of non-zero singular vectors in S, there exists $g \in N_G(S^*)$ such that $g(M^{\otimes 2})_2 \cap (B(1) \oplus B(2)) \neq 0$. This shows that $N_G(S^*)$ does not preserve B(3).

2.3 Elementary abelian 2-subgroup of order 2²⁷

Based on the exact sequence (2.2), we consider the following subgroup of $N_G(S^*)$:

$$A = \rho^{-1}(\langle \Delta(C), \sigma_U \rangle). \tag{2.4}$$

It follows from $S^* = \text{Ker } \rho \cong 2^{10}$, $C \cong 2^{16}$ and $|\sigma_U| = 2$ that $|A| = 2^{27}$. In this subsection, we prove that A is elementary abelian and is normal in both $N_G(S^*)$ and $C_G(z)$.

Let σ_V be an element in A with $\rho(\sigma_V) = \sigma_U$. Then $\sigma_V^2 \in S^*$; the order of σ_V is 2 or 4. Later, we will show in Corollary 2.10 that the order of σ_V is 2.

Lemma 2.7. (1) The subgroup $\rho^{-1}(\Delta(C))$ of A is an elementary abelian 2-group;

- (2) A is abelian;
- (3) A is normal in $N_G(S^*)$.

Proof. Since S^* acts on each $U^{\otimes 2}$ -submodule of V by a scalar and $A \subset C_G(S^*)$, we have $S^* \subset Z(A)$.

Let us show (1) and (2) by describing an explicit section of $\rho_{|A}$ in terms of τ -involutions. By Proposition 1.8 (9), C is generated by τ -involutions associate to Ising vectors in U. For $(c,1) \in C \times 1 \subset \rho(N_G(S^*))$, there exists an Ising vector $e \in U \otimes \mathbf{1}_U \subset V$ such that $\tau_{e|U^{\otimes 2}} = (c,1)$. Set $f = \sigma_V(e)$; $f \in \mathbf{1}_U \otimes U$ and it is orthogonal to e in V_2 , which shows that τ_f commutes with τ_e ; $\tau_e \tau_f$ is an involution in G and $\rho(\tau_e \tau_f) = (c,c)$; $\rho^{-1}(c,c)$ is elementary abelian. For $(c',1) \in C \times 1$, we obtain $\tau_{e'}$ and $\tau_{f'}$ by the same procedure. Since $[\tau_e, \tau_{f'}] = [\tau_{e'}, \tau_f] = 1$ and $[\tau_e, \tau_{e'}] = [\tau_f, \tau_{f'}] \in S^*$, we have $[\tau_e \tau_f, \tau_{e'} \tau_{f'}] = 1$; $\rho^{-1}(\langle (c,c), (c',c') \rangle)$ is elementary abelian. Moreover, these involutions generate an elementary abelian 2-subgroup whose image under ρ is $\Delta(C)$. Hence (1) holds. It follows from $\sigma_V(\tau_e \tau_f)\sigma_V^{-1} = \tau_f \tau_e = \tau_e \tau_f$ that $\sigma_V \in Z(A)$; (2) holds.

Let us show (3). Since A contains S^* , it suffices to show that $\rho(A) = \Delta(C) \times \langle \sigma_U \rangle$ is normal in $\rho(N_G(S^*)) = (C \wr \langle \sigma_U \rangle) \Delta(\operatorname{Aut} U)$ (cf. (2.2) and (2.4)). Clearly, $\rho(A)$ is normalized by $\Delta(\operatorname{Aut} U)$, and $\Delta(C)$ commutes with $C \wr \langle \sigma_U \rangle$. Moreover, for $(c, c') \in C \times C$, $(c, c')\sigma_U(c, c')^{-1} = (cc', cc')\sigma_U \in \rho(A)$. Hence $\rho(A)$ is normal in $\rho(N_G(S^*))$. Since A contains z, it is a subgroup of $C_G(z)$. In order to show that A is normal in $C_G(z)$, we consider the structure of A in $C_G(z)$. Recall that $\nu : C_G(z) \to \operatorname{Aut} V^+_{\mathrm{BW}_{32}}$ and $\eta : \operatorname{Aut} V^+_{\mathrm{BW}_{32}} \to \operatorname{Aut} \mathrm{BW}_{32}/\langle -1 \rangle$ are defined in (1.8) and (1.6), respectively, and that Ker $\nu = \langle z \rangle$, Ker $\eta = \operatorname{Hom}_{\mathbb{Z}}(\mathrm{BW}_{32}, \mathbb{Z}_2) \cong 2^{32}$, and Ker $\eta \circ \nu \cong 2^{1+32}_+$.

Lemma 2.8. (1) $\nu(S^*) \cap \text{Ker } \eta = \{ f \in \text{Hom}_{\mathbb{Z}}(BW_{32}, \mathbb{Z}_2) \mid f(BW_{16} \oplus BW_{16}) = 0 \} \cong 2^8.$

- (2) $\nu(A) \cap \text{Ker } \eta = \{ f \in \text{Hom}_{\mathbb{Z}}(BW_{32}, \mathbb{Z}_2) \mid f(BW_{32}[1]) = 0 \} \cong 2^{16}, \text{ where } BW_{32}[1] \text{ is the sublattice of } BW_{32} \text{ given in } (1.3).$
- (3) $\eta \circ \nu(A) = O_2(\operatorname{Aut} BW_{32}/\langle -1 \rangle) \cong 2^{10}.$
- (4) $\eta \circ \nu(S^*) = \langle (-1,1), -1 \rangle / \langle -1 \rangle$, where (-1,1) is the element in Aut BW₃₂ which sends $(x, y) \mapsto (-x, y)$ in the description (1.2) of BW₃₂.

Proof. (1) follows from $V^{S^*} = U^{\otimes 2}$ and Lemma 1.6 (2).

Let *L* be the sublattice of BW₃₂ satisfying $\nu(A) \cap \text{Ker } \eta = \{f \in \text{Hom}_{\mathbb{Z}}(\text{BW}_{32}, \mathbb{Z}_2) \mid f(L) = 0\}$ and $2\text{BW}_{32} \subset L$. Set $H = \nu \circ \rho^{-1}(\Delta(C \cap \text{Hom}_{\mathbb{Z}}(\text{BW}_{16}, \mathbb{Z}_2))) \cap \text{Ker } \eta$. It follows from (2.4) and $\rho^{-1}(\Delta(C))\sigma_V \cap \text{Ker } \eta = \emptyset$ that $\nu(A) \cap \text{Ker } \eta = \langle H, \nu(S^*) \cap \text{Ker } \eta \rangle$. By the definition of *H* and (1.9), for $f \in \text{Hom}_{\mathbb{Z}}(\text{BW}_{32}, \mathbb{Z}_2)$, $f \in H$ if and only if f(v) = 0 for all $v \in 2\text{BW}_{16}^* \oplus 2\text{BW}_{16}^* +$ $\{(v, v) \mid v \in \text{BW}_{16}^*\}$. By (1)

 $L = (BW_{16} \oplus BW_{16}) \cap (2BW_{16}^* \oplus 2BW_{16}^* + \{(v, v) \mid v \in BW_{16}^*\}) = BW_{32}[1].$

Hence by (1.3), we obtain $L = BW_{32}[1]$, which proves (2).

By (2) and $|\nu(A)| = 2^{26}$, we have $|\eta \circ \nu(A)| = 2^{10}$. By Lemma 2.7 (2), $\nu(A)$ is abelian, and hence it acts trivially on $\nu(A) \cap \text{Ker } \eta$. By (1.7) and (2), $\eta \circ \nu(A)$ acts trivially on BW₃₂[1]/2BW₃₂. By Lemma 1.1, $\eta \circ \nu(A)$ is a subgroup of $O_2(\text{Aut BW}_{32}/\langle -1 \rangle) \cong 2^{10}$. Comparing the orders of the groups, we have (3).

By (1), the order of $\eta \circ \nu(S^*)$ is 2. On the other hand, $U^{\otimes 2}$ is the subspace of $V^+_{BW_{16} \oplus BW_{16}}$ fixed by some lift of (-1, 1). Hence we have (4).

Proposition 2.9. The subgroup A is normal in $C_G(z)$.

Proof. Set $Q = \text{Ker } (\eta \circ \nu) \cap A$. By Lemmas 1.1 and 2.8 (2) and (1.7), Q is normal in $C_G(z)$. Let us show that $C_G(Q) = A$. Clearly, $A \subset C_G(Q)$. Since Q is maximal abelian in Ker $(\eta \circ \nu)$, we have Ker $(\eta \circ \nu) \cap C_G(Q) = Q = \text{Ker } (\eta \circ \nu) \cap A$.

Since $C_G(Q)$ acts trivially on $\nu(A) \cap \text{Ker } \eta$, so does $\eta \circ \nu(C_G(Q))$ on $BW_{32}[1]/2BW_{32}$ by Lemma 2.8 (2) and (1.7). By Lemmas 1.1 and 2.8 (3), $\eta \circ \nu(C_G(Q)) \subset \eta \circ \nu(A)$. Hence $C_G(Q) \subset A$. Thus $C_G(Q) = A$.

Corollary 2.10. (1) The normalizer $N_G(A)$ contains both $N_G(S^*)$ and $C_G(z)$.

- (2) Under the action of $N_G(A)$, B is irreducible.
- (3) The group A is an elementary abelian 2-group of order 2^{27} .

Proof. (1) follows from Lemma 2.7 (3) and Proposition 2.9, and (2) does from (1) and Proposition 2.6. Since $\eta \circ \nu(C_G(z))/O_2(\eta \circ \nu(C_G(z))) \cong \Omega^+(10, 2)$ acts irreducibility on $\eta \circ \nu(A) \cong 2^{10}$ and $\eta \circ \nu(\sigma_V) \neq 1$, we have $\sigma_V^2 = 1$. Hence (3) holds by Lemma 2.7 (1).

Proposition 2.11. The subgroup A is an irreducible $N_G(A)/A$ -module over \mathbb{F}_2 .

Proof. Let A_0 be a non-zero submodule of A for $N_G(A)/A$. By Corollary 2.10 (1), both $C_G(z)$ and $N_G(S^*)$ preserve A_0 .

If $\eta \circ \nu(A_0) \neq 0$, then by the action of $C_G(z)$ and Lemma 2.8 (3), we have $\eta \circ \nu(A_0) = \eta \circ \nu(A)$. By the action of $N_G(S^*)$, along with Lemma 2.8 (1) and (4), (Ker $\eta) \cap \nu(A_0) \neq 0$. By Lemmas 1.1 and 2.8 (2), $\nu(A) = \nu(A_0)$. By Ker $\nu = \langle z \rangle \subset S^*$, we have $A_0 = A$.

If $\eta \circ \nu(A_0) = 0$, then by the same argument above, we have $A_0 \cap (\text{Ker } \eta \circ \nu) = A \cap (\text{Ker } \eta \circ \nu)$. By the action of $N_G(S^*)$, along with Lemma 2.8 (3) and (4), we have $\eta \circ \nu(A_0) = \eta \circ \nu(A)$, and $A_0 = A$.

2.4 Graph on eigenspaces in the Griess algebra

In this subsection, we define a graph on the eigenspaces of A in $B = \omega_V^{\perp}$, which will be used to determine the group structure of $N_G(A)$.

Let $B = \bigoplus_{\chi \in A^*} B_{\chi}$ be the eigenspace decomposition of A in B.

Definition 2.12. For non-zero distinct eigenspaces B_1, B_2 , we say that B_1 is adjacent to B_2 if there exists $x_i \in B_i$ (i = 1, 2) such that $x_1 \cdot x_2 \neq 0$, where \cdot is the product in B.

Let Γ denote the graph on the vertex set $V\Gamma = \{B_{\chi} \mid B_{\chi} \neq 0\}$ defined by Definition 2.12. Since the product in the Griess algebra is commutative, Γ is undirected. For $g \in N_G(A)$, $g(B_{\chi}) = B_{\chi^g}$, where $\chi^g(a) = \chi(g^{-1}ag)$ for $a \in A$; $N_G(A)$ acts on Γ as an automorphism group.

Lemma 2.13. Let I be a subset of $V\Gamma$ and let H be a subgroup of $N_G(A)$ preserving I. Assume that $\bigoplus_{B_{\chi} \in I} B_{\chi}$ is irreducible for H. Then the following hold:

- (1) H is transitive on I;
- (2) For $B_1, B_2 \in I$, dim $B_1 = \dim B_2$.

Proof. Let I_0 be an orbit in I under the action of H. Then $\bigoplus_{B_{\chi} \in I_0} B_{\chi}$ is a submodule of B for H. By the irreducibility, we have $I_0 = I$, and (1) holds.

For $g \in H$ and $B_{\chi} \in V\Gamma$, dim $B_{\chi} = \dim B_{\chi^g}$. Hence (2) follows from (1).

2.5 Graph structure and the stabilizer of a vertex

In this subsection, we study the structure of the graph Γ defined in the previous subsection and the action of $N_G(A)/A$ on it.

Let μ denote the irreducible character of A defined by

$$\mu(a) = \begin{cases} 1 & \text{if } a \in \rho^{-1}(\Delta(C)), \\ -1 & \text{if } a \in \rho^{-1}(\Delta(C))\sigma_V. \end{cases}$$
(2.5)

Let us describe the eigenspace B_{μ} and its stabilizer in $N_G(A)$.

Lemma 2.14. The eigenspace B_{μ} is spanned by $\omega_U \otimes \mathbf{1}_U - \mathbf{1}_U \otimes \omega_U$, and dim $B_{\mu} = 1$.

Proof. Since $\rho^{-1}(\Delta(C))$ acts on $U^{\otimes 2}$ as a subgroup of Aut $U \times$ Aut U, it fixes both $\omega_U \otimes \mathbf{1}_U$ and $\mathbf{1}_U \otimes \omega_U$. Moreover, $\sigma_V(\omega_U \otimes \mathbf{1}_U - \mathbf{1}_U \otimes \omega_U) = -(\omega_U \otimes \mathbf{1}_U - \mathbf{1}_U \otimes \omega_U)$. Hence $\omega_U \otimes \mathbf{1}_U - \mathbf{1}_U \otimes \omega_U$ is an eigenvector of A with character μ .

By $S^* \subset \rho^{-1}(\Delta(C))$, $B_{\mu} \subset U^{\otimes 2} = \{u \in V \mid g(u) = u \; \forall g \in S^*\}$. It follows from $U_0 = \mathbb{C}\mathbf{1}_U$ and $U_1 = 0$ that $(U^{\otimes 2})_2 = U_2 \otimes \mathbf{1}_U \oplus \mathbf{1}_U \otimes U_2$. By Proposition 1.8 (8), $(U^{\otimes 2})^{\Delta(C)} = \mathbb{C}(\omega_U \otimes \mathbf{1}_U) \oplus \mathbb{C}(\mathbf{1}_U \otimes \omega_U)$. Considering the action of σ_V , we obtain this lemma. **Lemma 2.15.** Let $B_{\chi} \in V\Gamma$. Then the following hold:

- (1) B_{χ} is contained in some $U^{\otimes 2}$ -submodule of V, that is, $B_{\chi} \subset (M^{\otimes 2})_2$ for some $[M] \in S$;
- (2) Assume $\chi \neq \mu$. Then B_{χ} is adjacent to B_{μ} if and only if $B_{\chi} \subset (U^{\otimes 2})_2 \cap B$.

Proof. By $S^* \subset A$, B_{χ} is an eigenspace of S^* ; (1) holds.

Let us show (2). First, we assume that $B_{\chi} \subset (U^{\otimes 2})_2 \cap B$. Let $x \in B_{\chi}$ be a non-zero vector. Then $x = v \otimes \mathbf{1}_U + \mathbf{1}_U \otimes w$ for some $v, w \in U_2$ by $(U^{\otimes 2})_2 = U_2 \otimes \mathbf{1}_U \oplus \mathbf{1}_U \otimes U_2$. By

$$(\omega_U \otimes \mathbf{1}_U - \mathbf{1}_U \otimes \omega_U) \cdot x = 2(v \otimes \mathbf{1}_U - \mathbf{1}_U \otimes w) \neq 0,$$

 B_{μ} and B_{χ} are adjacent by Lemma 2.14.

Next, we assume that $B_{\chi} \subset (M^{\otimes 2})_2 \cap B$ and $M \neq U$. By Proposition 1.8 (3) and $(M^{\otimes 2})_2 \neq 0$, the lowest weight of M is 1. Hence any vector in B_{χ} has a form $\sum v_i \otimes w_i, v_i, w_i \in M_1$. By

$$(\omega_U \otimes \mathbf{1}_U - \mathbf{1}_U \otimes \omega_U) \cdot \sum v_i \otimes w_i = 0,$$

 B_{μ} and B_{χ} are not adjacent by Lemma 2.14.

Proposition 2.16. The stabilizer of B_{μ} in $N_G(A)$ is equal to $N_G(S^*)$.

Proof. By Lemmas 2.5 and 2.14, $N_G(S^*)$ is a subgroup of the stabilizer of B_{μ} in $N_G(A)$. Let $g \in N_G(A)$ with $g(B_{\mu}) = B_{\mu}$. Let I be a subset of all vertices in $V\Gamma$ adjacent to B_{μ} . Then g preserves $\bigoplus_{B_{\chi} \in I} B_{\chi}$ by the definition of Γ . By Lemma 2.15 (2), $B_{\mu} \oplus \bigoplus_{B_{\chi} \in I} B_{\chi} = (U^{\otimes 2})_2 \cap B$. Hence $g((U^{\otimes 2})_2) = (U^{\otimes 2})_2$, and $g(U^{\otimes 2}) = U^{\otimes 2}$ by Proposition 1.8 (2). It follows from Lemma 1.5 that $g \in N_G(S^*)$.

Proposition 2.17. The subgroup of $N_G(A)$ acting trivially on $V\Gamma$ is $C_G(A)$.

Proof. Clearly, $C_G(A)$ acts trivially on $V\Gamma$. Let g be an element in $N_G(A)$ acting on $V\Gamma$ trivially. Then $\chi^g = \chi$ for all $\chi \in A^*$ with $B_{\chi} \neq 0$. It follows from Lemma 1.7 that A acts faithfully on B. Hence, $\langle \chi \in \operatorname{Irr}(A) \mid B_{\chi} \neq 0 \rangle = A^*$, and g commutes with A.

Proposition 2.18. The subgroup A of G is self-centralizing, that is, $C_G(A) = A$.

Proof. Since A is abelian, $A \subset C_G(A)$. It follows from $S^* \subset A$ that $C_G(A) \subset C_G(S^*)$. It follows from $\sigma_V \in A$ and (2.3) that $\rho(C_G(A)) \subset \Delta(C) \times \langle \sigma_U \rangle$. Hence $C_G(A) \subset A$ by (2.4). \Box

Recall that dim B = 139503 and dim U = 2296 (cf. Lemmas 2.3 and 2.5). Combining Lemmas 2.13, 2.14, and 2.15 (2), Propositions 2.16, 2.17 and 2.18, we obtain the following:

Theorem 2.19. The graph Γ has 139503 vertices, and its valency is 4590. Moreover, $N_G(A)/A$ acts transitively on the vertex set, and the stabilizer of a vertex is isomorphic to $N_G(S^*)/A$.

2.6 Rank 3 property of the graph for the normalizer

By Proposition 2.16, the stabilizer of B_{μ} is $N_G(S^*)/A$. In this subsection, we show that Γ is rank 3 for $N_G(A)/A$, that is, there are three orbits in $V\Gamma$ under the action of $N_G(S^*)/A$.

Proposition 2.20. (1) The subgroup $N_G(S^*)/A$ is transitive on the subset of $V\Gamma$ consisting of all vertices adjacent to B_{μ} .

(2) The subgroup $N_G(S^*)/A$ is transitive on the subset of $V\Gamma \setminus \{B_\mu\}$ consisting of all vertices not adjacent to B_μ .

Proof. (1) Let I be the subset of $V\Gamma$ consisting of vertices adjacent to B_{μ} . Then $\bigoplus_{B_{\chi} \in I} B_{\chi}$ is a module for $N_G(S^*)$. By Lemmas 2.15, $\bigoplus_{B_{\chi} \in I} B_{\chi} = (U^{\otimes 2})_2 \cap B$, and by Lemma 2.5 it is irreducible for $N_G(S^*)$. Hence the transitivity of $N_G(S^*)$ on I follows from Lemma 2.13 (1).

(2) Let $B_{\chi} \in V\Gamma \setminus \{B_{\mu}\}$ not adjacent to B_{μ} . Then by Lemma 2.15 (1), $B_{\chi} \in (M^{\otimes 2})_2$ for some $[M] \in S \setminus \{[U]\}$. By (2.2) and (2.3), $N_G(S^*)/C_G(S^*) \cong \Omega^+(10, 2)$, and it is transitive on the set of non-zero singular vectors in $S = R(U) \cong \mathbb{F}_2^{10}$. By Proposition 1.8 (3), for $[M] \in S \setminus \{[U]\}$, $(M^{\otimes 2})_2 \neq 0$ if and only if [M] is non-zero singular.

Let us show that $C_G(S^*)$ acts transitively on the set of all vertices in $(M^{\otimes 2})_2$ for $M = V_{\text{BW}_{16}}^-$, which proves (2). Since BW₁₆ has no roots, $M_1 \cong \mathbb{C} \otimes_{\mathbb{Z}} \text{BW}_{16}$. Hence $(M^{\otimes 2})_2 = M_1^{\otimes 2} \cong$ $(\mathbb{C} \otimes_{\mathbb{Z}} \text{BW}_{16})^{\otimes 2}$ as vector spaces. By Lemma 1.6 (1) and the definition of z, Ker $\eta \circ \nu$ acts trivially on $(M^{\otimes 2})_2$ (see (1.6) and (1.8) for ν and η). By $C_G(S^*) \subset C_G(z)$, $C_G(S^*)$ acts on $(M^{\otimes 2})_2$ as $\eta \circ \nu(C_G(S^*))$. By (2.3), $\eta \circ \nu(\rho^{-1}(C \times C)) \subset \eta \circ \nu(C_G(S^*))$. By (1.10) and Lemma 2.8 (4), $\eta \circ \nu(\rho^{-1}(C \times C)) = O_2(\text{Aut BW}_{16}) * O_2(\text{Aut BW}_{16}) \cong 2^{1+8}_+$. Recall that M_1 is a 2⁴dimensional irreducible module for $O_2(\text{Aut BW}_{16}) \cong 2^{1+8}_+$. By Lemma 1.2, $M_1^{\otimes 2}$ is irreducible for $\eta \circ \nu(\rho^{-1}(C \times C))$. Hence by Lemma 2.13 (1), $C_G(S^*)$ acts transitively on the vertices in $(M^{\otimes 2})_2$.

Proposition 2.20 also shows that the action of $N_G(A)/A$ on $V\Gamma$ is primitive. Hence $N_G(S^*)/A$ is maximal in $N_G(A)/A$.

Theorem 2.21. The graph Γ is rank 3 for $N_G(A)/A$. Moreover, $N_G(A)$ is generated by $N_G(S^*)$ and $C_G(z)$.

2.7 Shape of the normalizer

In this subsection, we determine the shape of $N_G(A)$ by using the classification of permutation groups of rank 3.

As a corollary of Theorem 2.21, we have the following.

Corollary 2.22. The group $N_G(A)/A$ is a primitive permutation group of rank 3 of degree $139503 = 3 \cdot 7^2 \cdot 13 \cdot 73$.

Now, we recall that any primitive permutation group Q of rank 3 of finite degree n satisfies one of the following (see [Li87] and references therein):

- (i) Q is a subgroup of $E_0 \wr \mathbb{Z}_2$ and Q contains $E \times E$ as a normal subgroup, where E_0 is a 2-transitive group of degree n_0 and the socle E of E_0 is a simple group, and $n = n_0^2$;
- (ii) The socle of Q, the product of all the minimal normal subgroups of Q, is (non-abelian) simple;
- (iii) Q is an affine group, that is, the socle of Q is a vector space W, where $W \cong \mathbb{Z}_p^d$ for some prime p and $n = p^d$.

By the degree, $N_G(A)/A$ satisfies (ii). Since the stabilizer in $N_G(A)/A$ of a vertex in Γ has a shape $2^{16}: \Omega^+(10, 2)$ and its index is 139503, the order of $N_G(A)/A$ is $2^{36}\cdot 3^6\cdot 5^2\cdot 7^3\cdot 13\cdot 17\cdot 31\cdot 73$; the socle of $N_G(A)/A$ is isomorphic to neither an alternating group nor a sporadic group (cf. [CCNPW85]); the primitive permutation groups of rank 3 under (ii) were determined in [KL82] when the socle is a classical group, in [LS86] when the socle is a non-classical simple group of Lie type; $N_G(A)/A$ must be isomorphic to $E_6(2)$.

Proposition 2.23. The normalizer $N_G(A)$ has the shape $2^{27} \cdot E_6(2)$.

By Theorem 2.21 and Proposition 2.23 (cf. [LS86]), we obtain the following corollary:

Corollary 2.24. The graph Γ is isomorphic to the rank 3 graph on $E_6(2)/(2^{16}: \Omega^+(10, 2))$ of valency 4590, and its automorphism group is isomorphic to $E_6(2)$.

2.8 Full automorphism group

In this subsection, we determine the full automorphism group G of the VOA V.

First, we show the finiteness of G. The following is a modification of [Ti84, Theorem (iv)].

Proposition 2.25. Let \mathcal{H} be an algebraic group. Assume that \mathcal{H} contains commuting involutions z_1, z_2 such that $C_{\mathcal{H}}(z_1), C_{\mathcal{H}}(z_2)$ and $C_{\mathcal{H}}(z_1z_2)$ are finite. Then \mathcal{H} is finite.

Proof. Let \mathcal{H}^0 be the connected component of the identity in \mathcal{H} . Clearly, \mathcal{H}^0 is normal in \mathcal{H} . Let a be an involution in \mathcal{H} such that $C_{\mathcal{H}}(a)$ is finite. Then the involution $g \mapsto a^{-1}ga$ in Aut \mathcal{H}^0 has only finitely many fixed points. Let $L(\mathcal{H}^0)$ be the Lie algebra of \mathcal{H}^0 . Then the canonical group homomorphism ϕ from Aut (\mathcal{H}^0) to Aut $(L(\mathcal{H}^0))$ is injective.

We now suppose that $\mathcal{H}^0 \neq \{1\}$. Since $C_{\mathcal{H}}(a)$ is finite, $\phi(a)$ is an involution in Aut $L(\mathcal{H}^0)$ whose only fixed point is 0. This implies that $\phi(a) = -1$. In particular, $\phi(z_1) = \phi(z_2) = \phi(z_1 z_2) = -1$, which contradicts $\phi(z_1)\phi(z_2) = \phi(z_1 z_2)$. Hence $\mathcal{H}^0 = \{1\}$, and \mathcal{H} is discrete. Therefore \mathcal{H} is finite since it is an algebraic group.

By Proposition 2.1 $C_G(z)$ is finite. We now check that G satisfies the hypothesis of the proposition above.

Lemma 2.26. The abelian subgroup S^* of G contains commuting involutions z_1, z_2 such that z_1, z_2 and $z_1 z_2$ are conjugate to z. In particular, $C_G(z_1)$, $C_G(z_2)$ and $C_G(z_1 z_2)$ are finite.

Proof. By the natural correspondence between S^* and S with respect to the symplectic form, z corresponds to the non-zero singular vector $[V_{BW_{16}}^-] \in S$. The group $N_G(S^*)$ acts on S^* as a natural module for $\Omega^+(10, 2)$ by conjugation, and it is transitive on the set of non-zero singular vectors in S. Hence all involutions in S^* corresponding to non-zero singular vectors in S are conjugate to z. Since a 10-dimensional non-singular quadratic space over \mathbb{F}_2 contains a 2-dimensional totally singular subspace, we obtain this lemma.

It was shown in [DG02, Theorem 2.4] that the automorphism group of a finitely generated VOA is an algebraic group. By Lemma 1.7, V is finitely generated, and hence G is an algebraic group. Thus Proposition 2.25 and Lemma 2.26 imply the following:

Proposition 2.27. The automorphism group G of V is finite.

Next, we will show that A is normal in G.

Lemma 2.28. Let P be a Sylow 2-subgroup of $C_G(z)$. Then $Z(P) = \langle z \rangle$, and P is also a Sylow 2-subgroup of G.

Proof. It suffices to consider a specified Sylow 2-subgroup $P \cong 2^{1+32}_+ (Q_1.Q_2)$ of $C_G(z)$, where $Q_1 = O_2(\operatorname{Aut} BW_{32})$ and Q_2 is a Sylow 2-subgroup of Aut $BW_{32}/O_2(\operatorname{Aut} BW_{32})$. Then Q_2 and Q_1 act faithfully on Q_1 and $2^{1+32}/\langle z \rangle \cong BW_{32}/2BW_{32}$, respectively. Hence $Z(P) \subset$ $Z(2^{1+32}_+) = \langle z \rangle$, and $Z(P) = \langle z \rangle$. Let P_0 be a Sylow 2-subgroup of G containing P. Then $1 \neq Z(P_0) \subset Z(P) = \langle z \rangle$, and hence $Z(P_0) = Z(P)$. Thus $P_0 \subset C_G(z)$, and $P_0 = P$.

Proposition 2.29. Let F be a minimal normal subgroup of G. Then F is abelian, and F = A.

Proof. Set $J = F \cap N_G(A)$. By Proposition 2.11 A is irreducible as an $N_G(A)$ -module, and by Proposition 2.23 $N_G(A)/A$ is simple. Hence J must be $\{1\}$, A or $N_G(A)$.

First, we prove $A \subset F$. Suppose $A \not\subset F$. Then $J = \{1\}$ and $C_G(z) \cap F = \{1\}$. Let $x, y \in F$ with $x^z x^{-1} = y^z y^{-1}$. Then $(y^{-1}x)^z = (y^{-1})^z x^z = y^{-1}x$. By $C_G(z) \cap F = \{1\}$, we have $y^{-1}x = 1$, and y = x. Moreover, if $x \neq 1$ then $x^z x^{-1} \neq 1$. Hence $F = \{x^z x^{-1} \mid x \in F\}$, and $f^z = f^{-1}$ for all $f \in F$. By Lemma 2.26, there exist involutions $z_1, z_2 \in S^*$ such that z_1, z_2 and z_1z_2 are conjugate to z. The argument above shows that z_1, z_2, z_1z_2 induce the same automorphism of F of order 2, which is a contradiction.

Next, we suppose that F is not abelian. Then F is a direct product of isomorphic (nonabelian) simple groups, that is, $F = \prod_{i=1}^{l} F_i$, $F_i \cong F_j$ and F_i is simple for all $i, j; A = \prod_{i=1}^{l} (A \cap F_i)$ by $N_G(A) \cap F = A$. By $C_N(A) = A$, $A \cap F_i \neq \{1\}$ for all i. Hence $N_G(A)/A$ acts on $\{A \cap F_i\}$ as a permutation group of degree $d \leq 27$. Since $N_G(A)/A$ is simple and 73 | $|E_6(2)|$, d must be 1; F is simple.

We consider the following cases: (1) J = A; (2) $J = N_G(A)$.

(1) Suppose J = A. By Lemma 2.28, there exists a Sylow 2-subgroup P of G such that $A \subset P \subset C_G(z)$. Then $P \cap F = A$ is a Sylow 2-subgroup of F. By J = A, we have $N_F(A) = C_F(A) = A$. By [Go80, Theorem 4.3 in Chapter 7], F has a normal complement of A, which contradicts the simplicity of F.

(2) Suppose $J = N_G(A)$. This implies $N_G(A) \subset F$. It follows from $C_G(z) \subset N_G(A)$ that $C_F(z) = C_G(z)$. Let P be a Sylow 2-subgroup of $C_G(z)$. Then by Lemma 2.28 P is also a Sylow 2-subgroup of F. By Proposition 2.1, $|P| = 2^{63}$. By $N_G(A) \subset F$ and Proposition 2.23, |F| is divisible by 73. We now refer the classification of finite simple groups. If F is an alternating group, then by $|P| = 2^{63}$ it isomorphic to A_{66} or A_{67} , which contradicts that |F| is divisible by 73. By [CCNPW85] and $|P| = 2^{63}$, F is not a sporadic simple group. According to [AS76] and [GLS98, Section 4.5], $C_G(z) \cong 2^{1+32} \cdot 2^{10} \cdot \Omega^+(10, 2)$ can not be the centralizer of an involution in any finite simple group of Lie type. Thus (2) is impossible.

Therefore F is abelian and $A \subset F$. By Proposition 2.18, $A \subset F \subset C_G(A) = A$, and hence F = A.

Combining Propositions 2.23 and 2.29, we obtain the following main theorem:

Theorem 2.30. The automorphism group of the \mathbb{Z}_2 -orbifold VOA $\tilde{V}_{BW_{32}}$ has the shape $2^{27} \cdot E_6(2)$.

Acknowledgments. The author thanks Professor Robert L. Griess Jr. for helpful comments, Professor Masahiko Miyamoto for sending the unpublished manuscript [Mi], and Professor Ching Hung Lam for carefully reading an early version of this article. He also thanks the referee for many valuable suggestions.

References

- [AD04] T. Abe and C. Dong, Classification of irreducible modules for the vertex operator algebra V_L^+ : general case. J. Algebra **273** (2004), 657–685
- [ADL05] T. Abe, C. Dong and H. Li, Fusion rules for the vertex operator algebras $M(1)^+$ and V_L^+ , Comm. Math. Phys. **253** (2005), 171–219.
- [AS76] M. Aschbacher and G.M. Seitz, Involutions in Chevalley groups over fields of even order, Nagoya Math. J. 63 (1976), 1–91.
- [BW59] E.S. Barnes and G.E. Wall, Some extreme forms defined in terms of Abelian groups, J. Austral. Math. Soc. 1 (1959), 47–63

- [Bo86] R.E. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Nat'l. Acad. Sci. U.S.A.* 83 (1986), 3068–3071.
- [Bo92] R.E. Borcherds, Monstrous moonshine and monstrous Lie superalgebras, *Invent. Math.* **109** (1992), 405–444.
- [CCNPW85] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker and R.A. Wilson, Atlas of finite groups, Oxford University Press, Oxford, 1985.
- [CS99] J.H. Conway and N.J.A. Sloane, Sphere packings, lattices and groups, 3rd Edition, Springer, New York, 1999.
- [DGM96] L. Dolan, P. Goddard and P. Montague, Conformal field theories, representations and lattice constructions, Comm. Math. Phys. 179 (1996), 61–120.
- [Do93] C. Dong, Vertex algebras associated with even lattices, J. Algebra 161 (1993), 245–265.
- [DG98] C. Dong and R.L. Griess, Rank one lattice type vertex operator algebras and their automorphism groups, J. Algebra 208 (1998), 262–275.
- [DG02] C. Dong and R.L. Griess, Automorphism groups and derivation algebras of finitely generated vertex operator algebras. *Michigan Math. J.* **50** (2002), 227–239.
- [DG05] C. Dong and R.L. Griess, The rank two lattice type vertex operator algebras V_L^+ and their automorphism groups, *Michigan Math. J.* **53** (2005), 691–715.
- [DGH98] C. Dong, R.L. Griess, and G. Höhn, Framed vertex operator algebras, codes and Moonshine module, Comm. Math. Phys. 193 (1998), 407–448.
- [DLM00] C. Dong, H. Li, and G. Mason, Modular-invariance of trace functions in orbifold theory and generalized Moonshine, *Comm. Math. Phys.* 214 (2000), 1–56.
- [DM04] C. Dong and G. Mason, Rational vertex operator algebras and the effective central charge, Int. Math. Res. Not. (2004), 2989–3008.
- [DN99] C. Dong and K. Nagatomo, Automorphism groups and Twisted modules for lattice Vertex operator algebras, *Comtemp. Math.* 248 (1999), 117–133
- [FHL93] I. Frenkel, Y. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, Mem. Amer. Math. Soc. 104 1993.
- [FLM88] I. Frenkel, J. Lepowsky and A. Meurman, Vertex operator algebras and the Monster, Pure and Appl. Math., Vol.134, Academic Press, Boston, 1988.
- [Go80] D. Gorenstein, Finite groups, Chelsea Publishing Co., New York, 1980.
- [GLS98] D. Gorenstein, R. Lyons and R. Solomon, The classification of the finite simple groups, Number 3. Mathematical Surveys and Monographs, 40.3., American Mathematical Society, Providence, 1998.
- [Gr82] R.L. Griess, The friendly giant. Invent. Math. 69 (1982) 1–102.
- [Gr98] R.L. Griess, A vertex operator algebra related to E_8 with automorphism group O⁺(10, 2), Ohio State Univ. Math. Res. Inst. Publ. 7 (1998), 43–58.
- [Gr05] R.L. Griess, Pieces of 2^d : existence and uniqueness for Barnes-Wall and Ypsilanti lattices, Adv. Math. **196** (2005), 147–192, Corrections and additions, **211** (2007), 819–824.
- [Hö95] G. Höhn, Selbstduale Vertexoperatorsuperalgebren und das Babymonster, PhD thesis, Universität Bonn, 1995, Bonner Math. Schriften 286, 1996.

- [Hö08] G. Höhn, Conformal designs based on vertex operator algebras, Adv. Math. 217 (2008), 2301–2335.
- [HL95] Y.-Z Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra. III, J. Pure Appl. Algebra **100** (1995), 141–171.
- [KL82] W.M. Kantor and R.A. Liebler, The rank 3 permutation representations of the finite classical groups, Trans. Amer. Math. Soc. 271 (1982), 1–71.
- [LSY07] C. Lam, S. Sakuma and H. Yamauchi, Ising vectors and automorphism groups of commutant subalgebras related to root systems, *Math. Z.* 255 (2007), 597–626.
- [Li87] M.W. Liebeck, The affine permutation groups of rank three, *Proc. London Math. Soc.* (3) **54** (1987), 477–516.
- [LS86] M.W. Liebeck and J. Saxl, The finite primitive permutation groups of rank three, *Bull. London Math. Soc.* **18** (1986), 165–172.
- [Ma01] A. Matsuo, Norton's trace formulae for the Griess algebra of a vertex operator algebra with larger symmetry, *Comm. Math. Phys.* **224** (2001), 565–591.
- [MM00] A. Matsuo and M. Matsuo, The automorphism group of the Hamming code vertex operator algebra, J. Algebra **228** (2000), 204–226.
- [Mi96] M. Miyamoto, Griess algebras and conformal vectors in vertex operator algebras, J. Algebra 179 (1996), 523–548.
- [Mi] M. Miyamoto, Automorphism groups of \mathbb{Z}_2 -orbifold VOAs, unpublished paper, 1996.
- [Mi04] M. Miyamoto, A new construction of the Moonshine vertex operator algebra over the real number field, Ann. of Math. **159** (2004), 535–596.
- [Sh04] H. Shimakura, The automorphism group of the vertex operator algebra V_L^+ for an even lattice L without roots, J. Algebra **280** (2004), 29–57.
- [Sh06] H. Shimakura, The automorphism groups of the vertex operator algebras V_L^+ : general case, Math. Z. 252 (2006), 849–862.
- [Sh07] H. Shimakura, Lifts of automorphisms of vertex operator algebras in simple current extensions, Math. Z. 256 (2007), 491–508.
- [Sh11] H. Shimakura, An E_8 -approach to the moonshine vertex operator algebra, J. London Math. Soc. 83 (2011), 493–516.
- [Sh12] H. Shimakura, Classification of Ising vectors in the vertex operator algebra V_L^+ , Pacific J. Math. 258 (2012), 487–495.
- [Ti84] J. Tits, On R. Griess' "friendly giant", Invent. Math. 78 (1984), 491–499.