Minimax solutions for a problem with sign changing nonlinearity and lack of strict convexity

### Paola Magrone

Dipartimento di Architettura, Università degli Studi Roma Tre Via della Madonna dei Monti 40, Roma, Italia

#### Abstract

A result of existence of a nonnegative and a nontrivial solution is proved via critical point theorems for non smooth functionals. The equation considered presents a convex part and a nonlinearity which changes sign.<sup>1</sup>

# 1 Introduction and main results

Let us consider the problem

$$\begin{cases}
-\operatorname{div}(\Psi'(\nabla u)) = \lambda u + b(x)|u|^{p-2}u & \text{in } \Omega, \\
u = 0 & \text{on } \partial\Omega,
\end{cases}$$
(P)

where  $\lambda$  is a real parameter,  $\Omega$  is a bounded open subset of  $\mathbb{R}^N$ ,  $N \geq 2$ ,  $b(x) \in \overline{C}(\Omega)$  changes sign in  $\Omega$ . Finally  $2 , and we will assume that <math>\Psi : \mathbb{R}^N \to \mathbb{R}$  is a convex function of class  $C^1$  satisfying the following conditions:

<sup>&</sup>lt;sup>1</sup> AMS Subject Classification 35J65, 58E05; keywords: non strict convexity, sign changing, Linking theorem

$$(\Psi_1)$$
  $\lim_{\xi \to 0} \frac{\Psi(\xi)}{|\xi|^2} = \frac{1}{2};$ 

$$(\Psi_2) \quad \exists \mu > 0: \ \mu |\xi|^2 \le \Psi(\xi) \le \frac{1}{\mu} |\xi|^2 \quad \text{for every } \xi \in \mathbb{R}^N;$$

$$(\Psi_3) \quad \lim_{|\xi| \to \infty} \frac{\Psi'(\xi) \cdot \xi - 2\Psi(\xi)}{|\xi|^2} = 0;$$

Moreover the function b(x) has to be strictly positive in a non zero measure set, and the zero set must be "thin", in other words b(x) must satisfy the following conditions:

$$(b_1)$$
  $\Omega^+ := \{x \in \Omega : b(x) > 0\}$  is a nonempty open set

$$(b_2)$$
  $\Omega^0 := \{x \in \Omega : b(x) = 0\}$  has zero measure

Conditions  $(b_1)$  and  $(b_2)$  imply that  $b^+(x) = b(x) + b^-(x) \not\equiv 0$  and that, since b is continuous, the set  $\Omega^0$  is closed in  $\Omega$ .

Let us also denote by  $(\lambda_k)$  the eigenvalues of  $-\Delta$  with homogeneous Dirichlet boundary condition.

In the model case  $\Psi(\xi) = \frac{1}{2}|\xi|^2$ , there is a wide literature on problem  $(\mathcal{P})$ . To cite only some of the existing results, in [2] the authors found positive solutions to  $(\mathcal{P})$  in case that  $\lambda_1 < \lambda < \Lambda^*$ , with  $\Lambda^*$  suitably near to  $\lambda_1$ . In the following many other papers ([1], [2], [3], [5], [6]) were devoted to prove existence of (possibly infinitely many) solutions for  $\lambda \in [\lambda_1, \Lambda^*]$  or also for every  $\lambda$ , in case the nonlinearity satisfies some oddness assumption. A result concerning all  $\lambda$  different from the eigenvalues of the Laplacian under some quite general assumptions can be found in [11], while in [8] the authors proved a result of existence of a nontrivial solution (possibly changing sign) for every  $\lambda$ .

On the other hand, only a small literature is available when dealing with equations with a non strictly convex principal part. In this framework, in [7] the author applies non smooth variational methods in presence of subcritical, positive, nonlinearities; while using similar techniques a nonlinearity with critical growth was considered in [9].

The aim of this paper is to extend to the setting of non strictly convex functionals some of the results contained in [2] (existence of a positive solution for  $\lambda < \lambda_1$ ) and [8] (existence of a nontrivial solution for any  $\lambda$ .)

Problem  $(\mathcal{P})$  can be treated by variational techniques. Indeed, weak solutions u of  $(\mathcal{P})$  can be found as critical points of the  $C^1$  functional  $J: H_0^1(\Omega) \to \mathbb{R}$  defined as

$$J(u) = \int_{\Omega} \Psi(\nabla u) dx - \frac{\lambda}{2} \int_{\Omega} u^2 dx - \frac{1}{p} \int_{\Omega} b(x) |u|^p dx.$$
 (1.1)

The key point here is that, although  $\Psi$  shares some properties with this typical case, there is no assumption of strict convexity with respect to  $\xi$ .

For instance, one could consider

$$\Psi(\xi) = \psi(\xi_1) + \frac{1}{2} \sum_{j=2}^{N} \xi_j^2, \qquad (1.2)$$

where

$$\psi(t) = \begin{cases} \frac{1}{2}t^2 & \text{if } |t| < 1, \\ |t| - \frac{1}{2} & \text{if } 1 \le |t| \le 2, \\ \\ \frac{1}{2}|t|^2 - |t| + \frac{3}{2} & \text{if } |t| > 2. \end{cases}$$

If we look at the principal part of J as the energy stored in the deformation u, this means that the material has a plastic behavior when  $1 \leq |D_1 u| \leq 2$ . We refer the reader to [13, Chapter 6] for a discussion of several models of plasticity.

As shown in [7, 9], it may happen that Palais Smale sequences, even if bounded in  $H_0^1(\Omega)$ -norm, do not admit any subsequence which converges strongly in this norm. And there is no way to prevent the interaction between the area where  $\Psi$  loses strict convexity and the values of  $\nabla u$ . A possibile strategy is to look for compactness in a weaker norm  $(L^{2^*})$ .

Let us introduce the following notations: let  $k \geq 1$  be such that  $\lambda_k \leq \lambda < \lambda_{k+1}$ 

and let  $e_1, ..., e_k$  be eigenfunctions of  $-\Delta$  associated to  $\lambda_1, ..., \lambda_k$ , respectively. Finally, let  $E_- = \text{span}\{e_1, ..., e_k\}$  and  $E_+ = E_-^{\perp}$ . The main result of this paper are the following:

**Theorem 1.1.** Let  $N \geq 2$  and let  $\Psi : \mathbb{R}^N \to \mathbb{R}$  be a convex function of class  $C^1$  satisfying  $(\Psi_1), (\Psi_2)$ . Moreover let the function b(x) verify  $(b_1)$ . Then, for every  $\lambda \in ]0, \lambda_1[$ , problem  $(\mathcal{P})$  admits a nontrivial and nonnegative weak solution  $u \in H_0^1(\Omega)$ .

**Theorem 1.2.** Let  $N \geq 2$  and let  $\Psi : \mathbb{R}^N \to \mathbb{R}$  be a convex function of class  $C^1$  satisfying  $(\Psi_1)$ ,  $(\Psi_2)$  and let  $\lambda \geq \lambda_1$ . Moreover let the function b(x) verify  $(b_1)$ , and the following assumptions:

$$\int_{\Omega} b(x)|v|^p \ge 0 \qquad \forall v \in E_-. \tag{1.3}$$

$$\exists e \in E_{-}^{\perp} \setminus \{0\} : \int_{\Omega} b(x)|v|^p \, dx \ge C \int_{\Omega} |v|^p \, dx \qquad \forall v \in E_{-} \oplus span\{e\}. \quad (1.4)$$

Then problem  $(\mathcal{P})$  admits a nontrivial weak solution  $u \in H_0^1(\Omega)$ .

**Remark 1.3.** Arguing as in section 2 of [9] we can deduce the following properties for  $\Psi$ , up to modifying the constant  $\mu$ :

$$\Psi'(\xi) \cdot \xi \ge \mu |\xi|^2 \qquad \forall \xi \in \mathbb{R}^N,$$
(1.5)

$$|\Psi'(\xi)| \ge \mu|\xi| \qquad \forall \xi \in \mathbb{R}^N$$
 (1.6)

$$|\Psi'(\xi)| \le \frac{1}{\mu} |\xi| \qquad \forall \xi \in \mathbb{R}^N \tag{1.7}$$

Furthermore  $(\Psi_3)$  yields that  $\forall \sigma > 0, \exists M_{\sigma} \in \mathbb{R}$ :

$$\Psi'(\xi)\xi - 2\Psi(\xi) \le \sigma|\xi|^2 + M_{\sigma} \tag{1.8}$$

## 2 The variational framework

Let  $\Omega$  be a bounded open subset of  $\mathbb{R}^N$ ,  $N \geq 2$ , with Lipschitz boundary and let  $\lambda \in \mathbb{R}$ . Let us define the following functional  $J: H_0^1(\Omega) \to \mathbb{R}$ 

$$J(u) = \int_{\Omega} \Psi(\nabla u) \, dx - \frac{\lambda}{2} \int_{\Omega} u^2 \, dx - \frac{1}{p} \int_{\Omega} b(x) |u|^p \, dx.$$

By  $(\Psi_1)$ ,  $(\Psi_2)$  the functional J is of class  $C^1$  on  $H^1_0(\Omega)$ . We wish to apply variational methods to functional J, but, as already mentioned, it is well known that the Palais Smale (PS) condition for a functional which is not strictly convex is not satisfied on  $H^1_0(\Omega)$ . So it is convenient to extend the functional J to  $L^{2^*}$  with value  $+\infty$  outside  $H^1_0(\Omega)$ .

In other words we define the convex, lower semicontinuous functional (still denoted J)

$$J: L^{2^*}(\Omega) \longrightarrow [0, +\infty]$$

$$J(u) = \begin{cases} \int_{\Omega} \Psi(\nabla u) \, dx - \frac{\lambda}{2} \int_{\Omega} u^2 \, dx - \frac{1}{p} \int_{\Omega} b(x) |u|^p \, dx & \text{if } u \in H_0^1(\Omega), \\ +\infty & \text{if } u \in L^{2^*}(\Omega) \setminus H_0^1(\Omega) \end{cases}$$
(2.1)

This setting will allow us to recover PS condition.

This functional can be written as  $J = J_0 + J_1$ , where

$$J_0 = \int_{\Omega} \Psi(\nabla u) \, dx,$$

is proper, convex and l.s.c., while

$$J_1 = -\frac{\lambda}{2} \int_{\Omega} u^2 dx - \frac{1}{p} \int_{\Omega} b(x) |u|^p dx,$$

is of class  $C^1$ . We will use the following definitions ([12], [7]) of critical point and PS sequence for functionals of the type  $J = J_0 + J_1$ :

**Definition 2.1.** Let X be a real Banach space,  $u \in X$  is a critical point for J

if  $J(u) \in \mathbb{R}$  and  $-J'_1(u) \in \partial J_0$ , where  $\partial J_0$  is the subdifferential of  $J_0$  at u.

**Definition 2.2.** Let X be a real Banach space and let  $c \in \mathbb{R}$ . We say that  $u_k$  is a Palais Smale sequence at level c ( $(PS)_c$  sequence for short) for J if  $J(u_k) \to 0$  and there exists  $\alpha_k \in \partial J_0$  with  $(\alpha_k + J_1'(u_k)) \to 0$  in  $X^*$ .

The following proposition (see [7]) assures that the critical points of the extendend functional already defined gives the solutions of our problem.

**Proposition 2.3.** Let  $u \in L^{2^*}(\Omega, \mathbb{R}^N)$ . Then u is a critical point of J if and only if  $u \in H_0^1(\Omega)$  and u is a weak solution of  $(\mathcal{P})$ .

**Proof** Let  $v \in L^{2^*}$ . Then  $v \in \partial J_0$ , if and only if  $u \in H_0^1(\Omega)$  and

$$-div(\Psi'(\nabla u)) = v$$

that is a reformulation of definition 2.1.

Moreover we will apply the compactness result contained in [7], which we recall.

Let us define the functional  $\mathcal{E}:W^{1,2}_0(\Omega,\mathbb{R}^N)\to\mathbb{R}$  as

$$\mathcal{E}(u) = \int_{\Omega} \Psi(\nabla u) \, dx$$

**Theorem 2.4.** Assume that  $\Omega$  is bounded. If  $\{u_h\}$  is weakly convergent to u in  $W_0^{1,2}(\Omega,\mathbb{R}^N)$  with  $\mathcal{E}(\{u_h\}) \to \mathcal{E}(\{u\})$ , then u is strongly convergent to u in  $L^{2^*}(\Omega)$ .

# 3 Proof of main results

Since  $\Psi'(0) = 0$ , of course 0 is a solution of  $(\mathcal{P})$ . Therefore we are interested in nontrivial solutions. In order to find nonnegative solutions of  $(\mathcal{P})$ , we consider

the modified functional  $\overline{J}: L^{2^*}(\Omega) \to ]-\infty, +\infty]$  defined as

$$\overline{J}(u) = \begin{cases} \int_{\Omega} \Psi(\nabla u) dx - \frac{\lambda}{2} \int_{\Omega} (u^+)^2 dx - \frac{1}{p} \int_{\Omega} b(x) (u^+)^p dx & \text{if } u \in H_0^1(\Omega), \\ +\infty & \text{if } u \in L^{2^*}(\Omega) \setminus H_0^1(\Omega) \end{cases}$$

Of course,  $\overline{J}$  is also convex and lower semicontinuous.

**Proposition 3.1.** Let  $\Psi : \mathbb{R}^N \to \mathbb{R}$  be a convex function of class  $C^1$  satisfying  $(\Psi_2)$  with  $\mu > 0$ , and (1.6). Then each critical point  $u \in L^{2^*}$  of  $\overline{J}$  is a nonnegative solution of  $(\mathcal{P})$ .

**Proof** Since by Proposition 2.3 we already know that the critical points of J are solutions of our problem, it is only left to prove that the modified functional will give nonnegative solutions. By  $(\Psi_2)$  one has

$$\mu \int_{\Omega} |\nabla u^{-}|^{2} dx dx \le \int_{\Omega} \Psi'(\nabla u) \cdot (-\nabla u^{-}) dx dx =$$

$$= \lambda \int_{\Omega} u^{+}(-u^{-}) dx dx + \int_{\Omega} (u^{+})^{p-1}(-u^{-}) dx dx = 0,$$

whence the assertion.

sition 3.1.

**Remark 3.2.** From now on, to simplify notations, we will keep on using the functional J instead of  $\overline{J}$ , since it is understood what has been proved in Propo-

### Proof of Theorem 1.1

We aim to apply to J a nonsmooth version of Mountain Pass Theorem [12]. First of all, let us observe that, by  $(\Psi_1)$ , we have

$$\frac{\int_{\Omega} \Psi(\nabla u) \, dx}{\int_{\Omega} |\nabla u|^2 \, dx} \to \frac{1}{2} \quad \text{as } u \to 0 \text{ in } L^{2^*}.$$

Then, as in the case  $\Psi(\xi) = \frac{1}{2}|\xi|^2$  treated in [2, 8], we deduce that there exist  $\varrho > 0$  and  $\alpha > 0$  such that  $J(u) \geq \alpha$  whenever  $||u|| = \varrho$ . On the other hand,

there exists  $e \in L^{2^*}$  with  $e \ge 0$  a.e. in  $\Omega$  such that

$$\lim_{t \to +\infty} J(te) = -\infty \,,$$

again, this is proved in [2] in the case  $\Psi(\xi) = \frac{1}{2}|\xi|^2$ , but by  $(\Psi_2)$  the assertion is true also in our case.

By the Mountain Pass theorem, there exist a sequence  $(u_k)$  in  $L^{2^*}$  and a sequence  $(w_k)$  in  $L^{(2^*)'}(\Omega)$  strongly convergent to 0 such that (see definition 2.2)

$$\int_{\Omega} \Psi'(\nabla u_k)(\nabla v - \nabla u_k) \, dx \ge \lambda \int_{\Omega} u_k(v - u_k) \, dx + \int_{\Omega} b(x)|u_k|^{p-1}(v - u_k) \, dx$$
$$+ \int_{\Omega} w_k(v - u_k) \, dx \quad \forall v \in L^{(2^*)'} \quad (3.1)$$

Taking v = 0 and  $v = 2u_k$  as tests in the previous inequality yield

$$\int_{\Omega} \Psi'(\nabla u_k) \nabla u_k \, dx = \lambda \int_{\Omega} (u_k)^2 \, dx + \int_{\Omega} b(x) |u_k|^p \, dx + \int_{\Omega} w_k u_k \, dx \quad \forall v \in L^{(2^*)'}.$$
(3.2)

Furthermore also the following relation holds:

$$\lim_{k \to \infty} \left( \int_{\Omega} \Psi(\nabla u_k) \, dx - \frac{\lambda}{2} \int_{\Omega} (u_k)^2 \, dx - \frac{1}{p} \int_{\Omega} b(x) (u_k)^p \, dx \right) = c > \alpha. \tag{3.3}$$

Let us write the expression  $pJ(u_k) - J'(u_k)u_k$ , which is bounded by assumptions (3.2), (3.3):

$$p \int_{\Omega} \Psi(\nabla u_k) dx - \frac{p}{2} \lambda \int_{\Omega} (u_k)^2 dx - \int_{\Omega} b(x)(u_k)^p dx - \int_{\Omega} \Psi'(\nabla u_k) \cdot \nabla u_k dx$$
$$+ \lambda \int_{\Omega} (u_k)^2 dx + \int_{\Omega} b(x)(u_k)^p dx =$$
$$\int_{\Omega} (p-2)\Psi(\nabla u_k) dx + \int_{\Omega} \left[2\Psi(\nabla u_k) - \Psi'(\nabla u_k) \cdot \nabla u_k\right] dx - \lambda \left(\frac{p}{2} - 1\right) \int_{\Omega} (u_k)^2 dx =$$
$$(p-2)c - \int_{\Omega} w_k u_k dx + C \quad (3.4)$$

By (1.8) and  $(\Psi_2)$  one gets

$$\mu(p-2-\sigma) \int_{\Omega} |\nabla u_k|^2 dx - \lambda \left(\frac{p}{2}-1\right) \lambda \int_{\Omega} (u_k)^2 dx \le pc - \int_{\Omega} w_k u_k + C \quad (3.5)$$

SO

$$\mu(p-2-\sigma) \int_{\Omega} |\nabla u_k|^2 dx \le \lambda \left(\frac{p}{2} - 1\right) \int_{\Omega} (u_k)^2 dx + C \tag{3.6}$$

where the quantity  $(p-2-\sigma)$  is strictly positive since  $\sigma$  is arbitrarily small. Our aim is to prove the boundedness of the  $H_0^1$  norm of the Palais Smale sequences, so arguing by contradiction, let us assume that

$$||u_k|| \to \infty$$
 as  $k \to +\infty$ .

Dividing (3.3) by  $||u_k||^p$  yields

$$\liminf \left\{ \frac{p \int_{\Omega} \Psi(\nabla u_k)}{||u_k||^p} dx - \frac{\lambda p}{2} \frac{\int_{\Omega} (u_k)^2 dx}{||u_k||^p} dx - \frac{1}{p} \int_{\Omega} b(x) \left( \frac{u_k}{||u_k||} \right)^p dx \right\} = 0.$$

Since p > 2 and  $(\Psi_2)$  holds, the first two terms go to zero. So

$$\lim \sup \left( \int_{\Omega} b(x) \left( \frac{u_k}{||u_k||} \right)^p dx \right) = 0. \tag{3.7}$$

Since b is bounded, by Lebesgue dominated convergence Theorem we can take the limit and deduce that

$$\lim_{k} b(x) \left( \frac{u_k}{||u_k||} \right)^p = 0. \tag{3.8}$$

This yields that

$$\left(\frac{u_k}{||u_k||}\right) \to u_0$$

strongly in  $L^p$  and weakly in  $H_0^1(\Omega)$ . Arguing by contradiction let us suppose

that  $u_0 \equiv 0$ . Dividing (3.6) by  $||u_k||^2$  yields

$$\mu(p-2-2\sigma) \le \lambda \left(\frac{p}{2}-1\right) \frac{1}{||u_k||^2} \int_{\Omega} (u_k)^2 dx + \frac{C}{||u_k||^2}$$
 (3.9)

the right hand side goes to zero, which leads to a contradiction since  $p-2-2\sigma > 0$  and  $\mu > 0$ , so  $u_0$  must not be identically zero.

Now let  $\phi \in C_0^{\infty}(\Omega^+)$  be a compact support function,  $\phi \geq 0$  and  $\phi \not\equiv 0$ . Let us use the function  $t\phi v, v \in H_0^1(\Omega)$  as a test in 3.1.

$$\int_{\Omega^{+}} \Psi'(\nabla u_{k})(t\phi \nabla v + tv \nabla \phi - \nabla u_{k}) \ge \lambda \int_{\Omega^{+}} u_{k}(tv\phi - u_{k}) + \int_{\Omega^{+}} b(x)(u_{k})^{p-1}(tv\phi - u_{k})$$
$$\int_{\Omega^{+}} w_{k}(tv\phi - u_{k}) \quad \forall v \in H_{0}^{1}(\Omega)$$

Then let us divide the previous inequality by t and then let t go to  $+\infty$ :

$$\int_{\Omega^{+}} \Psi'(\nabla u_{k})(\phi \nabla v) + \Psi'(\nabla u_{k})v \nabla \phi \ge +\lambda \int_{\Omega^{+}} (u_{k})^{2} v \phi + \int_{\Omega^{+}} b^{+}(x)(u_{k})^{p-1} v \phi + \int_{\Omega^{+}} w_{k} v \phi \qquad \forall v \in H_{0}^{1}(\Omega) \quad (3.10)$$

On the other hand, if  $t \to -\infty$ , one gets the opposite inequality, so we can deduce that the equality holds in the last expression, that is

$$\int_{\Omega^{+}} \Psi'(\nabla u_{k})(\phi \nabla v) + \Psi'(\nabla u_{k})v \nabla \phi = 
+ \lambda \int_{\Omega^{+}} (u_{k})^{2} v \phi + \int_{\Omega^{+}} b^{+}(x)(u_{k})^{p-1} v \phi + \int_{\Omega} w_{k} v \phi \quad \forall v \in H_{0}^{1}(\Omega).$$
(3.11)

Now let us choose  $v = u_k$  and divide both handsides of (3.11) by  $||u_k||^p$ . It is easily seen that the terms containing  $\lambda$  and  $w_k$  go to 0 as  $k \to +\infty$ . Then

$$\int_{\Omega^+} \frac{\Psi'(\nabla u_k) \nabla u_k \phi}{||u_k||^p}$$

goes to 0 since p > 2 and (1.7) holds.

On the other hand, by (1.7), since p > 2 and  $\phi$  is of class  $C^{\infty}$  in  $\Omega^+$  bounded,

$$\frac{1}{||u_k||^p} \int_{\Omega^+} \Psi'(\nabla u_k) u_k \nabla \phi \le C \frac{||u_k||}{||u_k||^{p-1}} \frac{||u_k||_{L^2}}{||u_k||}$$

The term  $\frac{||u_k||_{L^2}}{||u_k||}$  is bounded, while  $\frac{||u_k||}{||u_k||^{p-1}}$  converges to 0.

By (3.11) We can conclude that

$$\int_{\Omega^+} \frac{1}{||u_k||^p} b^+(x) (u_k)^p \phi \mapsto 0 \text{ as } k \to \infty.$$

Applying Fatou's Lemma yields

$$\lim \inf \int_{\Omega^+} \frac{1}{||u_k||^p} b^+(x) (u_k)^p \phi \le 0$$

and since the integrand is nonnegative, this means that  $\frac{u_k^p}{||u_k||^p}$  must tend to 0 in  $\Omega^+$  as  $k \to \infty$ , but this is in contradiction with the fact that it was already proved that it converges to a nonzero function  $u_0$ .

Arguing in the same way, choosing now a compact support function  $\eta \in C_0^{\infty}(\Omega^-)$ , yields that  $\frac{u_k^p}{||u_k||^p} \to 0$  as  $k \to \infty$ , in  $\Omega^-$ .

This proves that  $||u_k||$  is bounded in  $H_0^1(\Omega)(\Omega^+ \cup \Omega^-)$ , and since  $\Omega^0$  is negligeable, this concludes this part of the proof. Then  $u_k$  admits a subsequence weakly converging in  $L^{2^*}$ .

According to (3.1) and taking v = u as a test function yields

$$\int_{\Omega} \Psi'(\nabla u_k)(\nabla u - \nabla u_k) dx \ge \lambda \int_{\Omega} u_k(u - u_k) dx + \int_{\Omega} b(x)(u_k)^{p-1} (u - u_k) dx + o(1)$$
(3.12)

so as  $k \to \infty$  the right hand-side terms go to zero, and we obtain

$$\liminf \int_{\Omega} \Psi'(\nabla u_k)(\nabla u - \nabla u_k) \, dx \ge 0. \tag{3.13}$$

On the other hand, by convexity

$$\int_{\Omega} \Psi(\nabla u) \, dx \ge \int_{\Omega} \Psi(\nabla u_k) \, dx + \int_{\Omega} \Psi'(\nabla u_k) (\nabla u - \nabla u_k) \, dx \tag{3.14}$$

So by (3.13) and (3.14)

$$\limsup \int_{\Omega} \Psi(\nabla u_{k}) \, dx \le \limsup \left( \int_{\Omega} \Psi(\nabla u) \, dx - \int_{\Omega} \Psi'(\nabla u_{k}) (\nabla u - \nabla u_{k}) \, dx \right)$$

$$\le \int_{\Omega} \Psi(\nabla u) \, dx - \liminf \int_{\Omega} \Psi'(\nabla u_{k}) (\nabla u - \nabla u_{k}) \, dx \le \int_{\Omega} \Psi(\nabla u) \, dx$$

$$(3.15)$$

By lower semicontinuity and convexity

$$\lim \inf \int_{\Omega} \Psi(\nabla u_k) \, dx \ge \int_{\Omega} \Psi(\nabla u) \, dx \tag{3.16}$$

We can conclude that

$$\int_{\Omega} \Psi(\nabla u_k) \, dx \to \int_{\Omega} \Psi(\nabla u) \, dx.$$

By Theorem 3.1  $u_k$  admits a subsequence strongly converging in  $L^{2^*}$ , which concludes the proof of PS condition and of Theorem 1.1.

#### Proof of Theorem 1.2

We are now concerned with the existence of (possibly sign-changing) nontrivial solutions u of  $(\mathcal{P})$ . Let  $(\lambda_k)$  denote the sequence of the eigenvalues of  $-\Delta$  with homogeneous Dirichlet condition, repeated according to multiplicity.

Since the case  $0 < \lambda < \lambda_1$  is already contained in Theorem 1.1, we may assume

12

that  $\lambda \geq \lambda_1$ . Let  $k \geq 1$  be such that  $\lambda_k \leq \lambda < \lambda_{k+1}, e_1, \dots, e_k$  are eigenfunctions of  $-\Delta$ , as defined in the introduction. Finally, let  $E_- = \operatorname{span}\{e_1, \dots, e_k\}$  and  $E_+ = E_-^{\perp}$ .

Consider the functional J defined in (2.1) We aim to apply the version of the Linking Theorem for convex functional presented by Szulkin in [12]. Since

$$\frac{\int_{\Omega} \Psi(\nabla u) \, dx}{\int_{\Omega} |\nabla u|^2 \, dx} \to \frac{1}{2} \qquad \text{as } u \to 0 \text{ in } H^1_0(\Omega),$$

as in the case  $\Psi(\xi) = \frac{1}{2}|\xi|^2$  treated in [8], we deduce that there exist  $\varrho > 0$  and  $\alpha > 0$  such that  $J(u) \ge \alpha$  whenever  $u \in E_+$  with  $||u|| = \varrho$ . On the other hand, there exists  $e \in H_0^1(\Omega) \setminus E_-$  such that

$$\lim_{\begin{subarray}{c} \|u\| \to \infty \\ u \in \mathbb{R} e \oplus E_- \end{subarray}} J(u) = -\infty \,,$$

Again, this is proved in [8] when  $\Psi(\xi) = \frac{1}{2}|\xi|^2$ , but by  $(\Psi_2)$  the assertion is true also in our case. Finally, it is clear that  $J(u) \leq 0$  for every  $u \in E_-$ .

By the Linking type theorem in [12] (Theorem 3.4), there exist a PS sequence  $(u_k)$  in  $H_0^1(\Omega)$  and we can continue, up to minor changes, as in the proof of Theorem 1.1 to prove that there exists a subsequence of  $(u_k)$  strongly converging in  $L^{2^*}$ . This concludes the proof of Theorem 1.2, since the nontriviality of the solution comes directly from the characterization of the critical level of the solution.

**Acknowledgment** The author thanks Prof. Marco Degiovanni for very helpful conversations.

## References

- [1] S. Alama and M. del Pino, Solutions of elliptic equations with indefinite nonlinearities via Morse theory and linking, Ann. I.H.P. Analyse non linearite, Vol. 13 (1996), 95–115.
- [2] S. Alama and G. Tarantello, On semilinear elliptic equations with indefinite nonlinearities, Calc. Var., Vol 1, (1993), 439–475.
- [3] S. Alama and G. Tarantello, Elliptic problems with nonlinearities indefinite in sign, J. Funct. An. Vol. 141, (1996), 159–215.
- [4] A. Ambrosetti and P.H. Rabinowitz, Dual variational methods in critical point theory and applications, J. Funct. Anal., Vol. 14 (1973), 349–381.
- [5] M. Badiale, Infinitely many solutions for some indefinite nonlinear elliptic problems, Comm. on Appl. Non. Anal., Vol. 3 (1996), 61–76.
- [6] M. Badiale and E. Nabana, A remark on multiplicity solutions for semilinear elliptic problems with indefinite nonlinearities, C.R. Acad. Sc. Paris, Vol. 323, serie 1 (1996), 151–156.
- [7] M. Degiovanni, Variational methods for functionals with lack of strict convexity, in "Nonlinear Equations: Methods, Models and Applications" (Bergamo, 2001), D. Lupo, C. Pagani and B. Ruf, eds., 127–139, Progress in Nonlinear Differential Equations and their Applications, Vol. 54, Birkhäuser, Boston, Inc., Boston, Ma, 2003.
- [8] M.Grossi, P.Magrone, M.Matzeu, Linking type solutions for elliptic equations with indefinite nonlinearities up to the critical growth, Discrete Contin. Dynam. Systems, Vol. 7, no 4, (2001), 703-718.
- [9] P.Magrone, An existence result for a problem with critical growth and lack of strict convexity, NoDEA Nonlinear Differential Equations Appl. Vol. 15, no 6, (2008), 717–728.
- [10] P.H. Rabinowitz, Minimax methods in critical point theory with applications to differential equations', "CBMS Regional Conference Series in

- Mathematics", Vol. 65, American Mathematical Society, Providence, R.I., 1986.
- [11] M. Ramos, S. Terracini and C. Troestler, Superlinear indefinite elliptic problems and Pohozaev type identities, J. Funct. Anal., Vol159, (1998), 596–628.
- [12] A. Szulkin, Minimax principles for lower semicontinuous functions and applications to nonlinear boundary value problems., Ann. Inst. H. Poincaré Anal. Non Linéaire, Vol. 3, no. 2, (1986), 77-109.
- [13] H.-C. Wu, Continuum mechanics and plasticity, CRC Series: Modern Mechanics and Mathematics, Chapman & Hall/CRC, Boca Raton, FL., (2005).