ANOTHER CHRISTOFFEL-DARBOUX FORMULA FOR MULTIPLE ORTHOGONAL POLYNOMIALS OF MIXED TYPE

GERARDO ARAZNIBARRETA AND MANUEL MAÑAS

ABSTRACT. An alternative expression for the Christoffel-Darboux formula for multiple orthogonal polynomials of mixed type is derived from the LU factorization of the moment matrix of a given measure and two sets of weights. We use the action of the generalized Jacobi matrix J, also responsible for the recurrence relations, on the linear forms and their duals to obtain the result.

1. Introduction

In this paper we address a natural question that arises from the LU factorization approach to multiple orthogonality [7]. The Gauss-Borel factorization of a Hankel matrix, which plays the role of a moment matrix, leads in the classical case to a natural description of algebraic facts regarding orthogonal polynomials on the real line (OPRL) such as recursion relations and Christoffel–Darboux formula. In that case we have a chain of orthogonal polynomials $\{P_l(x)\}_{l=0}^{\infty}$ of increasing degree l. In [7] we extended that approach to the multiple orthogonality scenario, and the Gauss–Borel factorization of an appropriate moment matrix leaded to sequences of families of multiple orthogonal polynomials in the real line (MOPRL), $\{Q_{[\vec{\nu}_1(l);\vec{\nu}_2(l-1)]}^{(II,a_1(l))}\}_{l=0}^{\infty}$ and $\{\bar{Q}_{[\vec{\nu}_2(l);\vec{\nu}_1(l-1)]}^{(I,a_1(l))}\}_{l=0}^{\infty}$. The recursion relations are relations constructed in terms of the linear forms in these sequences. However, the Christoffel-Darboux formula given in Proposition 4, that was re-deduced by linear algebraic means (Gauss-Borel factorization), was not expressed in terms of linear forms belonging to the mentioned sequences. This situation is rather different to the OPRL case, in that classical case the Christoffel-Darboux formula is expressed in terms of orthogonal polynomials in the sequence. The aim of this paper is to show that, within that scheme, we can deduce an alternative but equivalent MOPRL Christoffel–Darboux formula constructed in terms of linear forms in the sequences $\left\{Q_{[\vec{\nu}_1(l);\vec{\nu}_2(l-1)]}^{(\Pi,a_1(l))}\right\}_{l=0}^{\infty}$ and $\left\{\bar{Q}_{[\vec{\nu}_2(l);\vec{\nu}_1(l-1)]}^{(I,a_1(l))}\right\}_{l=0}^{\infty}$ as in OPRL situation.

1.1. Historical background. Simultaneous rational approximation starts back in 1873 when Hermite proved the transcendence of the Euler number e [17]. Later, K. Mahler delivered at the University of Groningen several lectures [20] where he settled down the foundations of this theory, see also [12] and [18]. Simultaneous rational approximation when expressed in terms of Cauchy transforms leads to multiple orthogonality of polynomials. Given an interval $\Delta \subset \mathbb{R}$ of the real line, let $\mathcal{M}(\Delta)$ denote all the finite positive Borel measures with support containing infinitely many points in Δ . Fix $\mu \in \mathcal{M}(\Delta)$, and let us consider a system of weights $\vec{w} = (w_1, \dots, w_n)$ on Δ , with $p \in \mathbb{N}$; i.e. w_1, \ldots, w_p being real integrable functions on Δ which does not change sign on Δ . Fix a multi-index $\vec{\nu} = (\nu_1, \dots, \nu_p) \in \mathbb{Z}_+^p$, $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$, and denote $|\vec{\nu}| = \nu_1 + \dots + \nu_p$. Then, there exist polynomials, A_1, \ldots, A_p not all identically equal to zero which satisfy the following orthogonality relations

(1)
$$\int_{\Delta} x^{j} \sum_{a=1}^{p} A_{a}(x) w_{a}(x) d\mu(x) = 0, \qquad \deg A_{a} \leq \nu_{a} - 1, \qquad j = 0, \dots, |\vec{\nu}| - 2.$$

Analogously, there exists a polynomial
$$B$$
 not identically equal to zero, such that
$$(2) \qquad \int_{\Delta} x^{j} B(x) w_{b}(x) \mathrm{d}\mu(x) = 0, \qquad \deg B \leq |\vec{\nu}|, \qquad j = 0, \dots, \nu_{b} - 1, \quad b = 1, \dots, p.$$

These are the so called multiple orthogonal polynomials of type I and type II, respectively, with respect to the combination $(\mu, \vec{w}, \vec{\nu})$ of the measure μ , the systems of weights \vec{w} and the multi-index $\vec{\nu}$. When p=1 both

¹⁹⁹¹ Mathematics Subject Classification. 33C45,42C05,15A23,37K10.

Key words and phrases. Multiple orthogonal polynomials, Christoffel-Darboux formula, moment matrices, Jacobi type matrices, Gaus-Borel factorization.

definitions coincide with standard orthogonal polynomials on the real line. Given a measure $\mu \in \mathcal{M}(\Delta)$ and a system of weights \vec{w} on Δ a multi-index $\vec{\nu}$ is called type I or type II normal if deg A_a must equal to $\nu_a - 1$, $a = 1, \ldots, p$, or deg B must equal to $|\vec{\nu}| - 1$, respectively. When for a pair (μ, \vec{w}) all the multi-indices are type I or type II normal, then the pair is called type I perfect or type II perfect respectively. Multiple orthogonal of polynomials have been employed in several proofs of irrationality of numbers. For example, in [10] F. Beukers shows that Apery's proof [9] of the irrationality of $\zeta(3)$ can be placed in the context of a combination of type I and type II multiple orthogonality which is called mixed type multiple orthogonality of polynomials. More recently, mixed type approximation has appeared in random matrix and non-intersecting Brownian motion theories, [11], [13], [19]. Sorokin [24] studied a simultaneous rational approximation construction which is closely connected with multiple orthogonal polynomials of mixed type. In [26] a Riemann–Hilbert problem was found that characterizes multiple orthogonality of type I and II, extending in this way the result previously found in [16] for standard orthogonality. In [13] mixed type multiple orthogonality was analyzed from this perspective.

1.2. **Perfect systems and MOPRL of mixed type.** In order to introduce multiple orthogonal polynomials of mixed type we consider two systems of weights $\vec{w}_1 = (w_{1,1}, \ldots, w_{1,p_1})$ and $\vec{w}_2 = (w_{2,1}, \ldots, w_{2,p_2})$ where $p_1, p_2 \in \mathbb{N}$, and two multi-indices $\vec{\nu}_1 = (\nu_{1,1}, \ldots, \nu_{1,p_1}) \in \mathbb{Z}_+^{p_1}$ and $\vec{\nu}_2 = (\nu_{2,1}, \ldots, \nu_{2,p_2}) \in \mathbb{Z}_+^{p_2}$ with $|\vec{\nu}_1| = |\vec{\nu}_2| + 1$. There exist polynomials A_1, \ldots, A_{p_1} , not all identically zero, such that $\deg A_s < \nu_{1,s}$, which satisfy the following relations

(3)
$$\int_{\Delta} \sum_{a=1}^{p_1} A_a(x) w_{1,a}(x) w_{2,b}(x) x^j d\mu(x) = 0, \quad j = 0, \dots, \nu_{2,b} - 1, \quad b = 1, \dots, p_2.$$

They are called mixed multiple-orthogonal polynomials with respect to the combination $(\mu, \vec{w}_1, \vec{w}_2, \vec{\nu}_1, \vec{\nu}_2)$ of the measure μ , the systems of weights \vec{w}_1 and \vec{w}_2 and the multi-indices $\vec{\nu}_1$ and $\vec{\nu}_2$. It is easy to show that finding the polynomials A_1, \ldots, A_{p_1} is equivalent to solving a system of $|\vec{\nu}_2|$ homogeneous linear equations for the $|\vec{\nu}_1|$ unknown coefficients of the polynomials. Since $|\vec{\nu}_1| = |\vec{\nu}_2| + 1$ the system always has a nontrivial solution. The matrix of this system of equations is the so called moment matrix, and the study of its Gauss–Borel factorization will be the cornerstone of this paper. Observe that when $p_1 = 1$ we are in the type II case and if $p_2 = 1$ in type I case. Hence in general we can find a solution of (3) where there is an $a \in \{1, \ldots, p_1\}$ such that $\deg A_a < \nu_{1,a} - 1$. When given a combination $(\mu, \vec{w}_1, \vec{w}_2)$ of a measure $\mu \in \mathcal{M}(\Delta)$ and systems of weights \vec{w}_1 and \vec{w}_2 on Δ if for each pair of multi-indices $(\vec{\nu}_1, \vec{\nu}_2)$ the conditions (3) determine that $\deg A_a = \nu_{1,a} - 1$, $a = 1, \ldots, p_1$, then we say that the combination $(\mu, \vec{w}_1, \vec{w}_2)$ is perfect. In this case we can determine a unique system of mixed type orthogonal polynomials (A_1, \ldots, A_{p_1}) satisfying (3) requiring for $a_1 \in \{1, \ldots, p_1\}$ that A_{a_1} monic. Following [13] we say that we have a type II normalization and denote the corresponding system of polynomials by $A_a^{(II,a_1)}$, $j = 1, \ldots, p_1$. Alternatively, we can proceed as follows, since the system of weights is perfect from (3) we deduce that

$$\int x^{\nu_{1,r_1}} \sum_{a=1}^{p_1} A_a(x) w_{1,a}(x) w_{2,b}(x) d\mu(x) \neq 0.$$

Then, we can determine a unique system of mixed type of multi-orthogonal polynomials $(A_1^{(I,a_2)}, \ldots, A_{p_2}^{(I,a_2)})$ imposing that

$$\int x^{\nu_{1,a_2}} \sum_{a=1}^{p_1} A_a^{(\mathbf{I},a_2)}(x) w_{1,a}(x) w_{2,b}(x) d\mu(x) = 1,$$

which is a type I normalization. We will use the notation $A_{[\vec{\nu}_1;\vec{\nu}_2],a}^{(\mathrm{II},a_1)}$ and $A_{[\vec{\nu}_1;\vec{\nu}_2],a}^{(\mathrm{I},a_2)}$ to denote these multiple orthogonal polynomials with type II and I normalizations, respectively.

A known illustration of perfect combinations $(\mu, \vec{w}_1, \vec{w}_2)$ can be constructed with an arbitrary positive finite Borel measure μ and systems of weights formed with exponentials:

(4)
$$(e^{\gamma_1 x}, \dots, e^{\gamma_p x}), \qquad \gamma_i \neq \gamma_j, \qquad i \neq j, \qquad i, j = 1, \dots, p,$$

or by binomial functions

(5)
$$((1-z)^{\alpha_1}, \dots, (1-z)^{\alpha_p}), \qquad \alpha_i - \alpha_j \notin \mathbb{Z}, \qquad i \neq j, \qquad i, j = 1, \dots, p.$$

or combining both classes, see [21]. Recently, in [15] the authors were able to prove perfectness for a wide class of systems of weights. These systems of functions, now called Nikishin systems, were introduced by E.M. Nikishin [21] and initially named MT-systems (after Markov and Tchebycheff).

1.3. Borel–Gauss factorization and multiple orthogonality of mixed type. A remainder. Orthogonal polynomials and the theory of integrable systems has been connected in several ways in the mathematical literature. We are particularly interested in the one based in the Gauss–Borel factorization that was developed in [1]-[5], and applied further in [6]-[8]. These papers set the basis for the method we use in this paper to get an alternative Christoffel–Darboux formula for MOPRL of mixed type.

In the following we extract from [7] the necessary material for the construction of the mentioned alternative Christoffel–Darboux formula. We introduce the moment matrix and recall how the Borel–Gauss factorization leads to multiple orthogonality. Then, we outline how the recursion relations appears by introducing a Jacobi type semi-infinite matrix and recall the reader the Chistoffel–Darboux formula [14, 13].

1.3.1. The moment matrix. Fixed a composition \vec{n}_{α} , $\alpha = 1, 2$, any given $l \in \mathbb{Z}_{+} := \{0, 1, 2, \ldots\}$, see [25], determines uniquely the following non-negative integers $k_{\alpha}(l) \in \mathbb{Z}_{+}$, $a_{\alpha}(l) \in \{1, 2, \ldots, p_{\alpha}\}$ and $r_{\alpha}(l)$ such that $0 \leq r_{\alpha}(l) < n_{\alpha,a_{\alpha}(l)}$ and

(6)
$$l = \begin{cases} k_{\alpha}(l)|\vec{n}_{\alpha}| + n_{\alpha,1} + \dots + n_{\alpha,a_{\alpha}(l)-1} + r_{\alpha}(l), & a_{\alpha}(l) \neq 1, \\ k_{\alpha}(l)|\vec{n}_{\alpha}| + r_{\alpha}(l), & a_{\alpha}(l) = 1. \end{cases}$$

We define now the monomial strings as vectors that may be understood as sequences of monomials according to the composition \vec{n}_{α} , $\alpha = 1, 2$, introduced previously.

$$\chi_{\alpha} := \begin{pmatrix} \chi_{\alpha,[0]} \\ \chi_{\alpha,[1]} \\ \vdots \\ \chi_{\alpha,[k]} \\ \vdots \end{pmatrix} \quad \text{where} \quad \chi_{\alpha,[k]} := \begin{pmatrix} \chi_{\alpha,[k],1} \\ \chi_{\alpha,[k],2} \\ \vdots \\ \chi_{\alpha,[k],a_{\alpha}} \\ \vdots \\ \chi_{\alpha,[k],p_{\alpha}} \end{pmatrix} \quad \text{and} \quad \chi_{\alpha,[k],a_{\alpha}} := \begin{pmatrix} x^{kn_{\alpha,a_{\alpha}}} \\ x^{kn_{\alpha,a_{\alpha}}+1} \\ \vdots \\ x^{kn_{\alpha,a_{\alpha}}+(n_{\alpha,a_{\alpha}}-1)} \end{pmatrix}.$$

In a similar manner for $\alpha = 1, 2$ we define the weighted monomial strings

$$\xi_{\alpha} := \begin{pmatrix} \xi_{\alpha,[0]} \\ \xi_{\alpha,[1]} \\ \vdots \\ \xi_{\alpha,[k]} \\ \vdots \end{pmatrix} \qquad \text{where} \qquad \qquad \xi_{\alpha,[k]} := \begin{pmatrix} w_{\alpha,1}\chi_{\alpha,[k],1} \\ w_{\alpha,2}\chi_{\alpha,[k],2} \\ \vdots \\ w_{\alpha,p_{\alpha}}\chi_{\alpha,[k],p_{\alpha}} \end{pmatrix}.$$

For any given $l \in \mathbb{Z}_+$ and $a_{\alpha} := 1, 2, \dots, p_{\alpha}$ we define

$$\nu_{\alpha,a_{\alpha}}(l) := \begin{cases} k_{\alpha}(l)|\vec{n}_{\alpha}| + n_{\alpha,a_{\alpha}} - 1, & a_{\alpha} < a_{\alpha}(l), \\ k_{\alpha}(l)|\vec{n}_{\alpha}| + r_{\alpha}(l), & a_{\alpha} = a_{\alpha}(l), \\ k_{\alpha}(l)|\vec{n}_{\alpha}| - 1, & a_{\alpha} > a_{\alpha}(l). \end{cases}$$

Notice that $\nu_{\alpha,a_{\alpha}}(l)$ is the hightest degree of all the monomials of type a_{α} up to the component $\chi_{\alpha}^{(l)}$ included, of the monomial string. Actually

$$\chi_{\alpha}^{(l)} = x^{\nu_{\alpha, a_{\alpha}(l)}}.$$

Given $l \geq 1$ and $a_{\alpha} = 1, \cdots, p_{\alpha}$ the + (-) associated integer is the smallest (largest) integer $l_{\{+,a_{\alpha}\}}$ ($l_{\{-,a_{\alpha}\}}$) such that $l_{\{+,a_{\alpha}\}} \geq l$ ($l_{\{-,a_{\alpha}\}} \leq l$) and $a(l_{\{+,a_{\alpha}\}}) = a_{\alpha}$ ($a(l_{\{-,a_{\alpha}\}}) = a_{\alpha}$). It can be shown that

$$l_{\{-,a_{\alpha}\}} := \begin{cases} k_{\alpha}(l)|\vec{n}_{\alpha}| + \sum_{i=1}^{a_{\alpha}} n_{\alpha,i} - 1, & a_{\alpha} < a_{\alpha}(l), \\ l, & a_{\alpha} = a_{\alpha}(l), \\ k_{\alpha}(l)|\vec{n}_{\alpha}| - \sum_{i=a_{\alpha}+1}^{p_{\alpha}} n_{\alpha,i} - 1, & a_{\alpha} > a_{\alpha}(l-1), \end{cases}$$

$$l_{\{+,a_{\alpha}\}} := \begin{cases} (k_{\alpha}(l)+1)|\vec{n}_{\alpha}| + \sum_{i=1}^{a_{\alpha}-1} n_{\alpha,i}, & a_{\alpha} < a_{\alpha}(l), \\ l, & a_{\alpha} = a_{\alpha}(l), \\ (k_{\alpha}(l)+1)|\vec{n}_{\alpha}| - \sum_{i=a_{\alpha}}^{p_{\alpha}} n_{\alpha,i}, & a_{\alpha} > a_{\alpha}(l). \end{cases}$$

Finally given the weighted monomials $\xi_{\vec{n}_{\alpha}}$, associated to the compositions \vec{n}_{α} , $\alpha = 1, 2$, we introduce the moment matrix in the following manner

Definition 1. The moment matrix is given by

(8)
$$g_{\vec{n}_1,\vec{n}_2} := \int \xi_{\vec{n}_1}(x)\xi_{\vec{n}_2}(x)^{\top} d\mu(x).$$

1.3.2. Multiple Orthogonality of mixed type.

Definition 2. For a given a perfect combination $(\mu, \vec{w}_1, \vec{w}_2)$ we define

(1) The Gauss-Borel factorization (also known as LU factorization) of a semi-infinite moment matrix g, determined by $(\mu, \vec{w}_1, \vec{w}_2)$, is the problem of finding the solution of

$$(9) g = S^{-1}\bar{S}, S = \begin{pmatrix} 1 & 0 & 0 & \cdots \\ S_{1,0} & 1 & 0 & \cdots \\ S_{2,0} & S_{2,1} & 1 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \in G_{-}, \bar{S}^{-1} = \begin{pmatrix} \bar{S}'_{0,0} & \bar{S}'_{0,1} & \bar{S}'_{0,2} & \cdots \\ 0 & \bar{S}'_{1,1} & \bar{S}'_{1,2} & \cdots \\ 0 & 0 & \bar{S}'_{2,2} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \in G_{+},$$

where $S_{i,j}, \bar{S}'_{i,j} \in \mathbb{R}$.

(2) In terms of these matrices we construct the polynomials

(10)
$$A_a^{(l)} := \sum_{i}' S_{l,i} x^{k_1(i)},$$

where the sum \sum' is taken for a fixed $a=1,\ldots,p_1$ over those i such that $a=a_1(i)$ and $i\leq l$. We also construct the dual polynomials

(11)
$$\bar{A}_b^{(l)} := \sum_{j}' x^{k_2(j)} \bar{S}'_{j,l},$$

where the sum \sum' is taken for a given b over those j such that $b = a_2(j)$ and $j \leq l$.

(3) Strings of linear forms and dual linear forms associated with multiple ortogonal polynomials and their duals are defined by

(12)
$$Q := \begin{pmatrix} Q^{(0)} \\ Q^{(1)} \\ \vdots \end{pmatrix} = S\xi_1, \qquad \bar{Q} := \begin{pmatrix} \bar{Q}^{(0)} \\ \bar{Q}^{(1)} \\ \vdots \end{pmatrix} = (\bar{S}^{-1})^{\top} \xi_2,$$

Then

Proposition 1. (1) The linear forms and their duals, introduced in Definition 2, are given by

(13)
$$Q^{(l)}(x) := \sum_{a=1}^{p_1} A_a^{(l)}(x) w_{1,a}(x), \qquad \bar{Q}^{(l)}(x) := \sum_{b=1}^{p_2} \bar{A}_b^{(l)}(x) w_{2,b}(x).$$

(2) The orthogonality relations

(14)
$$\int Q^{(l)}(x)w_{2,b}(x)x^k d\mu(x) = 0, \quad 0 \le k \le \nu_{2,b}(l-1) - 1, \quad b = 1, \dots, p_2,$$
$$\int \bar{Q}^{(l)}(x)w_{1,a}(x)x^k d\mu(x) = 0, \quad 0 \le k \le \nu_{1,a}(l-1) - 1, \quad a = 1, \dots, p_1,$$

are fulfilled.

(3) The following multiple bi-orthogonality relations among linear forms and their duals

(15)
$$\int Q^{(l)}(x)\bar{Q}^{(k)}(x)\mathrm{d}\mu(x) = \delta_{l,k}, \qquad l, k \ge 0,$$

hold.

(4) We have the following identifications

$$A_a^{(l)} = A_{[\vec{\nu}_1(l);\vec{\nu}_2(l-1)],a}^{(\mathrm{II},a_1(l))}, \qquad \qquad \bar{A}_b^{(l)} = A_{[\vec{\nu}_2(l);\vec{\nu}_1(l-1)],b}^{(\mathrm{II},a_1(l))},$$

in terms of multiple orthogonal polynomials of mixed type with two normalizations I and II, respectively.

1.3.3. Functions of the second kind. The Cauchy transforms of the linear forms (13) play a crucial role in the Riemann–Hilbert problem associated with the multiple orthogonal polynomials of mixed type [13]. Observe that the construction of multiple orthogonal polynomials performed so far is synthesized in the following strings of multiple orthogonal polynomials and their duals

(16)
$$\mathscr{A}_{a} := \begin{pmatrix} A_{a}^{(0)} \\ A_{a}^{(1)} \\ \vdots \end{pmatrix} = S\chi_{1,a}, \quad \bar{\mathscr{A}_{b}} := \begin{pmatrix} \bar{A}_{b}^{(0)} \\ \bar{A}_{b}^{(1)} \\ \vdots \end{pmatrix} = (\bar{S}^{-1})^{\top}\chi_{2,b},$$

for $a=1,\ldots,p_1$ and $b=1,\ldots,p_2$. In order to complete these formulae and in terms of $\chi_a^*:=z^{-1}\chi_a(z^{-1})$ let us introduce the following formal semi-infinite vectors

(17)
$$\mathscr{C}_{b} = \begin{pmatrix} C_{b}^{(0)} \\ C_{b}^{(1)} \\ \vdots \end{pmatrix} = \bar{S}\chi_{2,b}^{*}(z), \quad \bar{\mathscr{C}}_{a} = \begin{pmatrix} \bar{C}_{a}^{(0)} \\ \bar{C}_{a}^{(1)} \\ \vdots \end{pmatrix} = (S^{-1})^{\top}\chi_{1,a}^{*}(z),$$

for $a=1,\ldots,p_1$ and $b=1,\ldots,p_2$, that we call strings of second kind functions. These objects are actually Cauchy transforms of the linear forms $Q^{(l)}$, $l \in \mathbb{Z}_+$, whenever the series converge and outside the support of the measures involved.

Proposition 2. For each $l \in \mathbb{Z}_+$ the second kind functions can be expressed as follows

(18)
$$C_b^{(l)}(z) = \int_{\mathbb{R}} \frac{Q^{(l)}(x)w_{2,b}(x)}{z - x} d\mu(x), \quad z \in D_b^{(l)} \setminus \operatorname{supp}(w_{1,b}d\mu(x)),$$
$$\bar{C}_a^{(l)}(z) = \int_{\mathbb{R}} \frac{\bar{Q}^{(l)}(x)w_{1,a}(x)}{z - x} d\mu(x), \quad z \in \bar{D}_a^{(l)} \setminus \operatorname{supp}(w_{2,a}d\mu(x)).$$

1.3.4. Recursion relations, a Jacobi type matrix. The moment matrix has a Hankel type symmetry that implies the recursion relations and the Christoffel–Darboux formula. We consider the shift operators Υ_{α} defined by

$$(\Upsilon_{\alpha})_{l,j} := \delta_{j,(l+1)_{\{+,a_{\alpha}(l)\}}}$$

Wich satisfy the following relation

$$\Upsilon_{\alpha}\chi_{\alpha}(x) = x\chi_{\alpha}(x) \Longrightarrow \Upsilon_{\alpha}\xi_{\alpha}(x) = x\xi_{\alpha}(x)$$

In terms of these shift matrices we can describe the particular Hankel symmetries for the moment matrix

Proposition 3. The moment matrix q satisfies the Hankel type symmetry

$$\Upsilon_1 g = g \Upsilon_2^{\top}.$$

From this symmetry we see that the following is consistent

Definition 3. We define the matrices

$$J := S\Upsilon_1 S^{-1} = \bar{S}\Upsilon_2^\top \bar{S}^{-1} = J_+ + J_-, \qquad \qquad J_+ := (S\Upsilon_1 S^{-1})_+, \qquad \qquad J_- := (\bar{S}\Upsilon_2^\top \bar{S}^{-1})_-,$$

where the sub-indices + and -denote the upper triangular and strictly lower triangular projections.

The recursion relations follow immediately from the eigenvalue property

(21)
$$JQ(x) = xQ(x) \qquad \bar{Q}(x)^{\top} J = x\bar{Q}(x)^{\top}.$$

 $1.3.5. \ \ Christoffel-Darboux\ formula.$

Definition 4. The Christoffel–Darboux kernel is

(22)
$$K^{[l]}(x,y) := \sum_{k=0}^{l-1} Q^{(k)}(y)\bar{Q}^{(k)}(x)$$

In [14, 13] it was shown using a Riemann–Hilbert problem approach that

Proposition 4. For $l \ge \max(|\vec{n}_1|, |\vec{n}_2|)$ the following Christoffel–Darboux formula

$$(x-y)K^{[l]}(x,y) = \sum_{b=1}^{p_2} \bar{Q}^{(\text{II},b)}_{[\vec{\nu}_2(l-1)+\vec{e}_{2,b};\vec{\nu}_1(l-1)]}(x)Q^{(\text{I},b)}_{[\vec{\nu}_1(l-1);\vec{\nu}_2(l-1)-\vec{e}_{2,b}]}(y)$$

$$-\sum_{a=1}^{p_1} \bar{Q}^{(\text{I},a)}_{[\vec{\nu}_2(l-1);\vec{\nu}_1(l-1)-\vec{e}_{1,a}]}(x)Q^{(\text{II},a)}_{[\vec{\nu}_1(l-1)+\vec{e}_{1,a};\vec{\nu}_2(l-1)]}(y).$$

holds.

Here $\{\vec{e}_{i,a}\}_{a=1}^{p_i} \subset \mathbb{R}^{p_i}$ stands for the vectors in the respective canonical basis, i = 1, 2. In [7] it was given an algebraic proof of this statement not relying on analytic conditions. We refer the interested reader to [23] for a complete survey of the subject.

2. Alternative Christoffel–Darboux formula for multiple orthogonal polynomials of mixed type

The result of this paper is the following

Theorem 1. For $l \ge \max\{|\vec{n}_1|, |\vec{n}_2|\}$ the following Christofel–Darboux formula holds

$$(y-x)K^{[l]}(x,y) = \sum_{(i,j)\in\sigma_1[l]} \bar{Q}(x)^{(j)} J_{j,i} Q(y)^{(i)} - \sum_{(i,j)\in\sigma_2[l]} \bar{Q}(x)^{(j)} J_{j,i} Q(y)^{(i)}$$

where

$$\sigma_1[l] := \{l, \dots, (l)_{\{+, r_1(a_1(l)-1)\}}\} \times \{(l-1)_{\{-, r_1(a_1(l-1)+1)\}}, \dots, l-1\},$$

$$\sigma_2[l] := \{(l-1)_{\{-, r_2(a_2(l-1)+1)\}}, \dots, l-1\} \times \{l, \dots, (l)_{\{+, r_2(a_2(l)-1)\}}\}.$$

Proof. Splitting the eigenvalue property (21) into blocks we get

$$JQ(y) = yQ(y) \Longrightarrow J^{[l]}Q(y)^{[l]} + J^{[l,\geq l]}Q(y)^{[\geq l]} = yQ(y)^{[l]}$$
$$\bar{Q}(x)^{\top}J = x\bar{Q}(x)^{\top} \Longrightarrow [\bar{Q}(x)^{\top}]^{[l]}J^{[l]} + [\bar{Q}(x)^{\top}]^{[\geq l]}J^{[\geq l,l]} = x[\bar{Q}(x)^{\top}]^{[l]}$$

Multiply the first equation from the left by $[\bar{Q}(x)^{\top}]^{[l]}$ and the second one from the right by $Q(y)^{[l]}$ substract both results to obtain

$$\begin{split} [\bar{Q}(x)^{\top}]^{[l]}J^{[l,\geq l]}Q(y)^{[\geq l]} - [\bar{Q}(x)^{\top}]^{[\geq l]}J^{[\geq l,l]}Q(y)^{[l]} &= (y-x)[\bar{Q}(x)^{\top}]^{[l]} \cdot Q(y)^{[l]} \\ &= (y-x)K^{[l]}(x,y) \end{split}$$

After an brief study of the shape of J we realize that even though $J^{[l,\geq l]}$ has semi-infinte length rows, most of its elements are 0. Actually it only contains a finite number of nonzero entries that concentrate in the lower left corner of itself. The same reasoning applies to $J^{[\geq l,l]}$. This matrix has semi-infinite length columns but again it only contains a finite number of nonzero terms concentrated in the upper right corner of itself. Of course the

number of terms involved in this expression will depend on the value of [l]. To be more precise we proceed as follows. From the Euclidean division we know that for any positive integer $l \in \mathbb{Z}_+$ there exists unique integers $q_i, r_i, i = 1, 2$, the quotient and remainder, such that

$$l = q_i p_i + r_i,$$
 $0 \le r_i < p_i,$ $i = 1, 2.$

After a study of the shape of J we can state

Lemma 1. For $l \ge \max\{|\vec{n}_1|, |\vec{n}_2|\}$ the only nonzero elements of J along a given row or column are

$$J_{(l-1)_{\{-,r_1(a_1(l-1)+1)\}},l} \\ \vdots \\ \vdots \\ J_{l,(l-1)_{\{-,r_2(a_2(l-1)-1)\}}} \\ * \\ \cdots \\ * \\ J_{l,l} \\ * \\ \cdots \\ * \\ J_{l,(l+1)_{\{+,r_1(a_1(l+1)-1)\}}} \\ \vdots \\ * \\ J_{(l+1)_{\{+,r_2(a_2(l+1)-1)\}},l}$$

Using this Lemma we get the desired result and the proof is complete.

Example. In order to be more clear let us suppose that $p_1 = 3$ and $p_2 = 2$ with $\vec{n}_1 = (4,3,2)$ and $\vec{n}_2 = (3,2)$. The corresponding Jacobi type matrix has the following shape

where * denotes a non-necessarily null real number.

In our example $(p_1 = 3, p_2 = 2, \vec{n}_1 = (4, 3, 2) \text{ and } \vec{n}_2 = (3, 2))$ for [l] = [12] we have

$$(y-x)K^{[12]}(x,y) = \left[\sum_{i=8}^{11} \sum_{j=12}^{15} \bar{Q}(x)^{(i)} J_{i,j} Q(y)^{(j)}\right] - \left[\sum_{i=12}^{13} \sum_{j=9}^{11} \bar{Q}(x)^{(i)} J_{i,j} Q(y)^{(j)}\right]$$

2.1. Expressing the Jacobi type matrix in terms of factorization factors. As we have seen we can write J in terms of S or of \bar{S} , this means that for each term of J has two different expressions, giving relations between S with \bar{S} . We are not too concerned about these relations since what we want here is the most simple expression we can get for the elements of J. It is easy to realize that this is achieved if we use the expression involving S in order to calculate the upper part of J and the expression involving \bar{S} to calculate the lower part of it. Hence, for every $J_{l,k}$ we will have expressions in terms of the factorization matrices coefficients and the elements of their inverses –thus, in terms of the MOPRL and associated second kind functions. The only terms from the factorization matrices (or their inverses) that will be involved when calculating any $J_{l,k}$ are just those between the main diagonal and the $l - |n_1|$ diagonal (both included) of S and those between the main diagonal and the $l + |n_2|$ diagonal (both included) of S. And not even all of them. As we are about to see there are three different kinds of elements in J. The ones along the main diagonal, the ones along the immediate closest diagonals to the main one, and finally all the remaining diagonals. The recursion relation coefficients $J_{k,l}$ are ultimately related to the MOPRL and its associated second kind functions in the following way

Proposition 5. The elements of the recursion matrix J can be written in terms of products of the entries of the LU factorization matrices and its inverses as follows

$$\begin{split} J_{l,l} &= S_{l,(l-1)_{\{-,a_1(l)\}}} + S_{(l+1)_{\{+,a_1(l)\}},l}^{-1} + \sum_{\substack{a=1,\ldots,p_1\\ a\neq a_1(l)}} S_{l,(l-1)_{\{-,a\}}} S_{(l+1)_{\{+,a\}},l}^{-1}, \\ &= \bar{S}_{l,(l+1)_{\{+,a_2(l)\}}} \bar{S}_{l,l}^{-1} + \bar{S}_{l,l} \bar{S}_{(l-1)_{\{-,a_2(l)\}},l}^{-1} + \sum_{\substack{a=1,\ldots,p_2\\ a\neq a_2(l)}} \bar{S}_{l,(l+1)_{\{+,a\}}} \bar{S}_{(l-1)_{\{-,a\}},l}^{-1}, \\ J_{l,l+1} &= S_{(l+1)_{\{+,a_1(l)\}},l+1}^{-1} + \sum_{\substack{a=1,\ldots,p_1\\ a\neq a_1(l)}} S_{l,(l-1)_{\{-,a\}}} S_{(l+1)_{\{+,a\}},l+1}^{-1}, \\ J_{l+1,l} &= \bar{S}_{l+1,(l+1)_{\{+,a_2(l)\}}} \bar{S}_{l,l}^{-1} + \sum_{\substack{a=1,\ldots,p_2\\ a\neq a_2(l)}} \bar{S}_{l+1,(l+1)_{\{+,a\}}} \bar{S}_{(l-1)_{\{-,a\}},l}^{-1}, \\ J_{l,l+k} &= \sum_{\substack{a=r_1(a_1(l)-1)\\ a=r_1(a_1(l+k-1)+1)}}^{r_1(a_1(l)-1)} S_{l,(l-1)_{\{-,a_1\}}} S_{(l+1)_{\{+,a_1\}},l+k}^{-1} \quad 2 \leq k \leq (l+1)_{\{+,r_1(a_1(l+1)-1)\}} - l, \\ J_{l+k,l} &= \sum_{\substack{a=r_2(a_2(l)+k-1)+1)}}^{r_1(a_2(l)-1)} \bar{S}_{l+k,(l+1)_{\{+,a\}}} \bar{S}_{(l-1)_{\{-,a\}},l}^{-1}, \quad 2 \leq k \leq (l+1)_{\{+,r_2(a_2(l+1)-1)\}} - l. \end{split}$$

Where, for r, r' < p, we have used

$$\sum_{a=r}^{r'} X_a = \begin{cases} \sum_{a=r}^{r'} X_a, & r \le r', \\ \sum_{a=1}^{r'} X_a + \sum_{a=r}^{p} X_a, & r > r'. \end{cases}$$

ACKNOWLEDGEMENTS

GA thanks economical support from the Universidad Complutense de Madrid Program "Ayudas para Becas y Contratos Complutenses Predoctorales en España 2011". MM thanks economical support from the Spanish "Ministerio de Economía y Competitividad" research project MTM2012-36732-C03-01, Ortogonalidad y aproximacion; Teoria y Aplicaciones.

References

- [1] M. Adler and P. van Moerbeke, Group factorization, moment matrices and Toda lattices, International Mathematics Research Notices 12 (1997) 556-572.
- [2] M. Adler and P. van Moerbeke, Generalized orthogonal polynomials, discrete KP and Riemann–Hilbert problems, Communications in Mathematical Physics 207 (1999) 589-620.
- [3] M. Adler and P. van Moerbeke, Vertex operator solutions to the discrete KP hierarchy, Communications in Mathematical Physics 203 (1999) 185-210.
- [4] M. Adler and P. van Moerbeke, The spectrum of coupled random matrices, Annals of Mathematics 149 (1999) 921-976.
- [5] M. Adler and P. van Moerbeke, Darboux transforms on band matrices, weights and associated polynomials, International Mathematics Research Notices 18 (2001) 935-984.
- [6] C. Álvarez-Fernández, U. Fidalgo and M. Mañas, The multicomponent 2D Toda hierarchy: generalized matrix orthogonal polynomials, multiple orthogonal polynomials and Riemann–Hilbert problems. Inverse Problems 26 (2010) 055009 (17 pp).
- [7] C. Álvarez-Fernández, U. Fidalgo and M. Mañas, Multiple orthogonal polynomials of mixed type: Gauss-Borel factorization and the multi-component 2D Toda hierarchy. Advances in Mathematics 227 (2011) 1451-1525.
- [8] C. Álvarez-Fernández and M. Mañas, Orthogonal Laurent polynomials on the unit circle, extended CMV ordering and 2D Toda type integrable hierarchies, Advances in Mathematics 240 (2013) 132-193.
- [9] R. Apery. Irrationalite de $\zeta(2)$ et $\zeta(3)$, Astèrisque **61** (1979) 11-13.
- [10] F. Beukers, Padé approximation in number theory, Lecture Notes in Mathematics 888, Springer Verlag, Berlin, 1981, 90-99.
- [11] P.M. Bleher and A.B.J. Kuijlaars, Random matrices with external source and multiple orthogonal polynomials, International Mathematics Research Notices 2004 (2004), 109-129.
- [12] J. Coates, On the algebraic approximation of functions, I, II, III. Indagationes Mathematicae 28 (1966) 421-461.
- [13] E. Daems and A. B. J. Kuijlaars, Multiple orthogonal polynomials of mixed type and non-intersecting Brownian motions, Journal of Approximation Theory 146 (2007) 91-114.
- [14] E. Daems and A. B. J. Kuijlaars, A Christoffel-Darboux formula for multiple orthogonal polynomials, Journal of Approximation Theory 130 (2004) 188-200.
- [15] U. Fidalgo Prieto and G. López Lagomasino, Nikishin systems are perfect, Constructive Approximation 34 (2011) 297-356.
- [16] A. S. Fokas, A. R. Its and A. V. Kitaev, The isomonodromy approach to matrix models in 2D quatum gravity, Communications in Mathematical Physics (1992) 395-430.
- [17] Ch. Hermite, Sur la fonction exponentielle, C. R. Acad. Sci. Paris 77 (1873), 18-24, 74-79, 226-233, 285-293; reprinted in his Oeuvres, Tome III, Gauthier-Villars, Paris, 1912, 150-181.
- [18] H. Jager, A simultaneous generalization of the Padé table, I-VI, Indagationes Mathematicae 26 (1964), 193-249.
- [19] A. B. J. Kuijlaars, Multiple orthogonal polynomial ensembles, Contemporary Mathematics 507 (2010), 155–176
- [20] K. Mahler, Perfect systems, Compositio Mathematica 19 (1968), 95-166.
- [21] E. M. Nikishin, On simultaneous Padé approximants Matematicheskii Sbornik 113 (1980), 499-519 (Russian); English translation in Mathematics of the USSR-Sbornik 41 (1982), 409-425.
- [22] J. A. Shohat and J.D. Tamarkin, The problem of moments, American Mathematical Society (1943).
- [23] B. Simon, The Christoffel-Darboux Kernel, Proceedings of Symposia in Pure Mathematics 79: "Perspectives in Partial Differential Equations, Harmonic Analysis and Applications: A Volume in Honor of Vladimir G. Maz'ya's 70th Birthday", (2008) 295-336. arXiv:0806.1528
- [24] V.N. Sorokin, On simultaneous approximation of several linear forms, Vestnik Moskovskogo Universiteta. Seriya I. Matematika 1 (1983) 44-47.
- [25] R. P. Stanley, Enumerative combinatorics, Cambridge University Press, Cambridge (1998).
- [26] W. Van Assche, J. S. Geromino and A. B. J. Kuijlaars, Riemann-Hilbert problems for multiple orthogonal polynomials in: Bustoz et al (eds.), Special Functions 2000: Current Perspectives and Future Directions, Kluwer Academic Publishers, Dordrecht, 2001, pp 23-59.

DEPARTAMENTO DE FÍSICA TEÓRICA II (MÉTODOS MATEMÁTICOS DE LA FÍSICA), UNIVERSIDAD COMPLUTENSE DE MADRID, 28040-MADRID, SPAIN

 $E ext{-}mail\ address: gariznab@ucm.es}$

E-mail address: manuel.manas@ucm.es