

# Well-posedness of first order semilinear PDEs by stochastic perturbation

Christian Olivera <sup>1</sup>

*Departamento de Matemática, Universidade Estadual de Campinas,  
F. 54(19) 3521-5921 ? Fax 54(19) 3521-6094  
13.081-970 - Campinas - SP, Brazil. ; colivera@ime.unicamp.br*

## Abstract

We show that first order semilinear PDEs by stochastic perturbation are well-posedness for globally Holder continuous and bounded vector field, with an integrability condition on the divergence. This result extends the linear case presented in [2]. The proof is based on in the stochastic characteristics method and a version of the commuting Lemma.

**Key words:** Stochastic characteristic method, First order stochastic partial differential equations, Stochastic perturbation, commuting Lemma.

**MSC2000 subject classification:** 60H10 , 60H15.

## 1 Introduction

This work is motivated by the paper [2] where the linear equation

$$\begin{cases} du(t, x) + b(t, x)\nabla u(t, x)dt + \nabla u(t, x) \circ dB_t = 0, \\ u(0, x) = f(x) \in L^\infty(\mathbb{R}^d), \end{cases} \quad (1)$$

---

<sup>1</sup>Research partially supported by FAPESP 2012/18739-0.

has been studied, was proved existence and uniqueness of  $L^\infty$ - solutions for a globally Holder continuous and bounded vector field, with an integrability condition on the divergence, and where  $B_t = (B_t^1, \dots, B_t^d)$  is a standard Brownian motion in  $\mathbb{R}^d$ .

The aim of this paper is to investigate parts of this theory under the effect of nonlinear terms. Namely, we consider the semilinear SPDE

$$\begin{cases} du(t, x) + b(t, x)\nabla u(t, x) dt + F(t, x, u) dt + \nabla u(t, x) \circ dB_t = 0, \\ u(0, x) = f(x) \in L^\infty(\mathbb{R}^d). \end{cases} \quad (2)$$

We shall prove the existence and uniqueness of weak  $L^\infty$ - solutions for a globally Holder continuous and bounded vector field, with an integrability condition on the divergence. Moreover, we obtain a representation of the solution via stochastic flows. This is an example of nonlinear SPDE where the stochastic perturbation makes the equation well-posedness.

The fundamental tools used here is the stochastic characteristics method (see for example [1], [5] and [7]) and the version of the commuting Lemma presented in [2]. That is, we follow the strategy given in [2] in combination with the stochastic characteristics method.

The article is organized as follows: Section 2 we shall define the concept of weak  $L^\infty$ -solutions for the equation (2) and we shall prove existence for this class of solutions. In section 3, we shall show a uniqueness theorem for weak  $L^\infty$ -solutions.

Through of this paper we fix a stochastic basis with a  $d$ -dimensional Brownian motion  $(\Omega, \mathcal{F}, \{\mathcal{F}_t : t \in [0, T]\}, \mathbb{P}, (B_t))$ .

## 2 Existence of weak $L^\infty$ -solutions

Let  $T > 0$  be fixed. For  $\alpha \in (0, 1)$  define the space  $L^\infty([0, T], C^\alpha(\mathbb{R}^d))$  as the set of all bounded Borel functions  $f : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$  for which

$$[f]_{\alpha,T} = \sup_{t \in [0,T]} \sup_{x \neq y \in \mathbb{R}^d} \frac{|f(x) - f(y)|}{|x - y|}$$

We write the  $L^\infty([0, T], C^\alpha(\mathbb{R}^d, \mathbb{R}^d))$  for the space of all vector fields having components in  $L^\infty([0, T], C^\alpha(\mathbb{R}^d))$ .

We shall assume that

$$b \in L^\infty([0, T], C^\alpha(\mathbb{R}^d, \mathbb{R}^d)), \quad (3)$$

$$\text{Div } b \in L^p([0, T] \times \mathbb{R}^d) \text{ for } p > 2. \quad (4)$$

$$F \in L^1([0, T], L^\infty(\mathbb{R}^d \times \mathbb{R})) \quad (5)$$

and

$$F \in L^\infty([0, T] \times \mathbb{R}^d, LIP(\mathbb{R})). \quad (6)$$

## 2.1 Definition of weak $L^\infty$ -solutions

**Definition 2.1** *We assume (3), (4), (5) and (6). A weak  $L^\infty$ -solution of the Cauchy problem (2) is a stochastic process  $u \in L^\infty(\Omega \times [0, T] \times \mathbb{R}^d)$  such that, for every test function  $\varphi \in C_0^\infty(\mathbb{R}^d)$ , the process  $\int u(t, x)\varphi(x)dx$  has a continuous modification which is a  $\mathcal{F}_t$ -semimartingale and satisfies*

$$\begin{aligned} & \int u(t, x)\varphi(x)dx = \int f(x)\varphi(x) dx \\ & + \int_0^t \int b(s, x)\nabla\varphi(x)u(s, x) dx ds + \int_0^t \int \text{div } b(s, x)\varphi(x)u(s, x) dx ds \\ & + \int_0^t \int F(s, x, u)\varphi(x) dx ds + \sum_{i=0}^d \int_0^t \int D_i\varphi(x)u(s, x) dx \circ dB_s^i \end{aligned}$$

**Remark 2.1** We observe that a weak  $L^\infty$ -solution in the previous Stratonovich sense satisfies the Itô equation

$$\begin{aligned}
& \int u(t, x)\varphi(x)dx = \int f(x)\varphi(x) dx \\
& + \int_0^t \int b(s, x)\nabla\varphi(x)u(s, x) dx ds + \int_0^t \int \operatorname{div} b(s, x)\varphi(x)u(s, x) dx ds \\
& + \int_0^t \int F(s, x, u)\varphi(x) dx ds + \sum_{i=0}^d \int_0^t \int D_i\varphi(x)u(s, x) dx dB_s^i + \frac{1}{2} \int_0^t u(s, x)\Delta\varphi(x) dx ds
\end{aligned} \tag{7}$$

for every test function  $\varphi \in C_0^\infty(\mathbb{R}^d)$ . The converse is also true.

## 2.2 Existence of weak $L^\infty$ -solutions

**Lemma 2.1** Let  $f \in L^\infty(\mathbb{R}^d)$ . We assume (3), (4), (5) and (6). Then there exists a weak  $L^\infty$ -solution  $u$  of the SPDE (2).

**Proof:** Step 1 Assume that  $F \in L^1([0, T], C_b^\infty(\mathbb{R}^d \times \mathbb{R}))$  and  $f \in C_b^\infty(\mathbb{R}^d)$ . We take a mollifier regularization  $b_n$  of  $b$ . It is known (see [1], chapter 1) that there exist a unique classical solution  $u_n(t, x)$  of the SPDE (2), that written in weak Itô form is (7) with  $b_n$  in place of  $b$ . Moreover,

$$u_n(t, x) = Z_t^n(x, f(Y_t^n))$$

where  $Y_t^n$  is the inverse of  $X_t^n$ ,  $X_t^n(x)$  and  $Z_t^n(x, r)$  satisfy the following equations

$$X_t^n = x + \int_0^t b_n(s, X_s^n) ds + B_t, \tag{8}$$

and

$$Z_t^n = r + \int_0^t F(s, X_s^n(x), Z_s^n) ds. \tag{9}$$

According to theorem 5 of [2], see too remark 8, we have that

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[ \int_K \sup_{t \in [0, T]} |X_t^n - X_t| dx \right] = 0$$

and

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[ \int_K \sup_{t \in [0, T]} |DX_t^n - DX_t| dx \right] = 0$$

for any compact set  $K \subset \mathbb{R}^d$ , where  $X_t(x)$  verifies

$$X_t = x + \int_0^t b(s, X_s) ds + B_t. \quad (10)$$

Now, we denote

$$u(t, x) = Z_t(x, f(Y_t)),$$

$Y_t$  is the inverse of  $X_t$ ,

and

$$Z_t = r + \int_0^t F(s, X_s(x), Z_s) ds. \quad (11)$$

Then , we observe that

$$\begin{aligned} |u^n(t, x) - u(t, x)| &\leq |f(Y_t) - f(Y_t^n)| + \int_0^t |F(s, X_s^n, Z_s^n(f(Y_t^n))) - F(s, X_s, Z_s(f(Y_t)))| ds \\ &\leq |f(Y_t) - f(Y_t^n)| + C \int_0^t |Z_s^n(f(Y_t^n)) - Z_s(f(Y_t))| ds. \end{aligned}$$

From to theorem 5 of [2], see too remark 8, and the Lipchitz property of  $F$  we conclude that  $\lim_{n \rightarrow \infty} \mathbb{E}[\int_K \sup_{t \in [0, T]} |u_n(t, x) - u(t, x)|] = 0$  and  $u(t, x)$  is a weak  $L^\infty$ -solution of the SPDE (2).

**Step 2** Assume that  $F \in L^1([0, T], C_b^\infty(\mathbb{R}^d \times \mathbb{R}))$ . We a take a mollifier regularization  $f_n$  of  $f$  . By the last step  $u_n(t, x) = Z_t(x, f_n(Y_t))$  is a weak

$L^\infty$ -solution of the SPDE (2), that written in weak Itô form is (7) with  $f_n$  in place of  $f$ .

We have that any compact set  $K \subset \mathbb{R}^d$  and  $p \geq 1$

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup_{[0, T]} \int_K |f_n(X_t^{-1}) - f(X_t^{-1})|^p dx = \\ \lim_{n \rightarrow \infty} \sup_{[0, T]} \int_{X_t(K)} |f_n(x) - f(x)|^p JX_t(x) dx = 0 \end{aligned}$$

Then we have that

$$\lim_{n \rightarrow \infty} \sup_{[0, T]} \int_K |Z_t(x, f_n(Y_t)) - Z_t(x, f(Y_t))|^p dx = 0.$$

Thus  $u(t, x) = Z_t(x, f(Y_t))$  is a weak  $L^\infty$ -solution of the SPDE (2)

**Step 3** We take a mollifier regularization  $F_n$  of  $F$ . By the step 2, we have that  $u_n(t, x) = Z_t^n(x, f(Y_t))$  is a weak  $L^\infty$ -solution of the SPDE (2), and hold that  $Z_t^n(x, r)$  satisfies the equation (11) with  $F_n$  in place of  $F$ .

We observe that

$$\begin{aligned} |Z_t^n(x, r) - Z_t(x, r)| &\leq \int_0^t |F_n(t, X_s, Z_s^n) - F(t, X_s, Z_s)| ds \\ &\leq \int_0^t |F_n(t, X_s, Z_s^n) - F_n(t, X_s, Z_s)| ds + \int_0^t |F_n(t, X_s, Z_s) - F(t, X_s, Z_s)| ds \\ &\leq C \int_0^t |Z_s^n - Z_s| ds + \int_0^t |F_n(t, X_s, Z_s) - F(t, X_s, Z_s)| ds \end{aligned}$$

By the Gronwall Lemma we follow that

$$\lim_{n \rightarrow \infty} |Z_t^n(x, r) - Z_t(x, r)| = 0 \text{ uniformly in } t, x, r.$$

Then

$$\lim_{n \rightarrow \infty} |Z_t^n(x, f(Y_t)) - Z_t(x, f(Y_t))| = 0 \text{ uniformly in } t \text{ and } x.$$

Therefore, we conclude that  $u(t, x) = Z_t(x, f(Y_t))$  is a weak  $L^\infty$ -solution of the SPDE (2).

### 3 Uniqueness of weak $L^\infty$ -solutions

In this section, we shall present an uniqueness theorem for the SPDE (2) under similar conditions to the linear case, see theorem 20 of [2].

Let  $\varphi_n$  be a standard mollifier. We introduced the commutator defined as

$$\mathcal{R}_n(b, u) = (b\nabla)(\varphi_n * u) - \varphi_n * ((b\nabla)u)$$

We recall here the following version of the commutator lemma which is at the base of our uniqueness theorem.

**Lemma 3.1** *Let  $\phi_t$  be an  $C^1$ -diffeomorphism of  $\mathbb{R}^d$ . Assume  $b \in L^\infty_{loc}(\mathbb{R}^d, \mathbb{R}^d)$ ,  $divb \in L^1_{loc}(\mathbb{R}^d)$ ,  $u \in L^\infty_{loc}(\mathbb{R}^d)$ . Moreover, for  $d > 1$ , assume also  $J\phi^{-1} \in W^{1,1}_{loc}(\mathbb{R}^d)$ . Then for any  $\rho \in C^\infty_0(\mathbb{R}^d)$  there exists a constant  $C_\rho$  such that, given any  $R > 0$  such that  $supp(\rho \circ \phi^{-1}) \subset B(R)$ , we have :*

a) for  $d > 1$

$$\begin{aligned} & \left| \int \mathcal{R}_n(b, u)(\phi(x))\rho(x) dx \right| \\ & \leq C_\rho \|u\|_{L^\infty_{R+1}} \left[ \|divb\|_{L^1_{R+1}} \|J\phi^{-1}\|_{L^\infty_R} + \|b\|_{L^\infty_{R+1}} (\|D\phi^{-1}\|_{L^\infty_R} + \|DJ\phi^{-1}\|_{L^1_R}) \right] \end{aligned}$$

b) for  $d = 1$

$$\left| \int \mathcal{R}_n(b, u)(\phi(x))\rho(x) dx \right| \leq C_\rho \|u\|_{L^\infty_{R+2}} \|b\|_{W^{1,1}_{R+2}} \|J\phi^{-1}\|_{L^\infty_R}$$

**Proof:** See pp 28 of [2].

We are ready to prove our uniqueness result of weak  $L^\infty$ -solution to the Cauchy problem (2).

**Theorem 3.1** *Assume (3), (4), (5) and (6). Then, for every  $f \in L^\infty(\mathbb{R}^d)$  there exists an unique weak  $L^\infty$ -solution of the Cauchy problem (2).*

**Proof:**

Step 1( Itô-Ventzel-Kunita formula) Let  $u, v$  be are two weak  $L^\infty$ -solutions and  $\varphi_n$  be a standard mollifier. We put  $w = u - v$ , applying the Itô-Ventzel-Kunita formula (see Theorem 8.3 of [6] ) to  $F(y) = \int w(t, z)\varphi_n(y - z) dz$ , we obtain that

$$\int w(t, z)\varphi_n(X_s - z)dz$$

is equal to

$$\begin{aligned} & \int_0^t \int b(s, z)\nabla[\varphi_n(X_s - z)]w(s, z) dz ds + \int_0^t \int \operatorname{div} b(s, z)\varphi_n(X_s - z)u(s, z) dz ds + \\ & \int_0^t \int (F(s, z, u) - F(s, z, v))\varphi_n(X_s - z) dz ds + \sum_{i=1}^d \int_0^t \int w(s, z)D_i[\varphi_n(X_s - z)]dz \circ dB_s^i + \\ & \int_0^t \int (b\nabla)(w(s, \cdot) * \varphi_n)(X_s) ds - \sum_{i=1}^d \int_0^t \int w(s, z)D_i[\varphi_n(X_s - z)]dz \circ dB_s^i. \end{aligned}$$

Then

$$\begin{aligned} & \int w(t, z)\varphi_n(X_t - z)dz = \\ & \int_0^t \int (F(s, z, u) - F(s, z, v))\varphi_n(X_s - z) dz ds - \int_0^t \mathcal{R}_n(w, b)(X_s(x)) ds, \end{aligned}$$

where  $\mathcal{R}_n$  is the commutator defined above.

Step 2(  $\lim_{n \rightarrow \infty} \int_0^t \mathcal{R}_n(w, b)(X_s) ds = 0$ ) We argue as in [2]. We observe by Lemma 3.1 and the Lebesgue dominated theorem that

$$\lim_{n \rightarrow \infty} \int_0^t \int \mathcal{R}_n(w, b)(X_s)\rho(x) ds = 0$$

for all  $\rho \in C_0^\infty(\mathbb{R}^d)$ , for details see Theorem 20 of [2].

Step 3(  $w = 0$ ) We observe that



$$\lim_{n \rightarrow \infty} (w(t, \cdot) * \varphi_n)(\cdot) = w(t, \cdot)$$

, where the convergence is in  $L^1([0, T], L^1_{loc}(\mathbb{R}^d))$ . From the flow properties of  $X_t$ , see theorem 5 of [2], we obtain

$$\lim_{n \rightarrow \infty} (w(t, \cdot) * \varphi_n)(X_t) = w(t, X_t)$$

and

$$\lim_{n \rightarrow \infty} ((F(t, \cdot, u) - F(t, \cdot, v)) * \varphi_n)(X_t) =$$

$$(F(t, \cdot, X_t, u(t, \cdot, X_t)) - F(t, \cdot, X_t, v(t, \cdot, X_t))),$$

where the convergence is  $\mathbb{P}$  a.s in  $L^1([0, T], L^1_{loc}(\mathbb{R}^d))$ . Then by steps 1, 2 we have

$$w(t, X_t) = \int_0^t F(s, \cdot, X_s, u(t, \cdot, X_s)) - F(s, \cdot, X_s, v(t, \cdot, X_s)) ds.$$

Thus, for any compact set  $K \subset \mathbb{R}^d$  we obtain that

$$\int_K |w(t, X_t)| dx \leq \int_0^t \int_K |F(s, \cdot, X_s, u(t, \cdot, X_s)) - F(s, \cdot, X_s, v(t, \cdot, X_s))| dx ds.$$

$$\leq C \int_0^t \int_K |w(t, \cdot, X_s)| dx ds.$$

where  $C$  is constant related to the Lipchitz property of  $F$ . It follows

$$\int_K |w(t, X_t)| dx \leq C \int_0^t \int_K |w(t, \cdot, X_s)| dx ds.$$

and thus  $w(t, X_t) = 0$  by the Gronwall Lemma.

**Remark 3.1** *We observe that the unique solution  $u(t, x)$  has the representation  $u(t, x) = Z_t(x, f(X_t^{-1}))$ , where  $X_t$  and  $Z_t$  satisfy the equations (10) and (11) respectively.*

**Remark 3.2** *We mention that other variants of the theorem 3.1 can be proved. In fact, the step 2 is valid under other hypotheses, see corollary 23 of [2].*

**Remark 3.3** *We recall that relevant examples of non-uniqueness for the deterministic linear transport equation are presented in [2] and [3]. Currently we do not get a counter-example itself of the non-linear case. An interesting future work is to study if the nonlinear case may induce new pathologies.*

## References

- [1] Chow, P. L. 2007. Stochastic Partial Differential Equations, Chapman Hall/CRC.
- [2] Flandoli F., Gubinelli M. , Priola, E. 2010. Well-posedness of the transport equation by stochastic perturbation, Invent. Math., 180 (1): 1-53.
- [3] Flandoli,F. 2011. The Interaction Between Noise and Transport Mechanisms in PDEs, Milan j. Math, 79 (2) : 543-560.
- [4] Flandoli, F. 2009. Remarks on uniqueness and strong solutions to deterministic and stochastic differential equations, Metrika, 69 (2-3) : 101-123.
- [5] Kunita, H. 1984. First Order Stochastic Partial Differential Equations , in Proceedings of the Taniguchi International Symposium on Stochastic Analysis, North-Holland Mathematical Library, 249-269.

- [6] Kunita, H. 1982. Stochastic differential equations and stochastic flows of diffeomorphisms, Lectures Notes in Mathematics, 1097 : 143-303.
- [7] Kunita, H. 1990. Stochastic flows and stochastic differential equations, Cambridge University Press .