# Change of variables as a method to study general $\beta$ -models: bulk universality

M. Shcherbina, Institute for Low Temperature Physics Ukr.Ac.Sci, Kharkov, Ukraine. E-mail:shcherbi@ilt.kharkov.ua

#### Abstract

We consider  $\beta$  matrix models with real analytic potentials. Assuming that the corresponding equilibrium density  $\rho$  has a one-interval support (without loss of generality  $\sigma = [-2, 2]$ ), we study the transformation of the correlation functions after the change of variables  $\lambda_i \to \zeta(\lambda_i)$  with  $\zeta(\lambda)$  chosen from the equation  $\zeta'(\lambda)\rho(\zeta(\lambda)) = \rho_{sc}(\lambda)$ , where  $\rho_{sc}(\lambda)$  is the standard semicircle density. This gives us the "deformed"  $\beta$ -model which has an additional "interaction" term. Standard transformation with the Gaussian integral allows us to show that the "deformed"  $\beta$ -model may be reduced to the standard Gaussian  $\beta$ -model with a small perturbation  $n^{-1}h(\lambda)$ . This reduces most of the problems of local and global regimes for  $\beta$ -models to the corresponding problems for the Gaussian  $\beta$ -model with a small perturbation. In the present paper we prove the bulk universality of local eigenvalue statistics for both one-cut and multi-cut cases.

## **1** Introduction and main results

For any  $\beta > 0$  we consider the distribution in  $\mathbb{R}^n$  of the form

$$p_{n,\beta}(\bar{\lambda}) = Z_{n,\beta}^{-1}[V] e^{\beta H(\bar{\lambda})/2}, \qquad (1.1)$$

where H (Hamiltonian) and  $Z_n[\beta, V]$  (partition function) are

$$H(\bar{\lambda}) = -n \sum_{i=1}^{n} V(\lambda_i) + \sum_{i \neq j} \log |\lambda_i - \lambda_j|, \qquad (1.2)$$

$$V(\lambda) > (1+\varepsilon) \log(1+\lambda^2),$$
  

$$Z_n[\beta, V] = \int e^{\beta H(\bar{\lambda})/2} d\bar{\lambda}.$$
(1.3)

For any integrable function  $\Phi(\bar{\lambda})$  we denote its expectation by

$$\langle \Phi(\bar{\lambda}) \rangle_{V,n} = \int \Phi(\bar{\lambda}) p_{n,\beta}(\bar{\lambda}) d\bar{\lambda}.$$
(1.4)

The expectation is closely connected with the correlation functions

$$p_{n,\beta}^{(m)}(\lambda_1,...,\lambda_m) = \int_{\mathbb{R}^{n-l}} p_{n,\beta}(\bar{\lambda}) d\lambda_{m+1}...d\lambda_n.$$
(1.5)

It will be convenient below to use also the notation

$$\mathcal{N}_n[h] = \sum_{i=1}^n h(\lambda_i) \tag{1.6}$$

for the linear eigenvalue statistics, corresponding to the test function h.

For  $\beta = 1, 2, 4$  (1.1)-(1.3) is a joint eigenvalue distribution of real symmetric, hermitian and symplectic matrix models respectively.

Since the papers [3, 15] it is known that for any  $\beta > 0$  if V is a Hölder function, then

$$n^{-2}\log Z_{n,\beta}[V] = \frac{\beta}{2}\mathcal{E}[V] + O(\log n/n),$$
 (1.7)

where

$$\mathcal{E}[V] = \max_{m \in \mathcal{M}_1} \left\{ L[dm, dm] - \int V(\lambda)m(d\lambda) \right\} = \mathcal{E}_V(m^*), \tag{1.8}$$

and the maximizing measure  $m^*$  (called the equilibrium measure) has a compact support  $\sigma := \operatorname{supp} m^*$ . Here and below we denote

$$L[dm, dm] = \int \log |\lambda - \mu| dm(\lambda) dm(\mu), \qquad (1.9)$$
$$L[f](\lambda) = \int \log |\lambda - \mu| f(\mu) d\mu, \quad L[f, g] = (L[f], g),$$

where (.,.) is a standard inner product in  $L_2[\mathbb{R}]$ . The support  $\sigma$  can consist of one interval (one-cut case) and many intervals (multi-cut case). If V' is a Hölder function, then the equilibrium measure  $m^*$  has a density  $\rho$  (equilibrium density). The support  $\sigma$  and the density  $\rho$  are uniquely defined by the conditions:

$$\begin{aligned} v(\lambda) &:= 2 \int \log |\mu - \lambda| \rho(\mu) d\mu - V(\lambda) = \sup v(\lambda) := v^*, \quad \lambda \in \sigma, \\ v(\lambda) &\leq \sup v(\lambda), \quad \lambda \notin \sigma, \qquad \sigma = \sup\{\rho\}. \end{aligned} \tag{1.10}$$

Without loss of generality we will assume below that  $v^* = 0$ .

One of the most important questions of the theory of random matrices is the universality conjecture for the local eigenvalue statistics. According to this conjecture, e.g., in the bulk of the spectrum, the behavior of the scaled correlation functions (1.5)

$$p_{k,\beta}^{(n)}(\lambda_0 + x_1/(n\rho(\lambda_0)), ..., \lambda_0 + x_k/(n\rho(\lambda_0)))$$
(1.11)

in the limit  $n \to \infty$  is universal, i.e., does not depend on V and  $\lambda_0$  and depends only on  $\beta$ . The case  $\beta = 2$  is the simplest one, since for  $\beta = 2$  all correlation functions of (1.5) can be expressed in terms of the reproducing kernel of the system of polynomials orthogonal with a varying weight  $e^{-n\beta V}$  (see e.g. [17]). The orthogonal polynomial machinery, in particular, the Christoffel-Darboux formula and Christoffel's function simplify considerably the studies of marginal densities (1.5). This allows to study for  $\beta = 2$  the local eigenvalue statistics in many different cases: bulk of the spectrum, edges of the spectrum, special points, etc. (see [18], [20],[8],[4],[7],[16]).

For  $\beta = 1, 4$  the situation is more complicated. It was shown in [27] that all correlation functions can be expressed in terms of some 2 × 2- matrix kernels. But the representation is less convenient than that in the case  $\beta = 2$ . Therefore the universality conjecture for  $\beta = 1, 4$ was proven much later than for  $\beta = 2$ . There were a number of papers with improving results, first for monomials  $V = \lambda^{2m} + o(1)$ , (see [9], [10], [11]), then for arbitrary real analytic one-cut potentials (see [23], [24]) and finally for multi-cut real analytic potentials (see [25]).

Note, that for  $\beta = 1, 2, 4$  it was shown that the convergence of the scaled correlation functions (1.11) is uniform in  $(x_1, \ldots, x_k) \in S$ , where S is an arbitrary compact set in  $\mathbb{R}^k$ . There is also a more weak form of the universality, when the limit  $n \to \infty$  is taken after the integration of the correlation function of (1.11) with a smooth compactly supported function  $\phi(x_1, \ldots, x_k)$ . To prove universality in this form, it suffices to consider the limits of the expectations of the functions of the form

$$\Phi_k(\bar{\lambda};\lambda_0) = \prod_{j=1}^k \left( n^{-1} \sum_{i=1}^n n\phi_j \left( n\rho(\lambda_0)(\lambda_i - \lambda_0) \right) \right), \ \lambda_0 \in (-2 + \varepsilon, 2 - \varepsilon), \tag{1.12}$$

where  $\phi_j(x)$  (j = 1, ..., k) - are arbitrary smooth functions with compact supports.

In the series of recent papers [5, 12, 6] the bulk universality for any  $\beta > 0$  in the case of one-cut potentials of the generic behavior, possessing 4 derivatives, was proven in the form (see [6], Theorem 2.5):

$$\lim_{n \to \infty} (2n^{-1+\varepsilon})^{-1} \int_{-n^{-1+\varepsilon}}^{n^{-1+\varepsilon}} dt \langle \Phi_k(\bar{\lambda};\lambda_0+t) \rangle_{V,n} = \lim_{n \to \infty} \langle \Phi_k(\bar{\lambda};\lambda_0) \rangle_{*,n},$$

where  $\varepsilon$  is an arbitrary small number, and here and below we denote  $\langle \dots \rangle_{*,n}$  the expectation (1.4) for the Gaussian case  $V^*(\lambda) = \frac{1}{2}\lambda^2$ . Recall that the generic behavior of the potential V means that its equilibrium density has the form

$$\rho(\lambda) = \frac{1}{2\pi} P(\lambda) \Im X_{\sigma}^{1/2}(\lambda + i0), \quad \inf_{\lambda \in \sigma} |P(\lambda)| > 0, \quad X_{\sigma}(z) = z^2 - 4, \tag{1.13}$$

where we choose a branch of  $X_{\sigma}^{1/2}(z)$  such that  $X_{\sigma}^{1/2}(z) \sim z$ , as  $z \to +\infty$ . Moreover, the function v defined by (1.10) attains its maximum only if  $\lambda$  belongs to  $\sigma$ .

We recall also that for sufficiently smooth V the equilibrium density  $\rho$  always has the form (1.13) (see, e.g., [1]). For real analytic V the function P is also real analytic (1.13) and can be represented in the form

$$P(z) = \frac{1}{2\pi i} \oint_{\mathcal{L}} \frac{V'(z) - V'(\zeta)}{(z - \zeta) X_{\sigma}^{1/2}(\zeta)} d\zeta.$$
 (1.14)

Hence generic behavior just means that  $\rho$  has no zeros in the internal points of  $\sigma$  and behaves like square root near the edge points.

In the present paper we propose a different from [5, 12, 6] method, based on the analysis of the transformation of (1.1) under a smooth change of variables  $\lambda \to \zeta(\lambda)$ . For a good choice of  $\zeta$  (see (2.1)) we obtain that the partition function and all the correlation functions of (1.1) can be expressed in terms of the Hamiltonian

$$\tilde{H}^{(\zeta)}(\bar{\lambda}) = H_n^*(\bar{\lambda}) + \left(\frac{2}{\beta} - 1\right) \sum \log \zeta'(\lambda_j) + \sum_{k=1}^{\infty} \eta_k \left(\sum_j (\varphi_k(\lambda_j) - (\varphi_k, \rho_{sc}))\right)^2, \quad (1.15)$$

where  $H_n^*$  is the Hamiltonian of the form (1.1), corresponding to  $V^*(\lambda) = \lambda^2/2$ , and  $\{\eta_k\}_{k=1}^{\infty}$ and  $\{\varphi_k(\lambda)\}_{k=1}^{\infty}$  are eigenvalues and eigenvectors of the integral operator in  $L_2[\sigma_{\varepsilon}]$  ( $\sigma_{\varepsilon} = [-2 - \varepsilon, 2 + \varepsilon]$ ) with the kernel

$$L^{(\zeta)}(\lambda,\mu) := \log \left| \frac{\zeta(\lambda) - \zeta(\mu)}{\lambda - \mu} \right| = \sum_{k=1}^{\infty} \eta_k \varphi_k(\lambda) \varphi_k(\mu).$$
(1.16)

For sufficiently smooth  $\zeta(\lambda)$  the operator with this kernel is a compact operator with smooth eigenfunctions. The rate of convergence  $\eta_k \to 0, k \to \infty$ , depends on the number of derivatives of  $\zeta(\lambda)$ , e.g. for  $\zeta(\lambda) \in C_l[\sigma_{\varepsilon}]$ , we have  $\eta_k = o(k^{-l-1/2}), k \to \infty$  (see [14] Chapter

III, §10). Hence, restricting summation in (1.15) by M = M(n), we can provide that the remainder is  $o(n^{-2})$ , and so it does not contribute to the correlation functions. Then, using the Gaussian integration formula (see (2.4)) for each  $k = 1, \ldots M$ , we can "linearize" the terms under the summation and obtain that

$$\left\langle \Phi_k(\bar{\zeta}) \right\rangle_{V,n} = \left(\frac{\beta}{8\pi}\right)^{M/2} \int d\bar{u} e^{-\beta(\bar{u},\bar{u})/8} \left\langle \Phi_k(\zeta(\bar{\lambda})) e^{\beta\mathcal{N}_n[\dot{h}_u]/2} \right\rangle_{*,n} / I_n[\beta,\zeta], \tag{1.17}$$

where a "small perturbation" of  $V_* h_u(\lambda)$  (defined by (2.5)) depends linearly on the integration parameters  $\bar{u}$ , and  $I_n[\beta, \zeta]$  is the normalizing constant

$$I_n[\beta,\zeta] := \left(\frac{\beta}{8\pi}\right)^{M/2} \int d\bar{u} e^{-\beta(\bar{u},\bar{u})/8} \langle e^{\beta\mathcal{N}_n[\dot{h}_u]/2} \rangle_{*,n};$$

Note that a similar method was used in [26] in the multi-cut case in order to linearize the term which corresponds to the "interaction" between different intervals of the spectrum.

The analysis of  $\langle e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n}$  is based on the well-known result of [15], which we will use in the form, obtained in [26], Theorem 1.

**Theorem 1** Let V be a real analytic one-cut potential of generic behavior with  $\sigma = [-2, 2]$ , and h satisfy one of the two conditions:

(i) h is a real valued function with  $||h'||_2$ ,  $||h^{(6)}||_2 \le n^{1-\tilde{\delta}}$  (here and below  $||.||_2$  means the standard norm in  $L_2[-2-\varepsilon/2, 2+\varepsilon/2]$ , with some small  $\varepsilon > 0$ ),  $\tilde{\delta} > 0$  is any small constant; (ii) h is complex valued,  $(D_{\sigma}\Re h, \Re h) + (D_{\sigma}\Im h, \Im h) \le c_* \log n$  with some sufficiently

small  $c_*$ , and  $|h'||_2, ||h^{(6)}||_2 \le \log^s n$  with some s > 0.

Then we have

$$\langle e^{\beta \mathcal{N}_{n}[\dot{h}]/2} \rangle_{V,n} = \exp\left\{ (h, \nu_{\beta}) + \frac{\beta}{8} (\overline{D}_{\sigma}h, h) + n^{-\alpha} O\left( ||h'||_{2}^{3} \right) + n^{-\alpha} O\left( ||h^{(6)}||_{2}^{3} \right) \right\},$$
(1.18)

where  $\dot{h} := h - (\rho, h)$ , and  $\alpha = 1$  for the case (i) and  $\alpha = 1/2$  for the case (ii),

$$\overline{D}_{\sigma} = \frac{1}{2}(D_{\sigma} + D_{\sigma}^*), \quad D_{\sigma}h(\lambda) = \frac{X_{\sigma}^{-1/2}(\lambda)}{\pi^2} \int_{\sigma} \frac{h'(\mu)X_{\sigma}^{1/2}(\mu)d\mu}{(\lambda - \mu)}, \tag{1.19}$$

and  $D_{\sigma}^*$  is the adjoint operator to  $D_{\sigma}$  in  $L_2(\sigma)$ . We will use also the representation of  $D_{\sigma}$  obtained in [15]

$$(\overline{D}_{\sigma}h,h) = \sum_{k=1}^{\infty} kh_k^2, \quad h_k = \frac{2}{\pi} \int_0^{\pi} h(2\cos\theta)\cos k\theta d\theta.$$
(1.20)

A non positive measure  $\nu_{\beta}$  in (1.18) has the form

$$(h,\nu_{\beta}) := \left(1 - \frac{\beta}{2}\right) \left(\frac{1}{4}(h(-2) + h(2)) - \frac{1}{2\pi} \int_{\sigma} \frac{h(\lambda)d\lambda}{\sqrt{4 - \lambda^2}} - \frac{1}{2}(D_{\sigma}\log P, h)\right).$$

It will be important in what follows that  $\overline{D}_{\sigma}$  is a rank one perturbation of  $-L_{\sigma}^{-1}$ , where  $L_{\sigma}$  is the integral operator defined by the kernel  $\log |\lambda - \mu|$  for the interval  $\sigma$  (see [26]):

$$L_{\sigma}\bar{D}_{\sigma}v = -v + \pi^{-1}(v, X_{\sigma}^{-1/2})\mathbf{1}_{\sigma}.$$
 (1.21)

It is easy to understand that in view of (1.17) it suffices to prove that in the domain which gives non vanishing contribution in the integral (1.17) we have

$$\left|\frac{\langle \Phi_k e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n}}{\langle e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n}} - \langle \Phi_k \rangle_{*,n} \right| \le \tilde{\varepsilon}_n, \quad \tilde{\varepsilon}_n \to 0, \tag{1.22}$$

or, by another words, that for the Gaussian potential  $V^*$  the "small perturbation"  $n^{-1}h_u$ does not change correlation functions. We prove (1.22) in two steps. On the first step we replace  $h_u(\lambda)$  by some linear function  $c(h_u)\lambda$ , and on the second use the result of [28], Theorem 1, which (after "translation" on the langauge of correlation functions) states that for any sequence  $\lambda_0^{(n)}$  such that  $n^{2/3} ||\lambda_0^{(n)}| - 2| \to \infty$ , the integrated correlation functions  $\langle \Phi_k(\bar{\lambda}, \lambda_0^{(n)}) \rangle_{*,n}$  of (1.12) converges to some universal limit, depending only on  $\{\phi_j\}_{j=1}^k$ . This limit corresponds to the so-called  $Sine_\beta$  process, whose definition is not important here (see [28] for the precise definitions and results). We will use two simple corollaries from the above statement:

$$|\langle \Phi_k(\lambda,\lambda_0) \rangle_{*,n}| \le C_{\Phi},\tag{1.23}$$

$$|\langle \Phi_k(\lambda, \lambda_0 + t/n) \rangle_{*,n} - \langle \Phi_k(\lambda, \lambda_0) \rangle_{*,n}| \le \varepsilon_n \to 0, \quad n \to \infty, \tag{1.24}$$

where the first bound is uniform for  $\lambda_0 \in [-2 + \varepsilon, 2 - \varepsilon]$ , and the second relation is uniform in the same  $\lambda_0$  and  $|t| \leq n^{1-\delta}$  if  $\delta > 0$  is fixed. Note, that (1.23) and (1.24) become evident if we assume the contrary for some sequence of  $\lambda_0^{(n)}$  and obtain the contradiction with Theorem 1 of [28].

The method briefly described above gives the following result

**Theorem 2** Let V be a real analytic one-cut potential with  $\sigma = [-2, 2]$  of generic behavior and  $\lambda_0 \in [-2 + \varepsilon, 2 - \varepsilon]$  with any small  $\varepsilon > 0$ . Then for any  $k \ge 1$  and any  $\Phi_k(\bar{\lambda}, \lambda_0)$  of the form (1.12) with smooth  $\{\phi_j\}_{j=1}^k$  we have uniformly in  $\lambda_0 \in [-2 + \varepsilon, 2 - \varepsilon]$ 

$$\lim_{n \to \infty} \langle \Phi_k(\bar{\lambda}, \lambda_0) \rangle_{V,n} = \lim_{n \to \infty} \langle \Phi_k(\bar{\lambda}, 0) \rangle_{*,n}.$$
(1.25)

#### Remarks:

(i) The method of Theorem 2 can be generalized to the case of non-analytic V, for which P of (1.13) possesses 5 derivatives. The reasons to prove here Theorem 2 for real analytic V is that in this case  $|\eta_k| \sim e^{-kc}$ , thus we can take  $M = [\log^2 n]$  and do not care about  $M^p$  and the number of derivatives in the formulas. This allows to simplify the proof of Lemmas 2 and 3. Moreover, for P with 5 derivatives, the result (1.18) cannot be applied and we need to prove a new form of (1.18), which requires only  $2 + \varepsilon$  derivatives of h (in the sense of Sobolev spaces), but gives the bound only of the order  $n^{-\kappa}$  with small  $\kappa > 0$ , instead of  $n^{-1}$  in (1.18). All these technicalities make the proof less straightforward. Thus, the proof of Theorem 2 for non analytic potentials is postponed to the next paper, where the edge universality will be proved by the same method.

(ii) Examining the proof of Theorem 2, it is easy to see that to prove the edge universality for  $\lambda_0 = 2$ , it suffices just to replace  $\Phi_k$  of (1.12) by

$$\tilde{\Phi}_k(\bar{\lambda}, ) = \prod_{j=1}^k \left( n^{-1} \sum_{i=1}^n n\varphi_j \left( n^{2/3} \gamma_P(\lambda_i - 2) \right) \right), \ \gamma_P = P^{2/3}(2)$$

The only difference will be that instead of (1.23), (1.24) we need to use similar results of [22] on the existence of limits for the scaled correlation functions near the edge point for the Gaussian potential  $V^*$ .

(iii) The method proposed here works well for the global regime problems, e.g., for the proof of CLT for  $\mathcal{N}_n[h]$ , the expansion for  $Z_n[\beta, V]$  in  $n^{-k}$ , computations of the Hankel and the Toeplitz determinants etc. In particular, it simplifies considerably the proof of well-known results, (see, e.g., [15, 23, 24, 25, 2]) because it reduces their proofs to the case of the Gaussian potential  $V^*$  with a small perturbations  $\frac{1}{n}h$ .

Using the results of [26], Theorem 2 can be generalized to the multi-cut real analytic potentials of generic behavior.

**Theorem 3** Let V be a real analytic multi-cut potential with  $\sigma = \bigcup_{\alpha=1}^{q} \sigma_{\alpha}$  ( $\sigma_{\alpha} = [a_{\alpha}, b_{\alpha}]$ ) of generic behavior, which means that the correspondent equilibrium density  $\rho$  has the form (1.13), with  $X_{\sigma} = \prod_{\alpha=1}^{q} (z - a_{\alpha})(z - b_{\alpha})$ . Then, for any  $\lambda_0 \in \bigcup_{\alpha=1}^{q} [-a_{\alpha} + \varepsilon, b_{\alpha} - \varepsilon]$  (1.25) holds.

The paper is organized as follows. The proof of Theorem 2 modulo few auxiliary statements (see Lemmas 1-4) is given in Section 2. The proofs of Lemmas 1-4 are given in Section 3. The proof of Theorem 3 is given in Section 4.

# 2 Proof of Theorem 2

Take any *n*-independent small  $\varepsilon > 0$ . It is known (see [3]) that if we replace in the definition of the partition function and of the correlation functions the integration over  $\mathbb{R}$  by the integration  $\sigma_{\varepsilon/2}$ ,  $p_{n,\beta}^{(m)}$  and the new marginal densities  $p_{n,\beta}^{(m,\varepsilon)}$  for  $m = 1, 2, \ldots$  satisfy the inequalities

$$\sup_{\lambda_1,\dots,\lambda_m\in\sigma_{\varepsilon/2}} |p_{n,\beta}^{(m)}(\lambda_1,\dots,\lambda_m) - p_{k,\beta}^{(m,\varepsilon)}(\lambda_1,\dots,\lambda_m)| \le C_m e^{-n\beta d_{\varepsilon}}$$
$$Z_n[\beta,V] = Z_n^{(\varepsilon)}[\beta,V] (1 + e^{-n\beta d_{\varepsilon}}).$$

It is more convenient to consider the integration with respect to  $\sigma_{\varepsilon/2}$ , thus, starting from this moment it is assumed that this truncation is made, and below the integration without limits means the integration over  $\sigma_{\varepsilon/2}$ , but the superindex  $\varepsilon$  will be omitted.

#### 2.1 Change of variables in the one cut case

Let V be some smooth enough potential with the equilibrium density  $\rho$ , supp $\rho = [-2, 2]$ , and  $\zeta(\lambda) : \sigma_{\varepsilon} = [-2 - \varepsilon, 2 + \varepsilon] \rightarrow \sigma_{\varepsilon}$  be some smooth function such that  $\inf_{\sigma_{\varepsilon}} \zeta' > 0$ . Consider

$$H^{(\zeta)}(\bar{\lambda}) = -n \sum V(\zeta(\lambda_j)) + \sum_{i \neq j} \log |\zeta(\lambda_i) - \zeta(\lambda_j)| + \frac{2}{\beta} \sum \log \zeta'(\lambda_j).$$

It is evident that the correspondent partition function and all the marginal densities satisfy the relations

$$Z_{n,\beta}^{(\zeta)} := \int e^{\beta H^{(\zeta)}/2} d\bar{\lambda} = Z_n[\beta, V],$$
$$p_{n,\beta}^{(m,\zeta)}(\lambda_1, \dots, \lambda_m) := (Z_{n,\beta}^{(\zeta)})^{-1} \int e^{\beta H^{(\zeta)}/2} d\lambda_{m+1} \dots d\lambda_n$$
$$= p_{n,\beta}^{(m)}(\zeta(\lambda_1), \dots, \zeta(\lambda_m)).$$

On the other hand,

$$H^{(\zeta)}(\bar{\lambda}) = -n \sum V(\zeta(\lambda_j)) + \sum_{i \neq j} \log |\lambda_i - \lambda_j|$$
  
+ 
$$\sum_{i,j} \log \left| \frac{\zeta(\lambda_i) - \zeta(\lambda_j)}{\lambda_i - \lambda_j} \right| + (\frac{2}{\beta} - 1) \sum \log \zeta'(\lambda_j),$$

where we removed the condition  $i \neq j$  in the third sum and add the correspondent terms to the forth sum. Choose  $\zeta(\lambda)$  from the equation

$$\zeta'(\lambda) = \frac{\rho_{sc}(\lambda)}{\rho(\zeta(\lambda))}, \quad \zeta(-2) = -2, \quad \text{with} \quad \rho_{sc}(\lambda) = \frac{\sqrt{4-\lambda^2}}{2\pi}.$$
(2.1)

We will use this equation also in the form

$$\rho(\zeta(\lambda))\zeta'(\lambda) = \rho_{sc}(\lambda). \tag{2.2}$$

**Lemma 1**  $\zeta(\lambda)$  is a real analytic function in some  $\sigma_{\tilde{\varepsilon}}$ , and  $\zeta(2) = 2$ .

Without loss of generality we assume below that  $\sigma_{\tilde{\varepsilon}} = \sigma_{\varepsilon}$ .

For this choice of  $\zeta$  write

$$\sum_{i,j} L^{(\zeta)}(\lambda_i, \lambda_j) = \sum \eta_k \Big( \sum_j (\varphi_k(\lambda_j) - (\varphi_k, \rho_{sc})) \Big)^2 + 2n \sum_j \sum_k \eta_k \varphi_k(\lambda_j) (\varphi_k, \rho_{sc}) - n^2 \sum_k \eta_k (\varphi_k, \rho_{sc})^2 = \Delta(\bar{\lambda}) + 2n \sum_j \int L^{(\zeta)}(\lambda_j, \mu) \rho_{sc}(\mu) d\mu - n^2 \int L^{(\zeta)}(\lambda, \mu) \rho_{sc}(\lambda) \rho_{sc}(\mu) d\lambda d\mu,$$

where  $(f,g) := \int_{\sigma_{\varepsilon}} fg d\lambda$ . It is easy to see that for  $\lambda \in \sigma$  in view of (2.2)

$$2\int L^{(\zeta)}(\lambda,\mu)\rho_{sc}(\mu)d\mu = 2\int \log|\zeta(\lambda) - \zeta(\mu)|\rho_{sc}(\mu)d\mu - 2\int \log|\lambda - \mu|\rho_{sc}(\mu)d\mu \qquad (2.3)$$
$$= 2\int \log|\zeta(\lambda) - \zeta(\mu)|\rho(\zeta(\mu))\zeta'(\mu)d\mu - \frac{\lambda^2}{2} = V(\zeta(\lambda)) - \frac{\lambda^2}{2}.$$

On the other hand, the l.h.s. here is a real analytic function in  $\sigma_{\varepsilon}$  and the r.h.s. is also a real analytic function in  $\sigma_{\varepsilon}$ , hence (2.3) is valid for  $\lambda \in \sigma_{\varepsilon}$ . Similarly

$$\int L^{(\zeta)}(\lambda,\mu)\rho_{sc}(\lambda)\rho_{sc}(\mu)d\lambda d\mu = \mathcal{E}_{sc} - \mathcal{E}_V =: -\Delta \mathcal{E}.$$

Hence we finally obtain that our Hamiltonian for  $\bar{\lambda} \in \sigma_{\varepsilon}^{n}$  has the form (1.15)

### 2.2 "Linearization" of the quadratic terms in (1.15)

As it was mentioned in Section 1, in the case of real analytic  $\zeta$ , the eigenfunctions  $\{\varphi_k(\lambda)\}_{k=1}^{\infty}$  are analytic in the same domain as  $\zeta$ , and the eigenvalues  $|\eta_k| \leq e^{-kc}$ . Hence if we choose

 $M = [\log^2 n]$ , then the remainder of the sum in (1.15) will be less than any negative degree of n and will not have essential influence on the correlation functions. Write for any  $1 \le k \le M$ 

$$\exp\left\{\frac{\beta}{2}\eta_k\left(\sum_j(\varphi(\lambda_j) - (\varphi_k, \rho_{sc}))\right)^2\right\}$$

$$= \sqrt{\frac{\beta}{8\pi}} \int du_k \exp\left\{\frac{\beta}{2}\left(\sqrt{\eta_k}\left(\sum_j(\varphi_k(\lambda_j) - (\varphi_k, \rho_{sc}))\right)u_k - u_k^2/4\right)\right\},$$
(2.4)

where for  $k \in I_+ = \{k \leq M : \eta_k > 0\}$  we take an arithmetic square root, while for  $k \in I_- = \{k \leq M : \eta_k < 0\} \sqrt{\eta_k} = i\sqrt{|\eta_k|}$ . We will write  $\bar{u} = (u_1, \ldots, u_M)$ . Substituting this integrals in (1.15) and integrating first with respect to  $\bar{\lambda}$ , we get (1.17) with

$$h_{\bar{u}}(\lambda) = \sum_{k=1}^{M} \sqrt{\eta_k} \varphi_k(\lambda) u_k + \left(\frac{2}{\beta} - 1\right) \log \zeta'(\lambda), \quad \dot{h}_{\bar{u}}(\lambda) = h_{\bar{u}} - (h_{\bar{u}}, \rho_{sc}).$$
(2.5)

#### 2.3 Integration with respect to $\bar{u}$

The first our step is to get rid in (1.17) from the domain of  $\bar{u}$ , where  $\bar{u}$  is big, proving that the correspondent contribution in the integral (1.17) is small. Set

$$\begin{aligned} K_{jk}^{+} &= \eta_{j}^{1/2} \eta_{k}^{1/2} (\bar{D}_{\sigma} \varphi_{k}, \varphi_{j}), \ j, k \in I_{+}, \\ K_{jk}^{-} &= |\eta_{j}|^{1/2} |\eta_{k}|^{1/2} (\bar{D}_{\sigma} \varphi_{k}, \varphi_{j}), \ j, k \in I_{-}, \\ U_{1} &= \{ \bar{u} : (K^{+} \bar{u}, \bar{u}) + (K^{-} \bar{u}, \bar{u}) \leq k_{*} \log \varepsilon_{n}^{-1} \}, \end{aligned}$$
(2.6)

where  $\varepsilon_n$  is given by (1.24) and  $k_*$  is some absolute constant which will be chosen later.

**Lemma 2** There exists n-independent  $\delta > 0$  such that

$$K^+ < 1 - \delta. \tag{2.7}$$

Moreover, if  $U_1^c$  is a complement of  $U_1$  of (2.6), then

$$\left(\frac{\beta}{8\pi}\right)^{M/2} \int_{U_1^c} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8} |\langle \Phi_k e^{\beta \mathcal{N}_n[\dot{h}_u]/2} \rangle_{*,n}| \le \varepsilon_n^{\tau k_*}, \tag{2.8}$$

where  $\varepsilon_n$  is given by (1.24),  $k_*$  - by (2.6), and  $\tau > 0$  is some fixed number, depending on  $\delta$  in (2.7).

The proof of Lemma 2 is partially based on the following assertion

**Lemma 3** Let  $h(\lambda)$  be a real analytic function such that  $||h'(\lambda)||_2$ ,  $||h^{(16)}(\lambda)||_2 \le \log^s n$  with some n-independent positive s. Then

$$\left|\frac{\langle \Phi_k e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n}}{\langle e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n}} - \langle \Phi_k(\lambda) \rangle_{*,n} \right| \le C(n^{-\kappa} + \varepsilon_n), \tag{2.9}$$

where  $\varepsilon_n$  is the same as in (1.24), and  $\kappa > 0$ .

In particular, Lemma 3 and (1.23) for real h imply the bound which we need in the proof of Lemma 2:

$$\left| \langle \Phi_k e^{\beta \mathcal{N}_n[\dot{h}]/2} \rangle_{*,n} \right| \le C_{\Phi_k} \langle e^{\beta \mathcal{N}_n[\dot{h}]/2} \rangle_{*,n}.$$
(2.10)

Now let us prove (1.22). As it was mentioned above, for real  $h_u$  (1.22) follows from Lemma 3 (see (2.9)). To extend (2.9) to the complex valued  $h_u$ , we use the last lemma:

**Lemma 4** Let the analytic in  $t \in D = \{t : |t| \le \log^{1/2} \varepsilon_n^{-1}, \Im t \ge 0\}$  functions  $F_n$  satisfy two bounds:

$$|F_n(t)| \le C_1 \varepsilon_n e^{t^2/2}, \quad -\log^{1/2} \varepsilon_n^{-1} \le t \le \log^{1/2} \varepsilon_n^{-1}, \quad \varepsilon_n < 1,$$

$$|F_n(t)| \le C_2 e^{(\Re t)^2/2}, \quad t \in D.$$
(2.11)

Then the inequality

$$|F_n(t)| \le C\varepsilon_n^{1/2} |e^{t^2/2}|$$
 (2.12)

holds for  $t \in D' := \frac{1}{6}D$  with  $C = C_1^{3/4}C_2^{1/4}$ .

Denote

$$X_1 = \mathcal{N}_n[\Re h_{\bar{u}}], \quad X_2 = \mathcal{N}_n[\Im h_{\bar{u}}], \tag{2.13}$$
$$d_{11} = \frac{\beta}{4} (D\Re h_{\bar{u}}, \Re h_{\bar{u}}), \quad d_{12} = \frac{\beta}{4} (D\Re h_{\bar{u}}, \Im h_{\bar{u}}), \quad d_{22} = \frac{\beta}{4} (D\Im h_{\bar{u}}, \Im h_{\bar{u}}),$$

and use Lemma 4 for

$$F_n(t) = \frac{\langle \Phi_k(\bar{\lambda})e^{\beta(X_1 - X_2 d_{12}/d_{22} + tX_2/d_{22}^{1/2})/2} \rangle_{*,n}}{\langle e^{\beta(X_1 - X_2 d_{12}/d_{22})/2} \rangle_{*,n}} - e^{t^2/2} \langle \Phi_k(\bar{\lambda}) \rangle_{*,n}$$

Lemma 3, (1.18), and (1.23) guarantee that  $F_n(t)$  satisfy (2.11). Take  $t^* = id_{22}^{1/2} + d_{12}/d_{22}^{1/2}$ . For  $k_* \leq \frac{1}{6}$  in (2.6),  $t \in \frac{1}{6}D$ , since

$$k_* \log \varepsilon_n^{-1} \ge d_{11} + d_{22} \ge d_{22} + d_{12}^2/d_{22} = |id_{22}^{1/2} + d_{12}/d_{22}^{1/2}| = |t^*|.$$
(2.14)

In addition, by (1.18) and (2.13)

$$e^{t^2/2} \langle e^{\beta(X_1 - X_2 d_{12}/d_{22})/2} \rangle_{*,n} = e^{d_{11} + 2id_{12} - d_{22}} (1 + O(n^{-\kappa})),$$

hence Lemma 4 yields (cf (1.22))

$$\left| \langle \Phi_k(\bar{\lambda}) e^{\beta(X_1 + iX_2)/2} \rangle_{*,n} - e^{d_{11} + 2id_{12} - d_{22}} \langle \Phi_k(\bar{\lambda}) \rangle_{*,n} \right| \le C \varepsilon_n^{1/2} \left| e^{d_{11} + 2id_{12} - d_{22}} \right|.$$
(2.15)

Applying this inequality first for  $\Phi \equiv 1$ , we get that

$$\langle e^{\beta(X_1+iX_2)/2} \rangle_{*,n} = e^{d_{11}+2id_{12}-d_{22}}(1+O(\varepsilon_n^{-1/2})),$$

and then, substituting the last relation in (2.15), we obtain (1.22). Integrating (1.22) in  $U_1$  we complete the proof of Theorem 2.

# **3** Proofs of the auxiliary results

*Proof of Lemma 1.* The fact that  $\zeta(2) = 2$  follows from the relation (2.2) and the fact that

$$\int_{-2}^{2} \rho(\lambda) d\lambda = \int_{-2}^{2} \rho_{sc}(\lambda) d\lambda = 1.$$

The analyticity in all internal points of (-2, 2) follows from the analyticity of P (see (1.14)). Hence we are left to prove that  $\zeta(\lambda)$  is analytic in some neighborhood of  $\lambda = \pm 2$ . Consider, e.g.,  $\lambda = -2$ . To simplify formulas, we make the change  $x = \lambda + 2$ . Let us sick solution in the form

$$\zeta(x) + 2 = P_0^{-2/3} x(1 + \zeta_0(x)), \quad \zeta_0(0) = 0.$$

where  $P_0 := P(-2) \neq 0$ . Then

$$P(\zeta) = P_0(1 + \zeta \tilde{P}(\zeta)) = P_0(1 + x(1 + \zeta_0)P_1(\zeta_0, x))$$

 $(P_1(\zeta_0, x))$  is analytic in both variables) and (2.1) can be written as

$$(x\zeta_0)' = \frac{\sqrt{4-x}}{(1+x(1+\zeta_0)P_1(\zeta_0,x))\sqrt{(1+\zeta_0)(4-P_0^{-2/3}x(1+\zeta_0(x)))}} - 1$$
  
$$:= F(x,\zeta_0) = (xF_0(x)+\zeta_0F_1(x)+\zeta_0^2F_2(x,\zeta_0)),$$

where we used the fact that the r.h.s. of the first line is analytic in  $x, \zeta_0$  at the point (0,0), hence  $F_0, F_1, F_2$  are analytic at (0,0), moreover the r.h.s. is 0 at this point. Note that

$$F_0(x) = \frac{\partial F}{\partial \zeta_0}\Big|_{\zeta_0=0} = \frac{\partial \zeta}{\partial \zeta_0} \frac{\partial F}{\partial \zeta}\Big|_{\zeta_0=0} = x P_0^{-2/3} \frac{\partial}{\partial \zeta} \frac{\sqrt{x(4-x)}}{P(\zeta)\sqrt{4-\zeta^2}}\Big|_{\zeta_0=0}$$

can not be identically zero, if  $P(x) \neq 1$ . Moreover,  $F_1(x) = -\frac{1}{2} + xF_{11}(x)$ . Thus, the equation can be written in the form

$$\zeta_0' = -\frac{3}{2}\frac{\zeta_0}{x} + F_0(x) + \zeta_0 F_{11}(x) + \frac{\zeta_0^2}{x}F_2(x,\zeta_0), \quad F_0(x) = x^m F_{0,m}(x), \quad F_{0,m}(0) \neq 0,$$

where m could be 0 or any positive integer. It is evident that if we sick  $\zeta_0 = \sum_{k=1} \zeta_k x^k$ , then the equation above gives us the recursive system

$$(k + \frac{3}{2})\zeta_k = P_k(\zeta_1, \dots, \zeta_{k-1}),$$
 (3.1)

where  $P_k$  is a polynomial of  $\zeta_1, \ldots, \zeta_{k-1}$  with coefficients depending on the Taylor coefficients of  $F_0, F_{11}, F_2$ . This system always has a solution, the only problem is to check that the corresponding series is convergent, i.e. to find the upper bounds for  $|\zeta_k|$ . It is clear that if we replace all coefficients of  $P_k$  by something bigger, then the solution  $\zeta_k$  becomes bigger and similarly one can replace  $(k + \frac{3}{2}) \rightarrow 2$ . If  $F_0, F_{11}, F_2$  are analytic functions in  $x, \zeta_0$ for  $|x|, |\zeta_0| \leq \varepsilon_1$ , their Taylor coefficients are less than the corresponding coefficients of the functions  $Ax^m(\varepsilon_1 - 2x)^{-1}$ ,  $A(\varepsilon_1 - 2x)^{-1}$  and  $A(\varepsilon_1 - 2x)^{-1}(\varepsilon_1 - 2\zeta_0)^{-1}$ , where A is a sufficiently big number. Hence, the coefficients solving (3.1) are less than the coefficients of the solution of the algebraic equation

$$\frac{2\zeta_0}{x} = \frac{A(x^m + \zeta_0)}{(\varepsilon_1 - 2x)} + \frac{A\zeta_0^2}{x(\varepsilon_1 - 2\zeta_0)(\varepsilon_1 - 2x)}$$

One can easily check that the solution of this quadratic equation is an analytic function at x = 0, hence the coefficients solving (3.1) give us an analytic function at x = 0.

*Proof of Lemma 2.* We start from the technical proposition, whose proof is given after the proof of Lemma 2.

**Proposition 1** Set  $L_l^{(\zeta)}(x, y) := \frac{\partial^l}{\partial x^l} L^{(\zeta)}(x, y)$ , and denote  $L_l^{(\zeta)}$  the integral operator in  $L_2[\sigma_{\varepsilon}]$  with this kernel. Then

$$\sum |\eta_k| (D_\sigma \varphi_k, \varphi_k) \le C(\varepsilon) \Big( \operatorname{Tr} L_2^{(\zeta)} L_2^{(\zeta)*} + \operatorname{Tr} L_1^{(\zeta)} L_1^{(\zeta)*} \Big)^{1/2},$$
(3.2)

$$\sum |\eta_k| (\varphi_k^{(l)}, \varphi_k^{(l)})_{\varepsilon/2} \le C_l(\varepsilon) \Big( \operatorname{Tr} L_{2l+1}^{(\zeta)} L_{2l+1}^{(\zeta)*} + \operatorname{Tr} L_1^{(\zeta)} L_1^{(\zeta)*} \Big)^{1/2}.$$
(3.3)

Here and below we denote by  $(.,.)_{\varepsilon/2}$  the standard scalar product in  $L_2[\sigma_{\varepsilon/2}]$ .

The Schwartz inequality and (3.2) imply that  $K_+$  is a Hilbert-Schmidt matrix, hence, its eigenvalues  $\mu_k \to 0$ , and therefore  $\mu_k \leq \frac{1}{2}$  for all k except may be a finite set  $\{k_1, \ldots, k_\ell\} := I$ . Moreover, the definition of  $K^+$ , the standard properties of the operator norm and (1.21) yield for  $\varphi = \sum u_k \varphi_k$ 

$$(K^+u, u) = ((L_+^{(\zeta)})^{1/2} \overline{D}_{\sigma} (L_+^{(\zeta)})^{1/2} \varphi, \varphi) \le ||(L_+^{(\zeta)})^{1/2} \overline{D}_{\sigma} (L_+^{(\zeta)})^{1/2} ||$$
  
=  $||\overline{D}_{\sigma}^{1/2} L_+^{(\zeta)} \overline{D}_{\sigma}^{1/2} || \le ||\overline{D}_{\sigma}^{1/2} L_{\sigma} \overline{D}_{\sigma}^{1/2} || \le 1,$ 

where  $L_{+}^{(\zeta)}$  is a positive part of the operator  $L^{(\zeta)}$  and  $L_{\sigma}$  is defined in (1.21). Note that since the definition of  $D_{\sigma}$  includes projection  $\Pi$  on the interval  $\sigma$ , we can use the inequality  $\Pi L_{+}^{(\zeta)} \Pi \leq -L_{\sigma}$ . Hence,  $((L_{+}^{(\zeta)})^{1/2} \overline{D}_{\sigma} (L_{+}^{(\zeta)})^{1/2} \varphi, \varphi) = 1$  for some  $\varphi$  only if for  $\tilde{\varphi} = (L_{+}^{(\zeta)})^{1/2} \varphi$ we have

$$L_{+}^{(\zeta)}\overline{D}_{\sigma}\tilde{\varphi} = \tilde{\varphi} \Rightarrow (\overline{D}_{\sigma}L_{+}^{(\zeta)}\overline{D}_{\sigma}\tilde{\varphi},\tilde{\varphi}) = (\overline{D}_{\sigma}\tilde{\varphi},\tilde{\varphi}) = -(\overline{D}_{\sigma}L_{\sigma}\overline{D}_{\sigma}\tilde{\varphi},\tilde{\varphi}),$$
(3.4)

where the last equality follows from (1.21). On the other hand,  $\Pi L_{+}^{(\zeta)} \Pi \leq -L_{\sigma}$ , hence the above equality is possible only if  $L_{+}^{(\zeta)} \overline{D}_{\sigma} \tilde{\varphi} = L \overline{D}_{\sigma} \tilde{\varphi}$ . But  $\Pi L_{+}^{(\zeta)} \Pi \phi = -L_{\sigma} \phi$ , only if

$$\int_{\sigma} \log |\zeta(\lambda) - \zeta(\mu)|^{-1} \phi(\mu) \phi(\lambda) d\lambda d\mu = 0,$$

which contradicts to the positivity of the operator with the kernel  $\log |\zeta(\lambda) - \zeta(\mu)|^{-1}$ . Thus,  $\sup_{k \in I} \mu_k \leq 1 - \delta_1$  with  $\delta_1 > 0$ . Choosing  $\delta = \min\{\delta_1, \frac{1}{2}\}$  we obtain (2.7).

To prove (2.8), we denote

$$(A_0\bar{u},\bar{u}) := \sum_{k,j \le M} u_k u_j |\eta_k|^{1/2} |\eta_j|^{1/2} (\varphi_k^{(16)},\varphi_j^{(16)})_{\varepsilon/2}$$
(3.5)

(we need the 16th derivative here to control the 16th derivative of  $h_u$  in Lemma 3) and set

$$U_{2} = \{ \bar{u} : (A_{0}\bar{u}, \bar{u}) \leq \log^{2} n \land (K^{+}\bar{u}, \bar{u}) + (K^{-}\bar{u}, \bar{u}) \geq k_{*} \log \varepsilon_{n}^{-1} \},$$
(3.6)  

$$U_{3} = \{ \bar{u} : \log^{2} n \leq (A_{0}\bar{u}, \bar{u}) \leq n \log^{2} n \},$$
  

$$U_{4} = \{ \bar{u} : n \log^{2} n \leq (A_{0}\bar{u}, \bar{u}) \leq C_{*}n^{2} \},$$
  

$$U_{5} = \{ \bar{u} : C_{*}n^{2} \leq (A_{0}\bar{u}, \bar{u}) \}$$

with sufficiently large *n*-independent  $C_*$ . One can see easily that

$$U_1^c \subset U_2 \cup U_3 \cup U_4 \cup U_5,$$

Below we will often use the following evident statement

**Proposition 2** For any semi-infinite matrix A > 0 such that  $\sum_{i=1} A_{ii} < \infty$  and  $||A|| < 1-\delta$  $(\delta > 0)$ 

$$\left(\frac{\beta}{8\pi}\right)^{M/2} \int e^{-\beta(\bar{u},\bar{u})/8} e^{\beta \sum (A\bar{u},\bar{u})/8} d\bar{u} \le C \tag{3.7}$$

with some M-independent C.

In particular, (3.7) is true for  $(A\bar{u}, \bar{u}) = (K^+ \bar{u}, \bar{u}) + \tau (A_0 \bar{u}, \bar{u})$ , with sufficiently small  $\tau > 0$ , since we proved above that  $||K^+|| \le 1 - \delta$  and Proposition 1 guarantees that  $\sum_{i=1} A_{ii} < \infty$ . By the Schwartz inequality

By the Schwartz inequality,

$$\begin{aligned} |\Re h_u(\bar{\lambda})| &= \Big| \sum_{k \in I_+} u_k \sqrt{\eta}_k \varphi_k(\lambda) \Big| \le ||\bar{u}|| \Big| \sum_{k \in I_+} \eta_k |\varphi_k(\lambda)|^2 \Big| \\ &\le ||\bar{u}|| (L_+^{(\zeta)}(\lambda,\lambda))^{1/2} \le C ||\bar{u}||. \end{aligned}$$

Hence

$$\langle \Phi_k(\bar{\zeta})e^{\beta\mathcal{N}_n[\dot{h}_u]/2} \rangle_{*,n} \le (nC)^k e^{\beta \max_\lambda \{n|\Re h_u(\bar{\lambda})|\}/2} \le e^{nC_1(1+||\bar{u}||)},$$

where we used the trivial bound

$$|\Phi_k| \le (nC)^k. \tag{3.8}$$

Then, using the fact that the matrix  $A_0$  defined by the quadratic form  $(A_0\bar{u},\bar{u})$  is bounded (in view of Propositions 1), we have in  $U_5 ||\bar{u}||^2 ||A_0|| \ge (A_0\bar{u},\bar{u}) > n^2 C_*$ . Hence for sufficiently large  $C_*$  the integral

$$\int_{U_5} du e^{-\beta(\bar{u},\bar{u})/8} \langle \Phi_k(\bar{\zeta}) e^{\beta \mathcal{N}_n[\dot{h}_u]/2} \rangle_{*,n}$$
  
$$\leq \int_{||\bar{u}||^2 > n^2 C_*/||A_0||} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8} e^{nC(1+||\bar{u}||)} \leq e^{-Cn^2}.$$

For  $u \in U_4$ 

$$n^{-2} \int |\Re h_u^{(16)}(\lambda)|^2 d\lambda = n^{-2} (A_0 \bar{u}, \bar{u}) \le C_*.$$

Thus  $n^{-1}\Re h_u(\lambda)$  is a Hölder function for  $u \in U_4$ , and we can use the result of [3], according to which

$$Z_n[V^* - n^{-1}\Re h_u] \le \exp\left\{\frac{\beta n^2}{2} \max_{m \in \mathcal{M}_1^+[\sigma_{\varepsilon/2}]} \{L[m,m] - (m, V^* - n^{-1}h_u)\} + Cn\log n\right\},$$

where  $\mathcal{M}_1^+[\sigma_{\varepsilon/2}]$  is a set of positive unit measures with supports belonging to  $\sigma_{\varepsilon/2}$ . Since

$$-V^*(\lambda) \le -2L[\rho_{sc}](\lambda), \quad \lambda \in \sigma_{\varepsilon},$$

we have

$$\max_{m \in \mathcal{M}_{1}^{+}[\sigma_{\varepsilon}]} \{ L[m,m] - (m, V^{*} - n^{-1} \Re h_{u}) \}$$

$$\leq \max_{m \in \mathcal{M}_{1}^{+}[\sigma_{\varepsilon}]} \{ L[m,m] - (m, 2\mu_{\alpha}^{-1}L[\rho_{sc}] - n^{-1} \Re h_{u}) \}$$

$$\leq \max_{m \in \mathcal{M}_{1}[\sigma_{\varepsilon}]} \{ L[m,m] - (m, 2L[\rho_{sc}] - n^{-1} \Re h_{u}) \} =: E(\bar{u}).$$
(3.9)

Here  $\mathcal{M}_1[\sigma_{\varepsilon/2}]$  is a set of all signed unit measures with supports belonging to  $\sigma_{\varepsilon/2}$ . It is easy to see that, if we remove the condition of positivity of measures, then the maximum point  $\rho_1$  is uniquely defined by the conditions:

$$2L[\rho_1](\lambda) - 2L[\rho_{sc}](\lambda) - n^{-1}\Re h_u(\lambda) = \text{const}, \quad \lambda \in \sigma_{\varepsilon}, \quad \int_{\sigma_{\varepsilon}} \rho_1 = 1.$$

Hence  $\rho_1 = \rho_{sc} + \frac{1}{2}D_{\sigma_{\varepsilon}} \Re h_u$  and the r.h.s. of (3.9) takes the form

$$E(\bar{u}) = -L[\rho_{sc}, \rho_{sc}] + \frac{n^{-2}}{4}(\bar{D}_{\sigma_{\varepsilon}}\Re h_u, \Re h_u) + n^{-1}(\dot{\Re}h_u, \rho_{sc}).$$

But by the definition of  $\dot{h}_u$   $(\dot{h}_u, \rho_{sc}) = 0$ . Hence

$$E(\bar{u}) = -L[\rho_{sc}, \rho_{sc}] + \frac{n^{-2}}{4}(K_{\varepsilon}^{+}\bar{u}, \bar{u}) + O(n^{-1}\log n),$$

where  $K_{\varepsilon}^+$  is defined by the same way as  $K^+$  (see (2.6)), but with  $\bar{D}_{\sigma}$  replaced by  $\bar{D}_{\sigma_{\varepsilon/2}}$ . These relations and (3.8) yield

$$\langle \Phi_k(\bar{\zeta}) e^{\beta \mathcal{N}_n[h_u]/2} \rangle_{*,n} \leq (Cn)^k e^{\frac{\beta}{8}(K_{\varepsilon}^+ \bar{u}, \bar{u}) + O(n\log n)}$$

Then the Chebyshev inequality for sufficiently small  $\tau$  and (3.7) yield

$$\left(\frac{\beta}{8\pi}\right)^{M/2} \int_{U_4} e^{-\beta(\bar{u},\bar{u})/8} \langle \Phi_k(\bar{\zeta}) e^{\beta\mathcal{N}_n[\dot{h}_u]/2} \rangle_{*,n} d\bar{u}$$

$$\leq \left(\frac{\beta}{8\pi}\right)^{M/2} e^{O(n\log n)} \int e^{-\beta((I-K_{\varepsilon}^+)\bar{u},\bar{u})/8 + \tau((A_0\bar{u},\bar{u})-n\log^2 n)} d\bar{u} \leq e^{-\tau n\log^2 n/2}.$$

$$(3.10)$$

For  $u \in U_3$  (3.8) and (1.18) imply

$$\langle \Phi_k(\bar{\zeta}) e^{\beta \mathcal{N}_n[\dot{h}_u]/2} \rangle_{*,n} \leq (Cn)^k \exp\left\{ \frac{\beta}{4} (K^+ \bar{u}, \bar{u}) + (\bar{u}, \bar{q}) + C + Cn^{-1} ||\Re h_u^{(6)}||_2^3 + ||\Re h_u^{'}||_2^3 \right\}$$

$$\leq \exp\left\{ \frac{\beta}{4} (K^+ \bar{u}, \bar{u}) + n^{-1/3} (A_0 \bar{u}, \bar{u}) + C \right\},$$

$$(3.11)$$

where we used that the vector  $\bar{q} = (q_1, \ldots, q_M), q_k = \eta_k^{1/2} \beta(\bar{D}_\sigma \varphi_k, \zeta') + \eta_k^{1/2}(\nu_\beta, \varphi_k)$  is bounded (it is easy to check by the Schwartz inequality) and that

$$\begin{aligned} ||\Re h_u^{(l)}||_2^2 &\leq C_l (1+||\Re h_u^{(16)}||_2^2) = C(1+(A_0\bar{u},\bar{u})), \quad l=1,6, \\ n^{-1} (A_0\bar{u},\bar{u})^{3/2} &\leq \log n n^{-1/2} (A_0\bar{u},\bar{u}) \leq n^{-1/3} (A_0\bar{u},\bar{u}), \quad \bar{u} \in U_3. \end{aligned}$$

Then, similarly to (3.10), the Chebyshev inequality with sufficiently small  $\tau$  yields

$$\left(\frac{\beta}{8\pi}\right)^{M/2} \int_{U_3} e^{-\beta(\bar{u},\bar{u})/8} \langle \Phi_k(\bar{\zeta}) e^{\beta\mathcal{N}_n[\dot{h}_u]/2} \rangle_{*,n} d\bar{u}$$

$$\leq \left(\frac{\beta}{8\pi}\right)^{M/2} (Cn)^k \int e^{-\beta((I-K^+)\bar{u},\bar{u})/8 + (\bar{u},\bar{q}) + (\tau+n^{-1/3})(A_0\bar{u},\bar{u}) - \tau\log^2 n} du \leq e^{-\tau\log^2 n/2}$$

Finally, using the bound (2.10) for  $\bar{u} \in U_2$  and again the Chebyshev inequality with sufficiently small  $\tau$ , we obtain the bound, finishing the proof of the lemma:

$$\left(\frac{\beta}{8\pi}\right)^{M/2} \int_{U_2} e^{-\beta(\bar{u},\bar{u})/8} \left| \langle \Phi_k(\bar{\zeta}) e^{\beta\mathcal{N}_n[\dot{h}_u]/2} \rangle_{*,n} \right| d\bar{u} \qquad (3.12)$$

$$\leq C_{\Phi_k} \left(\frac{\beta}{8\pi}\right)^{M/2} \int \int d\bar{u} e^{-\beta((I-K^+)\bar{u},\bar{u})/8 + (\bar{u},\bar{q})} \\
\cdot e^{n^{-1/3}(A_0\bar{u},\bar{u}) + \tau((K^+\bar{u},\bar{u}) + (K^-\bar{u},\bar{u}) - k_*\log\varepsilon_n^{-1})} \leq C_3 e^{\tau k_*\log\varepsilon_n}.$$

Proof of Proposition 1. By (1.21)

$$\sum |\eta_k| (\bar{D}_{\sigma} \varphi_k, \varphi_k) = \sum |\eta_k| (\bar{D}_{\sigma}^2 \varphi_k, (-L_{\sigma}) \varphi_k)$$

$$\leq \left( \sum |\eta_k|^2 (\bar{D}_{\sigma}^2 \varphi_k, \bar{D}_{\sigma}^2 \varphi_k) \right)^{1/2} \left( \sum (L_{\sigma}^2 \varphi_k, \varphi_k) \right)^{1/2}$$

$$\leq C \left( \int dx dy \left( \sum \eta_k (\bar{D}_{\sigma}^2 \varphi_k) (x) \varphi_k(y) \right)^2 = C \int |\bar{D}_{\sigma}^2 L^{(\zeta)}(x, y)|^2 dx dy.$$
(3.13)

The last factor in the second line here is bounded since  $(-L_{\sigma})$  is a Hilbert-Schmidt operator. In addition, according to (1.20), for any h such that  $h^{(m)} \in L_2[\sigma]$ 

$$(\bar{D}_{\sigma}^{2m}h,h) = C \int_{0}^{\pi} \left| \frac{d^{m}}{d\theta^{m}} h(2\cos\theta) \right|^{2} d\theta = C \int_{-2}^{2} \left| \left( |X^{1/2}(x)| \frac{d}{dx} \right)^{m} h(x) \right|^{2} \frac{dx}{|X^{1/2}(x)|} \qquad (3.14)$$
$$\leq C \sum_{p=1}^{m} ||h^{(p)}||_{2}^{2} \leq \tilde{C}(||h^{(m)}||_{2} + ||h^{'}||_{2}).$$

Here we used that for  $p \leq m-1$ ,  $||h^{(p)}||_{\infty} \leq C(||h^{(p)}||_2 + ||h^{(p+1)}||_2)$ , hence the last integral in (3.14) is convergent. Applying (3.14) to the r.h.s. of (3.13) we obtain (3.2).

To obtain (3.3), we write

$$\int_{\sigma_{\varepsilon/2}} |h^{(l)}(x)|^2 dx = \int_{|\cos\theta| \le \frac{1+\varepsilon/2}{1+\varepsilon}} \left| \left( \frac{1}{\sin\theta} \frac{d}{d\theta} \right)^l h(2(1+\varepsilon)\cos\theta) \right|^2 \sin\theta d\theta$$
$$\le \tilde{C}_l(\varepsilon) \int_0^\pi \left| \left( \frac{d}{d\theta} \right)^l h(2(1+\varepsilon)\cos\theta) \right|^2 d\theta = C_l'(\varepsilon)(\bar{D}_{\sigma_\varepsilon}^{2l}h,h).$$

Hence, similarly to (3.13)-(3.14),

$$\sum |\eta_k| (\varphi_k^{(l)}, \varphi_k^{(l)})_{\varepsilon/2} \leq C_l'(\varepsilon) \sum |\eta_k| (\bar{D}_{\sigma_\varepsilon}^{2l} \varphi_k, \varphi_k) = \sum |\eta_k| (\bar{D}_{\sigma_\varepsilon}^{2l+1} \varphi_k, (-L_{\sigma_\varepsilon}) \varphi_k)$$
$$\leq C C_l'(\varepsilon) \Big( \sum |\eta_k|^2 (\bar{D}_{\sigma_\varepsilon}^{2l+1} \varphi_k, \bar{D}_{\sigma_\varepsilon}^{2l+1} \varphi_k) \Big)^{1/2} \leq C_l(\varepsilon) (\operatorname{Tr} L_{2l+1}^{(\zeta)} L_{2l+1}^{(\zeta)*} + \operatorname{Tr} L_1^{(\zeta)} L_1^{(\zeta)*})^{1/2}.$$

Proof of Lemma 3. The idea is to consider

$$V_{h} = \frac{\lambda^{2}}{2} + \frac{1}{n}h(\lambda) - c_{1}\frac{\lambda}{n} - c_{2}\frac{\lambda^{2}}{2n} := V_{*} + \frac{1}{n}\tilde{h}$$
(3.15)

with some appropriate  $c_1$  and  $c_2$  as a new potential and to apply to it the above procedure with the change of variables. But since it is possible only for the potentials whose support of the equilibrium measure is [-2, 2], we need to have two equalities:

$$\int_{\sigma} \frac{V_h'(\lambda)d\lambda}{X_{\sigma}^{1/2}(\lambda)} = 0, \quad \pi^{-1} \int_{\sigma} \frac{V_h'(\lambda)\lambda d\lambda}{X_{\sigma}^{1/2}(\lambda)} = 1.$$

Here the first equality is a necessary condition to have a bounded solution of the singular integral equation which can be obtained by the differentiation of (1.10), and the second equality provides the condition that the integral of the corresponding density  $\rho_h$  is 1. Thus we have to choose

$$c_1(h) = \pi^{-1} \int \frac{h'(\lambda)d\lambda}{X^{1/2}(\lambda)}, \quad c_2(h) = \pi^{-1} \int \frac{\lambda h'(\lambda)d\lambda}{2X^{1/2}(\lambda)}.$$

Solving equation (2.1) for  $V_h$ , we obtain uniformly in  $\lambda \in \sigma_{\varepsilon}$ 

$$\zeta_h = \lambda - \frac{1}{n}\tilde{\zeta}_h, \quad \tilde{\zeta}'_h(\lambda) = \frac{\rho_{\tilde{h}}(\lambda)}{\rho_{sc}(\lambda)} + O(n^{-1}||h''||_2^2). \tag{3.16}$$

where  $\rho_{\tilde{h}}$  is the equilibrium density, corresponding to  $\tilde{h}$ . According to (1.14),

$$\frac{\rho_{\tilde{h}}(\lambda)}{\rho_{sc}(\lambda)} = \int_{\sigma} \frac{\tilde{h}'(\mu)d\mu}{(\lambda-\mu)X_{\sigma}^{1/2}(\mu)},$$

hence, by the assumptions of the lemma  $||\zeta_h^{(14)}|| \leq C \log^s n$ . Then the correspondent compact operator kernel  $L^{(\zeta_h)}(\lambda,\mu)$  has the form

$$L^{(\zeta_h)}(\lambda,\mu) = \log \frac{\zeta_h(\lambda) - \zeta_h(\mu)}{\lambda - \mu} = \log \left(1 + \frac{1}{n} \frac{\tilde{\zeta}_h(\lambda) - \tilde{\zeta}_h(\mu)}{\lambda - \mu}\right) = \frac{1}{n} K_h(\lambda,\mu),$$

where  $K_h(\lambda,\mu) = K_h(\mu,\lambda)$  and for any  $\mu$   $K_h(.,\mu)$  is a real analytic function bounded by  $C \log^s n$ . Let  $\{\kappa_k, \varphi_{h,k}\}$  be eigenvalues and eigenvectors of  $K_h$ . Then we obtain that the *k*th correlation function of the Hamiltonian with the potential (3.15) at the point  $(\lambda_0 + x_1/n\rho(\lambda_0), \ldots, \lambda_0 + x_k/n\rho(\lambda_0))$  coincides with that at the point  $(\zeta_h(\lambda_0 + x_1/n\rho(\lambda_0)), \ldots, \zeta_h(\lambda_0 + x_k/n\rho(\lambda_0)))$  for the Hamiltonian (cf (1.15))

$$H^{(\zeta_h)}(\bar{\lambda}) = -n \sum \left(\frac{\lambda_i^2}{2} \left(1 - \frac{c_2}{n}\right) - \frac{c_1}{n} \lambda_i\right) + \sum_{i \neq j} \log |\lambda_i - \lambda_j|$$
  
+ 
$$\frac{1}{n} \sum_{k=1}^{\infty} \kappa_k \left(\sum_j \left((\varphi_{h,k}(\lambda_j) - (\varphi_{h,k}, \rho_{sc})\right)\right)^2 + \frac{1}{n} (\frac{2}{\beta} - 1) \sum \zeta^{(1)}(\lambda_i), \quad (3.17)$$

where  $\zeta_h^{(1)}(\lambda_i) = n \log(1 + n^{-1} \tilde{\zeta}'_h)$ . Taking  $M = [\log^2 n]$ , we obtain like before that we can restrict the summation above by k = M. Hence we get similarly to (2.4)

$$\frac{\langle \Phi_k e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n}}{\langle e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n}} = I_n^{-1}[\beta,\zeta_h] \left(\frac{\beta}{8\pi}\right)^{M/2} \int e^{-\beta(\bar{u},\bar{u})/8} d\bar{u} \langle \tilde{\Phi}_k e^{\beta \mathcal{N}_n[c_1\ell_1 + c_2\ell_2 + n^{-1/2}s_{\bar{u}}]/2} \rangle_{*,n} + o(1)$$
$$I_n[\beta,\zeta_h] := \left(\frac{\beta}{8\pi}\right)^{M/2} \int e^{-\beta(\bar{u},\bar{u})/8} d\bar{u} \langle e^{\beta \mathcal{N}_n[c_1\ell_1 + c_2\ell_2 + n^{-1/2}s_{\bar{u}}]/2} \rangle_{*,n},$$

where  $\ell_1(\lambda) = \lambda$ ,  $\ell_2(\lambda) = \lambda^2/2$ , and

$$s_{\bar{u}}(\lambda) = \sum_{k} u_k \sqrt{\kappa_k} \dot{\varphi}_{h,k}(\lambda) + n^{-1/2} (\frac{2}{\beta} - 1) \zeta^{(1)}(\lambda), \quad \tilde{\Phi}_k(\bar{\lambda}) := \Phi_k(\zeta_h(\bar{\lambda}), \lambda_0).$$

Then, changing variables once more

$$\lambda_i \to \zeta_c(\lambda) := (1 - c_2/n)^{1/2} \Big(\lambda_i - \frac{c_1}{n(1 - c_2/n)}\Big),$$

we obtain that

$$\frac{\langle \Phi_k e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n}}{\langle e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n}} = \left(\frac{\beta}{8\pi}\right)^{M/2} \int e^{-\beta(\bar{u},\bar{u})/8} d\bar{u} \langle \widehat{\Phi}_k e^{\beta \mathcal{N}_n[n^{-1/2}s_{\bar{u},c}]/2} \rangle_{*,n} / \bar{I}_n[\beta,\zeta_h] + o(1),$$
$$I_n[\beta,\zeta_h] := \left(\frac{\beta}{8\pi}\right)^{M/2} \int e^{-\beta(\bar{u},\bar{u})/8} d\bar{u} \langle e^{\beta \mathcal{N}_n[n^{-1/2}s_{\bar{u},c}]/2} \rangle_{*,n},$$

with  $\widehat{\Phi}_k(\lambda) = \widetilde{\Phi}_k(\zeta_c(\lambda))$  and  $s_{u,c}(\lambda) = s_u(\zeta_c(\lambda))$ . Represent  $\mathbb{R}^M = \widetilde{U}_1 \cup \widetilde{U}_2 \cup \widetilde{U}_3$ , where

$$\begin{split} \tilde{U}_1 &= \{ \bar{u} : (A_{h,0}\bar{u}, \bar{u}) \le \log^{2s+2} n \}, \\ \tilde{U}_2 &= \{ \bar{u} : \log^{2s+2} n \le (A_{h,0}\bar{u}, \bar{u}) \le C_* n \log^{4s} n \}, \\ \tilde{U}_3 &= \{ \bar{u} : C_* n \log^{4s} n \le (A_{h,0}\bar{u}, \bar{u}) \} \\ (A_{h,0})_{ij} &= |\kappa_i|^{1/2} |\kappa_j|^{1/2} (\varphi_{h,i}^{(6)}, \varphi_{h,j}^{(6)})_{\varepsilon/2}. \end{split}$$

Note, that Proposition 1 and the assumptions of Lemma 3 yield

$$\sum (A_{h,0})_{ii} \le C \log^{2s} n, \quad ||A_{h,0}|| \le C \log^{2s} n.$$
(3.18)

Repeating for  $\tilde{U}_3$  the argument used for  $U_5$  in Lemma 2, we get

$$||n^{-1/2}s_{u,c}||_{\infty} \le C_1 n^{-1/2} (A_{h,0}\bar{u},\bar{u})^{1/2} \le C_2 n^{-1/2} ||A_{h,0}|| \, ||\bar{u}|| \le C_3 n^{-1/2} \log^{2s} n ||\bar{u}||.$$

Hence

$$\left| \langle \Phi_k e^{\beta \mathcal{N}_n[n^{-1/2} s_{u,c}]/2} \rangle_{*,n} \right| \le (Cn)^k e^{\beta n ||n^{-1/2} s_{u,c}||_{\infty}/2} \le e^{c_1 n^{1/2} \log^{2s} n ||\bar{u}||}.$$

Then, since  $||\bar{u}|| \ge (A_{h,0}\bar{u},\bar{u})^{1/2}/||A_{h,0}||^{1/2} \ge C_* n \log^{3s} n$  in  $\tilde{U}_3$ , we have

$$\begin{split} & \int_{\tilde{U}_3} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8} \Big| \langle \Phi_k e^{\beta \mathcal{N}_n[h]/2} \rangle_{*,n} \Big| \\ & \leq \int_{||\bar{u}||^2 \ge C_* n \log^{3s} n} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\beta(\bar{u},\bar{u})/8 + c_1 n^{1/2} \log^{2s} n ||\bar{u}||} \le e^{-nc} d\bar{u} e^{-\alpha} d\bar{$$

For  $\bar{u} \in \tilde{U}_2$ 

$$||n^{-1/2} s_{u,c}^{(6)}||_2 \le C_* \log^{2s} n,$$

hence we can use here (1.18) and then the argument used for  $U_3$ , but replacing  $\tau$  in the Chebyshev inequality by  $\tau \log^{-2s} n$ . Since by (3.18) the matrix  $\tau \log^{-2s} n A_{0,h}$  for small  $\tau$  satisfy conditions of Proposition 2, we obtain that the integral in  $U_2$  is  $O(e^{-c \log^2 n})$ . Thus it suffices to study  $\bar{u} \in \tilde{U}_1$ . But here

$$\begin{aligned} \left| \langle \widehat{\Phi}_k(\bar{\lambda}) e^{\beta(\mathcal{N}_n[n^{-1/2}s_{\bar{u}}]/2)} \rangle_{*,n} - \langle \widehat{\Phi}_k(\bar{\lambda}) \rangle_{*,n} \right| &\leq \langle \widehat{\Phi}_k^2(\lambda) \rangle_{*,n}^{1/2} \langle \left| e^{n^{-1/2}\mathcal{N}[s_{\bar{u},c}]} - 1 \right|^2 \rangle_{*,n}^{1/2}, \\ &\leq C n^{-1/2} ((D_{\sigma} \Re s_{\bar{u},c}, \Re s_{\bar{u},c}) + (D_{\sigma} \Im s_{\bar{u},c}, \Im s_{\bar{u},c}))^{1/2} \leq n^{-\kappa}. \end{aligned}$$

Here we have used (1.23) which gives the bound for  $\langle \widehat{\Phi}_k^2 \rangle_{*,n}$  and (1.18) in the case (ii), according to which for  $\bar{u} \in \widetilde{U}_1$  we have for any bounded t

$$\langle e^{tn^{-1/2}\mathcal{N}[s_{\bar{u}}]} \rangle_{*,n} = e^{t^2n^{-1}(D_{\sigma}s_{\bar{u}},s_{\bar{u}})}(1+o(1)).$$

Since all  $\phi_j$  of  $\Phi_k$  are smooth and have finite supports, (1.23) imply that

$$\begin{split} \langle \widehat{\Phi}_k(\bar{\lambda};\lambda_0) \rangle_{*,n} = & \langle \Phi_k(\bar{\lambda} - n^{-1}(c_1 + c_2\lambda_0/2 - \widetilde{\zeta}_h(\lambda_0)) \rangle_{*,n} + O(n^{-\kappa}) \\ = & \langle \Phi_k(\bar{\lambda}) \rangle_{*,n} + O(\varepsilon_n) + O(n^{-\kappa}). \end{split}$$

Combining the above bounds, we get the assertion of Lemma 3.

#### 

Proof of Lemma 4. Introduce the analytic function

$$f_n(t) := C_1^{-1} e^{-t^2/2} \varepsilon_n^{-1} F_n(t), \quad t \in D.$$

Then

$$|f_n(t)| \le 1, \quad t \in \gamma = [-\log^{1/2} \varepsilon_n^{-1}, \log^{1/2} \varepsilon_n^{-1}].$$

 $|f_n(t)| \leq 1, \quad t \in T = [100] \quad e_n, \text{ for } e_n$ . Moreover,  $|f_n(t)| \leq C_2 C_1^{-1} \varepsilon_n^{-2}, t \in D$ . Then, by the theorem on two constants (see [13]), we conclude that

$$\log |f_n(t)| \le 2(1 - \omega(t; \gamma, D))(\log \varepsilon_n^{-1} + \log(C_2/C_1)^{1/2})$$

where  $\omega(t;\gamma,D)$  is the harmonic measure of the set  $\gamma$  with respect to the domain D at the point  $t \in D$ . It is well-known (see again [13]) that

$$\omega(t;\gamma,D) = 1 - \frac{2}{\pi} \Im \log \frac{1 + t/\log^{1/2} \varepsilon_n^{-1}}{1 - t/\log^{1/2} \varepsilon_n^{-1}}$$

Hence

$$1 - \omega(t; \gamma, D) \le \frac{6\Im t}{\pi \log^{1/2} \varepsilon_n^{-1}} \le \frac{1}{4}, \quad t \in \frac{\pi}{24} D = D',$$

and the above inequalities yield

$$\log |f_n(t)| \le \frac{1}{2} (\log \varepsilon_n^{-1} + \log (C_2/C_1)^{1/2}) \quad \Rightarrow \quad |f_n(t)| \le (C_2/C_1)^{1/4} \varepsilon_n^{-1/2}, \ t \in D'.$$

Then from the definition of  $f_n$  we obtain (2.12).

#### Proof of Theorem 3 4

Examining the proof of Theorem 2, one can see that its result can be reformulated as follows. For any real analytic *n*-independent one-cut V and real analytic  $h: ||h'||, ||h^{(6)}|| \leq \log^s n$  the inequalities hold uniformly in h:

$$\left|\frac{\langle \Phi_k(\bar{\lambda},\lambda_0)e^{\beta\mathcal{N}_n[h]/2}\rangle_{V,n}}{\langle e^{\beta\mathcal{N}_n[h]/2}\rangle_{V,n}}\right| \le \tilde{C}_{\Phi},\tag{4.1}$$

$$\left|\frac{\langle \Phi_k(\bar{\lambda},\lambda_0)e^{\beta\mathcal{N}_n[h]/2}\rangle_{V,n}}{\langle e^{\beta\mathcal{N}_n[h]/2}\rangle_{V,n}} - \langle \Phi_k(\bar{\lambda},0)\rangle_{*,n}\right| \le \tilde{\varepsilon}_n \to 0.$$
(4.2)

Note that  $C_{\Phi}$  and  $\tilde{\varepsilon}_n$  depend on  $C_{\Phi}$  and  $\varepsilon_n$  of (1.23), (1.24).

The proof of Theorem 3 is based on these two inequalities and on the results of [26]. Set

$$\mu_{\alpha} = \int_{\sigma_{\alpha}} \rho_{\alpha}(\lambda) d\lambda, \quad \rho_{\alpha} := \mathbf{1}_{\sigma_{\alpha}} \rho, \tag{4.3}$$

$$V_{\alpha}(\lambda) = \mathbf{1}_{\sigma_{\alpha,\varepsilon}}(\lambda)\mu_{\alpha}^{-1}\Big(V(\lambda) - 2\int_{\sigma\setminus\sigma_{\alpha}}\log|\lambda-\mu|\rho(\mu)d\mu\Big)$$
(4.4)

$$\bar{n} := (n_1, \dots, n_q), \quad |\bar{n}| := \sum_{\alpha=1}^q n_\alpha.$$
 (4.5)

It is easy to see that the potential  $V_{\alpha}$  and the equilibrium density  $\mu_{\alpha}^{-1}\rho_{\alpha}$  satisfy (1.10) and (1.13) in  $\sigma_{\alpha,\varepsilon}$ , hence  $V_{\alpha}$  is a real analytic potential of generic behavior in  $\sigma_{\alpha,\varepsilon}$  and so in each interval we can apply (1.18) and (4.1)-(4.2).

Assuming that  $\lambda_0 \in [a_1 + \varepsilon, b_1 - \varepsilon]$  and repeating the argument of Theorem 2 of [26], we obtain (cf(1.17))

$$\langle \Phi_k(\bar{\lambda}_1;\lambda_0) \rangle_{V,n} = \mathcal{I}_n^{-1}[V] \sum_{|\bar{n}|=n} \kappa_{\bar{n}} \left(\frac{\beta}{2\pi}\right)^{Mq} \int du e^{-\frac{\beta}{8}(u,u)}$$

$$\cdot \langle \Phi_k(\bar{\lambda}_1;\lambda_0) e^{\beta \mathcal{N}_{n_1}[\tilde{h}_1]/2} \rangle_{V_1,n_1} \prod_{\alpha=2}^q \langle e^{\beta \mathcal{N}_{n_\alpha}[\tilde{h}_\alpha]/2} \rangle_{V_\alpha,n_\alpha},$$

$$\mathcal{I}_n[V] = \sum_{|\bar{n}|=n} \kappa_{\bar{n}} \left(\frac{\beta}{2\pi}\right)^{Mq} \int du e^{-\frac{\beta}{8}(u,u)} \prod_{\alpha=1}^q \langle e^{\beta \mathcal{N}_{n_\alpha}[\tilde{h}_\alpha]/2} \rangle_{V_\alpha,n_\alpha},$$

$$(4.6)$$

where  $M = [\log^2 n], u := (u^{(1)}, u^{(2)}), \kappa_{\bar{n}}$  are some numbers,

$$\widetilde{h}_{\alpha}(\lambda) = (n_{\alpha} - n\mu_{\alpha})V_{\alpha} + \dot{s}^{(\alpha)}(u,\lambda), \qquad (4.7)$$

$$s^{(\alpha)}(u,\lambda) = \sum_{j,k,\alpha'} \left(\widehat{S}_{j,\alpha';k,\alpha}u^{(1)}_{j,\alpha'} + iS_{j,\alpha';k,\alpha}u^{(2)}_{j,\alpha'}\right)p^{(\alpha)}_{k}(\lambda), \qquad \dot{s}^{(\alpha)}(u,\lambda) = s^{(\alpha)}(u,\lambda) - \frac{n}{n_{\alpha}}\left(s^{(\alpha)}(u,.),\rho_{\alpha}\right).$$

Thus in each interval  $\sigma_{\alpha,\varepsilon}$  we are again in the situation of the one-cut analytic potential with a "small" perturbation  $\tilde{h}_{\alpha}$ . Here  $\{p_k^{(\alpha)}\}_{k=0}^M$  are polynomials on  $\sigma_{\varepsilon}$  of degree at most M and therefore in (1.18) we can use the bound, valid for any  $l = 1, \ldots, 6$  and for any  $\bar{u}$ 

$$(|\partial_{\lambda}^{(l)}s_{u}|, |\partial_{\lambda}^{(l)}s_{u}|) \leq (CM)^{4l} \sum_{\alpha} \left( (D_{\alpha}\Re s_{u}, \Re s_{u}) + (D_{\alpha}\Im s_{u}, \Im s_{u}) \right).$$
(4.8)

The bound follows from the inequality, valid for polynomials with degree not exceeding M:

$$\sup_{\deg p \le M} \frac{(p', p')}{(p, p)} \le CM^4,$$

which can be checked by expanding of an arbitrary polynomial in the sum of the Jacobi polynomials orthonormal on  $\sigma_{\varepsilon}$  without any weight.

The exact forms of positive matrices  $\widehat{S} = \{\widehat{S}_{j,\alpha';k,\alpha}\}_{j,k=1,\dots,M}$ , and  $S = \{S_{j,\alpha';k,\alpha}\}_{j,k=1,\dots,M}$ ,  $\alpha,\alpha'=1,\dots,q$ in (4.7) are not important for us. It will be important only that (4.11) is true.

Moreover, Lemma 2 of [26] implies

$$\mathcal{T}_{\bar{n}} := \kappa_{\bar{n}} \left(\frac{\beta}{2\pi}\right)^{Mq} \int du e^{-\frac{\beta}{8}(u,u)} \prod_{\alpha=1}^{q} \langle e^{\beta \mathcal{N}_{n\alpha}[\tilde{h}_{\alpha}]/2} \rangle_{V_{\alpha},n_{\alpha}} \le C e^{-c(\Delta n,\Delta n)}, \tag{4.9}$$

where  $\Delta n = (\Delta n_1, \dots, \Delta n_q)$ ,  $\Delta n_\alpha = n_\alpha - \mu_\alpha n$ , and  $\mu_\alpha$  were defined in (4.3). This relation and (4.1) yield that for our purposes it suffices to consider in (4.6) only those terms for which

$$(\Delta n, \Delta n) \le c_* \log \tilde{\varepsilon}_n^{-1} \tag{4.10}$$

with any *n*-independent  $c_*$ , hence the *u*-independent part of  $\tilde{h}_{\alpha}$  of (4.7) cannot be too big.

Let us again use (1.18) for  $\langle e^{\beta \mathcal{N}_{n_{\alpha}}[\tilde{h}_{\alpha}]/2} \rangle_{V_{\alpha},n_{\alpha}}$ ,  $\alpha = 1,\ldots,q$ . Similarly to the proof of Theorem 2, the key point is that after the application of (1.18) the real part of the correspondent quadratic form is negative definite, so the integral in  $\bar{u}$  is convergent. More precisely, Lemma 4 of [26] guarantees that there exists  $\delta > 0$  such that

$$\Re\left(\sum_{\alpha=1}^{q} (D_{\sigma_{\alpha}} s^{(\alpha)}, s^{(\alpha)})\right) \le (1-\delta)(\bar{u}, \bar{u}).$$

$$(4.11)$$

Moreover, it is shown (see again Lemma 4, the analog of Lemma 2 of the present paper) that if we define (cf (2.6))

$$U_1 = \{ u := (u^{(1)}, u^{(2)}) : \sum_{\alpha} |(D_{\alpha}\Im\dot{s}_{\alpha}, \Im\dot{s}_{\alpha})| \le k_* \log n \land (u^{(1)}, u^{(1)}) \le \log^4 n \},$$
(4.12)

then the integral over the complement in the r.h.s. of (4.6) is small. But if  $\log \tilde{\varepsilon}_n^{-1} \ll \log n$ , the domain  $U_1$  can be too big, because similarly to the proof of Theorem 2, we need to consider

$$U_0 = \{ u := (u^{(1)}, u^{(2)}) : \sum_{\alpha} (D_{\alpha} \Re s_{\alpha}, \Re s_{\alpha}) + (D_{\alpha} \Im s_{\alpha}, \Im s_{\alpha}) \le k_* \log \varepsilon_n^{-1} \},$$

To estimate the integral in  $U_1 \setminus U_0$ , one should use the Chebyshev inequality like in (3.12). We are left to prove the analog of (4.2) in  $U_0$ . For real  $s_u$  the bound is known because of (4.2), and for the complex s we obtain the bound from Lemma 4, repeating literally the argument used at the end of Section 2.

# References

- Albeverio, S., Pastur, L., Shcherbina, M.: On the 1/n expansion for some unitary invariant ensembles of random matrices. Commun. Math. Phys. 224, 271-305 (2001)
- Borot, G. and Guionnet, A.: Asymptotic expansion of β-matrix models in the one-cut regime. Comm. Math. Phys.317, 447-483 (2013)
- [3] Boutet de Monvel, A., Pastur L., Shcherbina M.: On the statistical mechanics approach in the random matrix theory. Integrated density of states. J. Stat. Phys. 79, 585-611 (1995)
- [4] Bleher, P., Its, A.: Double scaling limit in the random matrix model: the Riemann-Hilbert approach. Comm. Pure Appl. Math. 56, 433–516 (2003)
- [5] Bourgade, P., Erdös, L., Yau, H.-T.: Universality of General β-Ensembles. (http://arxiv.org/abs/1104.2272)
- [6] Bourgade, P., Erdös, L., Yau, H.-T.: Edge Universality of Beta Ensembles (http://arxiv.org/abs/1306.5728v1)
- [7] Claeys, T., Kuijalaars, A.B.J.: Universality of the double scaling limit in random matrix models. Comm. Pure Appl. Math. 59, 1573-1603 (2006)
- [8] Deift, P., Kriecherbauer, T., McLaughlin, K., Venakides, S., Zhou, X.: Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory. Commun. Pure Appl. Math. 52, 1335-1425 (1999)
- [9] Deift, P., Gioev, D.: Universality in random matrix theory for orthogonal and symplectic ensembles. Int. Math. Res. Papers. 2007, 004-116

- [10] Deift, P., Gioev, D.: Universality at the edge of the spectrum for unitary, orthogonal, and symplectic ensembles of random matrices. Comm. Pure Appl. Math. 60, 867-910 (2007)
- [11] Deift, P., Gioev, D., Kriecherbauer, T., Vanlessen, M.: Universality for orthogonal and symplectic Laguerre-type ensembles. J.Stat.Phys 129, 949-1053 (2007)
- [12] Erdos, L., Yau, H.-T. Gap Universality of Generalized Wigner and β-Ensembles (http://arxiv.org/abs/1211.3786v2)
- [13] Evgrafov, M.A.: Analytic Functions. Dover Pubns (1978)
- [14] Gohberg, I., Krein, M.G.: Introduction to the Theory of Linear Nonselfadjoint Operators American Mathematical Soc., 1969 - 378p
- [15] Johansson, K.: On fluctuations of eigenvalues of random Hermitian matrices. Duke Math. J. 91, 151-204 (1998)
- [16] Levin L., Lubinskky D.S.: Universality limits in the bulk for varying measures. Adv. Math. 219, 743-779 (2008)
- [17] M.L.Mehta, M.L.: Random Matrices. New York: Academic Press, (1991)
- [18] Pastur, L., Shcherbina, M.: Universality of the local eigenvalue statistics for a class of unitary invariant random matrix ensembles. J. Stat. Phys. 86, 109-147 (1997)
- [19] Kriecherbauer, T., Shcherbina, M.: Fluctuations of eigenvalues of matrix models and their applications. preprint arxive: math-ph/1003.6121
- [20] Pastur, L., Shcherbina, M.: Bulk universality and related properties of Hermitian matrix models. J.Stat.Phys. 130, 205-250 (2007)
- [21] Pastur, L., Shcherbina, M.: Eigenvalue Distribution of Large Random Matrices. Mathematical Survives and Monographs, V171, American Mathematical Society: Providence, Rhode Island (2011)
- [22] Ramìrez, J., Rider, B., and Viràg, B.: Beta ensembles, stochastic Airy spectrum, and a diffusion, J. Amer. Math. Soc. 24, 919944 (2011)
- [23] Shcherbina, M.: On Universality for Orthogonal Ensembles of Random Matrices Commun.Math.Phys. 285, 957-974 (2009)
- [24] Shcherbina, M.: Edge Universality for Orthogonal Ensembles of Random Matrices J.Stat.Phys 136, 35-50 (2009)
- [25] Shcherbina, M.: Orthogonal and symplectic matrix models: universality and other properties Commun.Math.Phys. 307, 761-790, (2011)
- [26] M.Shcherbina. Fluctuations of linear eigenvalue statistics of  $\beta$  matrix models in the multi-cut regime J.Stat.Phys., **151**, N 6, 1004-1034 (2013)
- [27] Tracy, C.A., Widom, H.: Correlation functions, cluster functions, and spacing distributions for random matrices. J.Stat.Phys. 92, 809-835 (1998)
- [28] Valkò, B., Viràg, B.: Continuum limits of random matrices and the Brownian carousel. Invent. Math. 177 (2009), no. 3, 463508.