

The probabilistic approach to limited packings in graphs

Andrei Gagarin*

Department of Computer Science, Royal Holloway, University of London
Egham, Surrey, TW20 0EX, UK

Vadim Zverovich†

Department of Mathematics and Statistics, University of the West of England
Bristol, BS16 1QY, UK

February 13, 2018

Abstract

We consider (closed neighbourhood) packings and their generalization in graphs. A vertex set X in a graph G is a k -limited packing if for any vertex $v \in V(G)$, $|N[v] \cap X| \leq k$, where $N[v]$ is the closed neighbourhood of v . The k -limited packing number $L_k(G)$ of a graph G is the largest size of a k -limited packing in G . Limited packing problems can be considered as secure facility location problems in networks.

In this paper, we develop a new probabilistic approach to limited packings in graphs, resulting in lower bounds for the k -limited packing number and a randomized algorithm to find k -limited packings satisfying the bounds. In particular, we prove that for any graph G of order n with maximum vertex degree Δ ,

$$L_k(G) \geq \frac{kn}{(k+1)^k \sqrt[k]{\binom{\Delta}{k}} (\Delta+1)}.$$

The problem of finding a maximum size k -limited packing is known to be NP -complete even in split or bipartite graphs.

Keywords: k -limited packings; the probabilistic method; lower and upper bounds; randomized algorithm

*e-mail: andrei.gagarin@rhul.ac.uk

†e-mail: vadim.zverovich@uwe.ac.uk

1 Introduction

We consider simple undirected graphs. If not specified otherwise, standard graph-theoretic terminology and notations are used (e.g., see [1, 2]). We are interested in the classical packings and packing numbers of graphs as introduced in [9], and their generalization, called limited packings and limited packing numbers, respectively, as presented in [6]. In the literature, the classical packings are often referred to under different names: for example, as (distance) 2-packings [9, 12], closed neighborhood packings [10] or strong stable sets [8]. They can also be considered as generalizations of independent (stable) sets which, following the terminology of [9], would be (distance) 1-packings.

Formally, a vertex set X in a graph G is a k -limited packing if for any vertex $v \in V(G)$,

$$|N[v] \cap X| \leq k,$$

where $N[v]$ is the closed neighbourhood of v . The k -limited packing number $L_k(G)$ of a graph G is the maximum size of a k -limited packing in G . In these terms, the classical (distance) 2-packings are 1-limited packings, and hence $\rho(G) = L_1(G)$, where $\rho(G)$ is the 2-packing number.

The problem of finding a 2-packing (1-limited packing) of maximum size is shown to be NP -complete by Hochbaum and Schmoys [8]. In [4], it is shown that the problem of finding a maximum size k -limited packing is NP -complete even for the classes of split and bipartite graphs.

Graphs usually serve as underlying models for networks. A number of interesting application scenarios of limited packings are described in [6], including network security, market saturation, and codes. These and others can be summarized as secure location or distribution of facilities in a network. In a more general sense, these problems can be viewed as (maximization) facility location problems to place/distribute in a given network as many resources as possible subject to some (security) constraints.

2-Packings (1-limited packings) are well-studied in the literature from the structural and algorithmic point of view (e.g., see [8, 9, 10, 11]) and in connection with other graph parameters (e.g., see [3, 7, 9, 10, 12]). In particular, several papers discuss connections between packings and dominating sets in graphs (e.g., see [3, 4, 6, 7, 10]). Although the formal definitions for packings and dominating sets may appear to be similar, the problems have a very different nature: one of the problems is a maximization problem not to break some (security) constraints, and the other is a minimization problem to satisfy some reliability requirements. For example, given a graph G , the definitions imply a simple inequality $\rho(G) \leq \gamma(G)$, where $\gamma(G)$ is the domination number of G (e.g., see [10]). However, the difference between $\rho(G)$ and $\gamma(G)$ can be arbitrarily large as illustrated in [3]: $\rho(K_n \times K_n) = 1$ for the Cartesian product of complete graphs, but $\gamma(K_n \times K_n) = n$.

In this paper, we develop the probabilistic method for k -limited packings in general and for 2-packings (1-limited packings) in particular. In Section 2 we present the probabilistic construction and use it to derive two lower bounds for the k -limited packing number $L_k(G)$. The construction implies a randomized algorithm to find k -limited packings satisfying the lower bounds. The algorithm and its analysis are presented in Section 3. Section 4 shows that one of the lower bounds is asymptotically sharp. Finally, Section 5 provides upper bounds for $L_k(G)$, e.g. in terms of the k -tuple domination number $\gamma_{\times k}(G)$.

Notice that the probabilistic construction and approach are different from the probabilistic constructions used for independent sets and deriving the lower bounds for the

independence number $\alpha(G)$ in [1], pp. 27–28, 91–92. In terms of packings, an independent set in a graph G is a distance 1-packing: for any two vertices in an independent set, the distance between them in G is greater than 1. To the best of our knowledge, the proposed probabilistic method is a new approach to work on packings and related maximization problems.

2 The probabilistic construction and lower bounds

Let $\Delta = \Delta(G)$ denote the maximum vertex degree in a graph G . Notice that $L_k(G) = n$ when $k \geq \Delta + 1$. We define

$$c_t = c_t(G) = \binom{\Delta}{t} \quad \text{and} \quad \tilde{c}_t = \tilde{c}_t(G) = \binom{\Delta + 1}{t}.$$

In what follows, we put $\binom{a}{b} = 0$ if $b > a$.

The following theorem gives a new lower bound for the k -limited packing number. It may be pointed out that the probabilistic construction used in the proof of Theorem 1 implies a randomized algorithm for finding a k -limited packing set, whose size satisfies the bound of Theorem 1 with a positive probability (see Algorithm 1 in Section 3).

Theorem 1 *For any graph G of order n with $\Delta \geq k \geq 1$,*

$$L_k(G) \geq \frac{kn}{\tilde{c}_{k+1}^{1/k} (1+k)^{1+1/k}}. \quad (1)$$

Proof: Let A be a set formed by an independent choice of vertices of G , where each vertex is selected with the probability

$$p = \left(\frac{1}{\tilde{c}_{k+1} (1+k)} \right)^{1/k}. \quad (2)$$

For $m = k, \dots, \Delta$, we denote

$$A_m = \{v \in A : |N(v) \cap A| = m\}.$$

For each set A_m , we form a set A'_m in the following way. For every vertex $v \in A_m$, we take $m - (k - 1)$ neighbours from $N(v) \cap A$ and add them to A'_m . Such neighbours always exist because $m \geq k$. It is obvious that

$$|A'_m| \leq (m - k + 1)|A_m|.$$

For $m = k + 1, \dots, \Delta$, let us denote

$$B_m = \{v \in V(G) - A : |N(v) \cap A| = m\}.$$

For each set B_m , we form a set B'_m by taking $m - k$ neighbours from $N(v) \cap A$ for every vertex $v \in B_m$. We have

$$|B'_m| \leq (m - k)|B_m|.$$

Let us construct the set X as follows:

$$X = A - \left(\bigcup_{m=k}^{\Delta} A'_m \right) - \left(\bigcup_{m=k+1}^{\Delta} B'_m \right).$$

It is easy to see that X is a k -limited packing in G . The expectation of $|X|$ is

$$\begin{aligned} \mathbf{E}[|X|] &\geq \mathbf{E} \left[|A| - \sum_{m=k}^{\Delta} |A'_m| - \sum_{m=k+1}^{\Delta} |B'_m| \right] \\ &\geq \mathbf{E} \left[|A| - \sum_{m=k}^{\Delta} (m-k+1)|A_m| - \sum_{m=k+1}^{\Delta} (m-k)|B_m| \right] \\ &= pn - \sum_{m=k}^{\Delta} (m-k+1)\mathbf{E}[|A_m|] - \sum_{m=k+1}^{\Delta} (m-k)\mathbf{E}[|B_m|]. \end{aligned}$$

Let us denote the vertices of G by v_1, v_2, \dots, v_n and the corresponding vertex degrees by d_1, d_2, \dots, d_n . We will need the following lemma:

Lemma 1 *If $p = \left(\frac{1}{\tilde{c}_{k+1}(1+k)} \right)^{1/k}$, then, for any vertex $v_i \in V(G)$,*

$$\binom{d_i}{m} (1-p)^{d_i-m} \leq \binom{\Delta}{m} (1-p)^{\Delta-m}. \quad (3)$$

Proof: The inequality (3) holds if $d_i = \Delta$. It is also true if $d_i < m$ because in this case $\binom{d_i}{m} = 0$. Thus, we may assume that

$$m \leq d_i < \Delta.$$

Now, it is easy to see that inequality (3) is equivalent to the following:

$$(1-p)^{\Delta-d_i} \geq \binom{d_i}{m} / \binom{\Delta}{m} = \frac{(\Delta-m)!/(d_i-m)!}{\Delta!/d_i!} = \prod_{i=0}^{\Delta-d_i-1} \frac{\Delta-m-i}{\Delta-i}. \quad (4)$$

Further, $\Delta \geq k$ implies $\frac{\Delta}{k} \leq \frac{\Delta-i}{k-i}$, where $0 \leq i \leq k-1$. Taking into account that $\Delta > 0$, we obtain

$$\left(\frac{\Delta}{k} \right)^k \leq \prod_{i=0}^{k-1} \frac{\Delta-i}{k-i} = c_k < \tilde{c}_{k+1}(1+k)$$

or

$$\frac{1}{\tilde{c}_{k+1}(1+k)} < \left(\frac{k}{\Delta} \right)^k.$$

Thus,

$$p^k < \left(\frac{k}{\Delta} \right)^k \quad \text{or} \quad p < \frac{k}{\Delta} \leq \frac{m}{\Delta}.$$

We have $p < \frac{m}{\Delta}$, which is equivalent to $1-p > \frac{\Delta-m}{\Delta}$. Therefore,

$$(1-p)^{\Delta-d_i} > \left(\frac{\Delta-m}{\Delta} \right)^{\Delta-d_i} \geq \prod_{i=0}^{\Delta-d_i-1} \frac{\Delta-m-i}{\Delta-i},$$

as required in (4). ■

Now we go on with the proof of Theorem 1. By Lemma 1,

$$\begin{aligned}
\mathbf{E}[|A_m|] &= \sum_{i=1}^n \mathbf{P}[v_i \in A_m] \\
&= \sum_{i=1}^n p \binom{d_i}{m} p^m (1-p)^{d_i-m} \\
&\leq p^{m+1} \sum_{i=1}^n \binom{\Delta}{m} (1-p)^{\Delta-m} \\
&= p^{m+1} (1-p)^{\Delta-m} c_m n
\end{aligned}$$

and

$$\begin{aligned}
\mathbf{E}[|B_m|] &= \sum_{i=1}^n \mathbf{P}[v_i \in B_m] \\
&= \sum_{i=1}^n (1-p) \binom{d_i}{m} p^m (1-p)^{d_i-m} \\
&\leq p^m \sum_{i=1}^n \binom{\Delta}{m} (1-p)^{\Delta-m+1} \\
&= p^m (1-p)^{\Delta-m+1} c_m n.
\end{aligned}$$

Taking into account that $c_{\Delta+1} = \binom{\Delta}{\Delta+1} = 0$, we obtain

$$\begin{aligned}
\mathbf{E}[|X|] &\geq pn - \sum_{m=k}^{\Delta} (m-k+1) p^{m+1} (1-p)^{\Delta-m} c_m n - \sum_{m=k+1}^{\Delta+1} (m-k) p^m (1-p)^{\Delta-m+1} c_m n \\
&= pn - \sum_{m=0}^{\Delta-k} (m+1) p^{m+k+1} (1-p)^{\Delta-m-k} c_{m+k} n \\
&\quad - \sum_{m=0}^{\Delta-k} (m+1) p^{m+k+1} (1-p)^{\Delta-m-k} c_{m+k+1} n \\
&= pn - \sum_{m=0}^{\Delta-k} (m+1) p^{m+k+1} (1-p)^{\Delta-m-k} n (c_{m+k} + c_{m+k+1}) \\
&= pn - p^{k+1} n \sum_{m=0}^{\Delta-k} (m+1) \tilde{c}_{m+k+1} p^m (1-p)^{\Delta-k-m}.
\end{aligned}$$

Furthermore,

$$\begin{aligned}
(m+1) \tilde{c}_{m+k+1} &= \binom{\Delta-k}{m} \frac{(m+1)! (\Delta+1)!}{(m+k+1)! (\Delta-k)!} \\
&\leq \binom{\Delta-k}{m} \frac{(\Delta+1)!}{(k+1)! (\Delta-k)!} = \binom{\Delta-k}{m} \tilde{c}_{k+1}.
\end{aligned}$$

We obtain

$$\begin{aligned}
\mathbf{E}[|X|] &\geq pn - p^{k+1}n \sum_{m=0}^{\Delta-k} \binom{\Delta-k}{m} \tilde{c}_{k+1} p^m (1-p)^{\Delta-k-m} \\
&= pn - p^{k+1}n \tilde{c}_{k+1} \\
&= pn(1 - p^k \tilde{c}_{k+1}) \\
&= \frac{kn}{\tilde{c}_{k+1}^{1/k} (1+k)^{1+1/k}}.
\end{aligned}$$

Since the expectation is an average value, there exists a particular k -limited packing of size at least $\frac{kn}{\tilde{c}_{k+1}^{1/k} (1+k)^{1+1/k}}$, as required. The proof of the theorem is complete. \blacksquare

The lower bound of Theorem 1 can be written in a simpler but weaker form as follows:

Corollary 1 *For any graph G of order n ,*

$$L_k(G) > \frac{kn}{e(1+\Delta)^{1+1/k}}.$$

Proof: It is not difficult to see that

$$\tilde{c}_{k+1} \leq \frac{(\Delta+1)^{k+1}}{(k+1)!}$$

and, using Stirling's formula,

$$(k!)^{1/k} > \left(\sqrt{2\pi k} \left(\frac{k}{e} \right)^k \right)^{1/k} = \sqrt[2k]{2\pi k} \frac{k}{e}.$$

By Theorem 1,

$$L_k(G) \geq \frac{kn ((k+1)!)^{1/k}}{(\Delta+1)^{1+1/k} (1+k)^{1+1/k}} > \frac{kn}{e(1+\Delta)^{1+1/k}} \times \frac{\sqrt[2k]{2\pi k} k}{1+k} > \frac{kn}{e(1+\Delta)^{1+1/k}}.$$

Note that $\frac{\sqrt[2k]{2\pi k} k}{1+k} = \frac{\sqrt[2k]{2\pi k}}{1+1/k} > 1$. The last inequality is obviously true for $k = 1$, while for $k \geq 2$ it can be rewritten in the equivalent form: $2\pi k > (1+1/k)^{2k} = e^2 - o(1)$. \blacksquare

3 Randomized algorithm

A pseudocode presented in Algorithm 1 explicitly describes a randomized algorithm to find a k -limited packing set, whose size satisfies bound (1) with a positive probability. Notice that Algorithm 1 constructs a k -limited packing by recursively removing unwanted vertices from the initially constructed set A . This is different from the probabilistic construction used in the proof of Theorem 1. The recursive removal of vertices from the set A may be more effective and efficient, especially if one tries to remove overall as few vertices as possible from A by maximizing intersections of the sets A'_m ($m = k, \dots, \Delta$) and B'_m ($m = k+1, \dots, \Delta$).

Algorithm 1 can be implemented to run in $O(n^2)$ time. To compute the probability $p = \frac{1}{\sqrt[k]{\binom{\Delta}{k}(\Delta+1)}}$, the binomial coefficient $\binom{\Delta}{k}$ can be computed by using the dynamic

programming and Pascal's triangle in $O(k\Delta) = O(\Delta^2)$ time using $O(k) = O(\Delta)$ memory. The maximum vertex degree Δ of G can be computed in $O(m)$ time, where m is the number of edges in G . Then p can be computed in $O(m + \Delta^2) = O(n^2)$ steps. It takes $O(n)$ time to find the initial set A . Computing the intersection numbers $r = |N(v) \cap A|$ and removing unwanted vertices of $N(v) \cap A$'s from A can be done in $O(n + m)$ steps. Finally, checking whether X is maximal or extending X to a maximal k -limited packing can be done in $O(n + m)$ time: try to add vertices of $V(G) - A$ to X recursively one by one, and check whether the addition of a new vertex $v \in V(G) - A$ to X violates the conditions of a k -limited packing for v or at least one of its neighbours in G with respect to $X \cup \{v\}$. Thus, overall Algorithm 1 takes $O(n^2)$ time, and, since $m = O(n^2)$ in general, it is linear in the graph size $(m + n)$ when $m = \theta(n^2)$.

Algorithm 1: Randomized k -limited packing

Input: Graph G and integer k , $1 \leq k \leq \Delta$.

Output: k -Limited packing X in G .

begin

 Compute $p = \left(\frac{1}{\bar{c}_{k+1}(1+k)}\right)^{1/k}$;

 Initialize $A = \emptyset$; /* Form a set $A \subseteq V(G)$ */

foreach *vertex* $v \in V(G)$ **do**

 | with the probability p , decide whether $v \in A$ or $v \notin A$;

end

/* Recursively remove redundant vertices from A */

foreach *vertex* $v \in V(G)$ **do**

 | Compute $r = |N(v) \cap A|$;

if $v \in A$ *and* $r \geq k$ **then**

 | remove any $r - k + 1$ vertices of $N(v) \cap A$ from A ;

end

if $v \notin A$ *and* $r > k$ **then**

 | remove any $r - k$ vertices of $N(v) \cap A$ from A ;

end

end

 Put $X = A$; /* A is a k -limited packing */

 Extend X to a maximal k -limited packing;

return X ;

end

Also, this randomized algorithm for finding k -limited packings in a graph G can be implemented in parallel or as a local distributed algorithm. As explained in [5], this kind of algorithms are especially important, e.g. in the context of ad hoc and wireless sensor networks. We hope that this approach can be also extended to design self-stabilizing or on-line algorithms for k -limited packings. For example, a self-stabilizing algorithm searching for maximal 2-packings in a distributed network system is presented in [11]. Notice that self-stabilizing algorithms are distributed and fault-tolerant, and use the fact that each

node has only a local view/knowledge of the distributed network system. This provides another motivation for efficient distributed search and algorithms to find k -limited packings in graphs and networks.

4 Sharpness of the bound of Theorem 1

We now show that the lower bound of Theorem 1 is asymptotically best possible for some values of k . Let $\delta = \delta(G)$ denote the minimum vertex degree in a graph G . The bound of Theorem 1 can be rewritten in the following form for $\Delta \geq k$:

$$L_k(G) \geq \frac{kn}{(k+1)\sqrt[k]{\binom{\Delta}{k}}(\Delta+1)}.$$

Combining this bound with the upper bound of Lemma 8 from [6], we obtain that for any connected graph G of order n with $\delta(G) \geq k$,

$$\frac{1}{\sqrt[k]{\binom{\Delta}{k}}(\Delta+1)} \times \frac{k}{k+1}n \leq L_k(G) \leq \frac{k}{k+1}n. \quad (5)$$

Notice that the upper bound in the inequality (5) is sharp (see [6]), so these bounds provide an interval of values for $L_k(G)$ in terms of k and Δ when $k \leq \delta$. For regular graphs, $\delta = \Delta$, and, when $k = \Delta$, we have

$$\frac{1}{\sqrt[k]{\binom{\Delta}{k}}(\Delta+1)} = \frac{1}{(k+1)^{1/k}} \longrightarrow 1 \quad \text{as } k \rightarrow \infty.$$

Therefore, Theorem 1 is asymptotically sharp in regular connected graphs for $k = \Delta$. A similar statement can be proved for the situation when $k = \Delta(1 - o(1))$.

Thus, the following result is true:

Theorem 2 *When n is large there exist graphs such that*

$$L_k(G) \leq \frac{kn}{\tilde{c}_{k+1}^{1/k} (1+k)^{1+1/k}} (1 + o(1)). \quad (6)$$

5 Upper bounds

As mentioned earlier, $\rho(G) = L_1(G) \leq \gamma(G)$. In [6], the authors provide several upper bounds for $L_k(G)$, e.g. $L_k(G) \leq k\gamma(G)$ for any graph G . Using the well-known bound (see e.g. [1])

$$\gamma(G) \leq \frac{\ln(\delta+1)+1}{\delta+1}n,$$

we obtain

$$L_k(G) \leq \frac{\ln(\delta+1)+1}{\delta+1}kn. \quad (7)$$

Even though this bound does not work well when k is ‘close’ to δ , it is very reasonable for small values of k .

We now prove an upper bound for the k -limited packing number in terms of the k -tuple domination number. A set X is called a k -tuple dominating set of G if for every vertex $v \in V(G)$, $|N[v] \cap X| \geq k$. The minimum cardinality of a k -tuple dominating set of G is the k -tuple domination number $\gamma_{\times k}(G)$. The k -tuple domination number is only defined for graphs with $\delta \geq k - 1$.

Theorem 3 For any graph G of order n with $\delta \geq k - 1$,

$$L_k(G) \leq \gamma_{\times k}(G). \quad (8)$$

Proof: We prove inequality (8) by contradiction. Let X be a maximum k -limited packing in G of size $L_k(G)$, and let Y be a minimum k -tuple dominating set in G of size $\gamma_{\times k}(G)$. We denote $B = X \cap Y$, i.e. $X = A \cup B$ and $Y = B \cup C$. Assume to the contrary that $L_k(G) > \gamma_{\times k}(G)$, thus $|A| > |C|$.

Since Y is k -tuple dominating set, each vertex of A is adjacent to at least k vertices of Y . Hence the number of edges between A and $B \cup C$ is as follows:

$$e(A, B \cup C) \geq k|A|.$$

Now, every vertex of C is adjacent to at most k vertices of X , because X is a k -limited packing set. Therefore, the number of edges between C and $A \cup B$ satisfies

$$e(C, A \cup B) \leq k|C|.$$

We obtain

$$e(C, A \cup B) \leq k|C| < k|A| \leq e(A, B \cup C),$$

i.e. $e(C, A \cup B) < e(A, B \cup C)$. By eliminating the edges between A and C , we conclude that

$$e(C, B) < e(A, B).$$

Now, let us consider an arbitrary vertex $b \in B$ and denote $s = |N(b) \cap A|$. Since $X = A \cup B$ is a k -limited packing set, we obtain $|N(b) \cap X| \leq k - 1$, and hence $|N(b) \cap B| \leq k - s - 1$. On the other hand, $Y = B \cup C$ is k -tuple dominating set, so $|N(b) \cap Y| \geq k - 1$. Therefore, $|N(b) \cap C| \geq s$. Thus, $|N(b) \cap C| \geq |N(b) \cap A|$ for any vertex $b \in B$. We obtain,

$$e(C, B) \geq e(A, B),$$

a contradiction. We conclude that $L_k(G) \leq \gamma_{\times k}(G)$. ■

Notice that it is possible to have $k = \Delta + 1$ in the statement of Theorem 3, which is not covered by Theorem 1. Then $\delta = \Delta$, which implies the graph is regular. However, $L_k(G) = \gamma_{\times k}(G) = n$ for $k = \delta + 1 = \Delta + 1$. In non-regular graphs, $\delta + 1 \leq \Delta$, and $k \leq \Delta$ to satisfy the conditions of Theorem 1 as well.

For $t \leq \delta$, we define

$$\delta' = \delta - k + 1 \quad \text{and} \quad \tilde{b}_t = \tilde{b}_t(G) = \binom{\delta + 1}{t}.$$

Using the upper bound for the k -tuple domination number from [5], we obtain:

Corollary 2 For any graph G with $\delta \geq k$,

$$L_k(G) \leq \left(1 - \frac{\delta'}{b_{k-1}^{1/\delta'}(1 + \delta')^{1+1/\delta'}}\right)n. \quad (9)$$

In some cases, Theorem 1 and Corollary 2 simultaneously provide good bounds for the k -limited packing number. For example, for a 40-regular graph G :

$$0.312n < L_{25}(G) < 0.843n.$$

References

- [1] N. Alon, J.H. Spencer, *The Probabilistic Method*, 2nd ed., John Wiley & Sons Inc., New York, 2000.
- [2] B. Bollobás, *Graph Theory: An Introductory Course*, Springer-Verlag, New York, 1979.
- [3] A.P. Burger, M.A. Henning, J.H. van Vuuren, On the ratios between packing and domination parameters of a graph, *Discrete Math.* **309** (2009) 2473–2478.
- [4] M.P. Dobson, V. Leoni, G. Nasini, The multiple domination and limited packing problems in graphs, *Inform. Process. Lett.* **111** (2011) 1108–1113.
- [5] A. Gagarin, A. Poghosyan, V.E. Zverovich, Randomized algorithms and upper bounds for multiple domination in graphs and networks, *Discrete Appl. Math.* **161** (2013) 604–611.
- [6] R. Gallant, G. Gunther, B. Hartnell, D. Rall, Limited packings in graphs, *Discrete Appl. Math.* **158**(12) (2010) 1357–1364.
- [7] M.A. Henning, C. Löwenstein, D. Rautenbach, Dominating sets, packings, and the maximum degree, *Discrete Math.* **311** (2011) 2031–2036.
- [8] D.S. Hochbaum, D.B. Schmoys, A best possible heuristic for the k -center problem, *Math. Oper. Res.* **10** (2) (1985) 180–184.
- [9] A. Meir, J.W. Moon, Relations between packing and covering numbers of a tree, *Pacific J. Math.* **61** (1975) 225–233.
- [10] R.R. Rubalcaba, A. Schneider, P.J. Slater, A survey on graphs which have equal domination and closed neighborhood packing numbers, *AKCE J. Graphs. Combin.* **3**(2) (2006) 93–114.
- [11] Z. Shi, A self-stabilizing algorithm to maximal 2-packing with improved complexity, *Inform. Process. Lett.* **112** (2012) 525–531.
- [12] J. Topp, L. Volkmann, On packing and covering numbers of graphs, *Discrete Math.* **96** (1991) 229–238.