THE MONGE-AMPÈRE CONSTRAINED ELASTIC THEORIES OF SHALLOW SHELLS

MARTA LEWICKA, L. MAHADEVAN AND MOHAMMAD REZA PAKZAD

ABSTRACT. Motivated by the degree of smoothness of constrained embeddings of surfaces in \mathbb{R}^3 , and by the recent applications to the elasticity of shallow shells, we rigorously derive the Γ -limit of 3-dimensional nonlinear elastic energy of a shallow shell of thickness h, where the depth of the shell scales like h^{α} and the applied forces scale like $h^{\alpha+2}$, in the limit when $h \to 0$. The main analytical ingredients are two independent results: a theorem on approximation of $W^{2,2}$ solutions of the Monge-Ampère equation by smooth solutions, and a theorem on the matching (in other words, continuation) of second order isometries to exact isometries.

1. Introduction

The mathematical theory of elastic shells must account for the deformation of thin elastic surfaces of non-zero curvature and the associated energetics. The subject thus brings together the differential geometry of surfaces with the theory of elasticity appropriately modified to account for the small aspect ratio of these slender elastic structures. From a practical engineering perspective, many approximate theories have been proposed for the mechanical behavior of elastic shells over the last 150 years [4]. The mathematical foundations for these theories has lagged behind but has recently been the focus of much attention [6, 5, 11] from two perspectives: (i) understanding the rigorous derivation of these theories teaches us about the limits of their practical applicability, and (ii) it also sheds light on the smoothness of allowable deformations with implications for the properties of elliptic operators that arise naturally in differential geometry. Both these questions can be couched in terms of the behavior of the elastic energy of the shell as a function of its aspect ratio, i.e. the ratio of its thickness to its curvature and/or lateral extent. For a flat sheet with no intrinsic curvature, the stretching energy per unit area of a thin sheet is proportional to its thickness, while the bending energy per unit area is proportional to the cube of its thickness. Thus, as the aspect ratio diminishes, the bending energy vanishes faster than the stretching or shearing energy. This leads to approximately isometric deformations of the sheet when it is subject to external loads, with a concomitant theory known as the Föppl-von Kármán theory [17] of elastic plates. For shallow shells with an intrinsic curvature, there is a new small parameter corresponding to the product of the intrinsic curvature and the thickness so that there are various possible distinguished limits associated with how small or large this parameter is independent of the aspect ratio of the shell. This leads to various theories

¹⁹⁹¹ Mathematics Subject Classification. 74K20, 74B20.

Key words and phrases. non-Euclidean plates, nonlinear elasticity, Gamma convergence, calculus of variations.

that penalize non-isometric deformations more or less depending on the relative magnitude of external loads that cause the shell to deform.

Over the recent past, we are beginning to get a good understanding of the limiting models and the types of isometries involved in various similar contexts (e.g. see the review and a conjecture in [24] for thin shells), but each separate case enjoys its own peculiarities related to the the geometry of the shell (hyperbolic, elliptic, degenerate or of mixed type), the order of the relevant approximate isometries on the shell, the linearity or it lack in the PDEs governing them (which become more complicated as the order increases) and their regularity. For shells of thickness h and depth h^{α} , subject to applied forces that scale with the shell depth as $h^{\alpha+2}$, as the thickness of the shell $h \to 0$, depending on the choice of α , various limiting theories arise. The regimes $\alpha = 1$ and $\alpha > 1$ can be treated in a similar manner as discussed in another context in [21] and are not of interest to us in this paper.

Here we consider the case when the shell is shallow and the forces weak, but not too weak, i.e. when $0 < \alpha < 1$, so that nearly isometric deformations might be expected, but as nonlinear constraints, rather than linear ones as appearing in the $\alpha \ge 1$ regimes. In the vanishing thickness limit, using the basic methods of Γ -convergence which were developed in this context in [10, 11], we derive a new thin film model (but see also previous results in [20, 21]). Our analysis corresponds to a situation where the second order infinitesimal isometries on a shallow shell reference configuration are given by the out-of-plane displacement $h^{\alpha}v_0$. In analogy with the results in [11] for plates with flat geometry $v_0 = 0$ in the energy scaling regime $2 < \beta < 4$, we recover a Monge-Ampère constraint det $\nabla^2 v = \det \nabla^2 v_0$ as a second order isometry constraint on the limiting displacements of Sobolev regularity $W^{2,2}$. This emergence of a "linearized curvature" constraint is natural from a mechanical point of view since the Laplacian of the strain which is proportional to the Laplacian of the stress for linear stress-strain relations characterizes local area changes and the isometry condition is softened to one that is effectively locally area-preserving.

Similarly to previously studied cases [11, 23], in order to complete the analysis, we need a result on the matching property, i.e. the continuation of second order isometries, on shallow shells, to exact isometries (under suitable regularity assumptions), and another result on approximating the $W^{2,2}$ solutions of the Monge-Ampère equation by smooth solutions. Contrary to [23, 14], the constraint here is a fully nonlinear one, and one cannot take advantage of the full degeneracy of det $\nabla^2 v = 0$ in [11, Theorems 7 and 10] (see also [32]). Nevertheless, we are able to prove the applicable versions of these results in the case det $\nabla^2 v_0 \equiv c > 0$. In what follows we start with a formulation of the problem (in Section 2) and discuss the various possible choices of α and the known corresponding limiting theories. In Sections 3 and 4, we present our main results, namely, the asymptotic behavior of the elastic energies when $h \to 0$, for the case $0 < \alpha < 1$, and put forward the corresponding matching property and density results as the main technical tools. In Sections 5-6, we present the details of our proofs. Section 7 is dedicated to the study of not necessarily convex $W^{2,2}$ solutions of the Monge-Ampère equation, using some key observations by Sverák from his unpublished manuscript [35]. These results are a necessary ingredient of the density result in Section 6.

2. Elastic shells of low curvature

Let $\Omega \subset \mathbb{R}^2$ be open, bounded and simply connected set. For a given out-of-plate displacement $v_0 \in W^{2,2}(\Omega,\mathbb{R}) \cap \mathcal{C}^2(\bar{\Omega})$, and for a given exponent $\alpha > 0$, consider a sequence of surfaces:

$$S_h = \phi_h(\Omega)$$
 where $\phi_h(x) = (x, h^{\alpha}v_0(x))$ $\forall x = (x_1, x_2) \in \Omega$,

and the family of thin plates $\Omega^h = \Omega \times (-h/2, h/2)$ and thin shallow shells $(S_h)^h$:

$$(2.1) (S_h)^h = \{ \tilde{\phi}_h(x, x_3); \ x \in \Omega, \ x_3 \in (-h/2, h/2) \} \forall 0 < h \ll 1.$$

Above, the Kirchhoff-Love extension $\tilde{\phi}_h:\Omega^h\to\mathbb{R}^3$ of the parametrization ϕ_h , is given by the formula:

(2.2)
$$\tilde{\phi}_h(x, x_3) = \phi_h(x) + x_3 \vec{n}^h(x) \qquad \forall (x, x_3) \in \Omega^h,$$

while the vector $\vec{n}^h(x)$ is the unit normal to S_h at $\phi_h(x)$:

$$\vec{n}^h(x) = \frac{\partial_1 \phi_h(x) \times \partial_1 \phi_h(x)}{|\partial_1 \phi_h(x) \times \partial_1 \phi_h(x)|} = \frac{1}{\sqrt{1 + h^{2\alpha} |\nabla v_0|^2}} \left(-h^\alpha \partial_1 v_0(x), -h^\alpha \partial_2 v_0(x), 1 \right) \qquad \forall x \in \Omega.$$

The thickness averaged elastic energy of a deformation u^h of $(S_h)^h$ is now given by:

(2.3)
$$I^{h}(u^{h}) = \frac{1}{h} \int_{(S_{h})^{h}} W(\nabla u^{h}) \qquad \forall u^{h} \in W^{1,2}((S_{h})^{h}, \mathbb{R}^{3}).$$

The energy density $W: \mathbb{R}^{3\times 3} \longrightarrow \overline{\mathbb{R}}_+$ above, in addition to being C^2 regular in a neighborhood of SO(3), is assumed to satisfy the normalization, frame indifference and nondegeneracy conditions:

(2.4)
$$\exists c > 0 \quad \forall F \in \mathbb{R}^{3 \times 3} \quad \forall R \in SO(3) \qquad W(R) = 0, \quad W(RF) = W(F), \\ W(F) > c \operatorname{dist}^{2}(F, SO(3)).$$

where $F = \nabla u$, is typically the deformation gradient associated with a mapping u. The following quadratic forms, generated by W, will be relevant in the subsequent analysis:

(2.5)
$$Q_3(F) = D^2W(\mathrm{Id})(F, F), \qquad Q_2(F_{tan}) = \min\{Q_3(\tilde{F}); \ \tilde{F} \in \mathbb{R}^{3\times 3}, \ (\tilde{F} - F)_{tan} = 0\}.$$

The form Q_3 is defined for all $F \in \mathbb{R}^{3\times 3}$, while Q_2 is defined on the 2×2 principal minors F_{tan} of such matrices. By (2.4), both forms Q_3 and all Q_2 are nonnegative definite and depend only on the symmetric parts of their arguments.

Let $f^h \in L^2((S_h)^h, \mathbb{R}^3)$ be a family of loads applied to the elastic shells under consideration. The total energy is then:

(2.6)
$$J^{h}(u^{h}) = \frac{1}{h} \int_{(S_{h})^{h}} W(\nabla u^{h}) - \frac{1}{h} \int_{(S_{h})^{h}} f^{h} u^{h} \qquad \forall u^{h} \in W^{1,2}((S_{h})^{h}, \mathbb{R}^{3}).$$

In what follows, we will make the simplifying assumptions:

(2.7)
$$f^{h} = (0, 0, h^{\alpha'} f \circ \tilde{\phi}_{h}^{-1})^{T}$$

for some $f \in L^2(\Omega)$, normalizes so that:

(2.8)
$$\int_{\Omega} f = 0 \quad \text{and} \quad \int_{\Omega} x f(x) \, dx = 0.$$

Heuristically, stronger forces ($\alpha' < \alpha + 2$) deform the shallow shell beyond the reference shape, while weaker forces ($\alpha' > \alpha + 2$) leave it undeformed, with an asymptotic behavior of displacements of lower order similar to that of a plate. The remaining case where the forces are tuned with the curvature of the mid-surface (shallowness), is given by the scaling regime $\alpha' = \alpha + 2$.

When $\alpha \geq 1$, by a simple change of variables, we see that:

$$J^{h}(u^{h}) = \frac{1}{h} \int_{\Omega^{h}} W((\nabla v^{h})(b^{h})^{-1}) \det b^{h} - \frac{1}{h} \int_{\Omega^{h}} h^{\alpha_{i}} f v_{3}^{h} \det b^{h},$$

with $v^h = u^h \circ \tilde{\phi}_h \in W^{1,2}(\Omega^h, \mathbb{R}^3)$ and $b^h = \nabla \tilde{\phi}_h$. Note that by the polar decomposition of positive definite matrices: $b^h = R(x, x_3)a^h$ for some $R(x, x_3) \in SO(3)$ and the symmetric tensor $a^h = \sqrt{(b^h)^T b^h}$. Therefore, for the isotropic energy W i.e. when:

(2.9)
$$W(FR) = W(F) \qquad \forall F \in \mathbb{R}^{3 \times 3} \quad \forall R \in SO(3)$$

one obtains:

(2.10)
$$I^{h}(u^{h}) = \frac{1}{h} \int_{\Omega^{h}} W\left((\nabla v^{h})(a^{h})^{-1} R(x, x_{3})^{-1}\right) \det b^{h} d(x, x_{3})$$
$$= \frac{1}{h} \int_{\Omega^{h}} W\left((\nabla v^{h})(a^{h})^{-1}\right) (1 + \mathcal{O}(h)) d(x, x_{3}),$$

hence reducing the problem to studying deformations of the flat plate Ω^h relative to the prestrain tensor a^h , see [20, 21] for a discussion of this topic.

When $\alpha = 1$, we derived in [21] the Γ -limit \mathcal{J}_{v_0} of the scaled energies $\frac{1}{h^4}I^h$. Namely, we showed that the energy of the almost minimizing deformations scales like:

$$\inf J^h \sim h^4$$
,

and that the Γ -limit (in the general, possibly non-isotropic case) is given by:

(2.11)
$$\mathcal{J}_{v_0}(w,v) = \frac{1}{2} \int_{\Omega} \mathcal{Q}_2 \left(\operatorname{sym} \nabla w + \frac{1}{2} \nabla v \otimes \nabla v - \frac{1}{2} \nabla v_0 \otimes \nabla v_0 \right) + \frac{1}{24} \int_{\Omega} \mathcal{Q}_2 \left(\nabla^2 v - \nabla^2 v_0 \right) - \int_{\Omega} f v,$$

which is the von Kármán-type functional defined for all out-of-plane displacements $v \in W^{2,2}(\Omega,\mathbb{R})$ and the in-plane displacements $w \in W^{1,2}(\Omega,\mathbb{R}^2)$. In analogy with the theory for flat plates with [11], due to the choice of scaling in (2.7), the limit energy is composed of two terms, corresponding to stretching and bending.

In the isotropic case (2.9), the Euler-Lagrange equations of \mathcal{J}_{v_0} are:

(2.12)
$$\begin{cases} \Delta^2 \Phi = -S(\det \nabla^2 v - \det \nabla^2 v_0) \\ B(\Delta^2 v - \Delta^2 v_0) = [v, \Phi] + f \end{cases}$$

where S is the Young modulus, B the bending stiffness, ν the Poisson ratio (given in terms of the Lamé constants μ and λ). A more involved version of the system (2.12) incorporating prestrain was first introduced in [26] using a thermoelastic analogy to growth, as a mathematical model of blooming activated by differential lateral growth from an initial non-zero

transverse displacement field v_0 (See also [25, 8]). See [21] for the rigorous derivation of that model.

When $\alpha > 1$, the bending energy takes over the stretching and hence the limiting theory is a variant of the linear elasticity as discussed in [11], yielding the Euler-Lagrange equations:

$$B(\Delta^2 v - \Delta^2 v_0) = f.$$

We now turn to the case of interest treated in this paper, namely $0 < \alpha < 1$.

3. The main results

Our first main result regards the matching of isometries on convex weakly shallow shells.

Theorem 3.1. Assume that Ω is simply connected and let $v_0 \in C^{2,\beta}(\bar{\Omega}, \mathbb{R})$ with det $\nabla^2 v_0 > c_0 > 0$. Let $v \in C^{2,\beta}(\bar{\Omega}, \mathbb{R})$ satisfy:

(3.1)
$$\det \nabla^2 v = \det \nabla^2 v_0 \quad in \ \Omega.$$

Then there exists a sequence $w_h \in \mathcal{C}^{2,\beta}(\bar{\Omega},\mathbb{R}^3)$ such that:

(3.2) $\forall h > 0 \quad \nabla (\mathrm{id} + hve_3 + h^2w_h)^T \nabla (\mathrm{id} + hve_3 + h^2w_h) = \nabla (\mathrm{id} + hv_0e_3)^T \nabla (\mathrm{id} + hv_0e_3)$ and $\sup \|w_h\|_{\mathcal{C}^{2,\beta}} < +\infty$.

We will explain the meaning of the "matching property" and put our analysis in a broader context in Section 4, while we give its proof in Section 5.

Our next main result concerns the density of regular solutions to the elliptic 2d Monge-Ampère equation.

Theorem 3.2. Let Ω be open, bounded, connected and star-shaped with respect to an interior ball $B \subset \Omega$. For a fixed constant $c_0 > 0$, define:

$$\mathcal{A} := \{ u \in W^{2,2}(\Omega); \det \nabla^2 u = c_0 \text{ a.e. in } \Omega \}.$$

Then $A \cap C^{\infty}(\bar{\Omega})$ is dense in A with respect to the strong $W^{2,2}$ norm.

We prove Theorem 3.2 by showing first the interior regularity of solutions and then by applying a simple scaling argument. The main difficulty to overcome is the absence of convexity assumptions on the $W^{2,2}$ solutions of the Monge-Ampère equation in our context. We first establish the convexity of elements of \mathcal{A} in a broader sense, combining some key observations by Šverák from his unpublished manuscript [35], with a theorem due to Iwaniec and Šverák in [16] (Theorem 7.4) on deformations with integrable dilatation in dimension 2. Some of the results in [35] are known within the community, but for completeness we write the proofs in full detail in Sections 6 and 7.

Our remaining results concern the variational limit of the elastic energy functionals (2.6). We first state the following lemma, whose proof is similar to [11, Theorem 2-i] and [22, Theorem 2.5], and hence it is omitted for brevity of presentation.

Lemma 3.3. Assume (2.7) and (2.8). Then:

(i) For every small h > 0 one has:

$$0 \ge \inf \left\{ \frac{1}{h^{2\alpha+2}} J^h(u^h); \ u^h \in W^{1,2}((S_h)^h, \mathbb{R}^3) \right\} \ge -C.$$

(ii) If $u^h \in W^{1,2}((S_h)^h, \mathbb{R}^3)$ is a minimizing sequence of $\frac{1}{h^{2\alpha+2}}J^h$, that is:

(3.3)
$$\lim_{h \to 0} \left(\frac{1}{h^{2\alpha+2}} J^h(y^h) - \inf \frac{1}{h^{2\alpha+2}} J^h \right) = 0,$$

then $\frac{1}{h^{2\alpha+2}}I^h(u^h)$ is bounded.

We now note, by a straightforward calculation, that a^h in (2.10), pertaining to the isotropic case (2.9), becomes:

$$a^{h} = \operatorname{Id} + \frac{1}{2}h^{2\alpha}(\nabla v_{0} \otimes \nabla v_{0})^{*} - h^{\alpha}x_{3}(\nabla^{2}v_{0})^{*} + o(h^{2\alpha}) + x_{3}o(h^{\alpha})$$

where the uniform quantities in $o(h^{2\alpha})$, $o(h^{\alpha})$ are independent of x_3 . Following the proof of Theorem 1.3 in [20], we actually obtain in the general (possibly non-isotropic) case:

Theorem 3.4. Assume that $u^h \in W^{1,2}((S_h)^h, \mathbb{R}^3)$ satisfies $I^h(u^h) \leq Ch^{2\alpha+2}$, where I^h is given in (2.3) and $0 < \alpha < 1$. Then there exists rotations $\bar{R}^h \in SO(3)$ and translations $c^h \in \mathbb{R}^3$ such that for the normalized deformations:

$$(3.4) y^h(x,t) = (\bar{R}^h)^T (u^h \circ \tilde{\phi}_h)(x,ht) - c^h : \Omega^1 \longrightarrow \mathbb{R}^3$$

defined by means of (2.2) on the common domain $\Omega^1 = \Omega \times (-1/2, 1/2)$ the following holds:

- (i) $y^h(x,t)$ converge in $W^{1,2}(\Omega^1,\mathbb{R}^3)$ to x
- (ii) the scaled displacements $V^h(x) = h^{-\alpha} \int_{-1/2}^{1/2} y^h(x,t) x$ dt converge (up to a subsequence) in $W^{1,2}(\Omega,\mathbb{R}^3)$ to $(0,0,v)^T$ where $v \in W^{2,2}(\Omega,\mathbb{R})$ and:

(3.5)
$$\det \nabla^2 v = \det \nabla^2 v_0.$$

(iii) Moreover: $\liminf_{h\to 0} h^{-(2\alpha+2)} J^h(u^h) \geq \mathcal{J}_{v_0}(v)$ where:

(3.6)
$$\mathcal{J}_{v_0}(v) = \frac{1}{24} \int_{\Omega} \mathcal{Q}_2(\nabla^2 v - \nabla^2 v_0) - \int_{\Omega} f v.$$

The constraint (3.5) in the assertion (ii) follows by observing that $h^{-\alpha} \operatorname{sym} \nabla V^h$ converges in $L^2(\Omega)$ to $F = \frac{1}{2}(\nabla v_0 \otimes \nabla v_0 - \nabla v \otimes \nabla v)$, and hence $F = \operatorname{sym} \nabla w$ for some $w : \Omega \to \mathbb{R}^2$. Consequently $\operatorname{curl}^T \operatorname{curl} F = 0$, which implies (3.5).

We conjecture that the linearized Kirchhoff-like energy (3.6) (3.5) is also the exact Γ limit of the rescaled energies $h^{-(2\alpha+2)}J^h(u^h)$ when Ω is simply connected. Note that this
result is established in [11] for the degenerate case $v_0 \equiv 0$ and hence our model can be
considered as a generalization of the linearized Kirchhoff model discussed in [11]. If Ω is
not simply connected, one must replace (3.5) with a more general variant which states that $\nabla v_0 \otimes \nabla v_0 - \nabla v \otimes \nabla v$ is a symmetric gradient.

We are able to prove our conjecture in the specific case when $\det \nabla^2 v_0$ is constant. Its proof relies on Theorems 3.1 and 3.2 and it will be given in Section 8. Namely, we have:

Theorem 3.5. Assume that Ω is open, bounded, connected and star-shaped with respect to an interior ball $B \subset \Omega$. Assume that:

$$\det \nabla^2 v_0 \equiv c_0 > 0 \quad in \ \Omega.$$

Fix $\alpha \in (0,1)$. Then, for every $v \in W^{2,2}(\Omega,\mathbb{R})$ with $\nabla^2 v = \nabla^2 v_0$, there exists a sequence of deformations $u^h \in W^{1,2}((S_h)^h,\mathbb{R}^3)$ such that:

- (i) The rescaled sequence $y^h(x,t) = u^h(x+h^{\alpha}v_0(x)e_3+ht\vec{n}^h(x))$ converges in $W^{1,2}(\Omega^1,\mathbb{R}^3)$ to x.
- (ii) The scaled displacements V^h as in (ii) Theorem 3.4 converge in $W^{1,2}$ to (0,0,v).
- (iii) $\lim_{h\to 0} h^{-(2\alpha+2)} J^h(u^h) = \mathcal{J}_{v_0}(v)$.

Remark 3.6. We expect that the $C^{2,\beta}$ scalar fields v are dense in the set of the $W^{2,2}$ fields with any prescribed, strictly positive but not necessarily constant det ∇^2 of $C^{0,\beta}$ regularity. With such a result at hand, it would follow that for all convex shells of sufficient regularity, the linearized Kirchhoff-type energy (3.6) is the rigorous variational limit on weakly shallow shells, in the same spirit as the matching and density of first order isometries on convex shells [23] resulted in that the only small slope theory for an elastic convex shell is the linear theory. The latter problems, when posed for surfaces of arbitrary geometry, are more difficult. One could hope to prove similar results for strictly hyperbolic surfaces S. In the general case, however, such problems reduce to the study of nonlinear PDEs of mixed types for which not so many suitable methods are at hand.

4. Remarks on the matching property in Theorem 3.1.

Remark 4.1. In [24], the authors put forward a conjecture regarding existence of infinitely many small slope shell theories (with no prestrain) each valid for a corresponding range of energy scalings. This conjecture is based on formal asymptotic expansions and it is in accordance with the previously obtained results for plates and shells. It predicts the form of the 2-dimensional limit energy functional, and identifies the space of admissible deformations as infinitesimal isometries of a given integer order N > 0 determined by the magnitude of the elastic energy. Hence, the influence of shells geometry on its qualitative response to an external force, i.e. the shells rigidity, is reflected in a hierarchy of functional spaces of isometries (and infinitesimal isometries) arising as constraints of the derived theories.

In certain cases, a given Nth order infinitesimal isometry can be modified by higher order corrections to yield an infinitesimal isometry of order M > N, a property to which we refer to by "matching property of infinitesimal isometries". This feature, combined with certain density results for spaces of isometries, cause the theories corresponding to orders of infinitesimal isometries between N and M, to collapse all into one and the same theory. The examples of such behavior are observed for plates [11], where any second order infinitesimal isometry can be matched to an exact isometry $(M = \infty)$, for convex shells [23], where any first order infinitesimal isometry satisfies the same property, and for non-flat developable surfaces [34, 14] where first order isometries can be matched to higher order isometries (see also [13]). The effects of these geometric properties on the elasticity of thin films are drastic. A plate whose boundary is at least partially free possesses three types of small-slope theories: the linear theory, the von Kármán theory and the linearized Kirchhoff theory, whereas the only small slope theory for a convex shell with free boundary is the

linear theory [23]: a convex shell transitions directly from the linear regime to the fully nonlinear bending one if the applied forces are adequately increased. In other words, while the von Kármán theory describes the buckling of thin plates at a body force magnitude of order thickness-cubed, the equivalent, variationally correct theory for buckling of elliptic shells is the purely nonlinear bending theory which comes only into effect when the body forces reach to a magnitude of order thickness-squared.

Remark 4.2. Writing $w_h = w_{h,tan} + w_h^3 e_3$ where $w_{h,tan}(x) \in \mathbb{R}^2$ and $w_h^3 \in \mathbb{R}$, equation (3.2) becomes:

(4.1)
$$\operatorname{Id} + h^{2}(2\operatorname{sym}\nabla w_{h,tan} + \nabla v \otimes \nabla v) + 2h^{3}\operatorname{sym}(\nabla v \otimes \nabla w_{h}^{3}) + h^{4}((\nabla w_{h,tan})^{T}\nabla w_{h,tan} + \nabla w_{h}^{3} \otimes \nabla w_{h}^{3}) = \operatorname{Id} + h^{2}\nabla v_{0} \otimes \nabla v_{0}.$$

Recall that (3.1) is equivalent to $\operatorname{curl}^T \operatorname{curl}(\nabla v \otimes \nabla v - \nabla v_0 \otimes \nabla v_0) = 0$, and hence to: $\nabla v \otimes \nabla v - \nabla v_0 \otimes \nabla v_0 = \operatorname{sym} \nabla w$ for some $w \in \mathcal{C}^{2,\beta}(\bar{\Omega},\mathbb{R}^2)$ since Ω is simply connected. Hence the constraint (3.1) is necessary and sufficient for matching the lowest order (h^2) terms in (4.1).

Our result states that actually it is possible to perturb w by an equibounded 3d displacement $w_h - w$ so that the full equality (4.1) holds. A natural way for proving this is by implicit function theorem. Indeed, this is how we proceed, and the ellipticity assumption $\det \nabla^2 v_0 > 0$ is precisely a sufficient condition for the invertibility of the implicit derivative $\mathcal{L}(p) : \mathcal{C}_0^{2,\beta}(\bar{\Omega},\mathbb{R}) \longrightarrow \mathcal{C}^{0,\beta}(\bar{\Omega},\mathbb{R}), \ \mathcal{L}(p) = -\cot \nabla^2 v : \nabla^2 p$ where p is the variation in w_h^3 . An extra argument for the uniform boundedness of $w_{h,tan}$ in $\mathcal{C}^{2,\beta}$ concludes the proof.

Remark 4.3. The condition (3.2) means that each deformation $u^h: S_h \to \mathbb{R}^3$ of a surface $S_h = \{x + hv_0(x)e_3; x \in \Omega\}$, given by $u^h(x + hv_0(x)e_3) = x + hv(x)e_3 + h^2w_h(x)$ is an isometry of S_h . In other words, the pull-back metrics from the Euclidean metric of S_h and of $u^h(S_h) = \{x + hv(x)e_3 + h^2w_h(x); x \in \Omega\}$ coincide. Hence Theorem 3.1 asserts that if two convex out-of-plane displacements of first order have the same determinants of Hessians, then they can be matched by a family of equibounded higher order displacements (the fields w_h) to be isometrically equivalent. When the parameter h is replaced by h^{α} , this result will be used in our context of shallow shells. For other results concerning matching of isometries see [11, Theorem 7], [23, Theorem 1.1], [14, Theorem 3.1] (which is comparable with [34, Lemma 3.3] and the remark which follows therein) and [13, Theorem 4.1].

5. The matching property on convex shallow shells: a proof of Theorem 3.1.

By a direct calculation, (3.2) is equivalent to:

- (5.1) $\nabla(\mathrm{id} + h^2 w_{h,tan})^T \nabla(\mathrm{id} + h^2 w_{h,tan}) = \mathrm{Id} + h^2 \nabla v_0 \otimes \nabla v_0 h^2 (\nabla v + \nabla z_h) \otimes (\nabla v + \nabla z_h),$ where $w_{h,tan} \in \mathcal{C}^{2,\beta}(\bar{\Omega},\mathbb{R}^2)$ and $z_h = h w_h^3 \in \mathcal{C}^{2,\beta}(\bar{\Omega},\mathbb{R})$ so that $w_h = w_{h,tan} + w_h^3 e_3$ is the required correction in (3.2).
- 1. We shall first find the formula for the Gaussian curvature of the 2d metric in the right hand side of (5.1):

$$(5.2) g_h(z_h) = \operatorname{Id} + h^2 \nabla v_0 \otimes \nabla v_0 - h^2 (\nabla v + \nabla z_h) \otimes (\nabla v + \nabla z_h).$$

Lemma 5.1. Let $v_0, v \in C^{2,\beta}(\bar{\Omega}, \mathbb{R})$ and consider the $C^{1,\beta}$ regular metrics on Ω of the type:

$$g = [g_{ij}]_{i,j=1,2} = \mathrm{Id} + h^2(\nabla v_0 \otimes \nabla v_0 - \nabla v_1 \otimes \nabla v_1).$$

Then, for any h > 0 small, the Gaussian curvature $\kappa(g)$ of g is $\mathcal{C}^{0,\beta}$ regular and it is given by the formula:

(5.3)
$$\kappa(g) = h^2 \left[\frac{\det(\nabla^2 v_0 - [\Gamma_{ij}^k \partial_k v_0]_{ij})}{\left(1 - h^2(g^{ij} \partial_i v_0 \partial_j v_0)\right)^2 \det g} - \frac{\left(1 - h^2(g^{ij} \partial_i v_0 \partial_j v_0)\right)^2}{\left(1 - h^2|\nabla v_1|^2\right)^2} \det \nabla^2 v_1 \right],$$

where the Christoffel symbols of g, the inverse of g, and its determinant are:

(5.4)
$$\Gamma_{ij}^{k} = \frac{1}{2} g^{kl} \left(\partial_{j} g_{il} + \partial_{i} g_{jl} - \partial_{j} g_{ij} \right),$$

(5.5)
$$g^{-1} = [g^{ij}] = \frac{1}{\det[g_{ij}]} \operatorname{cof}[g_{ij}],$$

$$\det g = 1 - h^4 |(\nabla v_0)^{\perp} \cdot \nabla v_1|^2 + h^2 (|\nabla v_0|^2 - |\nabla v_1|^2).$$

Proof. Assume first that v_0 and v_1 are in fact smooth. By Lemma 2.1.2 in [12], we have:

$$\kappa \left(\operatorname{Id} - h^2 \nabla v_1 \otimes \nabla v_1 \right) = -h^2 \frac{\det \nabla^2 v_1}{\left(1 - h^2 |\nabla v_1|^2 \right)^2}$$

$$\kappa \left(g - h^2 \nabla v_0 \otimes \nabla v_0 \right) = \frac{1}{\left(1 - h^2 (g^{ij} \partial_i v_0 \partial_j v_0) \right)^2} \left[\kappa(g) - \frac{h^2 \det (\nabla^2 v_0 - [\Gamma_{ij}^k \partial_k v_0]_{ij})}{\left(1 - h^2 (g^{ij} \partial_i v_0 \partial_j v_0) \right)^2 \det g} \right].$$

Since the two metrics above are equal, the formula (5.3) follows directly. The formula for $\det g$ is obtained by a direct calculation, via $\det(A+B) = \det A + \cot A : B + \det B$, valid for 2×2 matrices A, B.

In the general case when v_0, v_1 are only $C^{2,\beta}$ regular, one may approximate them by smooth sequences v_0^n, v_1^n . Then, each $\kappa_n = \kappa \left(\operatorname{Id} + h^2 (\nabla v_0^n \otimes v_0^n - \nabla v_1^n \otimes v_1^n) \right)$ is given by the formula in (5.3), and the sequence κ_n converges in $C^{0,\beta}$ to the right hand side in (5.3). On the other hand, κ_n converges in $\mathcal{D}'(\Omega)$ to $\kappa(g)$, which follows from the definition of Gauss curvature $\kappa = R_{1212}/\det g$. Hence the lemma is proven.

2. Applying Lemma 5.1 to $v_1 = v + z_h$, we now see that for small h, the Gauss curvature of metric $g_h(z_h)$ vanishes:

if and only if:

$$\Phi(h, z_h) = 0,$$

where:

$$\Phi(h,z) = (1 - h^2 |\nabla v + \nabla z|^2)^2 \det(\nabla^2 v_0 - [\Gamma_{ij}^k \partial_k v_0]_{ij}) - (1 - h^2 (g^{ij} \partial_i v_0 \partial_j v_0))^4 d(h,z) \det(\nabla^2 v + \nabla^2 z).$$

Here:

$$d(h,z) = 1 - h^4 |(\nabla v_0)^{\perp} \cdot \nabla (v+z)|^2 + h^2 (|\nabla v_0|^2 - |\nabla v + \nabla z|^2)$$

and Γ_{ij}^k and g^{ij} are given by (5.4) and (5.5) for the metric $g = \operatorname{Id} + h^2 \nabla v_0 \otimes \nabla v_0 - h^2 (\nabla v + \nabla z) \otimes (\nabla v + \nabla z)$. We shall consider:

$$\Phi: (-\epsilon, \epsilon) \times \mathcal{C}_0^{2,\beta}(\bar{\Omega}, \mathbb{R}) \longrightarrow \mathcal{C}^{0,\beta}(\bar{\Omega}, \mathbb{R})$$

and seek for solutions $z_h \in \mathcal{C}_0^{2,\beta}(\bar{\Omega},\mathbb{R})$ of (5.7) with zero boundary data. It is elementary to check that Φ is continuously Frechet differentiable at (0,0) and that

$$\Phi(0,0) = \det \nabla^2 v_0 - \det \nabla^2 v = 0.$$

Moreover, the partial Frechet derivative $\mathcal{L} = \partial \Phi / \partial z(0,0) : \mathcal{C}_0^{2,\beta}(\bar{\Omega},\mathbb{R}) \longrightarrow \mathcal{C}^{0,\beta}(\bar{\Omega},\mathbb{R})$ is a linear continuous operator of the form:

$$\forall z \in \mathcal{C}_0^{2,\beta} \qquad \mathcal{L}(z) = \lim_{\epsilon \to 0} \frac{1}{\epsilon} \Phi(0, \epsilon z) = \lim_{\epsilon \to 0} \frac{1}{\epsilon} \left(\det \nabla^2 v_0 - \det(\nabla^2 v + \epsilon \nabla^2 z) \right)$$
$$= \lim_{\epsilon \to 0} \frac{1}{\epsilon} \left(-\epsilon^2 \det \nabla^2 z - \epsilon \operatorname{cof} \nabla^2 v : \nabla^2 z \right) = -\operatorname{cof} \nabla^2 v : \nabla^2 z.$$

Clearly, \mathcal{L} is invertible to a continuous linear operator, because of the uniform ellipticity of the matrix field $\nabla^2 v$ which follows from the convexity assumption of det $\nabla^2 v = \det \nabla^2 v_0$ being strictly positive. Thus, invoking the implicit function theorem we obtain the solution operator:

$$\mathcal{Z}: (-\epsilon, \epsilon) \longrightarrow \mathcal{C}_0^{2,\beta}(\bar{\Omega}, \mathbb{R})$$

such that $z_h = \mathcal{Z}(h)$ satisfies (5.7). Moreover, \mathcal{Z} is differentiable at h = 0 and:

$$\mathcal{Z}'(0) = \mathcal{L}^{-1} \circ \left(\frac{\partial \Phi}{\partial h}(0,0)\right) = 0,$$

because:

$$\frac{\partial \Phi}{\partial h}(0,0) = \left(\operatorname{cof} \nabla^2 v_0 \right) : \left[\left(\frac{\partial}{\partial h} \Gamma_{ij}^k \right) \partial_k v_0 \right]_{ij} + \frac{\partial}{\partial h} \det \left[\Gamma_{ij}^k \partial_k v_0 \right]_{ij} - \left(\frac{\partial}{\partial h} d(0,0) \right) \det \nabla^2 v = 0.$$

Consequently:

(5.8)
$$||w_h^3||_{\mathcal{C}^{2,\beta}} = \frac{1}{h} ||z_h||_{\mathcal{C}^{2,\beta}} \to 0 \quad \text{as } h \to 0.$$

3. In conclusion, we have so far obtained a uniformly bounded sequence of $C_0^{2,\beta}$ out-of-plane displacements $w_h^3 = z_h/h$ such that the Gauss curvature (5.6) of the metric $g_h(z_h)$ in the right hand side of (5.1) is 0. By the result in [30] it follows that for each small h there exists exactly one (up to fixed rotations) orientation preserving isometric immersion $\phi_h \in C^2(\bar{\Omega}, \mathbb{R}^2)$ of $g_h(z_h)$:

(5.9)
$$\nabla \phi_h^T \nabla \phi_h = g_h(z_h) \quad \text{and} \quad \det \nabla \phi_h > 0.$$

What remains to be proven is that, in fact, $\phi_h = \mathrm{id} + h^2 w_{h,tan}$ with some $w_{h,tan}$ uniformly bounded in $\mathcal{C}^{2,\beta}(\bar{\Omega},\mathbb{R}^2)$.

It is a well known calculation (see [6, 30]) that (5.9) implies (is actually equivalent to):

(5.10)
$$\nabla^2 \phi_h - [\Gamma_{ij}^k \partial_k \phi_h]_{ij} = 0,$$

where Γ_{ij}^k are the Christoffel symbols (5.4) of the metric $g = g_h(z_h)$ in (5.2). By (5.9) $\|\nabla \phi_h\|_{L^{\infty}} \leq C$, and by (5.10) $\|\nabla^2 \phi_h\|_{L^{\infty}} \leq C$, hence $\|\phi_h\|_{\mathcal{C}^{2,\beta}} \leq C$. But Γ_{ij}^k are uniformly bounded (with respect to small h) in $\mathcal{C}^{0,\beta}$ so by (5.10) $\|\nabla^2 \phi_h\|_{\mathcal{C}^{0,\beta}} \leq C$ and thus:

$$\|\phi_h\|_{\mathcal{C}^{2,\beta}(\bar{\Omega},\mathbb{R}^2)} \leq C.$$

Note now that $\|\Gamma_{ij}^k\|_{\mathcal{C}^{0,\beta}} \leq Ch^2$ in view of the particular structure of the metrics $g_h(z_h)$. Hence, by (5.10):

$$\|\nabla^2 \phi_h\|_{\mathcal{C}^{0,\beta}} \le Ch^2.$$

Therefore, for some $A_h \in \mathbb{R}^{2 \times 2}$ we have:

We now prove that the matrix A_h in the inequality above can be chosen as a rotation and hence, without loss of generality, $A_h = \text{Id}$. For each $x \in \Omega$ there holds:

(5.13)
$$\operatorname{dist}(A_h, SO(3)) \le |A_h - \nabla \phi_h(x)| + \operatorname{dist}(\nabla \phi_h(x), SO(3)).$$

To evaluate the last term above, write: $\sqrt{\nabla \phi_h^T(x) \nabla \phi_h(x)} = QDQ^T$ for some $Q \in SO(3)$ and $D = \operatorname{diag}(\lambda_1, \lambda_2)$ with $\lambda_1, \lambda_2 > 0$. Since $\det \nabla \phi_h > 0$, it follows by polar decomposition theorem that:

$$\operatorname{dist}(\nabla \phi_h(x), SO(3)) = |\sqrt{\nabla \phi_h^T(x) \nabla \phi_h(x)} - \operatorname{Id}| \leq C|D - \operatorname{Id}|$$

$$= C \max_i \{|\lambda_i - 1|\} \leq C \max_i \{|\lambda_i^2 - 1|\} \leq C|D^2 - \operatorname{Id}|$$

$$= C|Q^T \nabla \phi_h^T(x) \nabla \phi_h(x) Q - \operatorname{Id}| \leq C|\nabla \phi_h^T \nabla \phi_h(x) - \operatorname{Id}| \leq Ch^2.$$

By the above and (5.13), (5.12) we see that $\operatorname{dist}(A_h, SO(3)) \leq Ch^2$. Hence, without loss of generality, $\|\nabla \phi_h - \operatorname{Id}\|_{\mathcal{C}^{1,\beta}} \leq Ch^2$ and:

$$\|\phi_h - \mathrm{id}\|_{\mathcal{C}^{2,\beta}} \le Ch^2.$$

Consequently, $\phi_h = \mathrm{id} + h^2 w_{h,tan}$ with $||w_{h,tan}||_{\mathcal{C}^{2,\beta}} \leq C$. This concludes the proof of Theorem 3.1, in view of (5.1) which is equivalent to (3.2).

Remark 5.2. The above proof is somewhat similar to [12, Theorem 4.1.1]. In analogy, note the similarity between the proof of the matching property in [23] and the Weyl problem by Nirenberg in [33].

6. Density and regularity for elliptic 2d Monge-Ampère equation: A proof of Theorem 3.2.

In this and the following sections, $\Omega \subset \mathbb{R}^2$ is a domain, i.e. an open, bounded and connected set. The main step towards proving Theorem 3.2 will be the following result in which we combine some key observations by Šverák from his unpublished manuscript [35], with a theorem due to Iwaniec and Šverák in [16] (Theorem 7.4) regarding deformations with integrable dilatation in dimension 2.

Theorem 6.1. Let $u \in W^{2,2}(\Omega)$ be such that:

(6.1) $\det \nabla^2 u = f \text{ in } \Omega, \text{ where } f: \Omega \to \mathbb{R}, f(x) \ge c_0 > 0 \quad \forall a.e. \ x \in \Omega.$

Then $u \in C^1(\Omega)$ and, modulo a global sign change, u is locally convex in Ω .

Example 6.2. Let B_1 be the unit disk in \mathbb{R}^2 and let $u \in \mathcal{C}^1(B_1)$ be given by:

$$u(x,y) = \begin{cases} x^2 e^{y^2/2} & \text{if } x \ge 0\\ -x^2 e^{y^2/2} & \text{otherwise.} \end{cases}$$

Note that u(0,y) = 0 and $\nabla u(0,y) = 0$ for all $y \in (-1,1)$. Indeed, we have $u_x = \pm 2xe^{y^2/2}$, $u_y = \pm yx^2e^{y^2/2}$, $u_{xx} = \pm 2e^{y^2/2}$, $u_{xy} = u_{yx} = \pm 2xye^{y^2/2}$, $u_{yy} = \pm (x^2e^{y^2/2} + y^2x^2e^{y^2/2})$ and $\Delta u = \pm e^{y^2/2}(2 + x^2 + y^2x^2)$, respectively for x > 0 and x < 0.

As a consequence $u \in W^{2,\infty}(B_1)$, u is strictly convex in $\{(x,y) \in B_1; x > 0\}$ and strictly concave in $\{(x,y) \in B_1; x < 0\}$. On the other hand $\det \nabla^2 u = 2x^2 e^{y^2} (1-y^2) \in \mathcal{C}^{\infty}(B_1)$ and it is positive if $x \neq 0$ and $y^2 < 1$. We right away note that $u \notin \mathcal{C}^2(B_1)$ although it solves the Monge-Ampère equation with smooth non-negative right hand side, a.e. in its domain.

Finally, our example shows that the assumption of strict positivity in Theorem 6.1 (and also 7.4 and Theorem 6.3 below) cannot be relaxed to det $\nabla^2 u > 0$ a.e., even assuming a better $W^{2,\infty}$ regularity for u.

As a consequence of Theorem 6.1 and of the monotonicity property by Vodopyanov and Goldstein which we quote in Theorem 7.6, we obtain:

Theorem 6.3. Let $f \in C^{k,\beta}(\Omega)$ be a positive function. Then any $W^{2,2}(\Omega)$ solution of $\det \nabla^2 u = f$, is $C^{k+2,\beta}$ regular, locally in Ω .

Proof. 1. We first note that u is a generalized Aleksandrov solution to (6.1). Since u is locally convex, it is twice differentiable in the classical sense a.e. in Ω , and its gradient agrees with f. By Lemma 2.3 in [37], the regular part of the Monge-Ampère measure μ_u equals $(\det \nabla^2 u)dx = fdx$. It suffices now to prove that there is no singular part of μ_u , i.e. that μ_u is absolutely continuous with respect to the Lebesgue measure dx.

Call $v = \nabla u$. By Theorem 6.1 we have $u \in \mathcal{C}^1(\Omega)$ and hence:

$$\mu_u(\omega) = |v(\omega)|$$
 for every Borel set $\omega \subset \Omega$.

We thus need to show that v satisfies Luzin's condition (N):

$$|v(\omega)| = 0$$
 $\forall \omega \subset \Omega; |\omega| = 0.$

The above claim follows directly from Theorem A in [27], in view of $v \in W^{1,2}(\Omega, \mathbb{R}^2)$ and the monotonicity property (7.3) of v due to Vodopyanov and Goldstein.

2. Since $f \in C^{0,\alpha}(\bar{\Omega})$, then Theorem 5.4 in [15] implies that u is locally $C^{2,\alpha}$. We note that this statement is the well-known result due to Caffarelli [2]. Indeed, fix $x_0 \in \Omega$. By Remark 3.2 in [37] which gives an elementary proof of a result by Aleksandrov and Heinz, the displacement u as above must be strictly convex in some $B(x_0, \epsilon)$. By adding an affine function to u, we may without loss of generality assume that u = 0 on the boundary of the convex set:

$$\Omega_0 = \{x \in \Omega; \ u(x) \le u(x_0) + \delta\} \subset B(x_0, \epsilon),$$

for a sufficiently small $\delta > 0$. Therefore, the statement of Theorem 5.4 in [15] can be directly applied. Once the $C^{2,\beta}$ regularity is established, the $C^{k+2,\beta}$ regularity follows as in Proposition 9.1 in [3].

Proof of Theorem 3.2. Without loss of generality we assume that Ω is starshaped with respect to $B = B(0, r) \subset \Omega$. Let $u \in \mathcal{A}$ and define $u_{\lambda}(x) = \frac{1}{\lambda^2} u(\lambda x)$ for $0 < \lambda < 1$. Then:

$$\det \nabla^2 u_{\lambda}(x) = c_0 \quad \forall a.e. \ x \in \Omega$$

and:

$$||u_{\lambda}||_{L^{2}(\Omega)} = \lambda^{-3} ||u||_{L^{2}(\lambda\Omega)}, \quad ||\nabla u_{\lambda}||_{L^{2}(\Omega)} = \lambda^{-2} ||u||_{L^{2}(\lambda\Omega)}, \quad ||\nabla^{2} u_{\lambda}||_{L^{2}(\Omega)} = \lambda^{-1} ||u||_{L^{2}(\lambda\Omega)}.$$

As a consequence, $u_{\lambda} \in \mathcal{A}$ for all $\lambda \in (0,1)$, and $u_{\lambda} \to u$ strongly in $W^{2,2}(\Omega)$ as $\lambda \to 1^-$.

So far we have used only the fact that Ω is starshaped with respect to the origin 0. Now, since Ω is star-shaped with respect to an open ball B, we have $\lambda \bar{\Omega} \subset \Omega$ for all $0 < \lambda < 1$. Hence, in view of Theorem 6.3, $u_{\lambda} \in \mathcal{C}^{\infty}(\bar{\Omega}) \cap \mathcal{A}$, which proves the claim.

7. Šverák's arguments: a proof of Theorem 6.1.

Definition 7.1. We say that a mapping $v \in C^0(\Omega, \mathbb{R}^2)$ is connectedly locally one-to-one iff it is locally one-to-one outside of a closed set $S \subset \Omega$ of measure zero, for which $\Omega \setminus S$ is connected.

Definition 7.2. Let $v \in W^{1,2}(\Omega, \mathbb{R}^2)$ and let $\det \nabla v \geq 0$ a.e. in Ω . We say that v has integrable dilatation iff:

$$\forall a.e. \ x \in \Omega \qquad |\nabla v|^2(x) \le K(x) \det \nabla v(x)$$

with some function $K \in L^1(\Omega)$.

For the proof of Theorem 6.1, the first result we propose is essentially a combination of arguments in Šverák's unpublished paper [35]. In this section, we will gather all the details of its proof.

Theorem 7.3 (Šverák). If $u \in W^{2,2}(\Omega)$ satisfies:

(7.1)
$$\det \nabla^2 u(x) > 0 \quad \forall a.e. \ x \in \Omega,$$

then $u \in C^1(\Omega)$. If additionally $v = \nabla u$ is connectedly locally one-to-one, then modulo a global sign change, u is locally convex in Ω . In particular, when Ω is convex then u is either convex or concave in the whole Ω .

We quote now the result, which will be crucial for the proof of Theorem 7.3:

Theorem 7.4 (Iwaniec and Šverák [16]). Let $v \in W^{1,2}(\Omega, \mathbb{R}^2)$ be as in Definition 7.2. Then there exists a homeomorphism $h \in W^{1,2}(\Omega', \Omega)$ and a holomorphic function $\varphi \in W^{1,2}(\Omega', \mathbb{R}^2 = \mathbb{C})$ such that:

$$v = \varphi \circ h^{-1}.$$

In particular, v is either constant or connectedly locally one-to-one, and in the latter case the singular set $S = h((\nabla \phi)^{-1}\{0\})$.

Without having Theorem 7.4 at hand, Šverák proved in [35] that if $u \in W^{2,2}(\Omega)$ satisfies $\det \nabla^2 u > 0$ a.e. in Ω , then there exists a closed set $S \subset \Omega$ of measure zero such that on each component of $\Omega \setminus S$, u is either locally convex or locally concave. In fact, the main step in the proof is to show that any such map is locally one-to-one outside a set of measure zero, which Šverák has achieved by using consequences of a version of Lemma 7.9 below and the classical degree theory.

Combining Theorem 7.3 with Theorem 7.4 one directly obtains:

Corollary 7.5. Let $u \in W^{2,2}(\Omega)$ satisfy (7.1) and be such that ∇u has integrable dilatation. Then $u \in C^1(\Omega)$ and modulo a global sign change, u is locally convex in Ω .

Theorem 6.1 is then, obviously, a particular case of the above corollary, where the displacement u trivially satisfies its assumptions. In the remaining part of this section, we will prove Theorem 7.3. We first remind a key result on the modulus of continuity of 2d deformations in $W^{1,2}$ with positive Jacobian:

Theorem 7.6 (Vodopyanov and Goldstein [39]). Assume that $v \in W^{1,2}(\Omega, \mathbb{R}^2)$ and that $\det \nabla v > 0$ a.e. in Ω . Then v is continuous in Ω , and for any $B(x, \delta) \subset B(x, R) \subset \Omega$ we have:

(7.2)
$$\operatorname{osc}_{B(x,\delta)} v \le \sqrt{2\pi} (\ln \frac{R}{\delta})^{-1/2} \|\nabla v\|_{L^2(B(x,R))}.$$

Proof. By a result of Vodopyanov and Goldstein [39] v is continuous (see also [28, 9, 36]). In fact, a key ingredient of this result is to show that ϕ is a monotone map, i.e. for $B_{\rho} = B(x, \rho)$:

$$(7.3) \operatorname{osc}_{B_{\rho}} v = \operatorname{osc}_{\partial B_{\rho}} v,$$

and hence v has the asserted modulus of continuity by [31, Theorem 4.3.4] (see also [28, 9]). We sketch the last part of the proof for the convenience of the reader. By Fubini's theorem v belongs to $W^{1,2}(\partial B_{\rho})$ for almost every $\rho \in (\delta, R)$. Hence the Morrey's theorem of embedding of $W^{1,2}$ into C^0 for the one-dimensional set ∂B_{ρ} yields:

$$\forall a.e. \ \rho \in (\delta, R) \qquad \operatorname{osc}_{B_{\delta}} v \leq \operatorname{osc}_{B_{\rho}} v = \operatorname{osc}_{\partial B_{\rho}} v \leq \sqrt{2\pi\rho} \Big(\int_{\partial B_{\rho}} |\nabla v|^2 \Big)^{1/2}.$$

To conclude, one squares both sides of the above inequality, divides by ρ and integrates from δ to R, in order to deduce (7.2).

Corollary 7.7. Assume that $v_n \in W^{1,2}(\Omega, \mathbb{R}^2)$ is a bounded sequence such that $\det \nabla v_n > 0$ a.e. in Ω . Then, up to a subsequence, v_n converges locally uniformly and also weakly in $W^{1,2}$ to a continuous mapping $v \in W^{1,2}(\Omega, \mathbb{R}^2)$ satisfying $\det \nabla v \geq 0$ a.e. in Ω .

Proof. The uniform convergence of a subsequence follows by Ascoli-Arzelá theorem in view of Theorem 7.6. Noting that $\det \nabla v = -\nabla v_1 \cdot \nabla^{\perp} v_2$, the Div-Curl Lemma implies then that the desired inequality is satisfied for the limit mapping v.

Corollary 7.8. Let $u_n \in W^{2,2}(\Omega)$ be a bounded sequence such that $\det \nabla^2 u_n \geq c_0 \in \mathbb{R}$ a.e. in Ω . Then, up to a subsequence, u_n converges weakly in $W^{2,2}$, as well as it converges locally uniformly together with its gradients, to a C^1 function $u \in W^{2,2}(\Omega, \mathbb{R}^2)$ satisfying $\det \nabla^2 u \geq c_0$ a.e. in Ω .

Proof. Let $v_n(x) = \nabla u(x) + (|c_0| + 1)^{1/2}x^{\perp}$, where $x^{\perp} = (x_1, x_2)^{\perp} = (-x_1, x_2)$. Clearly, $v_n \in W^{1,2}(\Omega, \mathbb{R}^2)$ and, since $\nabla^2 u_n$ is a symmetric matrix, we get:

$$\det \nabla v_n(x) = \det \nabla^2 u_n(x) + |c_0| + 1 > 0 \qquad \forall a.e. \ x \in \Omega.$$

The convergence assertion follows by Corollary 7.7. Again, the Div-Curl Lemma applied to sequence ∇u_n implies the desired inequality for the limit function u.

A consequence of Theorem 7.6 is the following assertion about $W^{2,2}$ functions whose Hessian determinants are uniformly controlled from below:

Lemma 7.9. Assume that $u \in W^{2,2}(\Omega)$ satisfies:

(7.4)
$$\det \nabla^2 u(x) \ge c_0 \in \mathbb{R} \quad \forall a.e. \ x \in \Omega.$$

Then $u \in C^1(\Omega)$. Moreover, if $x_0 \in \Omega$ is a Lebesgue point for $\nabla^2 u$, i.e. for some $A \in \mathbb{R}^{2 \times 2}_{sym}$:

(7.5)
$$\omega(r) := \frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} |\nabla^2 u - A|^2 dx \to 0 \quad as \quad r \to 0^+,$$

then for all $\epsilon > 0$ there exists $r_0 > 0$ such that:

(7.6)
$$\forall r < r_0 \quad \forall a \in D_r = \overline{B(x_0, r)} \qquad \|\nabla u(x) - \nabla u(a) - A(x - a)\|_{\mathcal{C}^0(D_r)} \le \frac{1}{2} \epsilon r, \\ \|u(x) - u(a) - \nabla u(a) \cdot (x - a) - \frac{1}{2} (x - a) \cdot A(x - a)\|_{\mathcal{C}^0(D_r)} \le \epsilon r^2.$$

Proof. Following Kirchheim [18] we set $v = \nabla u$ and we write $\phi(x) = v(x) + (|c_0| + 1)^{1/2}x^{\perp}$. Trivially $\phi \in W^{1,2}(\Omega, \mathbb{R}^2)$ and, as before:

$$\det \nabla \phi = \det \nabla^2 u + |c_0| + 1 > 0 \qquad \forall a.e. \ x \in \Omega.$$

Applying Theorem 7.6 to ϕ shows that v is continuous and so $u \in \mathcal{C}^1(\Omega)$.

In what follows, we assume without loss of generality that $x_0 = 0$, u(0) = 0 and $v(0) = \nabla u(0) = 0$ (otherwise it is sufficient to translate Ω and to modify u by its tangent map at 0). For r sufficiently small and for all $x \in B_2 = B(0, 2)$ we define:

$$v_r(x) := \frac{1}{r}v(rx), \qquad \phi_r(x) := v_r(x) + (|c_0| + 1)^{1/2}x^{\perp},$$

so that:

$$\forall x \in B_2 \qquad \nabla v_r(x) = \nabla v(rx) = \nabla^2 u(rx), \qquad \nabla \phi_r(x) = \nabla^2 u(rx) + (|c_0| + 1)^{1/2} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix},$$
$$\det \nabla \phi_r(x) = \det \nabla v_r(x) + (|c_0| + 1) > 0.$$

Since $\phi_r \in W^{1,2}(B_2, \mathbb{R}^2)$, we can apply (7.2) to $x \in B_1$ and $\delta < R = 1$, to obtain for $r < r_0$ small enough:

$$\operatorname{osc}_{B(x,\delta)} \phi_r \leq \sqrt{2\pi} \left(\ln \frac{1}{\delta}\right)^{-1/2} \left(\int_{B(x,1)} |\nabla \phi_r|^2 \right)^{1/2} \\
\leq \sqrt{2\pi} \left(\ln \frac{1}{\delta}\right)^{-1/2} \left(2(|m|+1)^{1/2} |B_1| + \left(\int_{B(x,1)} |\nabla^2 u(ry)|^2 \, dy \right)^{1/2} \right) \\
\leq C \ln \left(\frac{1}{\delta} \right)^{-1/2} \left((|m|+1)^{1/2} + \frac{1}{r} ||\nabla^2 u||_{B(0,2r)} \right) \\
\leq C \ln \left(\frac{1}{\delta} \right)^{-1/2},$$

where C = C(m, |A|) > 0. Above we used the fact that $B(x, 1) \subset B_2$ and that 0 is a Lebesgue point for $\nabla^2 u$. Now, given $\epsilon > 0$ we choose $\delta > 0$ such that:

$$\ln(\frac{1}{\delta})^{-1/2} < \epsilon/C.$$

Consequently:

$$\forall x, y \in D_1 \quad \forall r < r_0 \qquad |x - y| < \delta \implies |\phi_r(x) - \phi_r(y)| < \epsilon.$$

Since $v_r - \phi_r$ is a given linear deformation, we conclude that the family:

$$\mathcal{F} = \{ v_r : D_1 \to \mathbb{R}^2; \ r < r_0 \}$$

is equicontinuous. On the other hand:

$$\int_{D_1} \left| \nabla v_r - A \right|^2 = \pi \omega(r) \to 0 \quad \text{as } r \to 0.$$

Let $\tilde{v}_r = v_r - \int_{B_1} v_r$ and apply the Poincaré inequality to obtain that

$$\tilde{v}_r \to Ax$$
 in $W^{1,2}(\Omega, \mathbb{R}^2)$ as $r \to 0$.

Now, equicontinuity of \mathcal{F} and $v_r(0) = 0$ yield, by Arzelà-Ascoli theorem, that a subsequence of v_r (which we do not relabel) converges uniformly to a continuous function V on D_1 . Since $v_r - \tilde{v}$ is constant, we deduce that V(x) - Ax = c is constant too. But then, evaluating at 0 gives c = 0. Hence, v_r uniformly converges to Ax on D_1 .

Let us fix $\epsilon > 0$ and choose r_0 so that:

$$\forall r < r_0 \qquad \|v_r(x) - Ax\|_{\mathcal{C}^0(D_1)} \le \frac{1}{4}\epsilon.$$

This implies:

$$\|\nabla u(x) - Ax\|_{\mathcal{C}^0(D_r)} \le \frac{1}{4}\epsilon r.$$

Fixing $a \in D_r$ we get:

$$\|\nabla u(x) - \nabla u(a) - A(x-a)\|_{\mathcal{C}^0(D_r)} \le \frac{1}{2}\epsilon r,$$

giving the first estimate in (7.6). Since diam $D_r = 2r$, the second estimate follows.

We now prove a simple useful lemma whose statement we quote from [35]:

Lemma 7.10 ([35], Lemma 2). Let $u \in C^1(\bar{\Omega})$ and fix $a \in \Omega$. Suppose that:

(7.7)
$$u(x) \ge u(a) + \nabla u(a) \cdot (x - a) \quad \forall x \in \partial \Omega$$

and that $\nabla u(x) \neq \nabla u(a)$ for all $x \in \Omega \setminus \{a\}$. Then u has a supporting hyperplane at a, i.e. (7.7) holds for all $x \in \Omega$. In particular, if Ω is convex then:

$$(Cu)(a) = u(a)$$

where Cu denotes the convexification of the function u over $\bar{\Omega}$:

$$(Cu)(a) = \sup \{T(a); \ T: \Omega \to \mathbb{R} \ \text{is affine and} \ T(x) \le u(x) \ \forall x \in \bar{\Omega} \}.$$

Proof. Consider the tangent map $T(x) = u(a) + \nabla u(a) \cdot (x - a)$. We now claim that $T(x) \leq u(x)$ for all $x \in \bar{\Omega}$. For otherwise, the continuous function g(x) = u(x) - T(x) would assume a negative minimum on $\bar{\Omega}$ at some $c \in \Omega \setminus \{a\}$. Hence $\nabla g(c) = 0$, which is a contradiction with the second assumption as $\nabla u(c) = \nabla T(c) = \nabla u(a)$.

We are ready to prove the key theorem of this section:

Proof of Theorem 7.3.

1. The C^1 regularity of u is an immediate consequence of Theorem 7.6. Recall the properties of the singular set S from Definition 7.1. Since $\det \nabla^2 u > 0$ a.e. in Ω , modulo a global change of sign for u we can choose $x_0 \in \Omega \setminus S$ a Lebesgue point of $\nabla^2 u$ as in (7.5), such that the matrix A is positive definite. Hence there exists $\lambda > 0$ for which $\xi \cdot A\xi \geq \lambda |\xi|^2$ for all $\xi \in \mathbb{R}^2$. By Lemma 7.9 for all $r < r_0$ the estimate (7.6) holds true with $\epsilon = \frac{1}{4}\lambda$, and without loss of generality ∇u is also one-to-one on $D_r = \overline{B(x_0, r)} \subset \Omega \setminus S$. By (7.6) it follows that:

$$\forall x \in \partial D_r$$
 $u(x) - u(x_0) - \nabla u(x_0) \cdot (x - x_0) \ge \frac{1}{2}(x - x_0) \cdot A(x - x_0) - \varepsilon r^2 \ge \frac{\lambda}{4}r^2 > 0.$

In view of Lemma 7.10, u therefore admits a supporting hyperplane at x_0 on D_r .

2. Our next claim is that u is locally convex in $\Omega \setminus S$. Since $\Omega \setminus S$ is open and connected, it is also path-wise connected. Therefore, for a fixed $x \in \Omega \setminus S$, there exists a continuous path within $\Omega \setminus S$ connecting x and x_0 , which can be covered with a finite chain of open balls $B_i \subset \Omega \setminus S$, $i = 1, 2 \dots n$, such that $x_0 \in B_1$, $B_i \cap B_{i+1} \neq \emptyset$ and $x \in B_n$. We now need the following strong theorem due to J. Ball:

Theorem ([1], Theorem 1). Let $\Omega \subset \mathbb{R}^n$ be open and convex, and let $u \in \mathcal{C}^1(\Omega)$. The necessary and sufficient condition for u to be strictly convex on Ω is:

- (i) ∇u is locally one-to-one, and
- (ii) there exists a locally supporting hyperplane for u at some point of Ω :

$$\exists x_0 \in \Omega \quad \forall \rho > 0 \quad \forall x \in B(x_0, \rho) \qquad u(x) \ge u(x_0) + \nabla u(x_0) \cdot (x - x_0).$$

Applying this result consecutively to each ball B_i , we deduce that u is strictly convex on B_n , hence it is locally convex at x.

3. To finish the proof, we fix a direction in \mathbb{R}^2 and consider the family of straight lines parallel to that direction. For almost every such line L, the 1-dimensional Lebesgue measure of $L \cap S$ is zero and $u \in W^{2,2}(L \cap \Omega)$. Also, u is locally convex on $(L \cap \Omega) \setminus S$ in view of the

previously proven claim. We now state the following easy lemma to show that u is convex on connected components of $L \cap \Omega$:

Lemma. Let $I \subset \mathbb{R}$ be an open bounded interval, $\phi \in W^{2,1}(I)$ and assume that ϕ is locally convex on $I \setminus S$, where S is a set of measure 0. Then ϕ is convex on I.

The proof is elementary. Since $\phi \in \mathcal{C}^1(I)$ and ϕ is locally convex on a full measure open subset of I, we deduce that $\phi'' \geq 0$ a.e. in I. But this immediately implies that ϕ' is increasing in I, hence ϕ is globally convex.

We have previously shown that u is convex on connected components of $L \cap \Omega$, for almost all straight lines L in any direction. By continuity of ∇u , the same must hold, in fact, for all lines, by approaching any given line with a selected sequence of 'good' lines and passing to the limit in the convexity inequality. This implies that actually u is convex on any convex subset of Ω and the proof is done.

For completeness, we now note another corollary of Lemma 7.9 and Lemma 7.10:

Lemma 7.11. Let $u \in W^{2,2}(\Omega)$ satisfy (7.4). Assume that x_0 is a Lebesgue point for u with A in (7.5) being positive definite. Assume that ∇u is one-to-one in a neighborhood of x_0 . Then u is locally convex at x_0 .

Proof. There exists $\lambda > 0$ for which $\xi \cdot A\xi \geq \lambda |\xi|^2$ for all $\xi \in \mathbb{R}^2$. By Lemma 7.9 for all $r < r_0$ and all $a \in D_r = \overline{B(x_0, r)}$, estimate (7.6) holds true with $\epsilon = \lambda/4$. Without loss of generality, ∇u is one-to one on D_r and, by (7.6):

 $\forall a \in B(x_0, r/2) \quad \forall x \in \partial B(x_0, r/2)$

$$u(x) - u(a) - \nabla u(a) \cdot (x - a) \ge \frac{1}{2}(x - a) \cdot A(x - a) - \varepsilon r^2 \ge \frac{\lambda}{4}r^2 > 0.$$

The assumptions of Lemma 7.10 are satisfied and hence u(a) = (Cu)(a) for all $a \in B(x_0, r/2)$. The claim is proved.

Remark 7.12. In proving Theorem 6.1 we only used the conclusion of Theorem 7.4 that v is locally one-to-one on a connected set of full measure. Therefore, the assumptions of Theorem 6.1 could potentially be relaxed (as in Theorem 7.3), but not to det $\nabla^2 u > 0$ a.e. Indeed, let $v = \nabla u$ be as in Example 6.2. Then v is not of integrable dilatation because:

$$\frac{|\nabla v|^2}{\det \nabla v}(x,y) \ge \frac{2}{x^2(1-y^2)},$$

and also the singular set $S = \{(0, y); y \in (-1, 1)\}$ coincides with the vanishing set of v where v is obviously not locally one-to-one. On the other hand, Theorem 7.3 can be also applied to the cases where $\det \nabla^2 u \in \mathcal{C}^0(\Omega)$ is positive a.e. and $\Omega \setminus f^{-1}(0)$ is connected.

8. Recovery sequence: A proof of Theorem 3.5

It is enough to prove Theorem 3.5 for $v \in C^{2,\beta}(\bar{\Omega})$ satisfying $\det \nabla^2 v = \det \nabla^2 v_0$. In the general case of $v \in W^{2,2}(\Omega)$ satisfying the same constraint, the result follows then by a diagonal argument in view of the density property established in Theorem 3.2.

1. We now recall the useful change of variable $\tilde{\phi}_h \in \mathcal{C}^{1,\beta}(\Omega^h, \mathbb{R}^3)$ between thin plates $\Omega^h = \Omega \times (-h/2, h/2)$ and thin shallow shells $(S_h)^h$:

$$\tilde{\phi}_h(x, x_3) = x + h^{\alpha} v_0(x) e_3 + x_3 \vec{n}^h(x) \qquad \forall (x, x_3) \in \Omega,$$

where \vec{n}^h is the unit normal vector to the midsurface S_h , given as the image of the map $\phi_h(x) = x + h^{\alpha}v_0(x)e_3$:

$$\vec{n}^h(x) = \frac{\partial_1 \phi_h(x) \times \partial_2 \phi_h(x)}{|\partial_1 \phi_h(x) \times \partial_2 \phi_h(x)|} = \frac{1}{\sqrt{1 + h^{2\alpha} |\nabla v_0|^2}} \left(-h^{\alpha} (\nabla v_0)^* + e_3 \right).$$

By Theorem 3.1, there exists an equibounded sequence $w_h \in C^{2,\beta}(\bar{\Omega}, \mathbb{R}^3)$ such that the deformations $\xi_h(x) = x + h^{\alpha}v(x)e_3 + h^{2\alpha}w_h(x)$ are isometrically equivalent to $id + h^{\alpha}v_0e_3$:

(8.1)
$$\forall 0 < h \ll 1 \qquad (\nabla \xi_h)^T \nabla \xi_h = \nabla (\mathrm{id} + h^\alpha v_0 e_3)^T \nabla (\mathrm{id} + h^\alpha v_0 e_3).$$

Define now the recovery sequence $u^h \in \mathcal{C}^{1,\beta}((S_h)^h,\mathbb{R}^3)$ by:

$$v^{h}(x,x_{3}) = u^{h}(\tilde{\phi}_{h}(x,x_{3})) = \xi_{h}(x) + x_{3}\vec{N}^{h}(x) + \frac{x_{3}^{2}}{2}h^{\alpha}d^{h}(x),$$

where \vec{N}^h is the unit normal vector to the image surface $\xi_h(\Omega)$:

$$\vec{N}^h(x) = \frac{\partial_1 \xi_h(x) \times \partial_2 \xi_h(x)}{|\partial_1 \xi_h(x) \times \partial_2 \xi_h(x)|} = (-h^{\alpha} (\nabla v)^* + e_3) + \mathcal{O}(h^{2\alpha}),$$

while the 'warping' vector fields $d^h \in \mathcal{C}^{1,\beta}(\bar{\Omega},\mathbb{R}^3)$, approximating the effective warping $d \in \mathcal{C}^{0,\alpha}(\bar{\Omega},\mathbb{R}^3)$ are defined so that:

(8.2)
$$h^{\alpha} \|d^{h}\|_{\mathcal{C}^{1,\beta}} \leq C \quad \text{and} \quad \lim_{h \to 0} \|d^{h} - d\|_{L^{\infty}} = 0,$$
$$\mathcal{Q}_{2} (\nabla^{2} v_{0} - \nabla^{2} v) = \min \{ \mathcal{Q}_{3}(F); F \in \mathbb{R}^{3 \times 3}, F_{tan} = \nabla^{2} v_{0} - \nabla^{2} v \}$$
$$= \mathcal{Q}_{3} ((\nabla^{2} v_{0} - \nabla^{2} v)^{*} + \operatorname{sym}(d \otimes e_{3})).$$

For $F \in \mathbb{R}^{3\times 3}$, by F_{tan} we denote the principal 2×2 minor of F. Recall also that the quadratic form \mathcal{Q}_3 is given by $\mathcal{Q}_3(F) = D^2W(\mathrm{Id})(F,F)$.

2. Because of the first condition in (8.2), the statements in Theorem 3.5 (i), (ii) easily follow. In order to compute the energy limit in (iii), we write:

(8.3)
$$I^{h}(u^{h}) = \frac{1}{h} \int_{\Omega^{h}} W\left((\nabla v^{h})(b^{h})^{-1}\right) \det \nabla \tilde{\phi}_{h} = \frac{1}{h} \int_{\Omega^{h}} W\left(\sqrt{K^{h}}\right) \det b^{h},$$

where:

$$b^h = \nabla \tilde{\phi}^h$$

while the frame invariance of W justifies the second equality in (8.3) with:

$$K^h(x, x_3) = (b^h)^{-1,T} (\nabla v^h)^T (\nabla v^h) (b^h)^{-1}.$$

We will now compute the entries of the symmetric matrix field K^h , up to terms of order $o(h^{\alpha+1})$. In what follows we adopt the convention that all equalities hold modulo quantities which are uniformly $o(h^{\alpha+1})$. Call $M^h = (\nabla v^h)^T \nabla v^h$. Since:

$$\nabla_{tan}v^h = \nabla \xi_h + x_3 \nabla \vec{N}^h + o(h^{\alpha+1}), \qquad \partial_3 v^h = \vec{N}^h + x_3 h^\alpha d^h.$$

we obtain, in view of (8.1):

$$M_{tan}^{h} = \nabla (\mathrm{id} + h^{\alpha} v_{0} e_{3})^{T} \nabla (\mathrm{id} + h^{\alpha} v_{0} e_{3}) + 2x_{3} \mathrm{sym} \left((\nabla \xi_{h})^{T} \nabla \vec{N}^{h} \right)$$

$$= \mathrm{Id}_{2} + h^{2\alpha} \nabla v_{0} \otimes \nabla v_{0} - 2x_{3} h^{\alpha} \nabla^{2} v + o(h^{\alpha+1}),$$

$$M_{13,23}^{h} = (M^{h})_{13,23}^{T} = x_{3} h^{\alpha} d_{tan}^{h} + o(h^{\alpha+1}),$$

$$M_{33}^{h} = |\vec{N}^{h} + x_{3} h^{\alpha} d^{h}|^{2} = 1 + 2x_{3} h^{\alpha} d_{3}^{h} + o(h^{\alpha+1}).$$

Further, by a direct calculation, one obtains:

$$(b^h)_{tan} = \mathrm{Id}_2 - x_3 h^{\alpha} \nabla^2 v_0 + o(h^{\alpha+1}),$$

$$(b^h)_{13,23}^T = h^{\alpha} \nabla v_0 + o(h^{\alpha+1}), \qquad (b^h)_{13,23,33} = \vec{n}^h.$$

and the inverse matrix $(b^h)^{-1}$ has the following structure:

$$((b^h)^{-1})_{tan} = A + o(h^{\alpha+1}),$$

$$((b^h)^{-1})_{13,23} = h^{\alpha} A \nabla v_0 + o(h^{\alpha+1}), \qquad ((b^h)^{-1})_{13,23,33}^T = \vec{n}^h.$$

where the principal minor $A(x) \in \mathbb{R}^{2 \times 2}$ of $(b^h(x))^{-1}$ is the symmetric matrix:

(8.4)
$$A = \left(\mathrm{Id}_2 + h^{2\alpha} \nabla v_0 \otimes \nabla v_0 - x_3 h^{\alpha} \nabla^2 v_0 \right)^{-1}.$$

3. We now compute,

$$((b^{h})^{-1,T}M^{h})_{tan} = \operatorname{Id} + x_{3}h^{\alpha}A(\nabla^{2}v_{0} - 2\nabla^{2}v) + x_{3}h^{\alpha}\vec{n}_{tan}^{h} \otimes \vec{d}_{tan}^{h} + o(h^{\alpha+1})$$

$$= \operatorname{Id}_{2} + x_{3}h^{\alpha}(\nabla^{2}v_{0} - 2\nabla^{2}v) + o(h^{\alpha+1})$$

$$((b^{h})^{-1,T}M^{h})_{13,23}^{T} = h^{\alpha}(\operatorname{Id} + h^{2\alpha}\nabla v_{0} \otimes \nabla v_{0})A\nabla v_{0} + x_{3}h^{\alpha}d_{tan}^{h} + o(h^{\alpha+1})$$

$$= h^{\alpha}\nabla v_{0} + x_{3}h^{\alpha}d_{tan}^{h} + o(h^{\alpha+1})$$

$$((b^{h})^{-1,T}M^{h})_{13,23} = x_{3}h^{\alpha}d_{tan}^{h} + \vec{n}_{tan}^{h} + o(h^{\alpha+1})$$

$$((b^{h})^{-1,T}M^{h})_{33} = \vec{n}_{3}^{h} + 2x_{3}h^{\alpha}d_{3}^{h} + o(h^{\alpha+1}),$$

where we used that $A(\operatorname{Id} + h^{2\alpha}\nabla v_0 \otimes \nabla v_0 - 2x_3h^{\alpha}\nabla^2 v) = \operatorname{Id} + x_3h^{\alpha}A(\nabla^2 v_0 - 2\nabla^2 v)$ and that $h^{\alpha}(\operatorname{Id} + h^{2\alpha}\nabla v_0 \otimes \nabla v_0)A = h^{\alpha}(\operatorname{Id} + x_3h^{\alpha}\nabla^2 v_0A) = h^{\alpha}\operatorname{Id} + o(h^{\alpha+1})$. Consequently:

$$\begin{split} K_{tan}^{h} &= A + x_{3}h^{\alpha}(\nabla^{2}v_{0} - 2\nabla^{2}v) + \vec{n}_{tan}^{h} \otimes \vec{n}_{tan}^{h} + o(h^{\alpha+1}) \\ &= A \left(\operatorname{Id} + x_{3}h^{\alpha}(\nabla^{2}v_{0} - 2\nabla^{2}v) + \vec{n}_{tan}^{h} \otimes \vec{n}_{tan}^{h} + \frac{h^{4\alpha}|\nabla v_{0}|^{2}}{1 + h^{2\alpha}|\nabla v_{0}|^{2}} \nabla v_{0} \otimes \nabla v_{0} \right) + o(h^{\alpha+1}) \\ &= A \left(\operatorname{Id} + x_{3}h^{\alpha}\nabla^{2}v_{0} - 2x_{3}h^{\alpha}\nabla^{2}v + h^{2\alpha}\nabla v_{0} \otimes \nabla v_{0} \right) + o(h^{\alpha+1}) \\ &= \operatorname{Id}_{2} + 2x_{3}h^{\alpha}A(\nabla^{2}v_{0} - \nabla^{2}v) + o(h^{\alpha+1}) \\ &= \operatorname{Id}_{2} + 2x_{3}h^{\alpha}(\nabla^{2}v_{0} - \nabla^{2}v) + o(h^{\alpha+1}) \\ &= \operatorname{Id}_{2} + 2x_{3}h^{\alpha}(\nabla^{2}v_{0} - \nabla^{2}v) + o(h^{\alpha+1}) \\ &= h^{\alpha}A\nabla v_{0} + x_{3}h^{\alpha}d_{tan}^{h} + \vec{n}_{3}^{h}\vec{n}_{tan}^{h} + o(h^{\alpha+1}) \\ &= h^{\alpha}A\nabla v_{0} + x_{3}h^{\alpha}d_{tan}^{h} - \frac{h^{\alpha}}{1 + h^{2\alpha}|\nabla v_{0}|^{2}}\nabla v_{0} + o(h^{\alpha+1}) \\ &K_{33}^{h} = h^{2\alpha}\langle A\nabla v_{0}, \nabla v_{0}\rangle + 2x_{3}h^{\alpha}d_{3}^{h} + |\vec{n}_{3}^{h}|^{2} + o(h^{\alpha+1}) \\ &= 1 + h^{2\alpha}\langle A\nabla v_{0}, \nabla v_{0}\rangle + 2x_{3}h^{\alpha}d_{3}^{h} - \frac{h^{2\alpha}|\nabla v_{0}|^{2}}{1 + h^{2\alpha}|\nabla v_{0}|^{2}} + o(h^{\alpha+1}), \end{split}$$

where we used that $\vec{n}_{tan}^h \otimes \vec{n}_{tan}^h = \frac{h^{2\alpha}}{1 + h^{2\alpha} |\nabla v_0|^2} \nabla v_0 \otimes \nabla v_0$. Observe that:

$$\begin{split} h^{\alpha}A\nabla v_{0} - \frac{h^{\alpha}}{1 + h^{2\alpha}|\nabla v_{0}|^{2}}\nabla v_{0} &= \\ &= \frac{h^{\alpha}}{1 + h^{2\alpha}|\nabla v_{0}|^{2}}A\left((1 + h^{2\alpha}|\nabla v_{0}|^{2})\nabla v_{0} - \nabla v_{0} - h^{2\alpha}|\nabla v_{0}|^{2}\nabla v_{0}\right) + o(h^{\alpha+1}) \\ &= o(h^{\alpha+1}). \end{split}$$

Therefore, in fact:

$$K_{13,23}^h = x_3 h^{\alpha} d_{tan}^h + o(h^{\alpha+1}), \qquad K_{33}^h = 1 + 2x_3 h^{\alpha} d_3^h + o(h^{\alpha+1}).$$

Concluding, we get:

$$K^{h} = \mathrm{Id}_{3} + 2x_{3}h^{\alpha}((\nabla^{2}v_{0} - \nabla^{2}v)^{*} + \mathrm{sym}(d^{h} \otimes e_{3})) + o(h^{\alpha+1}).$$

4. Taylor expanding W at Id_3 and using (8.2) we now see that:

$$W(\sqrt{K^h}) = W(\mathrm{Id}_3 + x_3 h^{\alpha} ((\nabla^2 v_0 - \nabla^2 v)^* + \mathrm{sym}(d^h \otimes e_3)) + o(h^{\alpha+1}))$$

= $\frac{1}{2} x_3^2 h^{2\alpha} \mathcal{Q}_3 ((\nabla^2 v_0 - \nabla^2 v)^* + \mathrm{sym}(d^h \otimes e_3)) + o(h^{2\alpha+2}).$

Note that $\nabla \tilde{\phi}_h = 1 + \mathcal{O}(h^{\alpha})$. By this fact, recalling (8.3) and the convergence in (8.2), it follows that:

$$\lim_{h \to 0} \frac{1}{h^{2\alpha+2}} I^{h}(u^{h}) = \lim_{h \to 0} \frac{1}{h^{2\alpha+2}} \frac{1}{h} \int_{\Omega^{h}} W(\sqrt{K^{h}}) (1 + \mathcal{O}(h^{\alpha}))$$

$$= \lim_{h \to 0} \frac{1}{2h^{3}} \int_{\Omega^{h}} x_{3}^{2} \mathcal{Q}_{3} ((\nabla^{2} v_{0} - \nabla^{2} v)^{*} + \operatorname{sym}(d \otimes e_{3}))$$

$$= \lim_{h \to 0} \frac{1}{2h^{3}} (\int_{-h/2}^{h/2} x_{3}^{2} dx_{3}) \int_{\Omega} \mathcal{Q}_{2} (\nabla^{2} v_{0} - \nabla^{2} v) dx$$

$$= \frac{1}{24} \int_{\Omega} \mathcal{Q}_{2} (\nabla^{2} v_{0} - \nabla^{2} v) dx.$$

Since, clearly $v_3^h = h^{\alpha}v + \mathcal{O}(h^{2\alpha})$, we obtain:

$$\lim_{h \to 0} \frac{1}{h^{2\alpha+2}} \frac{1}{h} \int_{\Omega^h} h^{\alpha+2} f v_3^h \det b^h = \lim_{h \to 0} \frac{1}{h^{2\alpha+2}} \frac{1}{h} \int_{\Omega^h} h^{\alpha+2} f(h^{\alpha} v + \mathcal{O}(h^{2\alpha})) = \int_{\Omega} f v \, dx.$$

The proof of Theorem 3.5 is complete.

Acknowledgments. This project is based upon work supported by, among others, the National Science Foundation. M.L. is partially supported by the NSF grants DMS-0707275 and DMS-0846996, and by the Polish MN grant N N201 547438. L.M. is supported by the MacArthur Foundation, M.R.P. is partially supported by the NSF grants DMS-0907844 and DMS-1210258.

REFERENCES

- [1] J.M. Ball, Strict convexity, strong ellipticity and regularity in the calculus of variations, Math. Proc. Camb. Phil. Soc., 87, 501, (1980).
- [2] L.A. Caffarelli, Interior W^{2,p} estimates for solutions of the Monge-Ampère equation, Ann. of Math.
 (2) 131 (1990), no. 1, 135–150.
- [3] L.A. Caffarelli and X. Cabré, Fully nonlinear elliptic equations, American Mathematical Society Colloquium Publications, 43 AMS (1995).
- [4] C.R. Calladine, (1983) Theory of shell structures. Cambridge University Press, UK.
- [5] P.G. Ciarlet, Mathematical Elasticity, North-Holland, Amsterdam (1993).
- [6] P.G. Ciarlet, An introduction to differential geometry with applications to elasticity, Springer, Dordrecht, 2005.
- [7] G. Dal Maso, An introduction to Γ-convergence, Progress in Nonlinear Differential Equations and their Applications, 8, Birkhäuser, MA (1993).
- [8] Dervaux, J., Ciarletta, P. and Ben Amar, M. Morphogenesis of thin hyperelastic plates: a constitutive theory of biological growth in the Föppl-von Kármán limit. *Journal of the Mechanics and Physics of Solids*, **57**(3), 458–471, (2009).
- [9] I. Fonseca and W. Gangbo, Degree theory in analysis and applications, Oxford Univ. Press, 1995.
- [10] G. Friesecke, R. James and S. Müller, A theorem on geometric rigidity and the derivation of nonlinear plate theory from three dimensional elasticity, Comm. Pure. Appl. Math., 55 (2002), 1461–1506.
- [11] G. Friesecke, R. James and S. Müller, A hierarchy of plate models derived from nonlinear elasticity by gamma-convergence, Arch. Ration. Mech. Anal., 180 (2006), no. 2, 183–236.
- [12] Q. Han and J.-X. Hong, *Isometric embedding of Riemannian manifolds in Euclidean spaces*, Mathematical Surveys and Monographs, **130** American Mathematical Society, Providence, RI (2006).

- [13] P. Hornung, Continuation of infinitesimal bendings on developable surfaces and equilibrium equations for nonlinear bending theory of plates, Communications in Partial Differential Equations, 38, Issue 11, (2013), 1368–1408.
- [14] P. Hornung, M. Lewicka and M.R. Pakzad, Infinitesimal isometries on developable surfaces and asymptotic theories for thin developable shells, J. of Elasticity, (2012), doi 10.1007/s10659-012-9391-
- [15] Q. Huang, Sharp regularity results on second derivatives of solutions to the Monge-Ampère equation with VMO type data, Comm. Pure Appl. Math. 62 (2009), no. 5, 677–705.
- [16] T. Iwaniec and V. Šverák, On mappings with integrable dilatation, Proc. Amer. Math. Soc. 118, 181–188, (1993).
- [17] T. von Kármán, Festigkeitsprobleme im Maschinenbau, in Encyclopädie der Mathematischen Wissenschaften. Vol. IV/4, pp. 311-385, Leipzig, 1910.
- Kirchheim, Geometry andrigidity of microstructures, Habilitation University Leipzig, 2001 (see also: MPI-MIS Lecture Notes 16/2003sis, of http://www.mis.mpg.de/preprints/ln/index.html).
- [19] Koehl MAR, Silk WK, Liang HY, Mahadevan L (2008) How kelp produce blade shapes suited to different flow regimes: A new wrinkle. *Integ. and Comp. Biol.* 48, 834-851.
- [20] M. Lewicka, L. Mahadevan and M.R. Pakzad, The Föppl-von Kármán equations for plates with incompatible strains, Proceedings of the Royal Society A 467 (2011), 402-426.
- [21] M. Lewicka, L. Mahadevan and M.R. Pakzad, Models for elastic shells with incompatible strains, submitted.
- [22] M. Lewicka, M.G. Mora and M.R. Pakzad, Shell theories arising as low energy Γ-limit of 3d nonlinear elasticity, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5) Vol. IX (2010), 1–43.
- [23] M. Lewicka, M.G. Mora and M.R. Pakzad, *The matching property of infinitesimal isometries on elliptic surfaces and elasticity of thin shells*, Arch. Rational Mech. Anal. (2010).
- [24] M. Lewicka and M.R. Pakzad, *The infinite hierarchy of elastic shell models; some recent results and a conjecture*, accepted in Fields Institute Communications (2009).
- [25] H. Liang and L. Mahadevan, The shape of a long leaf, Proc. Nat. Acad. Sci. (2009).
- [26] H. Liang and L. Mahadevan, Growth, geometry and mechanics of the blooming lily, Proc. Nat. Acad. Sci., 108, 5516-21, (2011).
- [27] J.Malý and O. Martio, Lusin's condition (N) and mappings of the class W^{1,n}, J. reine angew. Math. 458 (1995), 19–36.
- [28] J. J. Manfredi, Weakly monotone functions, J. Geom. Anal. 4, no. 3, 393–402 (1994).
- [29] E.H. Mansfield, (1989) The bending and stretching of plates (Cambridge University Press), 2nd edn.
- [30] S. Mardare, On isometric immersions of a Riemannian space with little regularity, Anal. Appl. (Singap.) 2 (2004), no. 3, 193–226.
- [31] C.B. Morrey, Multiple integrals in the calculus of variations, Springer, 1966.
- [32] S. Müller and M.R. Pakzad, Regularity properties of isometric immersions, Math. Z., 251 (2005), no. 2, 313–331.
- [33] L. Nirenberg, The Weyl and Minkowski problems in differential geometry in the large, Comm. Pure Appl. Math., 6 (1953), 337–394.
- [34] B. Schmidt, Plate theory for stressed heterogeneous multilayers of finite bending energy, J. Math. Pures Appl. 9, 88 (2007), no. 1, 107–122.
- [35] V. Šverák, On regularity for the Monge-Ampère equation without convexity assumptions, preprint, Heriot-Watt University (1991).
- [36] V. Sverák, Regularity properties of deformations with finite energy, Arch. Rational Mech. Anal. 100, (1988) 105–127.
- [37] N.S. Trudinger and X.J. Wang, *The Monge-Ampère equation and its geometric applications*, Handbook of Geometric Analysis, International Press, (2008) Vol I, 467-524.
- [38] W.G. Van Doorn and U. Van Meeteren, Flower opening and closure: a review, J. Exp. Botany 54, 1801-1812 (2003).

[39] S.K. Vodopyanov and V.M. Goldstein, Quasiconformal mappings and spaces with generalized derivatives, Siberian Math. J. 17, 399–411, (1976), (Sibirskii Mat. Zh. 17, 515–531, (1976)).

Marta Lewicka, University of Pittsburgh, Department of Mathematics, 301 Thackeray Hall, Pittsburgh, PA 15260

L. Mahadevan, Harvard University, School of Engineering and Applied Sciences, and Department of Physics, Cambridge, MA 02138

Mohammad Reza Pakzad, University of Pittsburgh, Department of Mathematics, 301 Thackeray Hall, Pittsburgh, PA 15260

 $E ext{-}mail\ address: lewicka@pitt.edu, lm@seas.harvard.edu, pakzad@pitt.edu}$