Squared-Norm Empirical Process in Banach Space^{*†}

Vincent Q. Vu Department of Statistics The Ohio State University Columbus, OH vqv@stat.osu.edu Jing Lei Department of Statistics Carnegie Mellon University Pittsburgh, PA jinglei@andrew.cmu.edu

Abstract

This note extends a recent result of Mendelson on the supremum of a quadratic process to squared norms of functions taking values in a Banach space. Our method of proof is a reduction by a symmetrization argument and observation about the subadditivity of the generic chaining functional. We provide an application to the supremum of a linear process in the sample covariance matrix indexed by finite rank, positive definite matrices.

1 Introduction

Let F be a class of \mathbb{R} -valued functions on the probability space (Ω, \mathbb{P}) and $\{X_1, \ldots, X_n\}$ be independent, identically distributed random variables. Let $\|\cdot\|_{\psi_{\alpha}}$ be the Orlicz ψ_{α} norm for $\alpha \geq 1$ and γ_{α} be Talagrand's generic chaining complexity of F under the ψ_{α} metric. Mendelson has proved the following theorem:

Theorem 1.1 (Mendelson (2010)). If F is a symmetric class of mean-zero functions on (Ω, \mathbb{P}) then there exists absolute constants c_1 , c_2 , and c_3 such that for any $t \ge c_1$, with probability at least $1 - 2\exp(-c_2t^{2/5})$

$$\sup_{f \in F} \left| \frac{1}{n} \sum_{i=1}^{n} f^2(X_i) - \mathbb{E} f^2(X_i) \right| \le c_3 t \left\{ \frac{d_{\psi_1}(F)\gamma_2(F,\psi_2)}{\sqrt{n}} + \frac{\gamma_2^2(F,\psi_2)}{n} \right\},$$

where $d_{\psi_1}(F) = \sup_{f \in F} \|f(X_1)\|_{\psi_1}$.

The goal of this note is to extend the preceding theorem to a class G of functions that take values in a Banach space $(B, \|\cdot\|)$. We wish to bound

$$\sup_{g \in G} \left| \frac{1}{n} \sum_{i=1}^{n} \|g(X_i)\|^2 - \mathbb{E} \|g(X_i)\|^2 \right|$$
(1)

^{*}This note was first written in 2012 and a version has been available on the first author's website. It is now posted on arXiv to provide a universal identifier for citations elsewhere. It is not intended for publication in its current form, but a revision incorporating later work will be posted separately.

[†]This research was supported by NSF grants DMS-0903120 and BCS-0941518.

with a similar exponential tail probability bound. A special case is $g(X_1) = A^T X_1$, where X_1 is a random vector in \mathbb{R}^p and $A \in \mathcal{A}$, a class of $p \times k$ matrices. In this note we will assume that G is countable to avoid unnecessary measurability issues.

When $(B, \|\cdot\|) = (\mathbb{R}, |\cdot|)$, this expectation reduces to the one addressed by Theorem 1.1. Our goal is to bound the expectation in terms of complexity parameters of the class G. We will use a symmetrization argument to put (1) into a form suitable for Theorem 1.1. The next section contains an abstract technical result concerning subadditivity of the generic chaining γ_{α} functional (Talagrand 2005) that enables our reduction.

2 The Generic Chaining

Definition 2.1. Given a metric space (T, d) and A a subset of T, define the diameter of A to be

$$\Delta(A) := \sup_{t,u \in A} d(t,u) \,. \tag{2}$$

Definition 2.2. Given a metric space (T,d) and \mathcal{A} a partition of T, for $t \in T$, define $\mathcal{A}(t)$ to be the unique element of \mathcal{A} that contains t.

Definition 2.3 (Talagrand (2005)). Given a metric space (T, d), an admissible sequence is an increasing sequence $(\mathcal{A}_s)_{s\geq 0}$ of partitions of T such that $|\mathcal{A}_0| = 1$ and, for $s \geq 1$, $|\mathcal{A}_s| \leq 2^{2^s}$. For $\alpha \geq 1$, define the γ_{α} functional by

$$\gamma_{\alpha}(T,d) \coloneqq \inf \sup_{t \in T} \sum_{s \ge 0} 2^{s/\alpha} \Delta(\mathcal{A}_s(t)),$$

where the infimum is taken over all admissible sequences.

Lemma 2.1. If (T_1, d) and (T_2, d) are metric spaces, then

$$\gamma_{\alpha}(T_1 \cup T_2, d) \leq 3 \left[\Delta(T_1 \cup T_2) + \gamma_{\alpha}(T_1, d) + \gamma_{\alpha}(T_2, d) \right].$$

Moreover, if $T_1 \cap T_2 \neq \emptyset$, then

$$\gamma_{\alpha}(T_1 \cup T_2, d) \le 9 \big[\gamma_{\alpha}(T_1, d) + \gamma_{\alpha}(T_2, d) \big]$$

Proof. Let $(\mathcal{A}_s)_{s\geq 0}$ be an admissible sequence for T_1 such that

$$\sum_{s\geq 0} 2^{s/\alpha} \Delta(\mathcal{A}_s(t)) \leq 2\gamma_\alpha(T_1, d) \text{ for all } t \in T_1,$$
(3)

and $(\mathcal{B}_s)_{s\geq 0}$ an admissible sequence for T_2 such that

$$\sum_{s\geq 0} 2^{s/\alpha} \Delta(\mathcal{B}_s(t)) \leq 2\gamma_\alpha(T_2, d) \text{ for all } t \in T_2.$$
(4)

We define partitions C_s of $T_1 \cup T_2$ as follows. Let $C_0 = \{T_1 \cup T_2\}, C_1 = C_0$, and, for $s \ge 2$, let C_s be the partition consisting of the sets

$$A \cap (T_1 \setminus T_2) \text{ for } A \in \mathcal{A}_{s-2},$$

$$B \cap (T_2 \setminus T_1) \text{ for } B \in \mathcal{B}_{s-2}, \text{ and}$$

$$A \cap B \text{ for } A \in \mathcal{A}_{s-2} \text{ and } B \in \mathcal{B}_{s-2}$$

It is straightforward to check that $(C_s)_{s\geq 0}$ is an increasing sequence of partitions of $T_1 \cup T_2$, and that $|C_0| = |C_1| = 1$ and, for $s \geq 2$,

$$\begin{aligned} |\mathcal{C}_{s}| &\leq |\mathcal{A}_{s-2}| + |\mathcal{B}_{s-2}| + |\mathcal{A}_{s-2}| \times |\mathcal{B}_{s-2}| \\ &\leq 2^{2^{s-2}} + 2^{2^{s-2}} + 2^{2^{s-1}} \leq 2^{2^{s}}. \end{aligned}$$

Thus $(\mathcal{C}_s)_{s\geq 0}$ is an admissible sequence for $T_1 \cup T_2$. Let $t \in T_1 \cup T_2$. Since, for $s \geq 2$, every element of \mathcal{C}_s is a subset of an element of \mathcal{A}_{s-2} and/or \mathcal{B}_{s-2} ,

$$\begin{split} \sum_{s\geq 0} 2^{s/\alpha} \Delta(\mathcal{C}_s(t)) &\leq 3\Delta(T_1 \cup T_2) + \sum_{s\geq 2} 2^{s/\alpha} \Delta(\mathcal{C}_s(t)) \\ &\leq 3\Delta(T_1 \cup T_2) + \sum_{s\geq 0} 2^{s/\alpha} \Delta(\mathcal{A}_s(t)) + \sum_{s\geq 0} 2^{s/\alpha} \Delta(\mathcal{B}_s(t)) \\ &\leq 3 \left[\Delta(T_1 \cup T_2) + \gamma_\alpha(T_1, d) + \gamma_\alpha(T_2, d) \right], \end{split}$$

by (3) and (4). Then

$$\gamma_{\alpha}(T_1 \cup T_2, d) \leq \sup_{t \in T_1 \cup T_2} \sum_{s \geq 0} 2^{s/\alpha} \Delta(\mathcal{C}_s(t))$$
$$\leq 3 \left[\Delta(T_1 \cup T_2) + \gamma_{\alpha}(T_1, d) + \gamma_{\alpha}(T_2, d) \right]$$

For the "moreover" part, if $u \in T_1 \cap T_2$, then by the triangle inequality

$$\begin{split} \Delta(T_1 \cup T_2) &= \sup_{t_1, t_2 \in T_1 \cup T_2} d(t_1, t_2) \\ &\leq \sup_{t_1, t_2 \in T_1 \cup T_2} d(t_1, u) + d(t_2, u) \\ &\leq 2 \left[\Delta(T_1) + \Delta(T_2) \right] \\ &\leq 2 \left[\gamma_2(T_1, d) + \gamma_2(T_2, d) \right]. \end{split}$$

3 Symmetrization

The following theorem is our main result. Its proof is a symmetrization argument followed by an application of Theorem 1.1 to and Lemma 2.1.

Theorem 3.1. If X_1, X_2, \ldots, X_n are independent, identically distributed random variables, and G is a symmetric class of functions taking values in a Banach space $(B, \|\cdot\|)$, then there exists absolute constants c_1, c_2 , and c_3 such that for all $t \ge c_1$, with probability at least $1 - 2\exp(-c_2t^{2/5})$,

$$\sup_{g \in G} \left| \frac{1}{n} \sum_{i=1}^{n} \|g(X_i)\|^2 - \mathbb{E} \|g(X_i)\|^2 \right| \le c_3 t \left\{ \frac{\sup_{g \in G} \|\|g(X_1)\|\|_{\psi_1} \gamma_2(G,d)}{\sqrt{n}} + \frac{\gamma_2^2(G,d)}{n} \right\},$$
(5)

where d is the metric on G defined by

$$d(g_1, g_2) = \|\|g_1(X_1) - g_2(X_1)\|\|_{\psi_2} .$$
(6)

Proof. Let $\{\epsilon_1, \ldots, \epsilon_n\}$ be independent, identically distributed Rademacher random variables that are independent of $\{X_1, \ldots, X_n\}$. Define the function class

$$F = \left\{ f : f(x, \epsilon) = \epsilon \| g(x) \| \text{ for some } g \in G \right\},\$$

and let

$$F_0 = F \cup -F$$

 F_0 is a symmetric class of mean-zero functions by construction. So Theorem 1.1 implies that, for $t \ge c_1$, with probability at least $1 - 2 \exp(-c_2 t^{2/5})$,

$$\begin{split} \sup_{g \in G} & \left| \frac{1}{n} \sum_{i=1}^{n} \|g(X_{i})\|^{2} - \mathbb{E} \|g(X_{i})\|^{2} \right| \\ &= \sup_{f \in F_{0}} \left| \frac{1}{n} \sum_{i=1}^{n} f^{2}(X_{i}, \epsilon_{i}) - \mathbb{E} f^{2}(X_{i}, \epsilon_{i}) \right| \\ &\leq c_{3} t \left\{ \frac{\sup_{g \in G} \|\|g(X_{1})\|\|_{\psi_{1}} \gamma_{2}(F_{0}, \psi_{2})}{\sqrt{n}} + \frac{\gamma_{2}^{2}(F_{0}, \psi_{2})}{n} \right\}. \end{split}$$

The rest of the proof deals with showing that

$$\gamma_2(F_0,\psi_2) \le C\gamma_2(G,d)\,. \tag{7}$$

Apply Lemma 2.1 to

$$\gamma_2(F_0, \psi_2) \le 3 \left[\Delta(F \cup -F) + \gamma_2(F, \psi_2) + \gamma_2(-F, \psi_2) \right] \\= 3 \left[\Delta(F \cup -F) + 2\gamma_2(F, \psi_2) \right].$$
(8)

For any $f_1, f_2 \in F$ with corresponding $g_1, g_2 \in G$, we have by the triangle inequality that

$$\begin{aligned} \|f_1(X_1,\epsilon_1) - f_2(X_1,\epsilon_1)\|_{\psi_2} &= \|\|g_1(X_1)\| - \|g_2(X_1)\|\|_{\psi_2} \\ &\leq \|\|g_1(X_1) - g_2(X_1)\|\|_{\psi_2} \\ &=: d(g_1,g_2) \,. \end{aligned}$$

Then by Theorem 1.3.6 of Talagrand (2005),

$$\gamma_2(F,\psi_2) \le C\gamma_2(G,d) \tag{9}$$

for an absolute constant C > 0. Using the triangle inequality and the symmetry of G,

$$\begin{aligned} \Delta(F \cup -F) &= \max \left\{ \sup_{g_1, g_2 \in G} \| \|g_1(X_1)\| - \|g_2(X_1)\| \|_{\psi_2} , \sup_{g_1, g_2 \in G} \| \|g_1(X_1)\| + \|g_2(X_1)\| \|_{\psi_2} \right\} \\ &\leq \max \left\{ \sup_{g_1, g_2 \in G} d(g_1, g_2) , 2 \sup_{g \in G} \| \|g(X_1)\| \|_{\psi_2} \right\} \\ &= \max \left\{ \sup_{g_1, g_2 \in G} d(g_1, g_2) , \sup_{g \in G} \| \|g(X_1) - (-g(X_1))\| \|_{\psi_2} \right\} \\ &= \sup_{g_1, g_2 \in G} d(g_1, g_2) \\ &\leq \gamma_2(G, d) . \end{aligned}$$

Substituting the preceding inequality and (9) into (8) proves (7).

4 Linear Transformations

Let $\mathcal{A} \subset \mathbb{R}^{p \times k}$

$$G = \{g : g(x) = A^T x \text{ and } A \in \mathcal{A}\}.$$

Observe that

$$\frac{1}{n} \sum_{i=1}^{n} \|g(X_i)\|_2^2 = \frac{1}{n} \sum_{i=1}^{n} \langle X_i X_i^T, AA^T \rangle.$$

To apply Theorem 3.1 to this empirical process, we need to bound

$$\sup_{g \in G} \left\| \|g(X_1)\|_2 \right\|_{\psi_1} = \sup_{A \in \mathcal{A}} \left\| \|A^T X_1\|_2 \right\|_{\psi_1} \le \sup_{A \in \mathcal{A}} \left\| \|A^T X_1\|_2 \right\|_{\psi_2}$$
(10)

and

$$d(g_1, g_2) = \left\| \left\| g(X_1) - g_2(X_1) \right\|_2 \right\|_{\psi_2} = \left\| \left\| (A_1 - A_2)^T X_1 \right\|_2 \right\|_{\psi_2}.$$
 (11)

The next lemma allows us to relate the ψ_2 -norm on G to the Frobenius norm on \mathcal{A} .

Lemma 4.1. Let Z be a p-variate random vector and A be a $p \times k$ matrix. Then

$$\left\| \|A^{T}Z\|_{2} \right\|_{\psi_{2}} \leq \|A\|_{F} \|Z\|_{\psi_{2}},$$
(12)

where $||Z||_{\psi_2} \coloneqq \sup_{||u||_2 \leq 1} ||\langle Z, u \rangle||_{\psi_2}$.

Proof. Let a_1, \ldots, a_k denote the columns of A. Then

$$\left\| \|A^T Z\|_2^2 \right\|_{\psi_1} \le \sum_{i=1}^k \left\| |\langle a_i, Z \rangle|^2 \right\|_{\psi_1} = \sum_{i=1}^k \left\| \langle a_i, Z \rangle \right\|_{\psi_2}^2 \le \sum_{i=1}^k \|a_i\|_2^2 \|Z\|_{\psi_2}^2.$$

Thus,

$$\left\| \|A^T Z\|_2 \right\|_{\psi_2}^2 = \left\| \|A^T Z\|_2^2 \right\|_{\psi_1} \le \|A\|_F^2 \|Z\|_{\psi_2}^2 \,.$$

Applying Lemma 4.1 to eqs. (10) and (11) yields

$$\sup_{g \in G} \left\| \|g(X_1)\|_2 \right\|_{\psi_1} \le \sup_{A \in \mathcal{A}} \|X_1\|_{\psi_2} \|A\|_F$$

and

$$d(g_1, g_2) \le ||X_1||_{\psi_2} ||A_1 - A_2||_F.$$

Since $||A||_F = ||vec(A)||_2$, the majorizing measure theorem (Talagrand 2005, Theorem 2.1.1) implies that

$$\gamma_2(G,d) \le c \|X_1\|_{\psi_2} \mathbb{E} \sup_{A \in \mathcal{A}} \langle A, \mathcal{Z} \rangle,$$

where \mathcal{Z} is a $p \times k$ matrix with i.i.d. standard Gaussian entries and c > 0 is an absolute constant. Thus, we have the following corollary of Theorem 3.1. **Corollary 4.1.** Let $X_1, \ldots, X_n \in \mathbb{R}^p$ be i.i.d. mean 0 random vectors, $\Sigma = \mathbb{E}X_1 X_1^T$, and $\sigma = \sup_{\|u\|_2=1} \|\langle X_1, u \rangle\|_{\psi_2}$, and

$$S_n = \frac{1}{n} \sum_{i=1}^n X_i X_i^T$$

If $\mathcal{A} \subseteq \mathbb{R}^{p \times k}$ is symmetric, then there exist positive constants c_1 , c_2 , and c_3 such that for all $t \ge c_1$, with probability at least $1 - 2 \exp(-c_2 t^{2/5})$,

$$\sup_{A \in \mathcal{A}} |\langle S_n - \Sigma, AA^T \rangle| \le c_3 t \left\{ \frac{\sigma^2 \sup_{A \in \mathcal{A}} ||A||_F}{\sqrt{n}} \left(\mathbb{E} \sup_{A \in \mathcal{A}} \langle \mathcal{Z}, A \rangle \right) + \frac{\sigma^2}{n} \left(\mathbb{E} \sup_{A \in \mathcal{A}} \langle \mathcal{Z}, A \rangle \right)^2 \right\},$$

where Z is a $p \times k$ matrix with i.i.d. standard Gaussian entries. Furthermore, we have under the same condition, that

$$\mathbb{E} \sup_{A \in \mathcal{A}} |\langle S_n - \Sigma, AA^T \rangle| \le c \left\{ \frac{\sigma^2 \sup_{A \in \mathcal{A}} ||A||_F}{\sqrt{n}} \left(\mathbb{E} \sup_{A \in \mathcal{A}} \langle \mathcal{Z}, A \rangle \right) + \frac{\sigma^2}{n} \left(\mathbb{E} \sup_{A \in \mathcal{A}} \langle \mathcal{Z}, A \rangle \right)^2 \right\},\$$

with a universal constant c.

References

Mendelson, S. (2010). "Empirical Processes with a Bounded ψ_1 Diameter". In: Geometric and Functional Analysis 20.4, pp. 988–1027.

Talagrand, M. (2005). The Generic Chaining. Springer-Verlag.