

FPCB : a simple and swift strategy for mirror repeat identification

Bhardwaj Vikash*¹, Gupta Swapnil², Meena Sitaram and Sharma Kulbhusan*²

¹*Government College Sector 14 Gurgaon, Haryana, India.*

²*Division of Radiation Biosciences, Institute of Nuclear Medicine and Allied Sciences, Delhi -110054, India.*

Running Title: Three step strategy for mirror repeat identification.

*Corresponding authors:

1. Kulbhusan Sharma, Metabolic Cell Signaling Group

Division of Radiation Biosciences,

Institute of Nuclear Medicine and Allied Sciences, Delhi -110054, India

Phone: 91-11-23905186; Fax: 91-11-23919509

E-mail: kulsinmas@gmail.com

2. Vikash Bhardwaj, Government College Sector 14 Gurgaon, Haryana, India.

Email: vikashbhardwaj@ gmail.com.

ABSTRACT

After the recent advancement of sequencing strategies, mirror repeats have been found to be present in the gene sequence of many organisms and species. This presence of mirror repeats in most of the sequences indicates towards some important functional role of these repeats. However, a simple and quick strategy to search these repeats in a given sequence is not available. We in this manuscript have proposed a simple and swift strategy named as FPCB strategy to identify mirror repeats in a give sequence. The strategy includes three simple steps of downloading sequencing in FASTA format (F), making its parallel complement (PC) and finally performing a homology search with the original sequence (B). At least twenty genes were analyzed using the proposed study. A number and types of mirror repeats were observed. We have also tried to give

nomenclature to these repeats. We hope that the proposed FPCB strategy will be quite helpful for the identification of mirror repeats in DNA or mRNA sequence. Also the strategy may help in unravelling the functional role of mirror repeats in various processes including evolution.

INTRODUCTION

After completion of DNA sequencing, many different types of sequences able to have a regulatory role have been discovered. Among these sequences mirror repeats have been predicted to play an important roles like adoption of perfect or near-perfect homopurine or homopyrimidine mirror repeats into triple-helical H conformations (Spano *et al*, 2007) . A mirror repeat is for example 5'AGTTCATTACTTGA3'where the sequences AGTTCAT and TACTTGA are mirror repeats of each other (Fig 1). The repeats may or may not be separated by a spacer nucleotide sequences in between them. There are several computer programs available to develop to detect repeats and/or the associated secondary structure in DNA or RNA sequences. However, most of these programs have not been able to effectively find out mirror repeats.

The purpose of this work is to devise a simple strategy to find out a mirror repeat in a sequence. We have recently shown through carious techniques that parallel DNA synthesis is quite possible in PCR reaction when we introduced a parallel primer in the reaction (Bhardwaj *et al*, 2013). The interesting outcome in this reaction is that the final product resulting from parallel DNA PCR (PD-PCR) is the original template DNA in reverse orientation so that the initial 5' end becomes 3' end. We have shown in this work that if we align parallel complement of a nucleotide sequence to original nucleotide sequence and perform a blast analysis, mirror repeats present in original nucleotide sequence can be detected easily. We have analyzed twenty genes to validate our strategy. During this, we found the presence of various types of repeats in nature. We have tried to give nomenclature to these mirror repeats.

One of the problems in counting repeats is the fact that a single repeat can be counted many times if one does not define in some way a maximal repeat (Spano *et al*, 2007). When one considers mirror repeats, the definition of maximal repeat is less clear and must be clearly defined. We have also tried to define the same in this work. We are interested in finding the numbers and types of mirror repeats in model genome sequences. Interesting, no novel software is required for the proposed strategy. We just

have to follow three simple steps for the identification of mirror repeats in a given sequence.

We hope that this simple strategy will be helpful in identifying mirror repeats in a sequence. Also, the strategy will be helpful in understanding the role of these mirror repeats in a sequence and will also assist in understanding the events of evolution in terms of mirror repeats.




Figure 1: Mirror repeats : DNA mirror repeat (MR) is a sequence segment delimited on the basis of its containing a centre of symmetry on a single strand and identical terminal nucleotides. Picture in left depicts the mirror image of an animal whereas picture on right depicts an example of a nucleotide sequence mirror repeat.

MATERIALS and METHODS

1. Downloading sequence in FASTA format

The coding sequences for the gene of interest were downloaded in FASTA format using the following link: <http://www.ncbi.nlm.nih.gov>.

2. Making a Parallel complement

Once downloaded, the FASTA format of nucleotide sequences were pasted into Reverse Complement program (http://www.bioinformatics.org/sms/rev_comp.html) and DNA sequences were converted into its complement counterpart.

3. Mirror repeat search

Both FASTA format of Nucleotide sequence and its parallel complement were aligned for BLAST homology search using BLAST tool: (http://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=MegaBlast&PROGRAM=blastn&BLAST_PROGRAMS=megaBlast&PAGE_TYPE=BlastSearch&BLAST_SPEC=blast2seq&QUERY=&S

UBJECTS=). The program was optimized for somewhat similar sequences (blastn) which allows a word size down to seven bases. If the position number is exactly reversed in subject and query , it will be a mirror repeat.

4. Gene analysis and Nomenclature of mirror repeats

Twenty important mammalian genes were analysed using the proposed study. The nomenclature was proposed for different types of mirror repeats observed during the analysis.

RESULTS

A simple strategy named **FPCB (FASTA-parallel complement-BLAST)** was devised to find out mirror repeats in various human mRNAs. The coding sequences for the gene of interest were downloaded in FASTA format using <http://www.ncbi.nlm.nih.gov>. In order to obtain the reverse complement sequence, the sequence in FASTA format was pasted into the following program (http://www.bioinformatics.org/sms/rev_comp.html). The final output of reverse complement program was selected for homology search with the parental sequence. Complement sequence was aligned with original sequence was submitted for BLAST analysis. The FPCB strategy has been depicted in Figure 2.




Figure 2: FPCB Strategy to find out mirror repeats. The strategy is based on three simple steps reflected in the name assigned FPCB. F stands for the first step of

downloading gene coding sequence in FASTA format. PC stands for formation of parallel complement of the gene of interest. The third and final step of the strategy is BLAST alignment analysis depicted by the word B.

The strategy was applied for twenty genes. Various genes selected for the FPCB analysis has been shown in table 1. During this analysis, a number and types of mirror repeats were observed. We have proposed the nomenclature of these mirror repeats (Fig.3). A typical mirror repeat has been shown in Fig.3 (i). Depending on the number of spacers at centre of symmetry of the mirror repeats, these were named as single spacer mirror repeats (SSMR), double spacer mirror repeats (DSMR) and multi spacer mirror repeats (MSMR) (Fig.3, ii, iii and iv respectively). Tandem mirror repeats and continuous di-mirror repeats were also observed (Fig.3, v and vi respectively). In some of the genes, few rare and interesting mirror repeats were observed. These were Continuous overlapping mirror repeats (COMR) and mirror repeats with simple tandem repeats (MRSMR).




Figure 3: Nomenclature and analysis of various types of mirror repeats observed after the use of FPCB Strategy. At least eight different types of mirror repeats were obtained after FPCB analysis in different genes. The strategy is based on three simple steps reflected in the name assigned FPCB.

Various important human genes were arbitrarily selected. Some of these genes included p53, BRCA 1, COX-2, NFkB, TGF beta etc. (Table 1). We included only mammalian genes in this study. However, future studies revealing mirror repeats in another organisms using our proposed FPCB strategy may be of interest and may also reveal various novel types of mirror repeats.

S.No.	Gene/CDS	Sequence Reference	No of MRs	Mirror repeats
1	P53	GenBank: AB082923.1	3	1. CCAAAGAAGAAACC 2. GGAACTCAAGG 3. ACCTGAAGTCCA
2	EIF2A	NCBI Reference Sequence NM_032025.3	2	1. ACAATTTTAACA 2. TTAACAACAATT These two mirror repeats are part of one continuous sequence ACAATTTTAACAACAATT
3	STAT3	GenBank: AJ012463.1	3	1. AGATCGGCTAGA 2. TCACTTTCCT 3. CTATCTCTATC
4	TNF-alpha	NCBI Reference Sequence X02910.1	3	1. CCTCATCTACTCC 2. CCAGAGGGAGACC 3. CCTCATCTACTCC No 2 and No 3 Mirror repeats are part of one continuous sequence CCAGAGGGAGACCCAGAGGG
5	Tgf-beta	GenBank: M60316.1	5	1. TGTGAGGGGGAGTGT 2. CGCCCGCGCCCGC 3. GCTACCACCATCG 4. CCGACTTCAGCC 5. GTCCGGGCCTG
6	Human protein kinase C theta	GenBank: L07032.1	2	1. ACATGTTTTGTACA 2. GACCTTCCAG

7	CCR5) mRNA	GenBank: U54994.1	1	1. GAGAAG AAGAG
8	AKT1	NCBI Reference Sequence NM_005163.2	2	1. AGGAGG AGGAGGA 2. GAGGAG GAGGAG These two mirror repeats are part of one sequence AGGAGGAGGAGGAG, It contains AGG Tandem repeat and GAG Tandem repeat.
9	FOXP2	NCBI Reference Sequence NG_007491.2	2	1. CTCTCA CACTCTC 2. CCCAGG GACCC
10	MTOR	NCBI Reference Sequence NM_004958.3	4	1. CGCCACC ACCGC 2. ACCTTCT TCTTCCA 3. ACTACAA ACATCA 4. AGAAGA AGAAGA
11	Wnt7a	GenBank: U53476.1	1	1. GCTACGG CA T CG
12	Ubiquitin	GenBank: M26880.1	3	1. CGAGGT G GGAGC Same sequence is repeated three times
13	BRCA1	GenBank: L78833.1	5	1. AAGAGA A AGAA A AGAA AGAGAA 2. AAATGA AC AGACA AGTAAA 3. TGGATT CA AACT TAGGT 4. AAAGATA A ATAGAAA 5. AAACCG T GCCAAA
14	NF-kappa-B	GenBank: BC051765.	6	1. CTGGGAG A GGGTC 2. AAAGTT A TTGAAA 3. AAGAACA A AGAA 4. GGAGG C GGAGG 5. AACGT A TGCAA 6. GACAGT G ACAG
15	COX2	GenBank: AY462100.	3	1. CGCGGT C CTGGCGC 2. ACAAC T ATCAACA 3. TGCAAT A ACGT

16	PTEN	GenBank: U93051.1	5	<ol style="list-style-type: none"> 1. ATGATGTAGTA 2. AGGTTTTTGGA 3. AAATTTTAAA 4. TTCATGTACTT 5. TTATAGATATT <p>First two mirror repeats are part of one continuous strand</p> <p>ATGATGTAGTAAGGTTTTTGGA</p>
17	HIF1A	NCBI Reference Sequence: NM_001530.3	7	<ol style="list-style-type: none"> 1. AAAACAGTGACAAAA 2. TTCAGCACGACTT 3. GAAGACACAGAAG 4. GAAAAGAAAAG 5. CAGATTTAGAC 6. CCCTATATCCC 7. ACATTATTACA
18	PCK1	NCBI Reference Sequence: NM_002591.3	2	<ol style="list-style-type: none"> 1. GGTCACAACTGG 2. TGGCTTTTTCGGT
19	Ceruloplasmin	GenBank: M13699.1	2	<ol style="list-style-type: none"> 1. TCCTGGGTCCT 2. TAGTTTTTGAT
20	Caspase 8	GenBank: BC068050.	2	<ol style="list-style-type: none"> 1. GTCTCACTCTG 2. CCTCCGCCTCC

Table 1 Various types of mirror repeats observed in various Human mRNA sequence using FPCB strategy. The sequence shown in red is mirror image of sequence shown in blue.

The exact role of mirror repeats has not been understood up till date. However, as most of the genes analyzed using FPCB were found to have the presence of these repeats, a major contribution of these repeats even in the evolutionary process may be possible. Future studies regarding the detailed understanding of the functional relevance of the mirror repeats may reveal some interesting findings. We hope that our proposed FPCB strategy will be helping in these

References :

1. Bhardwaj et al, 2013, <http://arxiv.org/abs/1309.3658>
2. Spano et al, 2007; <http://arxiv.org/abs/0705.2143v1>.
3. <http://www.ncbi.nlm.nih.gov>.
4. http://www.bioinformatics.org/sms/rev_comp.html.
5. http://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=MegaBlast&PROGRAM=blastn&BLAST_PROGRAMS=megaBlast&PAGE_TYPE=BlastSearch&BLAST_SPEC=blast2seq&QUERY=&SUBJECTS=

Supplementary Data

1) Homo sapiens mRNA for P53, complete cds

GenBank: AB082923.1

GenBank Graphics

```
>gi|23491728:64-1245 Homo sapiens mRNA for P53, complete cds
ATGGAGGAGCCGCGAGTCAGATCCTAGCGTCGAGCCCCCTCTGAGTCAGGAAACATTTTCAGACCTATGGA
AACTACTTCTGAAAACAACGTTCTGTCCCCCTTGCCGTCCCAAGCAATGGATGATTTGATGCTGTCCCC
GGACGATATTGAACAATGGTTCACTGAAGACCCAGGTCCAGATGAAGCTCCAGAAATGCCAGAGGCTGCT
CCCCGCGTGGCCCCCTGCACCAGCAGCTCCTACACCGGCGGCCCTGCACCAGCCCCCTCTGGCCCCCTGT
CATCTTCTGTCCCTTCCCAGAAAACCTACCAGGGCAGCTACGGTTTCCGTCTGGGCTTCTTGCAATTCTGG
GACAGCCAAGTCTGTGACTTGCACGTACTCCCCTGCCCTCAACAAGATGTTTTGCCAACTGGCCAAGACC
TGCCCTGTGCAGCTGTGGGTTGATTCCACACCCCCGCCCCGACCCGCGTCCGCGCCATGGCCATCTACA
AGCAGTCACAGCACATGACGGAGGTTGTGAGGCGCTGCCCCACCATGAGCGCTGCTCAGATAGCGATGG
TCTGGCCCCCTCTCAGCATCTTATCCGAGTGGAAGGAAATTTGCGTGTGGAGTATTTGGATGACAGAAAC
ACTTTTCGACATAGTGTGGTGGTGCCCTATGAGCCGCTGAGGTTGGCTCTGACTGTACCACCATCCACT
ACAACACTACATGTGTAACAGTTTCTGCATGGGCGGCATGAACCGGAGGCCATCCTCACCATCATCACACT
GGAAGACTCCAGTGGAATCTACTGGGACGGAACAGCTTTGAGGTGCATGTTTGTGCCCTGTCTGGGAGA
GACCGGCGCACAGAGGAAGAGAATCTCCGCAAGAAAGGGGAGCCTCACCACGAGCTGCCCCAGGGAGCA
CTAAGCGAGCACTGTCCAACAACACCAGCTCCTCTCCCCAGCCAAAGAAGAAACCCTGGATGGAGAATA
TTTACCCCTTCAGATCCGTGGGCGTGAGCGCTTCGAGATGTTCCGAGAGCTGAATGAGGCCTTGGAACTC
AAGGATGCCCAGGCTGGGAAGGAGCCAGGGGGAGCAGGGCTCACTCCAGCCACCTGAAGTCCA AAAAGG
GTCAGTCTACCTCCCGCCATAAAAACTCATGTTCAAGACAGAAGGGCCTGACTCAGACTGA
```

Parallel complement Homo sapiens mRNA for P53

```
TACCTCCTCGGCGTCAGTCTAGGATCGCAGCTCGGGGGAGACTCAGTCCTTTGTAAAAGTCTGGATA
CCTTTGATGAAGGACTTTTGTGCAAGACAGGGGGAACGGCAGGGTTCGTTACCTACTAACTACG
ACAGGGGCCTGCTATAACTTGTACCAAGTGACTTCTGGGTCCAGGTCTACTTCGAGGGTCTTACGG
```

TCTCCGACGAGGGGCGCACCGGGGACGTGGTCGTCGAGGATGTGGCCGCCGGGGACGTGGTTCGG
GGGAGGACCGGGGACAGTAGAAGACAGGGAAGGGTCTTTTGGATGGTCCCGTCGATGCCAAAGGC
AGACCCGAAGAACGTAAGACCCTGTCCGGTTCAGACACTGAACGTGCATGAGGGGACGGGAGTTGTT
CTACAAAACGGTTGACCGGTTCTGGACGGGACACGTGACACCCAACTAAGGTGTGGGGGCGGGC
CGTGGGCGCAGGCGCGGTACCGGTAGATGTTTCGTCAGTGTGTCGTACTGCCTCCAACACTCCGCGA
CGGGGGTGGTACTCGCGACGAGTCTATCGCTACCAGACCGGGGAGGAGTCGTAGAATAGGCTCAC
CTTCCTTTAAACGCACACCTCATAAACCTACTGTCTTTGTGAAAAGCTGTATCACACCACCACGGGAT
ACTCGGCGGACTCCAACCGAGACTGACATGGTGGTAGGTGATGTTGATGTACACATTGTCAAGGAC
GTACCCGCCGTACTTGGCCTCCGGGTAGGAGTGGTAGTAGTGACCTTCTGAGGTCACCATTAGAT
GACCCTGCCTTGTGAAACTCCACGTACAAACACGGACAGGACCCTCTCTGGCCGCGTGTCTCCTTCT
CTTAGAGGCGTTCTTTCCCTCGGAGTGGTGCTCGACGGGGTCCCTCGTGATTGCTCGTGACAGG
TTGTTGGTTCGAGGAGAGGGGTCGGTTTCTTCTTTGGTGACCTACCTCTTATAAAGTGGGAAGTCT
AGGCACCCGCACTCGCGAAGCTCTACAAGGCTCTCGACTTACTCCGGAACCTTGAGTTCCTACGGGT
CCGACCCTTCTCGGTCCCCCTCGTCCCGAGTGAGGTCGGTGGACTTCAGGTTTTTCCAGTCAGAT
GGAGGGCGGTATTTTTTGGAGTACAAGTCTGTCTTCCCGACTGAGTCTGACT

Alignment statistics for match #1

Score	Expect	Identities	Gaps	Strand
26.5 bits(28)	0.014	14/14(100%)	0/14(0%)	Plus/Minus

Query	952	CCAAAGAAGAAACC	965
Sbjct	965	CCAAAGAAGAAACC	952

Range 2: 1103 to 1114 [Graphics](#) [Next Match](#) [Previous Match](#) [First Match](#)

Alignment statistics for match #2

Score	Expect	Identities	Gaps	Strand
22.9 bits(24)	0.17	12/12(100%)	0/12(0%)	Plus/Minus

Query	1103	ACCTGAAGTCCA	1114
--------------	-------------	---------------------	-------------

| | | | | | | | | |

Sbjct 1114 ACCTGAAGTCCA 1103

Range 3: 1044 to 1054 [Graphics](#) [Next Match](#) [Previous Match](#) [First Match](#)

Alignment statistics for match #3

Score	Expect	Identities	Gaps	Strand
21.1 bits(22)	0.60	11/11(100%)	0/11(0%)	Plus/Minus

Query 1044 GGAACTCAAGG 1054

| | | | | | | | | |

Sbjct 1054 GGAACTCAAGG 1044

2) Homo sapiens eukaryotic translation initiation factor 2A, 65kDa (EIF2A), mRNA

NCBI Reference Sequence: NM_032025.3

[GenBank](#) [Graphics](#)

>gi|83656780:17-1774 Homo sapiens eukaryotic translation initiation factor 2A, 65kDa (EIF2A), mRNA

```

ATGGCGCCGTCCACGCCGCTCTTGACAGTCCGAGGATCAGAAGGACTGTACATGGTGAATGGACCACCAC
ATTTTACAGAAAGCACAGTGTTCCTCAAGGGAATCTGGGAAGAATTGCAAAGTCTGTATCTTTAGTAAGGA
TGGGACCTTGTTCCTGGGGCAATGGAGAAAAAGTAAATATTATCAGTGTCACTAACAAGGGACTACTG
CACTCCTTCGACCTCCTGAAGGCAGTTTGCCTTGAATTCTCACCCAAAAATACTGTCCCTGGCAACGTGGC
AGCCTTACACTACTTCTAAAGATGGCACAGCTGGGATACCCAACCTACAACCTTTATGATGTGAAAACTGG
GACATGTTTGAAATCTTTTCATCCAGAAAAAATGCAAAATTTGGTGTCCATCCTGGTCAGAAGATGAAACT
CTTTGTGCCCCGAATGTAAACAATGAAGTTCACCTTCTTTGAAAAACAACAATTTTAACACAATTGCAAATA
AATTGCATTTGCAAAAAATTAATGATTTTGTATTATCACCTGGACCCCAACCATACAAGGTGGCTGTCTA
TGTTCCAGGAAGTAAAGGTGCACCTTCATTTGTTAGATTATATCAGTACCCCAACTTTGCTGGACCTCAT
GCAGCTTTAGCTAATAAAAAGTTTCTTTAAGGCAGATAAAGTTACAATGCTGTGGAATAAAAAAGCTACTG
CTGTGTTGGTAATAGCTAGCACAGATGTTGACAAGACAGGAGCTTCCTACTATGGAGAACAAACTCTACA
CTACATTGCAACAAATGGAGAAAGTGCTGTAGTGCAATTACCAAAAAATGGCCCCATTTATGATGTAGTT
TGGAATTCTAGTTCTACTGAGTTTTTGTGCTGTATATGGTTTTATGCCTGCCAAAGCGACAATTTTCAACT
TGAAATGTGATCCTGTATTTGACTTTGGAAGTGGTCCTCGTAATGCAGCCTACTATAGCCCTCATGGACA
TATATTAGTATTAGCTGGATTTGGAAATCTGAGGGGACAAATGGAAGTGTGGGATGTGAAAACTACAAA

```

CTTATTTCTAAACCGGTGGCTTCTGATTCTACATATTTTGCTTGGTGCCCGGATGGTGAGCATATTTTAA
CAGCTACATGTGCTCCCAGGTTACGGGTTAATAATGGATACAAAATTTGGCATTATACTGGCTCTATCTT
GCACAAGTATGATGTGCCATCAAATGCAGAATTATGGCAGGTTTCTTGGCAGCCATTTTGGATGGAATA
TTTCCAGCAAAAACAATAACTTACCAAGCAGTTCCAAGTGAAGTACCCAATGAGGAACCTAAAGTTGCAA
CAGCTTATAGACCCCCAGCTTTAAGAAATAAACCAATCACCAATTCCAAATTCATGAAGAGGAACCACC
TCAGAATATGAAACCACAATCAGGAAACGATAAGCCATTATCAAAAACAGCTCTTAAAAATCAAAGGAAG
CATGAAGCTAAGAAAGCTGCAAAGCAGGAAGCAAGAAGTGACAAGAGTCCAGATTTGGCACCTACTCCTG
CCCCACAGAGCACACCACGAAACACTGTCTCTCAGTCAATTTCTGGGGACCCTGAGATAGACAAAAAAT
CAAGAACCTAAAGAAGAAAAGTAAAAGCAATCGAAACAACTGAAAAGAACAAAGCAGCAACTGGAAAACAGCTA
GAAAAAATCAGTTGGAGAAAATTCAGAAAGAAACAGCCCTTCTCCAGGAGCTGGAAGATTTGGAATTGG
GTATTTAA

Parallel Complement

TACCGCGGCAGGTGCGGCGAGAAGTGTGAGGCTCCTAGTCTTCTGACATGTACCACTTACCTGGTGGTGTA
AATGTCTTTTCGTGTCAAAAGGTTCCCTTAGACCCTTCTAACGTTTCAGACATAGAAATCATTCTACCTGGA
ACAAACGGACCCCGTTACCTCTTTTTCATTTATAATAGTCACAGTGATTGTTCCCTGATGACGTGAGGAAGCTG
GAGGACTTCCGTCAAACGGAACCTAAGAGTGGGTTTTATGACAGGACCGTTGCACCGTCGGAATGTGATGA
AGATTTCTACCGTGTGACCCCTATGGGTTGGATGTTGAAATACTACACTTTTGACCCTGTACAACTTTAGAAA
GTAGGTCTTTTTTACGTTTTAACCACAGGTAGGACCAGTCTTCTACTTTGAGAAACACGGGCGTTACAATTGT
TACTTCAAGTGAAGAACTTTTGTGTTAAATTTGTGTTAACGTTTATTTAACGTAAACGTTTTTTAATTACTAA
AACATAATAGTGGACCTGGGGTTGGTATGTTCCACCGACAGATACAAGGTCCTTCATTTCCACGTGGAAGTAA
ACAATCTAATATAGTCATGGGTTGAAACGACCTGGAGTACGTGAAATCGATTATTTCAAAGAAATTCGT
CTATTTCAATGTTACGACACCTATTTTTTCGATGACGACACAACCATTATCGATCGTGTCTACAACCTGTTCTGT
CCTCGAAGGATGATACCTCTGTTTGAGATGTGATGTAACGTTGTTTACCTCTTTCACGACATCACGTTAATGG
TTTTTACCAGGGTAAATACTACATCAAACCTAAGATCAAGATGACTCAAACACGACATATAACAAAATACG
GACGGTTTCGCTGTTAAAAGTTGAACTTTACACTAGGACATAAACTGAAACCTTGACCAGGAGCATTACGTGCG
GATGATATCGGGAGTACCTGTATATAATCATAATCGACCTAAACCTTTAGACTCCCCTGTTTACCTTCACACCCT
ACACTTTTTGATGTTTGAATAAAGATTTGGCCACCGAAGACTAAGATGTATAAAACGAACCACGGGCCTACCA
CTCGTATAAAATTTGTCGATGTACACGAGGGTCCAATGCCAATTATTACCTATGTTTTAAACCGTAATATGACC
GAGATAGAACGTGTTACTACTACACGGTAGTTTACGTCTTAATACCGTCCAAAGAACCGTCGGTAAAAACCTA
CCTATAAAGGTCGTTTTTGTATTGAATGGTTCGTCAAGGTTCACTTCATGGGTTACTCCTTGGATTTCAACGT
TGTCGAATATCTGGGGTTCGAAATCTTTATTTGGTTAGTGGTTAAGGTTAACGTACTTCTCCTTGGTGGAGT
CTTATACTTTGGTGTAGTCTTTGCTATTCGGAATAGTTTTTGTGAGAATTTTTAGTTTCTTCTGACTTCGA
TTCTTTGACGTTTTGTCCTTCGTTCTTCACTGTTCTCAGGTCTAAACCGTGGATGAGGACGGGGTGTCTCGTG
TGGTGCTTTGTGACAGAGAGTCAGTTAAAGACCCCTGGGACTCTATCTGTTTTTTAGTTCTTGGATTTCTTCTT
TGACTTTGCTTAGCTTGTGACTTTCTGTTGTCGTTGACCTTTTGTGATCTTTTTTTAGTCAACCTCTTTAAG
TCTTTCTTGTGCGGAAGAGGTCCTCGACCTTCTAAACCTTAACCCATAAAT

Range 1: 467 to 478 [Graphics](#) [Next Match](#) [Previous Match](#)

Alignment statistics for match #1

Score Expect Identities Gaps Strand
22.9 bits(24) 0.38 12/12(100%) 0/12(0%) Plus/Minus

Query 467 ACAATTTTAACA 478
|||||||
Sbjct 478 ACAATTTTAACA 467

Range 2: 454 to 467[Graphics](#)[Next Match](#)[Previous Match](#)[First Match](#)

Alignment statistics for match #2

Score Expect Identities Gaps Strand
21.1 bits(22) 1.3 13/14(93%) 0/14(0%) Plus/Minus

Query 643 AATAAAAGTTTCTT 656
|| |||||
Sbjct 467 AACAAAAGTTTCTT 454

Range 3: 473 to 483[Graphics](#)[Next Match](#)[Previous Match](#)[First Match](#)

Score Expect Identities Gaps Strand
21.1 bits(22) 1.3 11/11(100%) 0/11(0%) Plus/Minus

Query 473 TTAACACAATT 483
|||||||
Sbjct 483 TTAACACAATT 473

Range 4: 643 to 656[Graphics](#)[Next Match](#)[Previous Match](#)[First Match](#)

Score Expect Identities Gaps Strand

21.1 bits(22) 1.3 13/14(93%) 0/14(0%) Plus/Minus

Query	454	TTCTTTGAAAACAA	467
Sbjct	656	TTCTTTGAAAATAA	643