METRIC MEASURE SPACES SUPPORTING GAGLIARDO-NIRENBERG INEQUALITIES: VOLUME NON-COLLAPSING AND RIGIDITIES

ALEXANDRU KRISTÁLY

ABSTRACT. Let $(M, \mathsf{d}, \mathsf{m})$ be a metric measure space which satisfies the Lott-Sturm-Villani curvature-dimension condition $\mathsf{CD}(K, n)$ for some $K \ge 0$ and $n \ge 2$, and a lower n-density assumption at some point of M. We prove that if $(M, \mathsf{d}, \mathsf{m})$ supports the Gagliardo-Nirenberg inequality or any of its limit cases $(L^p$ -logarithmic Sobolev inequality or Faber-Krahn-type inequality), then a global non-collapsing n-dimensional volume growth holds, i.e., there exists a universal constant $C_0 > 0$ such that $\mathsf{m}(B_x(\rho)) \ge C_0 \rho^n$ for all $x \in M$ and $\rho \ge 0$, where $B_x(\rho) = \{y \in M : \mathsf{d}(x, y) < \rho\}$. Due to the quantitative character of the volume growth estimate, we establish several rigidity results on Riemannian manifolds with non-negative Ricci curvature supporting Gagliardo-Nirenberg inequalities by exploring a quantitative Perelman-type homotopy construction developed by Munn (J. Geom. Anal., 2010). Further rigidity results are also presented on some reversible Finsler manifolds.

Contents

1. Introduction	1
1.1. Recalling optimal Gagliardo-Nirenberg inequalities on normed spaces	1
1.2. Statement of main results	3
2. Volume non-collapsing via Gagliardo-Nirenberg inequalities	6
2.1. Cases $\alpha > 1 \& 0 < \alpha < 1$: usual Gagliardo-Nirenberg inequalities	7
2.2. Limit case I ($\alpha \to 1$): L^p -logarithmic Sobolev inequality	12
2.3. Limit case II ($\alpha \to 0$): Faber-Krahn-type inequality	15
3. Rigidity results in smooth settings	16
3.1. Gagliardo-Nirenberg inequalities on Riemannian manifolds with Ricci ≥ 0	17
3.2. Gagliardo-Nirenberg inequalities on Finsler manifolds with $n-\text{Ricci} \ge 0$	18
References	21

1. INTRODUCTION

An important role in the theory of geometric functional inequalities is played by the Gagliardo-Nirenberg interpolation inequality and its limit cases. The present paper is devoted to the study of Gagliardo-Nirenberg inequalities on metric measure spaces; to be more precise, we shall

- (a) establish quantitative volume non-collapsing properties of metric measure spaces satisfying the Lott-Sturm-Villani curvature-dimension condition CD(K, n) for some $K \ge 0$ and $n \ge 2$, in the presence of a Gagliardo-Nirenberg inequality or one of its limit cases $(L^p-\text{logarithmic Sobolev inequality})$;
- (b) provide *rigidity* results in the framework of Riemannian and Finsler manifolds with nonnegative Ricci curvature which support (*almost*) optimal Gagliardo-Nirenberg inequalities by using the volume non-collapsing property from (a) and a quantitative homotopy construction due to Munn [17] and Perelman [22].

In §1.1 we recall the optimal Gagliardo-Nirenberg inequalities on normed spaces which play a comparison role in our investigations; in §1.2 we present the main results of the paper.

1.1. Recalling optimal Gagliardo-Nirenberg inequalities on normed spaces. The optimal Gagliardo-Nirenberg inequality in the Euclidean case has been obtained by Del Pino and Dolbeault [7] for a certain range of parameters by using symmetrization arguments. By using mass transportation argument, Cordero-Erausquin, Nazaret and Villani [6] extended the results from [7] to prove optimal Gagliardo-Nirenberg inequalities on arbitrary normed spaces. In the sequel, we recall the main theorems from [6] and some related results.

²⁰⁰⁰ Mathematics Subject Classification. Primary 53C23; Secondary 35R06, 53C60.

Key words and phrases. Gagliardo-Nirenberg inequality, L^p -logarithmic Sobolev inequality, Faber-Krahn-type inequality, metric measure space, CD(K, n) condition, volume non-collapsing, rigidity.

ALEXANDRU KRISTÁLY

Let $\|\cdot\|$ be an arbitrary norm on \mathbb{R}^n ; without loss of generality, we may assume that the Lebesgue measure of the unit ball in $(\mathbb{R}^n, \|\cdot\|)$ is the volume of the *n*-dimensional Euclidean unit ball $\omega_n = \pi^{\frac{n}{2}} \Gamma(\frac{n}{2}+1)^{-1}$. The dual norm $\|\cdot\|_*$ of $\|\cdot\|$ is given by $\|x\|_* = \sup_{\|y\|\leq 1} x \cdot y$ where '.' is the Euclidean inner product. Let $p \in [1, n)$ and $L^p(\mathbb{R}^n)$ be the Lebesgue space of order p. As usual, we consider the Sobolev spaces

$$\dot{W}^{1,p}(\mathbb{R}^n) = \{ u \in L^{p^*}(\mathbb{R}^n) : \nabla u \in L^p(\mathbb{R}^n) \}$$

and

$$W^{1,p}(\mathbb{R}^n) = \{ u \in L^p(\mathbb{R}^n) : \nabla u \in L^p(\mathbb{R}^n) \},\$$

where $p^* = \frac{pn}{n-p}$ and ∇ is the gradient operator. On account of the Finslerian duality (see also §3.2), if $u \in \dot{W}^{1,p}(\mathbb{R}^n)$, the norm of ∇u is defined by

$$\|\nabla u\|_{L^p} = \left(\int_{\mathbb{R}^n} \|\nabla u(x)\|_*^p dx\right)^{1/p},$$

where dx is the Lebesgue measure on \mathbb{R}^n .

Fix $n \ge 2$, $p \in (1, n)$ and $\alpha \in (0, \frac{n}{n-p}] \setminus \{1\}$; for every $\lambda > 0$, let

$$h_{\alpha,p}^{\lambda}(x) = (\lambda + (\alpha - 1) \|x\|^{p'})_{+}^{\frac{1}{1-\alpha}}, \ x \in \mathbb{R}^{n}, ^{1}$$

where $p' = \frac{p}{p-1}$ is the conjugate to p, and $r_+ = \max\{0, r\}$ for $r \in \mathbb{R}$. The following optimal Gagliardo-Nirenberg inequalities are known on normed spaces:

Theorem A. [see [6, Theorem 4]] Let $n \ge 2$, $p \in (1, n)$ and $\|\cdot\|$ be an arbitrary norm on \mathbb{R}^n .

• If $1 < \alpha \leq \frac{n}{n-p}$, then

$$\|u\|_{L^{\alpha p}} \le \mathcal{G}_{\alpha,p,n} \|\nabla u\|_{L^{p}}^{\theta} \|u\|_{L^{\alpha(p-1)+1}}^{1-\theta}, \ \forall u \in \dot{W}^{1,p}(\mathbb{R}^{n}),$$
(1.1)

where

$$\theta = \frac{p^{\star}(\alpha - 1)}{\alpha p(p^{\star} - \alpha p + \alpha - 1)},\tag{1.2}$$

and the best constant

$$\mathcal{G}_{\alpha,p,n} = \left(\frac{\alpha - 1}{p'}\right)^{\theta} \frac{\left(\frac{p'}{n}\right)^{\frac{\theta}{p} + \frac{\theta}{n}} \left(\frac{\alpha(p-1)+1}{\alpha-1} - \frac{n}{p'}\right)^{\frac{1}{\alpha p}} \left(\frac{\alpha(p-1)+1}{\alpha-1}\right)^{\frac{\theta}{p} - \frac{1}{\alpha p}}}{\left(\omega_n \mathsf{B}\left(\frac{\alpha(p-1)+1}{\alpha-1} - \frac{n}{p'}, \frac{n}{p'}\right)\right)^{\frac{\theta}{n}}}$$

is achieved by the family of functions $h_{\alpha,p}^{\lambda}$, $\lambda > 0$;

• If $0 < \alpha < 1$, then

$$\|u\|_{L^{\alpha(p-1)+1}} \le \mathcal{N}_{\alpha,p,n} \|\nabla u\|_{L^p}^{\gamma} \|u\|_{L^{\alpha p}}^{1-\gamma}, \ \forall u \in \dot{W}^{1,p}(\mathbb{R}^n),$$
(1.3)

where

$$\gamma = \frac{p^{\star}(1-\alpha)}{(p^{\star}-\alpha p)(\alpha p+1-\alpha)},\tag{1.4}$$

and the best constant

$$\mathcal{N}_{\alpha,p,n} = \left(\frac{1-\alpha}{p'}\right)^{\gamma} \frac{\left(\frac{p'}{n}\right)^{\frac{\gamma}{p}+\frac{\gamma}{n}} \left(\frac{\alpha(p-1)+1}{1-\alpha} + \frac{n}{p'}\right)^{\frac{\gamma}{p}-\frac{1}{\alpha(p-1)+1}} \left(\frac{\alpha(p-1)+1}{1-\alpha}\right)^{\frac{1}{\alpha(p-1)+1}}}{\left(\omega_n \mathsf{B}\left(\frac{\alpha(p-1)+1}{1-\alpha}, \frac{n}{p'}\right)\right)^{\frac{\gamma}{n}}}$$

is achieved by the family of functions $h_{\alpha,p}^{\lambda}$, $\lambda > 0$. Hereafter, $\mathsf{B}(\cdot, \cdot)$ is the Euler beta-function.

¹The function $h_{\alpha,p}^{\lambda}$ is positive everywhere for $\alpha > 1$ while $h_{\alpha,p}^{\lambda}$ has always a compact support for $\alpha < 1$.

The borderline case $\alpha = \frac{n}{n-p}$ (thus $\theta = 1$) reduces to the *optimal Sobolev inequality*, see Aubin [3] and Talenti [26] in the Euclidean case, and Alvino, Ferone, Lions and Trombetti [1] for normed spaces. Furthermore, inequalities (1.1) and (1.3) degenerate to the *optimal L^p-logarithmic Sobolev inequality* whenever $\alpha \to 1$ (called also as the entropy-energy inequality involving the Shannon entropy), while (1.3) reduces to a *Faber-Krahn-type inequality* whenever $\alpha \to 0$, respectively. More precisely, one has

Theorem B. Let $n \ge 2$, $p \in (1, n)$ and $\|\cdot\|$ be an arbitrary norm on \mathbb{R}^n .

• Limit case I ($\alpha \rightarrow 1$) [see [9, Theorem 1.1]²]: One has

$$\mathbf{Ent}_{dx}(|u|^p) = \int_{\mathbb{R}^n} |u|^p \log |u|^p dx \le \frac{n}{p} \log \left(\mathcal{L}_{p,n} \|\nabla u\|_{L^p}^p\right), \ \forall u \in W^{1,p}(\mathbb{R}^n), \ \|u\|_{L^p} = 1,$$
(1.5)
where the best constant
$$n (n-1)^{p-1} (n-1)^{-\frac{p}{n}}$$

$$\mathcal{L}_{p,n} = \frac{p}{n} \left(\frac{p-1}{e} \right)^{p-1} \left(\omega_n \Gamma \left(\frac{n}{p'} + 1 \right) \right)^{-\frac{1}{2}}$$

is achieved by the family of functions

$$l_p^{\lambda}(x) = \lambda^{\frac{n}{pp'}} \omega_n^{-\frac{1}{p}} \Gamma\left(\frac{n}{p'} + 1\right)^{-\frac{1}{p}} e^{-\frac{\lambda}{p} \|x\|^{p'}}, \lambda > 0;$$

• Limit case II $(\alpha \rightarrow 0)$ [see [6, p. 320]]: One has

$$\|u\|_{L^{1}} \leq \mathcal{F}_{p,n} \|\nabla u\|_{L^{p}} |\mathrm{supp}(u)|^{1-\frac{1}{p^{*}}}, \ \forall u \in \dot{W}^{1,p}(\mathbb{R}^{n})$$
(1.6)

and the best constant

$$\mathcal{F}_{p,n} = \lim_{\alpha \to 0} \mathcal{N}_{\alpha,p,n} = n^{-\frac{1}{p}} \omega_n^{-\frac{1}{n}} (p'+n)^{-\frac{1}{p'}}$$

is achieved by the family of functions

$$f_p^{\lambda}(x) = \lim_{\alpha \to 0} h_{\alpha,p}^{\lambda}(x) = (\lambda - \|x\|^{p'})_+, \ x \in \mathbb{R}^n,$$

where supp(u) stands for the support of u and |supp(u)| is its Lebesgue measure.

1.2. Statement of main results. As we already pointed out, the primordial purpose of the present paper is to establish fine topological properties of metric measure spaces curved in the sense of Lott-Sturm-Villani which support Gagliardo-Nirenberg-type inequalities. In fact, the metric spaces we are working on are supposed to satisfy the curvature-dimension condition CD(K, n) for some $K \ge 0$ and $n \ge 2$, introduced by Lott and Villani [15] and Sturm [24, 25]; see §2 for its formal definition.

1.2.1. Volume non-collapsing on metric measure spaces. Let $(M, \mathsf{d}, \mathsf{m})$ be a metric measure space (with a strictly positive Borel measure m) and $\operatorname{Lip}_0(M)$ be the space of Lipschitz functions with compact support on M. For $u \in \operatorname{Lip}_0(M)$, let

$$|\nabla u|_{\mathsf{d}}(x) := \limsup_{y \to x} \frac{|u(y) - u(x)|}{\mathsf{d}(x, y)}, \ x \in M.$$

$$(1.7)$$

Note that $x \mapsto |\nabla u|_{\mathsf{d}}(x)$ is Borel measurable on M for $u \in \operatorname{Lip}_0(M)$.

As before, let $n \ge 2$ be an integer, $p \in (1, n)$ and $\alpha \in (0, \frac{n}{n-p}] \setminus \{1\}$. Throughout this section we assume that the lower *n*-density of the measure **m** at a point $x_0 \in M$ is unitary, i.e.,

$$(\mathbf{D})_{x_0}^n: \liminf_{\rho \to 0} \frac{\mathsf{m}(B_{x_0}(\rho))}{\omega_n \rho^n} = 1,$$

where $B_x(r) = \{ y \in M : d(x, y) < r \}.$

Throughout the whole paper, we shall keep the notations from Theorems A and B (i.e., the four best constants from the Gagliardo-Nirenberg inequalities on normed spaces and the numbers θ and γ from (1.2) and (1.4), respectively); the Lebesgue spaces L^p are defined on the measure space (M, m) . We now are the position to state our quantitative, globally non-collapsing volume growth results:

²Gentil [9] proved an optimal L^p -logarithmic Sobolev inequality for even, q-homogeneous (q > 1), strictly convex functions $C : \mathbb{R}^n \to [0, \infty)$. In our case, $C(x) = \frac{\|x\|^{p'}}{n'}$.

ALEXANDRU KRISTÁLY

Theorem 1.1. [Gagliardo-Nirenberg inequalities] Let $(M, \mathsf{d}, \mathsf{m})$ be a proper metric measure space which satisfies the curvature-dimension condition $\mathsf{CD}(K, n)$ for some $K \ge 0$ and $n \ge 2$. Let $p \in (1, n)$ and assume that $(\mathbf{D})_{x_0}^n$ holds for some $x_0 \in M$. Then the following statements hold:

(i) If $1 < \alpha \leq \frac{n}{n-p}$ and the inequality

$$\|u\|_{L^{\alpha p}} \le \mathcal{C}\||\nabla u|_{\mathsf{d}}\|_{L^{p}}^{\theta}\|u\|_{L^{\alpha(p-1)+1}}^{1-\theta}, \ \forall u \in \operatorname{Lip}_{0}(M)$$
(GN1) ^{α, p}

holds for some $\mathcal{C} \geq \mathcal{G}_{\alpha,p,n}$, then K = 0 and

$$\mathsf{m}(B_x(\rho)) \ge \left(\frac{\mathcal{G}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\theta}} \omega_n \rho^n \quad for \ all \ x \in M \ and \ \rho \ge 0.$$

(ii) If $0 < \alpha < 1$ and the inequality

$$\|u\|_{L^{\alpha(p-1)+1}} \le \mathcal{C}\||\nabla u|_{\mathsf{d}}\|_{L^{p}}^{\gamma}\|u\|_{L^{\alpha p}}^{1-\gamma}, \ \forall u \in \operatorname{Lip}_{0}(M)$$
(GN2)^{*a,p*}

holds for some $\mathcal{C} \geq \mathcal{N}_{\alpha,p,n}$, then K = 0 and

$$\mathsf{m}(B_x(\rho)) \ge \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}} \omega_n \rho^n \quad for \ all \ x \in M \ and \ \rho \ge 0$$

In the limit case $\alpha \to 1$, we can state

Theorem 1.2. $[L^p-\text{logarithmic Sobolev inequality}]$ Under the same assumptions as in Theorem 1.1, if

$$\operatorname{Ent}_{d\mathsf{m}}(|u|^p) = \int_M |u|^p \log |u|^p d\mathsf{m} \le \frac{n}{p} \log \left(\mathcal{C} \| |\nabla u|_{\mathsf{d}} \|_{L^p}^p \right), \ \forall u \in \operatorname{Lip}_0(M), \ \|u\|_{L^p} = 1 \qquad (\mathbf{LS})_{\mathcal{C}}^p$$
for some $\mathcal{C} \ge \mathcal{C}$, then $K = 0$ and

holds for some $C \geq \mathcal{L}_{p,n}$, then K = 0 and

$$\mathsf{m}(B_x(\rho)) \ge \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}} \omega_n \rho^n \quad for \ all \ x \in M \ and \ \rho \ge 0.$$

In the remaining limit case $\alpha \to 0$, one can prove

Theorem 1.3. [Faber-Krahn-type inequality] Under the same assumptions as in Theorem 1.1, if

$$\|u\|_{L^1} \le \mathcal{C} \||\nabla u|_{\mathsf{d}}\|_{L^p} \mathsf{m}(\operatorname{supp}(u))^{1-\frac{1}{p^\star}}, \ \forall u \in \operatorname{Lip}_0(M)$$
(**FK**)^{*p*}_{*C*}

holds for some $C \geq \mathcal{F}_{p,n}$, then K = 0 and

$$\mathsf{m}(B_x(\rho)) \ge \left(\frac{\mathcal{F}_{p,n}}{\mathcal{C}}\right)^n \omega_n \rho^n \text{ for all } x \in M \text{ and } \rho \ge 0.$$

Some remarks are in order.

Remark 1.1. (a) The proofs of Theorems 1.1-1.3 are synthetic where we shall exploit some basic features of metric measure spaces satisfying the CD(K, n) condition (such as generalized Bonnet-Myers and Bishop-Gromov comparison inequalities) and direct constructions. Although the lines of the proofs of these results are similar, our arguments require different technics, deeply depending on the shape of certain test functions whose profiles come from the family of extremals in normed spaces (cf. Theorems A & B). Note that instead of the CD(K, n) condition it is enough to consider the slightly weaker measure contraction property MCP(K, n), see Ohta [20].

(b) The case p = 2 and $\alpha = \frac{n}{n-2}$ $(n \ge 3)$ is contained in Kristály and Ohta [12], where the authors studied Caffarelli-Kohn-Nirenberg inequalities on metric measure spaces. We notice that the roots of Theorem 1.1 (i) on Riemannian manifolds with non-negative Ricci curvature can be found in do Carmo and Xia [8], Ledoux [13] and Xia [28].

(c) The generalized Bishop-Gromov inequality and density assumption $(\mathbf{D})_{x_0}^n$ imply $\mathsf{m}(B_{x_0}(\rho)) \leq \omega_n \rho^n$ for all $\rho \geq 0$. In particular, the latter inequality and the conclusions of Theorems 1.1-1.3 imply the Ahlfors n-regularity at the point x_0 ; therefore, the Hausdorff dimension of (M, d) is precisely n.

(d) $(\mathbf{D})_{x_0}^n$ clearly holds for every point x_0 on *n*-dimensional Riemannian and Finsler manifolds endowed with the canonical Busemann-Hausdorff measure.

1.2.2. Applications: rigidity results in smooth settings. Having fine volume growth estimates in Theorems 1.1-1.3, important rigidity results can be deduced in the context of Riemannian and Finsler manifolds supporting Gagliardo-Nirenberg-type inequalities.

In order to state such results, let (M, g) be an *n*-dimensional complete Riemannian manifold with non-negative Ricci curvature $(n \ge 2)$ endowed with its canonical volume form dv_q . Let $\alpha_{MP}(k,n) \in$ (0,1] be the so-called Munn-Perelman constant for every k = 1, ..., n, see Munn [17]. In fact, based on the double induction argument of Perelman [22], Munn determined explicit lower bounds for the volume growth in terms of the constant $\alpha_{MP}(k,n)$ which guarantee the triviality of the k-th homotopy group $\pi_k(M)$ of (M, g); see details in §3.

For sake of simplicity, we restrict here our attention to the L^p -logarithmic Sobolev inequality $(\mathbf{LS})_{\mathcal{C}}^p$ on (M, g) by proving that once $\mathcal{C} > 0$ is closer and closer to the optimal Euclidean constant $\mathcal{L}_{p,n}$, the manifold (M, g) approaches topologically more and more to the Euclidean space \mathbb{R}^n .

Theorem 1.4. Let (M,g) be an *n*-dimensional complete Riemannian manifold with non-negative Ricci curvature $(n \ge 2)$ and assume the L^p -logarithmic Sobolev inequality $(\mathbf{LS})^p_{\mathcal{C}}$ holds on (M, g) for some $p \in (1, n)$ and C > 0. Then the following assertions hold:

- (i) $\mathcal{C} \geq \mathcal{L}_{p,n}$;
- (ii) The order of the fundamental group $\pi_1(M)$ is bounded above by $\left(\frac{\mathcal{C}}{\mathcal{L}_{p,n}}\right)^{\frac{n}{p}}$;
- (iii) If $\mathcal{C} < \alpha_{MP}(k_0, n)^{-\frac{p}{n}} \mathcal{L}_{p,n}$ for some $k_0 \in \{1, ..., n\}$ then $\pi_1(M) = ... = \pi_{k_0}(M) = 0;$
- (iv) If $\mathcal{C} < \alpha_{MP}(n,n)^{-\frac{p}{n}} \mathcal{L}_{p,n}$ then M is contractible;
- (v) $\mathcal{C} = \mathcal{L}_{p,n}$ if and only if (M, g) is isometric to the Euclidean space \mathbb{R}^n .

Remark 1.2. (a) Theorem 1.4 (v) answers an open question of Xia [29] for generic $p \in (1, n)$. For p = 2 the latter equivalence is well known by using sharp analytic estimates for the heat kernel on complete Riemannian manifolds with non-negative Ricci curvature; see Bakry, Concordet and Ledoux [4], Ni [18], and Li [14]. Details are presented in $\S3.1$ (see Remark 3.1).

(b) The conclusion $\mathcal{C} \geq \mathcal{L}_{p,n}$ in Theorem 1.4 (i) is in a perfect concordance with the assumption of Theorem 1.2. Analogous statements hold for the other Gagliardo-Nirenberg inequalities.

(c) Similar results to Theorem 1.4 can be stated also for Gagliardo-Nirenberg inequalities $(GN1)_{\mathcal{C}}$ and $(\mathbf{GN2})_{\mathcal{C}}$, and Faber-Krahn inequality $(\mathbf{FK})_{\mathcal{C}}$ with trivial modifications. In particular, we have:

Corollary 1.1. [Optimality vs. flatness] Let (M, g) be an $n \geq 2$ -dimensional complete Riemannian manifold with non-negative Ricci curvature. The following statements are equivalent:

- (i) $(\mathbf{GN1})_{\mathcal{G}_{\alpha,p,n}}^{\alpha,p}$ holds on (M,g) for some $p \in (1,n)$ and $\alpha \in (1, \frac{n}{n-p}]$; (ii) $(\mathbf{GN2})_{\mathcal{N}_{\alpha,p,n}}^{\alpha,p}$ holds on (M,g) for some $p \in (1,n)$ and $\alpha \in (0,1)$;
- (iii) $(\mathbf{LS})_{\mathcal{L}_{p,n}}^p$ holds on (M,g) for some $p \in (1,n)$;
- (iv) $(\mathbf{FK})_{\mathcal{F}_{p,n}}^{\tilde{p},n}$ holds on (M,g) for some $p \in (1,n)$;
- (v) (M,q) is isometric to the Euclidean space \mathbb{R}^n .

Remark 1.3. (a) The equivalence (i) \Leftrightarrow (v) in Corollary 1.1 is precisely the main result of Xia [28].

(b) A similar rigidity result to Corollary 1.1 can be stated on reversible Finsler manifolds endowed with the natural Busemann-Hausdoff measure dV_F of (M, F); roughly speaking, we can replace the notions 'Riemannian' and 'Euclidean' in Corollary 1.1 by the notions 'Berwald' and 'Minkowski', respectively (see Theorem 3.2). The latter notions will be introduced in $\S 3.2$.

Notations. When no confusion arises, $\|\cdot\|_{L^p}$ abbreviates: (a) $\|\cdot\|_{L^p(M,dm)}$ on the metric measure space $(M, \mathsf{d}, \mathsf{m})$; (b) $\|\cdot\|_{L^p(M, dv_q)}$ on the Riemannian manifold (M, g) where dv_g stands for the canonical Riemannian measure on (M, g); (c) $\|\cdot\|_{L^p(M, dV_F)}$ on the Finsler manifold (M, F) where dV_F denotes the Busemann-Hausdoff measure on (M, F); and (d) $\|\cdot\|_{L^p(\mathbb{R}^n, dx)}$ on the Euclidean/normed space \mathbb{R}^n where dx is the usual Lebesgue measure, respectively. When A is not the whole space we are working on, we shall use the notation $||u||_{L^p(A)}$ for the L^p -norm of the function $u: A \to \mathbb{R}$.

ALEXANDRU KRISTÁLY

2. Volume non-collapsing via Gagliardo-Nirenberg inequalities

Before the presentation of the proofs of Theorems 1.1-1.3, we recall for completeness some notions and results from Lott and Villani [15] and Sturm [24, 25], which are indispensable in our arguments.

Let $(M, \mathsf{d}, \mathsf{m})$ be a metric measure space, i.e., (M, d) is a complete separable metric space and m is a locally finite measure on M endowed with its Borel σ -algebra. In the sequel, we assume that the measure m on M is strictly positive, i.e., $\operatorname{supp}[\mathsf{m}] = M$. As usual, $\mathcal{P}_2(M, \mathsf{d})$ is the L^2 -Wasserstein space of probability measures on M, while $\mathcal{P}_2(M, \mathsf{d}, \mathsf{m})$ will denote the subspace of m -absolutely continuous measures. $(M, \mathsf{d}, \mathsf{m})$ is said to be proper if every bounded and closed subset of M is compact.

For a given number $N \ge 1$, the *Rényi entropy functional* $S_N(\cdot|\mathbf{m}) : \mathcal{P}_2(M, \mathsf{d}) \to \mathbb{R}$ with respect to the measure \mathbf{m} is defined by $S_N(\mu|\mathbf{m}) = -\int_M \rho^{-\frac{1}{N}} d\mu$, ρ being the density of μ^c in $\mu = \mu^c + \mu^s = \rho \mathbf{m} + \mu^s$, where μ^c and μ^s represent the absolutely continuous and singular parts of $\mu \in \mathcal{P}_2(M, \mathsf{d})$, respectively. Let $K, N \in \mathbb{R}$ be two numbers with $K \ge 0$ and $N \ge 1$. For every $t \in [0, 1]$ and $s \ge 0$, let

$$\tau_{K,N}^{(t)}(s) = \begin{cases} +\infty, & \text{if } Ks^2 \ge (N-1)\pi^2; \\ t^{\frac{1}{N}} \left(\sin\left(\sqrt{\frac{K}{N-1}}ts\right) / \sin\left(\sqrt{\frac{K}{N-1}}s\right) \right)^{1-\frac{1}{N}}, & \text{if } 0 < Ks^2 < (N-1)\pi^2; \\ t, & \text{if } Ks^2 = 0. \end{cases}$$

We say that $(M, \mathsf{d}, \mathsf{m})$ satisfies the curvature-dimension condition $\mathsf{CD}(K, N)$ if for each $\mu_0, \mu_1 \in \mathcal{P}_2(M, \mathsf{d}, \mathsf{m})$ there exists an optimal coupling γ of μ_0, μ_1 and a geodesic $\Gamma : [0, 1] \to \mathcal{P}_2(M, \mathsf{d}, \mathsf{m})$ joining μ_0 and μ_1 such that

$$S_{N'}(\Gamma(t)|\mathsf{m}) \leq -\int_{M \times M} [\tau_{K,N'}^{(1-t)}(\mathsf{d}(x_0,x_1))\rho_0^{-\frac{1}{N'}}(x_0) + \tau_{K,N'}^{(t)}(\mathsf{d}(x_0,x_1))\rho_1^{-\frac{1}{N'}}(x_1)]d\gamma(x_0,x_1)$$

for every $t \in [0,1]$ and $N' \geq N$, where ρ_0 and ρ_1 are the densities of μ_0 and μ_1 with respect to m. Clearly, when K = 0, the above inequality reduces to the the geodesic convexity of $S_{N'}(\cdot|\mathbf{m})$ on the L^2 -Wasserstein space $\mathcal{P}_2(M, \mathsf{d}, \mathsf{m})$.

It is well known that CD(K, n) holds on a complete Riemannian manifold (M, g) endowed with the Riemannian volume element dv_g if and only if its Ricci curvature $\geq K$ and $\dim(M) \leq n$.

Let $B_x(r) = \{y \in M : d(x, y) < r\}$. In the sequel we shall exploit properties which are resumed in the following results.

Theorem 2.1. (see [25]) Let (M, d, m) be a metric measure space with strictly positive measure m satisfying the curvature-dimension condition CD(K, N) for some $K \ge 0$ and N > 1. Then every bounded set $S \subset M$ has finite m-measure and the metric spheres $\partial B_x(r)$ have zero m-measures. Moreover, one has:

- (i) [Generalized Bonnet-Myers theorem] If K > 0, then $M = \text{supp}[\mathsf{m}]$ is compact and has diameter less than or equal to $\sqrt{\frac{N-1}{K}}\pi$.
- (ii) [Generalized Bishop-Gromov inequality] If K = 0, then for every R > r > 0 and $x \in M$,

$$\frac{\mathsf{m}(B_x(r))}{r^N} \ge \frac{\mathsf{m}(B_x(R))}{R^N}$$

Lemma 2.1. Let (M, d, m) be a metric measure space which satisfies the curvature-dimension condition CD(0, n) for some $n \ge 2$. If

$$\ell_{\infty}^{x_0} := \limsup_{\rho \to \infty} \frac{\mathsf{m}(B_{x_0}(\rho))}{\omega_n \rho^n} \ge a \tag{2.1}$$

for some $x_0 \in M$ and a > 0, then

$$\mathsf{m}(B_x(\rho)) \ge a\omega_n \rho^n, \ \forall x \in M, \ \rho \ge 0.$$

Proof. Let us fix $x \in M$ and $\rho > 0$; then we have

$$\frac{\mathsf{m}(B_x(\rho))}{\omega_n \rho^n} \geq \limsup_{r \to \infty} \frac{\mathsf{m}(B_x(r))}{\omega_n r^n} \qquad [\text{Bishop} - \text{Gromov inequality}]$$

$$\geq \limsup_{r \to \infty} \frac{\mathsf{m}(B_{x_0}(r - \mathsf{d}(x_0, x)))}{\omega_n r^n} \qquad [B_x(r) \supset B_{x_0}(r - \mathsf{d}(x_0, x))]$$

$$= \limsup_{r \to \infty} \left(\frac{\mathsf{m}(B_{x_0}(r - \mathsf{d}(x_0, x)))}{\omega_n (r - \mathsf{d}(x_0, x))^n} \cdot \frac{(r - \mathsf{d}(x_0, x))^n}{r^n} \right)$$

$$= \ell_{\infty}^{x_0}$$

$$\geq a, \qquad [cf. (2.1)]$$

which concludes the proof.

We are now in the position to prove our volume non-collapsing results.

2.1. Cases $\alpha > 1$ & $0 < \alpha < 1$: usual Gagliardo-Nirenberg inequalities. In this subsection we present the proof of Theorem 1.1 by distinguishing two cases:

Proof of Theorem 1.1 (i): the case $1 < \alpha \leq \frac{n}{n-p}$. In this part we follow the line of [12]; the proof is divided into several steps. We clearly may assume that $\mathcal{C} > \mathcal{G}_{\alpha,p,n}$ in $(\mathbf{GN1})^{\alpha,p}_{\mathcal{C}}$; indeed, if $\mathcal{C} = \mathcal{G}_{\alpha,p,n}$ we can consider the subsequent arguments for $\mathcal{C} := \mathcal{G}_{\alpha,p,n} + \varepsilon$ with small $\varepsilon > 0$ and then take $\varepsilon \to 0^+$.

STEP 1 (K = 0). If we assume that K > 0 then the generalized Bonnet-Myers theorem (see Theorem 2.1 (i)) implies that M is compact and $\mathsf{m}(M)$ is finite. Taking the constant map $u(x) = \mathsf{m}(M)$ in $(\mathbf{GN1})^{\alpha,p}_{\mathcal{C}}$ as a test function, one gets a contradiction. Therefore, K = 0.

STEP 2 (*ODE from the optimal Euclidean Gagliardo-Nirenberg inequality I*). We consider the optimal Gagliardo-Nirenberg inequality (1.1) in the particular case when the norm is precisely the Euclidean norm $|\cdot|$. After a simple rescaling, one can see that the function $x \mapsto (\lambda + |x|^{p'})^{\frac{1}{1-\alpha}}, \lambda > 0$, is a family of extremals in (1.1); therefore, we have the following first order ODE

$$\left(\frac{1-\alpha}{\alpha(p-1)+1}h'_G(\lambda)\right)^{\frac{1}{\alpha p}} = \mathcal{G}_{\alpha,p,n}\left(\frac{p'}{\alpha-1}\right)^{\theta} \left(h_G(\lambda) + \frac{\alpha-1}{\alpha(p-1)+1}\lambda h'_G(\lambda)\right)^{\frac{\theta}{p}} h_G(\lambda)^{\frac{1-\theta}{\alpha(p-1)+1}}, \quad (2.2)$$

where $h_G: (0,\infty) \to \mathbb{R}$ is given by

$$h_G(\lambda) = \int_{\mathbb{R}^n} \left(\lambda + |x|^{p'}\right)^{\frac{\alpha(p-1)+1}{1-\alpha}} dx, \ \lambda > 0.$$

For further use, we shall represent the function h_G in two different ways, namely

$$h_G(\lambda) = \omega_n \frac{n}{p'} \mathsf{B}\left(\frac{\alpha(p-1)+1}{\alpha-1} - \frac{n}{p'}, \frac{n}{p'}\right) \lambda^{\frac{\alpha(p-1)+1}{1-\alpha} + \frac{n}{p'}} = \int_0^\infty \omega_n \rho^n f_G(\lambda, \rho) d\rho,$$
(2.3)

where

$$f_G(\lambda,\rho) = p' \frac{\alpha(p-1)+1}{\alpha-1} \left(\lambda + \rho^{p'}\right)^{\frac{\alpha p}{1-\alpha}} \rho^{p'-1}.$$
(2.4)

STEP 3 (*Differential inequality from* $(\mathbf{GN1})^{\alpha,p}_{\mathcal{C}}$). By the generalized Bishop-Gromov inequality (see Theorem 2.1 (ii)) and hypothesis $(\mathbf{D})^n_{x_0}$ one has that

$$\frac{\mathsf{m}(B_{x_0}(\rho))}{\omega_n \rho^n} \le \liminf_{r \to 0} \frac{\mathsf{m}(B_{x_0}(r))}{\omega_n r^n} = 1, \ \rho > 0.$$
(2.5)

Inspired by the form of h_G , we consider the function $w_G: (0, \infty) \to \mathbb{R}$ defined by

$$w_G(\lambda) = \int_M \left(\lambda + \mathsf{d}(x_0, x)^{p'}\right)^{\frac{\alpha(p-1)+1}{1-\alpha}} d\mathsf{m}(x), \ \lambda > 0.$$

By using the layer cake representation, it follows that w_G is well-defined and of class C^1 ; indeed,

$$w_{G}(\lambda) = \int_{0}^{\infty} \mathsf{m}\left(\left\{x \in M : \left(\lambda + \mathsf{d}(x_{0}, x)^{p'}\right)^{\frac{\alpha(p-1)+1}{1-\alpha}} > t\right\}\right) dt$$

$$= \int_{0}^{\infty} \mathsf{m}(B_{x_{0}}(\rho)) f_{G}(\lambda, \rho) d\rho \qquad [\text{change } t = \left(\lambda + \rho^{p'}\right)^{\frac{\alpha(p-1)+1}{1-\alpha}} \text{ and see (2.4)}]$$

$$\leq \int_{0}^{\infty} \omega_{n} \rho^{n} f_{G}(\lambda, \rho) d\rho \qquad [\text{see (2.5)}]$$

$$= h_{G}(\lambda),$$

thus

$$0 < w_G(\lambda) \le h_G(\lambda) < \infty, \ \lambda > 0.$$
(2.6)

For every $\lambda > 0$ and $k \in \mathbb{N}$, we consider the function $u_{\lambda,k} : M \to \mathbb{R}$ defined by

$$u_{\lambda,k}(x) = (\min\{0, k - \mathsf{d}(x_0, x)\} + 1)_+ \left(\lambda + \max\{\mathsf{d}(x_0, x), k^{-1}\}^{p'}\right)^{\frac{1}{1-\alpha}}$$

Note that since $(M, \mathsf{d}, \mathsf{m})$ is proper, the set $\operatorname{supp}(u_{\lambda,k}) = \overline{B_{x_0}(k+1)}$ is compact. Consequently, $u_{\lambda,k} \in \operatorname{Lip}_0(M)$ for every $\lambda > 0$ and $k \in \mathbb{N}$; thus we can apply these functions in $(\mathbf{GN1})_{\mathcal{C}}^{\alpha,p}$, i.e.,

$$\|u_{\lambda,k}\|_{L^{\alpha p}} \leq \mathcal{C} \||\nabla u_{\lambda,k}|_{\mathsf{d}}\|_{L^p}^{\theta} \|u_{\lambda,k}\|_{L^{\alpha(p-1)+1}}^{1-\theta}$$

Moreover,

$$\lim_{k \to \infty} u_{\lambda,k}(x) = \left(\lambda + \mathsf{d}(x_0, x)^{p'}\right)^{\frac{1}{1-\alpha}} =: u_{\lambda}(x).$$

By using the dominated convergence theorem, it turns out from the above inequality that u_{λ} also verifies $(\mathbf{GN1})_{\mathcal{C}}^{\alpha,p}$, i.e.,

$$\|u_{\lambda}\|_{L^{\alpha p}} \leq \mathcal{C} \||\nabla u_{\lambda}|_{\mathsf{d}}\|_{L^{p}}^{\theta} \|u_{\lambda}\|_{L^{\alpha(p-1)+1}}^{1-\theta}.$$
(2.7)

The non-smooth chain rule gives that

$$|\nabla u_{\lambda}|_{\mathsf{d}}(x) = \frac{p'}{\alpha - 1} \left(\lambda + \mathsf{d}(x_0, x)^{p'} \right)^{\frac{\alpha}{1 - \alpha}} \mathsf{d}(x_0, x)^{p' - 1} |\nabla \mathsf{d}(x_0, \cdot)|_{\mathsf{d}}(x), \ x \in M.$$
(2.8)

Since $d(x_0, \cdot)$ is 1-Lipschitz (therefore, $|\nabla d(x_0, \cdot)|_d(x) \le 1$ for all $x \in M$), due to (2.7), (2.8) and the form of the function w_G , we obtain the differential inequality

$$\left(\frac{1-\alpha}{\alpha(p-1)+1}w_G'(\lambda)\right)^{\frac{1}{\alpha p}} \le \mathcal{C}\left(\frac{p'}{\alpha-1}\right)^{\theta} \left(w_G(\lambda) + \frac{\alpha-1}{\alpha(p-1)+1}\lambda w_G'(\lambda)\right)^{\frac{\theta}{p}} w_G(\lambda)^{\frac{1-\theta}{\alpha(p-1)+1}}.$$
 (2.9)

STEP 4 (Comparison of w_G and h_G near the origin). We claim that

$$\lim_{\lambda \to 0^+} \frac{w_G(\lambda)}{h_G(\lambda)} = 1.$$
(2.10)

By hypothesis $(\mathbf{D})_{x_0}^n$, for every $\varepsilon > 0$ there exists $\rho_{\varepsilon} > 0$ such that

$$\mathsf{m}(B_{x_0}(\rho)) \ge (1-\varepsilon)\omega_n \rho^n \text{ for all } \rho \in [0, \rho_\varepsilon].$$
(2.11)

By (2.11), one has that

$$w_{G}(\lambda) = \int_{0}^{\infty} \mathsf{m}(B_{x_{0}}(\rho)) f_{G}(\lambda, \rho) d\rho$$

$$\geq (1 - \varepsilon) \int_{0}^{\rho_{\varepsilon}} \omega_{n} \rho^{n} f_{G}(\lambda, \rho) d\rho = (1 - \varepsilon) \lambda^{\frac{\alpha(p-1)+1}{1-\alpha} + \frac{n}{p'}} \int_{0}^{\rho_{\varepsilon} \lambda^{-\frac{1}{p'}}} \omega_{n} \rho^{n} f_{G}(1, \rho) d\rho.$$

Thus, by the representation (2.3) of h_G and a change of variables, it turns out that

$$\liminf_{\lambda \to 0^+} \frac{w_G(\lambda)}{h_G(\lambda)} \ge (1-\varepsilon) \liminf_{\lambda \to 0^+} \frac{\int_0^{\rho_\varepsilon \lambda^{-\frac{1}{p'}}} \omega_n \rho^n f_G(1,\rho) d\rho}{\int_0^\infty \omega_n \rho^n f_G(1,\rho) d\rho} = 1-\varepsilon$$

The above inequality (with $\varepsilon > 0$ arbitrary small) combined with (2.6) proves the claim (2.10).

STEP 5 (Global comparison of w_G and h_G). We now claim that

$$w_G(\lambda) \ge \left(\frac{\mathcal{G}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\theta}} h_G(\lambda) = \tilde{h}_G(\lambda), \ \lambda > 0.$$
(2.12)

Since we assumed that $C > \mathcal{G}_{\alpha,p,n}$, by (2.10) one has

$$\lim_{\lambda \to 0^+} \frac{w_G(\lambda)}{\tilde{h}_G(\lambda)} = \left(\frac{\mathcal{C}}{\mathcal{G}_{\alpha,p,n}}\right)^{\frac{n}{\theta}} > 1$$

Therefore, there exists $\lambda_0 > 0$ such that for every $\lambda \in (0, \lambda_0)$, one has $w_G(\lambda) > h_G(\lambda)$.

By contradiction to (2.12), we assume that there exists $\lambda^{\#} > 0$ such that $w_G(\lambda^{\#}) < \tilde{h}_G(\lambda^{\#})$. If $\lambda^* = \sup\{0 < \lambda < \lambda^{\#} : w_G(\lambda) = \tilde{h}_G(\lambda)\}$, then $0 < \lambda_0 \leq \lambda^* < \lambda^{\#}$. In particular,

$$w_G(\lambda) \le \tilde{h}_G(\lambda), \ \forall \lambda \in [\lambda^*, \lambda^{\#}].$$

The latter relation and the differential inequality (2.9) imply that for every $\lambda \in [\lambda^*, \lambda^{\#}]$,

$$\left(\frac{1-\alpha}{\alpha(p-1)+1}w_G'(\lambda)\right)^{\frac{1}{\alpha\theta}} \le \mathcal{C}^{\frac{p}{\theta}}\left(\frac{p'}{\alpha-1}\right)^p \left(\tilde{h}_G(\lambda) + \frac{\alpha-1}{\alpha(p-1)+1}\lambda w_G'(\lambda)\right) \tilde{h}_G(\lambda)^{\frac{(1-\theta)p}{\theta(\alpha(p-1)+1)}}.$$
 (2.13)

Moreover, since $\tilde{h}_G(\lambda) = \left(\frac{\mathcal{G}_{\alpha,p,b}}{\mathcal{C}}\right)^{\overline{\theta}} h_G(\lambda)$, the ODE in (2.2) can be equivalently transformed for every $\lambda > 0$ into the equation

$$\left(\frac{1-\alpha}{\alpha(p-1)+1}\tilde{h}'_G(\lambda)\right)^{\frac{1}{\alpha\theta}} = \mathcal{C}^{\frac{p}{\theta}}\left(\frac{p'}{\alpha-1}\right)^p \left(\tilde{h}_G(\lambda) + \frac{\alpha-1}{\alpha(p-1)+1}\lambda\tilde{h}'_G(\lambda)\right)\tilde{h}_G(\lambda)^{\frac{(1-\theta)p}{\theta(\alpha(p-1)+1)}}.$$
 (2.14)

For $\lambda > 0$ fixed we introduce the increasing function $j_G^{\lambda} : (0, \infty) \to \mathbb{R}$ defined by

$$j_G^{\lambda}(t) = \left(\frac{\alpha - 1}{\alpha(p - 1) + 1}t\right)^{\frac{1}{\alpha\theta}} + \mathcal{C}^{\frac{p}{\theta}}\left(\frac{p'}{\alpha - 1}\right)^p \frac{\alpha - 1}{\alpha(p - 1) + 1}\lambda \tilde{h}_G(\lambda)^{\frac{(1 - \theta)p}{\theta(\alpha(p - 1) + 1)}}t.$$

Relations (2.13) and (2.14) can be rewritten into

$$j_G^{\lambda}(-w_G'(\lambda)) \le \mathcal{C}^{\frac{p}{\theta}} \left(\frac{p'}{\alpha-1}\right)^p \tilde{h}_G(\lambda)^{1+\frac{(1-\theta)p}{\theta(\alpha(p-1)+1)}} = j_G^{\lambda}(-\tilde{h}_G'(\lambda)), \ \forall \lambda \in [\lambda^*, \lambda^{\#}],$$

which implies that

$$-w'_G(\lambda) \le -\tilde{h}'_G(\lambda), \ \forall \lambda \in [\lambda^*, \lambda^{\#}],$$

i.e., the function $\tilde{h}_G - w_G$ is non-increasing in $[\lambda^*, \lambda^{\#}]$. In particular, $0 < (\tilde{h}_G - w_G)(\lambda^{\#}) \le (\tilde{h}_G - w_G)(\lambda^*) = 0$, a contradiction. This concludes the proof of (2.12).

STEP 6 (Asymptotic volume growth estimate w.r.t. x_0). We claim that

$$\ell_{\infty}^{x_0} := \limsup_{\rho \to \infty} \frac{\mathsf{m}(B_{x_0}(\rho))}{\omega_n \rho^n} \ge \left(\frac{\mathcal{G}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\theta}}.$$
(2.15)

By assuming the contrary, there exists $\varepsilon_0 > 0$ such that for some $\rho_0 > 0$,

$$\frac{\mathsf{m}(B_{x_0}(\rho))}{\omega_n \rho^n} \le \left(\frac{\mathcal{G}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\theta}} - \varepsilon_0, \ \forall \rho \ge \rho_0$$

By (2.12) and from the latter relation, we have for every $\lambda > 0$ that

$$0 \leq w_{G}(\lambda) - \left(\frac{\mathcal{G}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\theta}} h_{G}(\lambda)$$

$$= \int_{0}^{\infty} \left(\frac{\mathsf{m}(B_{x_{0}}(\rho))}{\omega_{n}\rho^{n}} - \left(\frac{\mathcal{G}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\theta}}\right) \omega_{n}\rho^{n} f_{G}(\lambda,\rho)d\rho$$

$$\leq \left(1 + \varepsilon_{0} - \left(\frac{\mathcal{G}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\theta}}\right) \int_{0}^{\rho_{0}} \omega_{n}\rho^{n} f_{G}(\lambda,\rho)d\rho - \varepsilon_{0} \int_{0}^{\infty} \omega_{n}\rho^{n} f_{G}(\lambda,\rho)d\rho$$

By using (2.3), a suitable rearrangement of the terms in the above relation shows that

$$\varepsilon_0 \frac{n}{p'} \mathsf{B}\left(\frac{\alpha(p-1)+1}{\alpha-1} - \frac{n}{p'}, \frac{n}{p'}\right) \lambda^{1+\frac{n}{p'}} \le \frac{p'}{n+p'} \left(1 + \varepsilon_0 - \left(\frac{\mathcal{G}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\theta}}\right) \frac{\alpha(p-1)+1}{\alpha-1} \rho_0^{n+p'}.$$

If we take the limit $\lambda \to +\infty$ in the last estimate, we obtain a contradiction. Thus, the claim (2.15) is proved and it remains to apply Lemma 2.1, which concludes the proof of Theorem 1.1 (i).

Proof of Theorem 1.1 (ii): the case $0 < \alpha < 1$. We shall invoke some of the arguments from the proof of Theorem 1.1 (i), emphasizing that subtle differences arise due to the 'dual' nature of the Gagliardo-Nirenberg inequalities $(\mathbf{GN1})_{\mathcal{C}}^{\alpha,p}$ and $(\mathbf{GN2})_{\mathcal{C}}^{\alpha,p}$, respectively. As before, we may assume that the inequality $(\mathbf{GN2})_{\mathcal{C}}^{\alpha,p}$ holds with $\mathcal{C} > \mathcal{N}_{\alpha,p,n}$. STEP 1. The fact that K = 0 works similarly as in Theorem 1.1 (i).

STEP 2. Since $x \mapsto \left(\lambda^{p'} - |x|^{p'}\right)_{+}^{\frac{1}{1-\alpha}}$ is an extremal function in (1.3) for every $\lambda > 0$, we obtain the ODE

$$h_N(\lambda)^{\frac{1}{\alpha(p-1)+1}} = \mathcal{N}_{\alpha,p,n}\left(\frac{p'}{1-\alpha}\right)^{\gamma} \left(-h_N(\lambda) + \frac{1-\alpha}{p'(\alpha(p-1)+1)}\lambda h'_N(\lambda)\right)^{\frac{\gamma}{p}} \times \left(\frac{1-\alpha}{p'(\alpha(p-1)+1)}\lambda^{1-p'}h'_N(\lambda)\right)^{\frac{1-\gamma}{\alpha p}},$$
(2.16)

where the function $h_N: (0, \infty) \to \mathbb{R}$ is defined by

$$h_N(\lambda) = \int_{\mathbb{R}^n} \left(\lambda^{p'} - |x|^{p'} \right)_+^{\frac{\alpha(p-1)+1}{1-\alpha}} dx, \ \lambda > 0$$

It is clear that h_N is well-defined, of class C^1 and can be represented as

$$h_N(\lambda) = \omega_n \frac{n}{p'} \mathsf{B}\left(\frac{\alpha(p-1)+1}{1-\alpha} + 1, \frac{n}{p'}\right) \lambda^{\frac{\alpha p p'}{1-\alpha} + n+p'} = \int_0^\lambda \omega_n \rho^n f_N(\lambda, \rho) d\rho,$$

where

$$f_N(\lambda,\rho) = p' \frac{\alpha(p-1)+1}{1-\alpha} \left(\lambda^{p'} - \rho^{p'}\right)^{\frac{\alpha_p}{1-\alpha}} \rho^{p'-1}, \text{ for every } \lambda > 0 \text{ and } \rho \in (0,\lambda).$$
(2.17)

STEP 3. Let $w_N: (0,\infty) \to \mathbb{R}$ be the function defined by

$$w_N(\lambda) = \int_M \left(\lambda^{p'} - \mathsf{d}(x_0, x)^{p'}\right)_+^{\frac{\alpha(p-1)+1}{1-\alpha}} d\mathsf{m}(x), \ \lambda > 0.$$

where $x_0 \in M$ is from $(\mathbf{D})_{x_0}^n$. By the layer cake representation and relations (2.5) and (2.17), w_N is well-defined, positive, of class C^1 and

$$0 < w_N(\lambda) = \int_0^\lambda \mathsf{m}(B_{x_0}(\rho)) f_N(\lambda, \rho) d\rho \le \int_0^\lambda \omega_n \rho^n f_N(\lambda, \rho) d\rho = h_N(\lambda) < \infty, \ \lambda > 0.$$
(2.18)

Since $u_{\lambda} = \left(\lambda^{p'} - \mathsf{d}(x_0, \cdot)^{p'}\right)_{+}^{\frac{1}{1-\alpha}}$ is a Lipschitz function on M with compact support $\overline{B_{x_0}(\lambda)}$, it belongs to $\operatorname{Lip}_0(M)$. Therefore, we may apply u_{λ} in $(\mathbf{GN2})_{\mathcal{C}}^{\alpha,p}$; a similar reasoning as in (2.8) leads to the differential inequality

$$w_N(\lambda)^{\frac{1}{\alpha(p-1)+1}} \leq \mathcal{C}\left(\frac{p'}{1-\alpha}\right)^{\gamma} \left(-w_N(\lambda) + \frac{1-\alpha}{p'(\alpha(p-1)+1)}\lambda w'_N(\lambda)\right)^{\frac{\gamma}{p}} \times \left(\frac{1-\alpha}{p'(\alpha(p-1)+1)}\lambda^{1-p'}w'_N(\lambda)\right)^{\frac{1-\gamma}{\alpha p}}, \ \lambda > 0.$$
(2.19)

STEP 4. For an arbitrarily fixed $\varepsilon > 0$, let $\rho_{\varepsilon} > 0$ from (2.11). If $0 < \lambda < \rho_{\varepsilon}$, one has that

$$w_N(\lambda) = \int_0^\lambda \mathsf{m}(B_{x_0}(\rho)) f_N(\lambda, \rho) d\rho \ge (1 - \varepsilon) \int_0^\lambda \omega_n \rho^n f_N(\lambda, \rho) d\rho = (1 - \varepsilon) h_N(\lambda).$$

Consequently, the latter relation together with (2.18) implies that

$$\lim_{\lambda \to 0^+} \frac{w_N(\lambda)}{h_N(\lambda)} = 1.$$
(2.20)

STEP 5. We shall prove that

$$w_N(\lambda) \ge \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}} h_N(\lambda) = \tilde{h}_N(\lambda), \ \lambda > 0.$$
 (2.21)

By (2.20) one has

$$\lim_{\lambda \to 0^+} \frac{w_N(\lambda)}{\tilde{h}_N(\lambda)} = \left(\frac{\mathcal{C}}{\mathcal{N}_{\alpha,p,n}}\right)^{\frac{n}{\gamma}} > 1,$$

which implies the existence of a number $\lambda_0 > 0$ such that $w_N(\lambda) > \tilde{h}_N(\lambda)$ for every $\lambda \in (0, \lambda_0)$.

We assume by contradiction that there exists $\lambda^{\#} > 0$ such that $w_N(\lambda^{\#}) < \tilde{h}_N(\lambda^{\#})$. If $\lambda^* = \sup\{0 < \lambda < \lambda^{\#} : w_N(\lambda) = \tilde{h}_N(\lambda)\}$, then $0 < \lambda_0 \le \lambda^* < \lambda^{\#}$ and

$$w_N(\lambda) \le \tilde{h}_N(\lambda), \ \forall \lambda \in [\lambda^*, \lambda^\#].$$
 (2.22)

For every $\lambda > 0$, let $j_N^{\lambda} : \left(\frac{p'(\alpha(p-1)+1)}{(1-\alpha)\lambda}, \infty\right) \to \mathbb{R}$ be the function defined by

$$j_N^{\lambda}(t) = \mathcal{C}\left(\frac{p'}{1-\alpha}\right)^{\gamma} \left(-1 + \frac{1-\alpha}{p'(\alpha(p-1)+1)}\lambda t\right)^{\frac{\gamma}{p}} \left(\frac{1-\alpha}{p'(\alpha(p-1)+1)}\lambda^{1-p'}t\right)^{\frac{1-\gamma}{\alpha p}}.$$

It is clear that j_N^{λ} is well-defined, positive and increasing. A direct computation yields that both values $(\log w_N)'(\lambda) = \frac{w'_N(\lambda)}{w_N(\lambda)}$ and $(\log \tilde{h}_N)'(\lambda) = \frac{\tilde{h}'_N(\lambda)}{\tilde{h}_N(\lambda)}$ are greater than $\frac{p'(\alpha(p-1)+1)}{(1-\alpha)\lambda}$ for every $\lambda > 0$. Taking into account (1.4), we have

$$\frac{1}{\alpha(p-1)+1} - \frac{\gamma}{p} - \frac{1-\gamma}{\alpha p} = -\frac{\gamma}{n};$$

therefore, if we divide the inequality (2.19) by $w_N(\lambda)^{\frac{\gamma}{p}+\frac{1-\gamma}{\alpha p}}$, we obtain that

$$w_N(\lambda)^{-\frac{\gamma}{n}} \le j_N^\lambda \left((\log w_N)'(\lambda) \right), \ \forall \lambda > 0.$$
(2.23)

In a similar manner, by $\tilde{h}_N(\lambda) = \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}} h_N(\lambda)$ and relation (2.16), we have that

$$\tilde{h}_N(\lambda)^{-\frac{\gamma}{n}} = j_N^\lambda \left((\log \tilde{h}_N)'(\lambda) \right), \ \forall \lambda > 0.$$
(2.24)

Thus, by (2.22), (2.23) and (2.24), it turns out that

$$j_N^{\lambda}\left((\log \tilde{h}_N)'(\lambda)\right) = \tilde{h}_N(\lambda)^{-\frac{\gamma}{n}} \le w_N(\lambda)^{-\frac{\gamma}{n}} \le j_N^{\lambda}\left((\log w_N)'(\lambda)\right), \ \forall \lambda \in [\lambda^*, \lambda^{\#}].$$

Since the inverse of j_N^{λ} is also increasing, it follows that $(\log \tilde{h}_N)'(\lambda) \leq (\log w_N)'(\lambda)$ for every $\lambda \in [\lambda^*, \lambda^{\#}]$. Therefore, the function $\lambda \mapsto \log \frac{\tilde{h}_N(\lambda)}{w_N(\lambda)}$ is non-increasing in the interval $[\lambda^*, \lambda^{\#}]$. In particular, it follows that

$$0 < \log \frac{\tilde{h}_N(\lambda^{\#})}{w_N(\lambda^{\#})} \le \log \frac{\tilde{h}_N(\lambda^*)}{w_N(\lambda^*)} = 0,$$

a contradiction, which proves the validity of the claim (2.21).

STEP 6. We shall prove that

$$\limsup_{\rho \to \infty} \frac{\mathsf{m}(B_{x_0}(\rho))}{\omega_n \rho^n} \ge \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}}.$$
(2.25)

By contradiction, we assume that there exists $\varepsilon_0 > 0$ such that for some $\rho_0 > 0$,

$$\frac{\mathsf{m}(B_{x_0}(\rho))}{\omega_n \rho^n} \le \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}} - \varepsilon_0, \ \forall \rho \ge \rho_0.$$

The above inequality and (2.21) imply that for every $\lambda > \rho_0$,

$$0 \leq w_{N}(\lambda) - \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}} h_{N}(\lambda) = \int_{0}^{\lambda} \left(\frac{\mathsf{m}(B_{x_{0}}(\rho))}{\omega_{n}\rho^{n}} - \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}}\right) \omega_{n}\rho^{n} f_{N}(\lambda,\rho)d\rho$$
$$\leq \left(1 + \varepsilon_{0} - \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}}\right) \int_{0}^{\rho_{0}} \omega_{n}\rho^{n} f_{N}(\lambda,\rho)d\rho - \varepsilon_{0} \int_{0}^{\lambda} \omega_{n}\rho^{n} f_{N}(\lambda,\rho)d\rho.$$

Reorganizing the latter estimate, it follows that for every $\lambda > 0$,

$$\varepsilon_0 \frac{n}{p'} \mathsf{B}\left(\frac{\alpha(p-1)+1}{1-\alpha}+1, \frac{n}{p'}\right) \lambda^{n+p'} \le \frac{p'}{n+p'} \left(1+\varepsilon_0 - \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}}\right) \frac{\alpha(p-1)+1}{1-\alpha} \rho_0^{n+p'}$$

Once we let $\lambda \to \infty$, we get a contradiction. Therefore, (2.25) holds and Lemma 2.1 yields that

$$\frac{\mathsf{m}(B_x(\rho))}{\omega_n \rho^n} \ge \left(\frac{\mathcal{N}_{\alpha,p,n}}{\mathcal{C}}\right)^{\frac{n}{\gamma}}, \ \forall x \in M, \ \rho > 0,$$

which concludes the proof of Theorem 1.1 (ii).

2.2. Limit case I ($\alpha \to 1$): L^p -logarithmic Sobolev inequality. In this subsection we shall provide the proof of Theorem 1.2. We shall assume that $\mathcal{C} > \mathcal{L}_{p,n}$ in $(\mathbf{LS})^p_{\mathcal{C}}$.

STEP 1. As in the previous proofs, we obtain that K = 0; the only difference is that we shall consider $u(x) = \mathsf{m}(M)^{-1/p}$ as a test function in $(\mathbf{LS})^p_{\mathcal{C}}$, in order to fulfil the normalization assumption $||u||_{L^p} = 1$.

STEP 2. Since the functions l_p^{λ} ($\lambda > 0$) in Theorem B are extremals in (1.5), once we plug them we obtain a first order ODE of the form

$$-\log h_L(\lambda) + \lambda \frac{h'_L(\lambda)}{h_L(\lambda)} = \frac{n}{p} \log \left(-\mathcal{L}_{p,n} \left(\frac{p'}{p} \right)^p \lambda^p \frac{h'_L(\lambda)}{h_L(\lambda)} \right), \ \lambda > 0,$$
(2.26)

where $h_L: (0, \infty) \to \mathbb{R}$ is defined by

$$h_L(\lambda) = \int_{\mathbb{R}^n} e^{-\lambda |x|^{p'}} dx$$

For later use, we recall that h_L can be represented alternatively by

$$h_L(\lambda) = \frac{2\pi^{\frac{n}{2}}}{p'\lambda^{\frac{n}{p'}}} \cdot \frac{\Gamma\left(\frac{n}{p'}\right)}{\Gamma\left(\frac{n}{2}\right)} = \lambda p'\omega_n \int_0^\infty e^{-\lambda\rho^{p'}} \rho^{n+p'-1} d\rho = \lambda^{-\frac{n}{p'}} p'\omega_n \int_0^\infty e^{-t^{p'}} t^{n+p'-1} dt.$$
(2.27)

STEP 3. Let $w_L: (0,\infty) \to \mathbb{R}$ be defined by

$$w_L(\lambda) = \int_M e^{-\lambda \mathsf{d}(x_0, x)^{p'}} d\mathsf{m}(x).$$

where $x_0 \in M$ is the element from hypothesis $(\mathbf{D})_{x_0}^n$. Note that w_L is well-defined, positive and differentiable. Indeed, by the layer cake representation, for every $\lambda > 0$ we obtain that

$$w_{L}(\lambda) = \int_{0}^{\infty} \mathsf{m}\left(\left\{x \in M : e^{-\lambda \mathsf{d}(x_{0},x)^{p'}} > t\right\}\right) dt = \int_{0}^{1} \mathsf{m}\left(\left\{x \in M : e^{-\lambda \mathsf{d}(x_{0},x)^{p'}} > t\right\}\right) dt$$
$$= \lambda p' \int_{0}^{\infty} \mathsf{m}(B_{x_{0}}(\rho)) e^{-\lambda \rho^{p'}} \rho^{p'-1} d\rho \qquad [\text{change } t = e^{-\lambda \rho^{p'}}]$$
$$\leq \lambda p' \omega_{n} \int_{0}^{\infty} e^{-\lambda \rho^{p'}} \rho^{n+p'-1} d\rho \qquad [\text{see } (2.5)]$$
$$= h_{L}(\lambda) < +\infty.$$

Let us consider the family of functions $\tilde{u}_{\lambda}: M \to \mathbb{R} \ (\lambda > 0)$ defined by

$$\tilde{u}_{\lambda}(x) = \frac{e^{-\frac{\lambda}{p}\mathsf{d}(x_0,x)^{p'}}}{w_L(\lambda)^{\frac{1}{p}}}, \ x \in M.$$

It is clear that $\|\tilde{u}_{\lambda}\|_{L^{p}} = 1$ and as in the proof of Theorem 1.1 (i), the function \tilde{u}_{λ} can be approximated by elements from $\operatorname{Lip}_{0}(M)$; in fact, \tilde{u}_{λ} can be used as a test function in $(\mathbf{LS})^{p}_{\mathcal{C}}$. Thus, plugging \tilde{u}_{λ} into the inequality $(\mathbf{LS})^{p}_{\mathcal{C}}$, applying the non-smooth chain rule and the fact that $|\nabla \mathsf{d}(x_{0}, \cdot)|_{\mathsf{d}}(x) \leq 1$ for every $x \in M$, it yields

$$-\log w_L(\lambda) + \lambda \frac{w'_L(\lambda)}{w_L(\lambda)} \le \frac{n}{p} \log \left(-\mathcal{C}\left(\frac{p'}{p}\right)^p \lambda^p \frac{w'_L(\lambda)}{w_L(\lambda)} \right), \ \lambda > 0.$$
(2.28)

STEP 4. We prove that

$$\lim_{\lambda \to +\infty} \frac{w_L(\lambda)}{h_L(\lambda)} = 1.$$
(2.29)

For a fixed $\varepsilon > 0$, let $\rho_{\varepsilon} > 0$ from (2.11). Then one has

$$w_{L}(\lambda) = \lambda p' \int_{0}^{\infty} \mathsf{m}(B_{x_{0}}(\rho)) e^{-\lambda \rho^{p'}} \rho^{p'-1} d\rho \ge \lambda p'(1-\varepsilon)\omega_{n} \int_{0}^{\rho_{\varepsilon}} e^{-\lambda \rho^{p'}} \rho^{n+p'-1} d\rho$$
$$= \lambda^{-\frac{n}{p'}} p'(1-\varepsilon)\omega_{n} \int_{0}^{\rho_{\varepsilon}\lambda^{\frac{1}{p'}}} e^{-t^{p'}} t^{n+p'-1} dt. \qquad [\text{change } t = \lambda^{\frac{1}{p'}} \rho]$$

Therefore, by the third representation of h_L (see (2.27)) it turns out that

$$\liminf_{\lambda \to +\infty} \frac{w_L(\lambda)}{h_L(\lambda)} \ge 1 - \varepsilon.$$

The arbitrariness of $\varepsilon > 0$ together with Step 3 implies the validity of (2.29).

STEP 5. We claim that

$$w_L(\lambda) \ge \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}} h_L(\lambda) =: \tilde{h}_L(\lambda), \ \lambda > 0.$$
(2.30)

n

Since $\mathcal{C} > \mathcal{L}_{p,n}$, by (2.29) it follows that

$$\lim_{\lambda \to +\infty} \frac{w_L(\lambda)}{\tilde{h}_L(\lambda)} = \left(\frac{\mathcal{C}}{\mathcal{L}_{p,n}}\right)^{\frac{n}{p}} > 1.$$

Consequently, there exists $\tilde{\lambda} > 0$ such that $w_L(\lambda) > \tilde{h}_L(\lambda)$ for all $\lambda > \tilde{\lambda}$. If we introduce the notations

$$W(\lambda) = \log w_L(\lambda)$$
 and $\dot{H}(\lambda) = \log h_L(\lambda), \ \lambda > 0,$

the latter relation implies that

$$W(\lambda) > \hat{H}(\lambda), \ \forall \lambda > \hat{\lambda},$$
 (2.31)

while relations in (2.28) and (2.26) can be rewritten in terms of W and H as

$$-W(\lambda) + \lambda W'(\lambda) \le \frac{n}{p} \log\left(-\mathcal{C}\left(\frac{p'}{p}\right)^p \lambda^p W'(\lambda)\right), \ \lambda > 0,$$
(2.32)

and

$$-\tilde{H}(\lambda) + \lambda \tilde{H}'(\lambda) = \frac{n}{p} \log \left(-\mathcal{C}\left(\frac{p'}{p}\right)^p \lambda^p \tilde{H}'(\lambda) \right), \ \lambda > 0.$$
(2.33)

Claim (2.30) is proved once we show that $W(\lambda) \geq \tilde{H}(\lambda)$ for all $\lambda > 0$. By contradiction, we assume there exists $\lambda^{\#} > 0$ such that $W(\lambda^{\#}) < \tilde{H}(\lambda^{\#})$. Due to (2.31), $\lambda^{\#} < \tilde{\lambda}$. On the one hand, let $\lambda^* = \inf\{\lambda > \lambda^{\#} : W(\lambda) = \tilde{H}(\lambda)\}$. In particular,

$$W(\lambda) \le \tilde{H}(\lambda), \ \forall \lambda \in [\lambda^{\#}, \lambda^*].$$
 (2.34)

On the other hand, if we introduce for every $\lambda > 0$ the function $j_L^{\lambda} : (0, \infty) \to \mathbb{R}$ by

$$j_L^{\lambda}(t) = \frac{n}{p} \log\left(\mathcal{C}\left(\frac{p'}{p}\right)^p \lambda^p t\right) + \lambda t, \ t > 0,$$

relations (2.32) and (2.33) become

$$-W(\lambda) \le j_L^{\lambda}(-W'(\lambda)) \text{ and } -\tilde{H}(\lambda) = j_L^{\lambda}(-\tilde{H}'(\lambda)), \ \lambda > 0$$

By the above relations and (2.34) it yields that

$$j_L^{\lambda}(-\tilde{H}'(\lambda)) = -\tilde{H}(\lambda) \le -W(\lambda) \le j_L^{\lambda}(-W'(\lambda)), \ \forall \lambda \in [\lambda^{\#}, \lambda^*].$$

Since j_L^{λ} is increasing, it follows that $W - \tilde{H}$ is a non-increasing function on $[\lambda^{\#}, \lambda^*]$, which implies

$$0 = (W - \tilde{H})(\lambda^*) \le (W - \tilde{H})(\lambda^{\#}) < 0,$$

a contradiction. This completes the proof of (2.30).

STEP 6. We claim that

$$\limsup_{\rho \to \infty} \frac{\mathsf{m}(B_{x_0}(\rho))}{\omega_n \rho^n} \ge \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}}.$$
(2.35)

By assuming the contrary, there exists $\varepsilon_0 > 0$ such that for some $\rho_0 > 0$,

$$\frac{\mathsf{m}(B_{x_0}(\rho))}{\omega_n \rho^n} \le \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}} - \varepsilon_0, \ \forall \rho \ge \rho_0.$$

Combining the latter relation with (2.30) and (2.27), for every $\lambda > 0$ we obtain that

$$0 \leq w_{L}(\lambda) - \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}} h_{L}(\lambda)$$

$$\leq \lambda p' \int_{0}^{\rho_{0}} \mathsf{m}(B_{x_{0}}(\rho)) e^{-\lambda \rho^{p'}} \rho^{p'-1} d\rho + \lambda p' \omega_{n} \left(\left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}} - \varepsilon_{0}\right) \int_{\rho_{0}}^{\infty} e^{-\lambda \rho^{p'}} \rho^{n+p'-1} d\rho$$

$$-\lambda p' \omega_{n} \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}} \int_{0}^{\infty} e^{-\lambda \rho^{p'}} \rho^{n+p'-1} d\rho.$$

Rearranging the above inequality, by virtue of (2.5) it follows for every $\lambda > 0$ that

$$\varepsilon_0 \int_0^\infty e^{-\lambda \rho^{p'}} \rho^{n+p'-1} d\rho \le \left(1 - \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}} + \varepsilon_0\right) \int_0^{\rho_0} e^{-\lambda \rho^{p'}} \rho^{n+p'-1} d\rho$$

Due to (2.27), the latter inequality implies

$$\varepsilon_0 \frac{1}{p' \lambda^{1+\frac{n}{p'}}} \Gamma\left(\frac{n}{p'}+1\right) \le \left(1 - \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}} + \varepsilon_0\right) \frac{\rho_0^{n+p'}}{n+p'}, \ \lambda > 0.$$

Now, letting $\lambda \to 0^+$ we arrive to a contradiction. Therefore, the proof of (2.35) is concluded. Thus, Lemma 2.1 gives that

$$\frac{\mathsf{m}(B_x(\rho))}{\omega_n \rho^n} \ge \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}}, \ \forall x \in M, \ \rho > 0,$$

concluding the proof of Theorem 1.2.

2.3. Limit case II ($\alpha \to 0$): Faber-Krahn-type inequality. In this part we sketch the proof of Theorem 1.3. Similarly as before, we assume that $C > \mathcal{F}_{p,n}$.

STEP 1. Analogously to Theorem 1.1 (i), it follows that K = 0.

STEP 2. The function $x \mapsto (\lambda^{p'} - |x|^{p'})_+$ being extremal in (1.6) for every $\lambda > 0$, a direct computation shows that

$$h_F(\lambda) = \mathcal{F}_{p,n} p' \left(-h_F(\lambda) + \frac{1}{p'} \lambda h'_F(\lambda) \right)^{\frac{1}{p}} \left(\frac{1}{p'} \lambda^{1-p'} h'_F(\lambda) \right)^{1-\frac{1}{p^{\star}}}, \qquad (2.36)$$

where $h_F: (0, \infty) \to \mathbb{R}$ is given by

$$h_F(\lambda) = \int_{\mathbb{R}^n} \left(\lambda^{p'} - |x|^{p'} \right)_+ dx, \ \lambda > 0.$$

STEP 3. Let $x_0 \in M$ from $(\mathbf{D})_{x_0}^n$. Since $u_{\lambda} = \left(\lambda^{p'} - \mathsf{d}(x_0, \cdot)^{p'}\right)_+ \in \operatorname{Lip}_0(M)$, we may insert u_{λ} into $(\mathbf{FK})_{\mathcal{C}}^p$ obtaining

$$\|u_{\lambda}\|_{L^{1}} \leq \mathcal{C} \||\nabla u_{\lambda}|_{\mathsf{d}}\|_{L^{p}} \mathsf{m}(\operatorname{supp}(u_{\lambda}))^{1-\frac{1}{p^{\star}}}.$$
(2.37)

First, we observe that

$$|\nabla u_{\lambda}|_{\mathsf{d}}(x) = p'\mathsf{d}(x_0, x)^{p'-1} |\nabla \mathsf{d}(x_0, \cdot)|_{\mathsf{d}}(x) \le p'\mathsf{d}(x_0, x)^{p'-1}, \ \forall x \in B_{x_0}(\lambda)$$

while $|\nabla u_{\lambda}|_{\mathsf{d}}(x) = 0$ for every $x \notin B_{x_0}(\lambda)$. Moreover, since the spheres have zero m-measures (see Theorem 2.1), we have that

$$\mathsf{m}(\mathrm{supp}(u_{\lambda})) = \mathsf{m}(\overline{B_{x_0}(\lambda)}) = \mathsf{m}(B_{x_0}(\lambda)).$$

We now introduce the function $w_F: (0,\infty) \to \mathbb{R}$ given by

$$w_F(\lambda) = \int_M \left(\lambda^{p'} - \mathsf{d}(x_0, x)^{p'}\right)_+ d\mathsf{m}(x), \ \lambda > 0.$$

Due to the layer cake representation, one has

$$w_{F}(\lambda) = \int_{B_{x_{0}}(\lambda)} \left(\lambda^{p'} - \mathsf{d}(x_{0}, x)^{p'}\right) d\mathsf{m}(x) = \lambda^{p'} \mathsf{m}(B_{x_{0}}(\lambda)) - \int_{B_{x_{0}}(\lambda)} \mathsf{d}(x_{0}, x)^{p'} d\mathsf{m}(x)$$

$$= \lambda^{p'} \mathsf{m}(B_{x_{0}}(\lambda)) - \int_{0}^{\lambda^{p'}} \mathsf{m}\left(\{x \in B_{x_{0}}(\lambda) : \mathsf{d}(x_{0}, x)^{p'} > t\}\right) dt$$

$$= \lambda^{p'} \mathsf{m}(B_{x_{0}}(\lambda)) - p' \int_{0}^{\lambda} \left(\mathsf{m}(B_{x_{0}}(\lambda)) - \mathsf{m}(B_{x_{0}}(\rho))\right) \rho^{p'-1} d\rho \quad [\text{change } t = \rho^{p'}]$$

$$= p' \int_{0}^{\lambda} \mathsf{m}(B_{x_{0}}(\rho)) \rho^{p'-1} d\rho.$$

Therefore,

$$||u_{\lambda}||_{L^1} = w_F(\lambda), \ \mathsf{m}(\mathrm{supp}(u_{\lambda})) = \mathsf{m}(B_{x_0}(\lambda)) = \frac{1}{p'} \lambda^{1-p'} w'_F(\lambda),$$

and

$$\||\nabla u_{\lambda}|_{\mathsf{d}}\|_{L^{p}} \leq p' \left(\int_{B_{x_{0}}(\lambda)} \mathsf{d}(x_{0}, x)^{p'} d\mathsf{m}(x)\right)^{\frac{1}{p}} = p' \left(-w_{F}(\lambda) + \frac{1}{p'} \lambda w'_{F}(\lambda)\right)^{\frac{1}{p}}.$$

Consequently, inequality (2.37) takes the form

$$w_F(\lambda) \le \mathcal{C}p'\left(-w_F(\lambda) + \frac{1}{p'}\lambda w'_F(\lambda)\right)^{\frac{1}{p}} \left(\frac{1}{p'}\lambda^{1-p'}w'_F(\lambda)\right)^{1-\frac{1}{p^*}}, \ \lambda > 0,$$

which is formally (2.19) if $\alpha \to 0$ since due to (1.4), $\lim_{\alpha\to 0} \gamma = 1$ and $\lim_{\alpha\to 0} \frac{1-\gamma}{\alpha p} = 1 - \frac{1}{p^*}$. Therefore, we may proceed as in the proof of Theorem 1.1 (ii) (Steps 4-6), proving that

$$\lim_{\lambda \to 0^+} \frac{w_F(\lambda)}{h_F(\lambda)} = 1,$$
$$w_F(\lambda) \ge \left(\frac{\mathcal{F}_{p,n}}{\mathcal{C}}\right)^n h_F(\lambda), \ \forall \lambda > 0$$

and finally

$$\frac{\mathsf{m}(B_x(\rho))}{\omega_n \rho^n} \ge \left(\frac{\mathcal{F}_{p,n}}{\mathcal{C}}\right)^n, \ \forall x \in M, \ \rho > 0,$$

which concludes the proof of Theorem 1.3.

3. RIGIDITY RESULTS IN SMOOTH SETTINGS

As a starting point, we need an Aubin-Hebey-type result (see [3] and [11]) for Gagliardo-Nirenberg inequalities which is valid on generic Riemannian manifolds.

Lemma 3.1. Let (M, g) be a complete n-dimensional Riemannian manifold and C > 0. The following statements hold:

- (i) If $(\mathbf{GN1})^{\alpha,p}_{\mathcal{C}}$ holds on (M,g) for some $p \in (1,n)$ and $\alpha \in (1,\frac{n}{n-p}]$ then $\mathcal{C} \geq \mathcal{G}_{\alpha,p,n}$;
- (ii) If $(\mathbf{GN2})^{\alpha,p}_{\mathcal{C}}$ holds on (M,g) for some $p \in (1,n)$ and $\alpha \in (0,1)$ then $\mathcal{C} \geq \mathcal{N}_{\alpha,p,n}$;
- (iii) If $(\mathbf{LS})^p_{\mathcal{C}}$ holds on (M, g) for some $p \in (1, n)$ then $\mathcal{C} \geq \mathcal{L}_{p,n}$;
- (iv) If $(\mathbf{FK})^p_{\mathcal{C}}$ holds on (M,g) for some $p \in (1,n)$ then $\mathcal{C} \geq \mathcal{F}_{p,n}$.

Proof. (i) By contradiction, we assume that $(\mathbf{GN1})_{\mathcal{C}}^{\alpha,p}$ holds on (M,g) for some $p \in (1,n)$, $\alpha \in (1, \frac{n}{n-p}]$, and $\mathcal{C} < \mathcal{G}_{\alpha,p,n}$. Let $x_0 \in M$ be fixed arbitrarily. For every $\varepsilon > 0$, there exists a local chart (Ω, ϕ) of M at the point x_0 and a number $\delta > 0$ such that $\phi(\Omega) = B_0(\delta) = \{\tilde{x} \in \mathbb{R}^n : |\tilde{x}| < \delta\}$ and the components $g_{ij} = g_{ij}(x)$ of the Riemannian metric g on (Ω, ϕ) satisfy

$$(1-\varepsilon)\delta_{ij} \le g_{ij} \le (1+\varepsilon)\delta_{ij} \tag{3.1}$$

in the sense of bilinear forms. Since $(\mathbf{GN1})_{\mathcal{C}}^{\alpha,p}$ is valid, relation (3.1) shows that for every $\varepsilon > 0$ small enough, there exists $\delta_{\varepsilon} > 0$ and $\mathcal{C}_{\varepsilon} \in (\mathcal{C}, \mathcal{G}_{\alpha,p,n})$ such that for every $\delta \in (0, \delta_{\varepsilon})$ and $v \in \operatorname{Lip}_0(B_0(\delta))$,

$$\|v\|_{L^{\alpha p}(B_{0}(\delta),dx)} \leq C_{\varepsilon} \|\nabla v\|_{L^{p}(B_{0}(\delta),dx)}^{\theta} \|v\|_{L^{\alpha(p-1)+1}(B_{0}(\delta),dx)}^{1-\theta}.$$
(3.2)

Let us fix $u \in \operatorname{Lip}_0(\mathbb{R}^n)$ arbitrarily and set $v_{\lambda}(x) = \lambda^{\frac{n}{p}} u(\lambda x), \lambda > 0$. For $\lambda > 0$ large enough, one has $v_{\lambda} \in \operatorname{Lip}_0(B_0(\delta))$. If we plug in v_{λ} into (3.2), by using the scaling properties

$$\|\nabla v_{\lambda}\|_{L^{p}(B_{0}(\delta),dx)} = \lambda \|\nabla u\|_{L^{p}(\mathbb{R}^{n},dx)} \quad \text{and} \quad \|v_{\lambda}\|_{L^{q}(B_{0}(\delta),dx)} = \lambda^{\frac{n}{p}-\frac{n}{q}} \|u\|_{L^{q}(\mathbb{R}^{n},dx)}, \quad \forall q > 0,$$

$$(3.3)$$

and the form of the number θ (see (1.2)), it follows that

$$\|u\|_{L^{\alpha p}(\mathbb{R}^n, dx)} \leq \mathcal{C}_{\varepsilon} \|\nabla u\|_{L^p(\mathbb{R}^n, dx)}^{\theta} \|u\|_{L^{\alpha(p-1)+1}(\mathbb{R}^n, dx)}^{1-\theta}$$

If we insert the extremal function $h_{\alpha,p}^{\lambda}$ of the optimal Gagliardo-Nirenberg inequality on \mathbb{R}^n ($\alpha > 1$) into the latter relation, Theorem A yields that $\mathcal{G}_{\alpha,p,n} \leq \mathcal{C}_{\varepsilon}$, a contradiction.

The proofs of (ii) (iii) and (iv) are analogous to (i), taking into account in addition to (3.3) that

$$\operatorname{Ent}_{dx}(|v_{\lambda}|^{p}) = \operatorname{Ent}_{dx}(|u|^{p}) + n \|u\|_{L^{p}}^{p} \log \lambda,$$

and

$$\mathcal{H}^{n}(\operatorname{supp}(v_{\lambda})) = \lambda^{-n} \mathcal{H}^{n}(\operatorname{supp}(u)),$$

respectively.

CES

3.1. Gagliardo-Nirenberg inequalities on Riemannian manifolds with Ricci ≥ 0 . Before presenting the proofs of Theorem 1.4 and Corollary 1.1, we recall some results from Munn [17].

To do this, let (M, g) be an $n \geq 2$ -dimensional complete Riemannian manifold with non-positive Ricci curvature endowed with its canonical volume element dv_g . The asymptotic volume growth of (M, g) is defined by

$$\operatorname{AVG}_{(M,g)} = \lim_{r \to \infty} \frac{\operatorname{Vol}_g(B_x(r))}{\omega_n r^n}.$$

By Bishop-Gromov comparison theorem it follows that $AVG_{(M,g)} \leq 1$ and this number is independent of the point $x \in M$.

Given $k \in \{1, ..., n\}$, let us denote by $\delta_{k,n} > 0$ the smallest positive solution to the equation $10^{k+2}C_{k,n}(k)s\left(1+\frac{s}{2k}\right)^k = 1$ in variable s, where

$$C_{k,n}(i) = \begin{cases} 1 & \text{if } i = 0, \\ 3 + 10C_{k,n}(i-1) + (16k)^{n-1}(1 + 10C_{k,n}(i-1))^n & \text{if } i \in \{1, \dots, k\} \end{cases}$$

We now consider the smooth, bijective and increasing function $h_{k,n}: (0, \delta_{k,n}) \to (1, \infty)$ defined by

$$h_{k,n}(s) = \left[1 - 10^{k+2} C_{k,n}(k) s \left(1 + \frac{s}{2k}\right)^k\right]^{-1}$$

For every s > 1, let

$$\beta(k,s,n) = \begin{cases} 1 - \left[1 + \frac{s^n}{[h_{1,n}^{-1}(s)]^n}\right]^{-1} & \text{if } k = 1, \\ \max\left\{\beta(1,s,n), \beta(i,1 + \frac{h_{k,n}^{-1}(s)}{2k},n) : i = 1, \dots, k-1\right\} & \text{if } k \in \{2,\dots,n\} \end{cases}$$

Note that the constant $\beta(k, s, n)$, which is used to prove the Perelman's maximal volume lemma, denotes the minimum volume growth of (M, g) needed to guarantee that any continuous map $f : \mathbb{S}^k \to B_x(\rho)$ has a continuous extension $g : \mathbb{D}^{k+1} \to B_x(c\rho)$, where $\mathbb{D}^{k+1} = \{y \in \mathbb{R}^{k+1} : |y| \leq 1\}$ and $\mathbb{S}^k = \partial \mathbb{D}^{k+1}$, see [17, Definition 3.3]. Finally, the Munn-Perelman constant is defined as

$$\alpha_{MP}(k,n) = \inf_{s \in (1,\infty)} \beta(k,s,n).$$

By construction, $\alpha_{MP}(k, n)$ is non-decreasing in k; for numerical values of $\alpha_{MP}(k, n)$ one can consult [17, Appendix A].

Proof of Theorem 1.4. Let (M, g) be an *n*-dimensional complete Riemannian manifold with nonnegative Ricci curvature $(n \ge 2)$ and assume the L^p -logarithmic Sobolev inequality $(\mathbf{LS})^p_{\mathcal{C}}$ holds on (M, g) for some $p \in (1, n)$ and $\mathcal{C} > 0$.

(i) It follows from Lemma 3.1 (iii), i.e., $C \geq \mathcal{L}_{p,n}$.

(ii) Anderson [2] and Li [14] stated that if there exists $c_0 > 0$ such that $\operatorname{Vol}_g(B_x(\rho)) \ge c_0 \omega_n \rho^n$ for every $\rho > 0$, then (M, g) has finite fundamental group $\pi_1(M)$ and its order is bounded above by c_0^{-1} . Thus it remains to apply Theorem 1.2.

(iii) Assume that $\mathcal{C} < \alpha_{MP}(k_0, n)^{-\frac{p}{n}} \mathcal{L}_{p,n}$ for some $k_0 \in \{1, ..., n\}$. By Theorem 1.2, we have that

$$\operatorname{AVG}_{(M,g)} = \lim_{r \to \infty} \frac{\operatorname{Vol}_g(B_x(r))}{\omega_n r^n} \ge \left(\frac{\mathcal{L}_{p,n}}{\mathcal{C}}\right)^{\frac{n}{p}} > \alpha_{MP}(k_0, n) \ge \dots \ge \alpha_{MP}(1, n).$$

By Munn [17, Theorem 1.2], it follows that $\pi_1(M) = ... = \pi_{k_0}(M) = 0$.

(iv) If $\mathcal{C} < \alpha_{MP}(n,n)^{-\frac{p}{n}} \mathcal{L}_{p,n}$, then $\pi_1(M) = \dots = \pi_n(M) = 0$, which implies the contractibility of M, see e.g. Luft [16].

(v) If $C = \mathcal{L}_{p,n}$ then by Theorem 1.2 and the Bishop-Gromov volume comparison theorem follows that $\operatorname{Vol}_g(B_x(\rho)) = \omega_n \rho^n$ for every $x \in M$ and $\rho > 0$. The equality in Bishop-Gromov theorem implies that (M, g) is isometric to the Euclidean space \mathbb{R}^n . The converse trivially holds.

Remark 3.1. In the study of heat kernel bounds on an n-dimensional complete Riemannian manifold (M, g) with non-negative Ricci curvature, the logarithmic Sobolev inequality

$$\operatorname{Ent}_{dv_g}(u^2) \le \frac{n}{2} \log \left(C \|\nabla_g u\|_{L^2(M, dv_g)}^2 \right), \ \forall u \in C_0^\infty(M), \ \|u\|_{L^2} = 1,$$
(3.4)

plays a central role, C > 0. In fact, (3.4) is equivalent to an upper bound of the heat kernel $p_t(x,y)$ on M, i.e.,

$$\sup_{x,y \in M} p_t(x,y) \le \tilde{C}t^{-\frac{n}{2}}, \ t > 0,$$
(3.5)

for some C > 0. According to Theorem B (from §1.1), the optimal constant in (3.4) in the Euclidean space \mathbb{R}^n is given by $C = \mathcal{L}_{n,2} = \frac{2}{n\pi e}$; this scale invariant form on \mathbb{R}^n can be deduced by Gross [10] logarithmic Sobolev inequality

$$\operatorname{Ent}_{d\gamma_n}(u^2) \le 2 \|\nabla u\|_{L^2(\mathbb{R}^n, d\gamma_n)}^2, \ \forall u \in C_0^{\infty}(\mathbb{R}^n), \ \|u\|_{L^2(\mathbb{R}^n, d\gamma_n)} = 1,$$

where the canonical Gaussian measure γ_n has the density $\delta_n(x) = (2\pi)^{-\frac{n}{2}} e^{-\frac{|x|^2}{2}}$, $x \in \mathbb{R}^n$, see Weissler [27]. Sharp estimates on the heat kernel shows that on a complete Riemannian manifold (M, g)with non-negative Ricci curvature the L^2 -logarithmic Sobolev inequality (3.4) holds with the optimal Euclidean constant $C = \mathcal{L}_{n,2} = \frac{2}{n\pi e}$ if and only if (M,g) is isometric to \mathbb{R}^n , cf. Bakry, Concordet and Ledoux [4], Ni [18], and Li [14]. In this case, $\tilde{C} = (4\pi)^{-\frac{n}{2}}$ in (3.5).

In particular, Theorem 1.4 (v) gives a positive answer to the open problem of C. Xia [29] concerning the validity of the optimal L^p -logarithmic Sobolev inequality for generic $p \in (1, n)$ in the same geometric context as above. Xia's formulation was deeply motivated by the lack of sharp L^p -estimates $(p \neq 2)$ for the heat kernel on Riemannian manifolds with non-negative Ricci curvature.

Similar results to Theorem 1.4 can be stated for the other three Gagliardo-Nirenberg-type inequalities; here we formulate one for $(\mathbf{GN1})^{\alpha,p}_{\mathcal{C}}$, the other two inequalities are left to the reader.

Theorem 3.1. Let (M, q) be an n-dimensional complete Riemannian manifold with non-negative Ricci curvature $(n \geq 2)$ and assume the $(\mathbf{GN1})^{\alpha,p}_{\mathcal{C}}$ holds on (M,g) for some $p \in (1,n), \alpha \in (1, \frac{n}{n-p}]$ and C > 0. Then the following assertions hold:

- (i) $\mathcal{C} \geq \mathcal{G}_{\alpha,p,n}$;
- (ii) The order of the fundamental group $\pi_1(M)$ is bounded above by $\left(\frac{\mathcal{C}}{\mathcal{G}_{\alpha,p,n}}\right)^{\frac{n}{\theta}}$;
- (iii) If $\mathcal{C} < \alpha_{MP}(k_0, n)^{-\frac{\theta}{n}} \mathcal{G}_{\alpha, p, n}$ for some $k_0 \in \{1, ..., n\}$ then $\pi_1(M) = ... = \pi_{k_0}(M) = 0;$
- (iv) If $\mathcal{C} < \alpha_{MP}(n,n)^{-\frac{\theta}{n}} \mathcal{G}_{\alpha,p,n}$ then M is contractible; (v) $\mathcal{C} = \mathcal{G}_{\alpha,p,n}$ if and only if (M,g) is isometric to the Euclidean space \mathbb{R}^n .

3.2. Gagliardo-Nirenberg inequalities on Finsler manifolds with n-Ricci > 0. Let M be a connected *n*-dimensional C^{∞} -manifold and $TM = \bigcup_{x \in M} T_x M$ be its tangent bundle. The pair (M, F) is called a *reversible Finsler manifold* if a continuous function $F: TM \longrightarrow [0, \infty)$ satisfies the conditions:

- (a) $F \in C^{\infty}(TM \setminus \{0\});$
- (b) F(x,tv) = |t|F(x,v) for all $t \in \mathbb{R}$ and $(x,v) \in TM$;

(b) P(x, v) = |v|P(x, v) for all $v \in \mathbb{R}$ and $(x, v) \in PM$, (c) the $n \times n$ matrix $g_{ij}(x, v) = \frac{1}{2} \frac{\partial^2 (F^2)}{\partial v^i \partial v^j}(x, v)$ is positive definite for all $(x, v) \in TM \setminus \{0\}$. Here $v = \sum_{i=1}^{n} v^i \frac{\partial}{\partial x^i}$, and we shall denote by g_v the inner product on $T_x M$ induced by the above form. If $g_{ij}(x) = g_{ij}(x, v)$ is independent of v then (M, F) is called *Riemannian manifold*. A *Minkowski* space consists of a finite dimensional vector space V and a Minkowski norm which induces a Finsler metric on V by translation, i.e., F(x, v) is independent of x. A Finsler manifold (M, F) is called a locally Minkowski space if every point in M admits a local coordinate system (x^i) on its neighborhood such that F(x, v) depends only on v and not on x.

We consider on the pull-back bundle π^*TM the *Chern connection*, see Bao, Chern and Shen [5, Theorem 2.4.1]. The coefficients of the Chern connection are denoted by Γ^i_{ik} , which are instead of the well-known Christoffel symbols from Riemannian geometry. A Finsler manifold is of Berwald type if the coefficients $\Gamma_{ij}^k(x,v)$ in natural coordinates are independent of v. It is clear that Riemannian manifolds and (locally) Minkowski spaces are Berwald spaces. The Chern connection induces in a natural manner on π^*TM the curvature tensor R, see Bao, Chern and Shen [5, Chapter 3]. By means of the connection, we also have the covariant derivative $D_v u$ of a vector field u in the direction $v \in T_x M$. Note that $v \mapsto D_v u$ is not linear. A vector field u = u(t) along a curve σ is parallel if $D_{\dot{\sigma}} u = 0$. A C^{∞} curve $\sigma : [0, a] \to M$ is a geodesic if $D_{\dot{\sigma}} \dot{\sigma} = 0$. Geodesics are considered to be parametrized proportionally to arc-length. The Finsler manifold is complete if every geodesic segment can be extended to \mathbb{R} . For a C^{∞} -curve $\sigma : [0, l] \longrightarrow M$, its integral length is given by $L_F(\sigma) := \int_0^l F(\sigma(t), \dot{\sigma}(t)) dt$. Define the distance function $d_F : M \times M \longrightarrow [0, \infty)$ by

$$d_F(x_1, x_2) = \inf_{\sigma} L_F(\sigma),$$

where σ runs over all C^{∞} -curves from x_1 to x_2 . Geodesics locally minimize d_F -distances.

Let $u, v \in T_x M$ be two non-collinear vectors and $S = \operatorname{span}\{u, v\} \subset T_x M$. By means of the curvature tensor R, the flag curvature of the flag $\{S, v\}$ is defined by

$$K(\mathcal{S};v) = \frac{g_v(R(U,V)V,U)}{g_v(V,V)g_v(U,U) - g_v(U,V)^2}$$

where $U = (v; u), V = (v; v) \in \pi^* TM$. If (M, F) is Riemannian, the flag curvature reduces to the well known sectional curvature.

Let $v \in T_x M$ be such that F(x, v) = 1 and let $\{e_i\}_{i=1,...,n}$ with $e_n = v$ be a basis for $T_x M$ such that $\{(v; e_i)\}_{i=1,...,n}$ is an orthonormal basis for π_*TM . Let $\mathcal{S}_i = \operatorname{span}\{e_i, v\}, i = 1, ..., n - 1$. The *Ricci curvature* Ric: $TM \to \mathbb{R}$ is defined by $\operatorname{Ric}(cv) = c^2 \sum_{i=1}^{n-1} K(\mathcal{S}_i; v)$ for every c > 0.

Let (M, F) be an *n*-dimensional complete Finsler manifold and let m be an arbitrarily positive smooth measure on M; such a manifold is viewed as a regular metric measure space and we denote it by (M, F, m). Let $v \in T_x M$ be such that F(x, v) = 1 and let

$$\Upsilon(v) = \log\left(\frac{\operatorname{vol}_{g_v}(\mathbf{B}(0,1))}{\mathsf{m}_x(\mathbf{B}(0,1))}\right),\,$$

where vol_{g_v} and m_x denote the Lebesgue measures on $T_x M$ induced by g_v and m , respectively, while $\mathrm{B}(0,1) = \{y \in T_x M : F(x,y) < 1\}$ is the unit tangent ball at $T_x M$. The latter relation can be rewritten into the more familiar form $\mathsf{m}_x(\mathrm{B}(0,1)) = e^{-\Upsilon(v)} \operatorname{vol}_{g_v}(\mathrm{B}(0,1))$. We introduce the notation

$$\partial_{v}\Upsilon = \frac{d}{dt}\Upsilon(\dot{\sigma}(t))\big|_{t=0},\tag{3.6}$$

where $\sigma : (-\varepsilon, \varepsilon) \to M$ is the geodesic with $\sigma(0) = x$ and $\dot{\sigma}(0) = v$. We say that the space (M, F, m) has n-Ricci curvature bounded below by $K \in \mathbb{R}$ if and only if $\operatorname{Ric}(v) \geq K$ and $\partial_v \Upsilon = 0$ for every $v \in T_x M$ such that F(x, v) = 1, see Ohta [19, Theorem 1.2] and Ohta and Sturm [21, Definition 5.1]. Note that a Berwald space endowed with the Busemann-Hausdorff measure m_{BH} (and inducing the volume form dV_F) verifies the property $\partial_v \Upsilon \equiv 0$, see Shen [23, Propositions 2.6 & 2.7].

The polar transform of F is defined for every $(x, \alpha) \in T^*M$ by

$$F^*(x,\alpha) = \sup_{v \in T_x M \setminus \{0\}} \frac{\alpha(v)}{F(x,v)}.$$
(3.7)

Note that, for every $x \in M$, the function $F^*(x, \cdot)$ is a Minkowski norm on $T^*_r M$.

If $u \in \text{Lip}_0(M)$, then relation (1.7) can be interpreted as

$$|\nabla u|_{d_F}(x) = F^*(x, Du(x)) \text{ for a.e. } x \in M,$$
(3.8)

where $Du(x) \in T_x^*(M)$ is the distributional derivative of u at $x \in M$, see Ohta and Sturm [21]. In particular, if (M, F) = (M, g) is a Riemannian manifold, then $|\nabla u|_{d_g} = |\nabla_g u|$, where d_g is the distance function on (M, g), ∇_g is the Riemannian gradient on (M, g), and $|\cdot|$ is the norm coming from the Riemannian metric g, respectively.

Although a slightly more general result can be proved, we present an application on Berwald spaces (M, F) endowed with the canonical Busemann-Hausdorff measure m_{BH} (and its induced volume form dV_F), by exploring the results of Cordero-Erausquin, Nazaret and Villani [6] and Gentil [9] (see Theorems A & B).

Theorem 3.2. [Optimality vs. flatness] Let (M, F) be an *n*-dimensional complete reversible Berwald space with non-negative Ricci curvature. The following statements are equivalent:

- (i) $(\mathbf{GN1})_{\mathcal{G}_{\alpha,p,n}}^{\alpha,p}$ holds on (M,F) for some $p \in (1,n)$ and $\alpha \in (1,\frac{n}{n-p}]$; (ii) $(\mathbf{GN2})_{\mathcal{N}_{\alpha,p,n}}^{\alpha,p}$ holds on (M,F) for some $p \in (1,n)$ and $\alpha \in (0,1)$;
- (iii) $(\mathbf{LS})^p_{\mathcal{L}_{p,n}}$ holds on (M, F) for some $p \in (1, n)$;
- (iv) $(\mathbf{FK})_{\mathcal{F}_{n,n}}^{p}$ holds on (M, F) for some $p \in (1, n)$;
- (v) (M, F) is isometric to an *n*-dimensional Minkowski space.

Proof. We divide the proof into two parts.

 $(i) \lor (ii) \lor (iii) \lor (iv) \Rightarrow (v)$. Note that the Busemann-Hausdorff measure m_{BH} satisfies the *n*-density assumption for every $x \in M$, i.e.,

$$\lim_{\rho \to 0} \frac{\mathsf{m}_{BH}(B_x(\rho))}{\omega_n \rho^n} = 1,$$

see Shen [23, Lemma 5.2]. Since (M, F) is a Berwald space (thus $\partial_v \Upsilon \equiv 0$ for every $v \in T_x M, x \in M$, see (3.6), the non-negativity of the Ricci curvature on (M, F) coincides with the non-negativity of the *n*-Ricci curvature on $(M, d_F, \mathsf{m}_{BH})$, thus the metric measure space $(M, d_F, \mathsf{m}_{BH})$ satisfies the curvature-dimension condition CD(0,n), see Ohta [19]. Moreover, the completeness of (M,F) via Hopf-Rinow theorem implies that the $(M, d_F, \mathsf{m}_{BH})$ is proper. Applying now any of the Theorems 1.1, 1.2 or 1.3 (according to which of the assumptions (i), (ii), (iii) or (iv) is satisfied), it yields that

$$\mathsf{m}_{BH}(B_x(\rho)) \ge \omega_n \rho^n$$
 for all $x \in M, \ \rho \ge 0$.

By the generalized Bishop-Gromov theorem on Finsler manifolds and the n-density property we also have the reverse inequality, thus

$$\mathbf{m}_{BH}(B_x(\rho)) = \omega_n \rho^n \text{ for all } x \in M, \ \rho \ge 0.$$
(3.9)

The latter relation immediately implies that the flag curvature on (M, F) is identically zero, see Ohta [19, Theorem 7.3], and Kristály and Ohta [12, Theorem 3.3]. Due to Bao, Chern and Shen [5, Section 10.5]), every Berwald space with zero flag curvature is necessarily a locally Minkowski space. By (3.9) it follows that (M, F) is actually isometric to a Minkowski space.

 $(\mathbf{v}) \Rightarrow (\mathbf{i}) \land (\mathbf{i}\mathbf{i}) \land (\mathbf{i}\mathbf{i}\mathbf{i}) \land (\mathbf{i}\mathbf{v})$. Let us fix an arbitrary norm $\|\cdot\|$ on \mathbb{R}^n , and let $\Phi: (M, F) \rightarrow (\mathbb{R}^n, \|\cdot\|)$ be an isometry. Then

$$F(x,y) = \|d\Phi_x(y)\|, \ x \in M, y \in T_x M,$$

and a simple computation based on the definition of the polar transform (see (3.7)) gives

$$F^*(x,\alpha) = \|\alpha d\Phi_{\Phi(x)}^{-1}\|_*, \ x \in M, \alpha \in T^*_x M.$$
(3.10)

If we consider the change of variables $\tilde{x} = \Phi(x)$, relations (3.8) and (3.10) imply

$$|\nabla v|_{d_F}(x) = F^*(x, Dv(x)) = \|(D(v \circ \Phi^{-1})(\tilde{x}))\|_*, \ v \in C_0^\infty(M), \ x \in M.$$
(3.11)

Thus, for every $v \in C_0^{\infty}(M)$, $p \in (1, n)$ and q > 0, we have

$$\begin{split} \|D(v \circ \Phi^{-1})\|_{L^{p}(\mathbb{R}^{n}, d\tilde{x})} &= \left(\int_{\mathbb{R}^{n}} \|(D(v \circ \Phi^{-1})(\tilde{x}))\|_{*}^{p} d\tilde{x}\right)^{\frac{1}{p}} = \left(\int_{M} (|\nabla v|_{d_{F}}(x))^{p} dV_{F}(x)\right)^{\frac{1}{p}} \\ &= \||\nabla v|_{d_{F}}\|_{L^{p}(M, dV_{F})}, \end{split}$$

 $\operatorname{Ent}_{d\tilde{x}}(|v \circ \Phi^{-1}|^p) = \operatorname{Ent}_{dV_F}(|v|^p)$ and $||v \circ \Phi^{-1}||_{L^q} = ||v||_{L^q}$.

It remains to apply the results of Cordero-Erausquin, Nazaret and Villani [6] and Gentil [9] (cf. Theorems A & B) for $u = v \circ \Phi^{-1}$. Acknowledgments. The author is grateful to the Universität Bern for the warm hospitality where this work has been initiated. He thanks Professor Zoltán M. Balogh and Professor Cédric Villani for stimulating conversations on the L^p -logarithmic Sobolev inequality and Professor Michel Ledoux for suggesting the study of the whole family of Gagliardo-Nirenberg inequalities. Research supported by a grant of the Romanian National Authority for Scientific Research, CNCS-UEFISCDI, "Symmetries in elliptic problems: Euclidean and non-Euclidean techniques", project no. PN-II-ID-PCE-2011-3-0241, and by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences.

References

- A. Alvino, V. Ferone, P.-L. Lions, G. Trombetti, Convex symmetrization and applications. Ann. Inst. H. Poincaré Anal. Non Linéaire 14 (1997), no. 2, 275–293.
- [2] M. Anderson, On the topology of complete manifold of nonnegative Ricci curvature. Topology 3 (1990), 41–55.
- [3] T. Aubin, Problèmes isopérimétriques et espaces de Sobolev. J. Differential Geometry 11 (1976), no. 4, 573–598.
- [4] D. Bakry, D. Concordet, M. Ledoux, Optimal heat kernel bounds under logarithmic Sobolev inequalities. ESAIM Probab. Statist. 1 (1995/97), 391–407.
- [5] D. Bao, S. S. Chern, Z. Shen, Introduction to Riemann-Finsler Geometry, Graduate Texts in Mathematics, 200, Springer Verlag, 2000.
- [6] D. Cordero-Erausquin, B. Nazaret, C. Villani, A mass-transportation approach to sharp Sobolev and Gagliardo-Nirenberg inequalities. Adv. Math. 182 (2004), no. 2, 307–332.
- [7] M. Del Pino, J. Dolbeault, The optimal Euclidean L^p-Sobolev logarithmic inequality. J. Funct. Anal. 197 (2003) 151–161.
- [8] M. P. do Carmo, C. Xia, Complete manifolds with non-negative Ricci curvature and the Caffarelli-Kohn-Nirenberg inequalities. Compos. Math. 140 (2004), 818–826.
- [9] I. Gentil, The general optimal L^p-Euclidean logarithmic Sobolev inequality by Hamilton-Jacobi equations. J. Funct. Anal. 202 (2003), no. 2, 591–599.
- [10] L. Gross, Logarithmic Sobolev inequalities. Amer. J. Math. 97 (1975) 1061–1083.
- [11] E. Hebey, Nonlinear analysis on manifolds: Sobolev spaces and inequalities. Courant Lecture Notes in Mathematics, 5. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999.
- [12] A. Kristály, S. Ohta, Caffarelli-Kohn-Nirenberg inequality on metric measure spaces with applications. Math. Ann. 357 (2013), no. 2, 711–726.
- [13] M. Ledoux, On manifolds with non-negative Ricci curvature and Sobolev inequalities. Comm. Anal. Geom. 7 (1999), no. 2, 347–353.
- [14] P. Li, Large time behavior of the heat equation on complete manifolds with nonnegative Ricci curvature. Ann. of Math. (2) 124 (1986), no. 1, 1–21.
- [15] J. Lott, C. Villani, Ricci curvature for metric measure spaces via optimal transport. Ann. of Math. 169 (3) (2009), 903–991.
- [16] E. Luft, On contractible open topological manifolds. Invent. Math. 4 (1967), 192–201.
- [17] M. Munn, Volume growth and the topology of manifolds with nonnegative Ricci curvature. J. Geom. Anal. 20 (2010), no. 3, 723–750.
- [18] L. Ni, The entropy formula for linear heat equation. J. Geom. Anal. 14 (2004), no. 1, 87–100.
- [19] S. Ohta, Finsler interpolation inequalities. Calc. Var. Partial Differential Equations 36 (2009) 211–249.
- [20] S. Ohta, On the measure contraction property of metric measure spaces. Comment. Math. Helv. 82 (2007), 805–828.
- [21] S. Ohta, K.-T. Sturm, Heat flow on Finsler manifolds. Comm. Pure Appl. Math. 62 (2009), no. 10, 1386–1433.
- [22] G. Perelman, Manifolds of positive Ricci curvature with almost maximal volume. J. Am. Math. Soc. 7 (1994), 299–305.
- [23] Z. Shen, Volume comparison and its applications in Riemann-Finsler geometry. Adv. Math. 128 (1997), 306–328.
- [24] K.-T. Sturm, On the geometry of metric measure spaces. I, Acta Math. 196 (1) (2006) 65–131.
- [25] K.-T. Sturm, On the geometry of metric measure spaces. II, Acta Math. 196 (1) (2006) 133–177.
- [26] G. Talenti, Best constant in Sobolev inequality. Ann. Mat. Pura Appl. 110 (1976), 353–372.
- [27] F. B. Weissler, Logarithmic Sobolev inequalities for the heat-diffusion semigroup. Trans. Amer. Math. Soc. 237 (1978), 255–269.
- [28] C. Xia, The Gagliardo-Nirenberg inequalities and manifolds of non-negative Ricci curvature. J. Funct. Anal. 224 (2005), no. 1, 230–241.
- [29] C. Xia, Sobolev type inequalities on complete Riemannian manifolds, http://www.official.kotaroy.com/meeting/symposium/abstract.bak/02-xia.pdf

DEPARTMENT OF ECONOMICS, BABEŞ-BOLYAI UNIVERSITY, 400591 CLUJ-NAPOCA, ROMANIA & INSTITUTE OF AP-PLIED MATHEMATICS, ÓBUDA UNIVERSITY, 1034 BUDAPEST, HUNGARY

E-mail address: alex.kristaly@econ.ubbcluj.ro; alexandrukristaly@yahoo.com