Sleep, Neuroengineering and Dynamics

Jens Christian Claussen · Ulrich G. Hofmann

Cognitive Neurodynamics 6 (3), 211-214 (2012) DOI: 10.1007/s11571-012-9204-2 Received: 23.04.2012

Cognitive Neurodynamics 6 (3), 211-214 (2012)

Abstract Modeling of consciousness-related phenomena and neuroengineering are fields that are rapidly growing together. We review recent approaches and developments and point out some promising directions of future research: Understanding the dynamics of consciousness states and associated oscillations, pathological oscillations as well as their treatment by stimulation, neuroprosthetics and brain-computer-interface approaches, and stimulation approaches that probe, influence and strengthen memory consolidation. In all these fields, computational models connect theory, neurophysiology and neuroengineering research and pave a way towards medical applications.

 $\label{eq:computational neuroscience} \begin{array}{l} \textbf{Keywords} \ sleep \ \cdot \ neuroengineering \ \cdot \ dynamics \ \cdot \ computational neuroscience \ \cdot \ brain-computer interface \ \cdot \ neuroproschetics \ \cdot \ pathological \ oscillations \ \cdot \ plasticity \ \cdot \ consciousness \end{array}$

Retweeting John von Neumann's words from the 1950s, "All stable processes we shall predict. All unstable processes we shall control.", it becomes immediate how computational neuroscience might form a basis for novel

Jens Christian Claussen University of Luebeck Institute for Neuro- and Bioinformatics D-23538 Lübeck E-mail: claussen@inb.uni-luebeck.de

Ulrich Hofmann Neuro-Electronic Systems University Clinic Freiburg D-79095 Freiburg E-mail: ulrich.hofmann@uniklinik-freiburg.de engineering approaches in neural medicine. Brain modeling and neural engineering are, in footsteps of these words, aiming at understanding the brain and its emerging phenomena on the scientific side, and interacting with the brain – also with the perspective of medical treatment – from the engineering science side. Both sides of the theory-experiment coin have always been connected as, e.g., electrodes are used for data acquisition as well as to influence neural systems, be it only by well-defined input pulses of a measurement protocol [Bi and Poo 2001], [Siegelbaum and Kandel 1991].

This roadmap, however, persistently is a difficult enterprise for various reasons. The brain is a highly nonlinear dynamical system and remains a challenge for data analysis, theoretical modeling, and large-scale computation [Singer 1999], [Kantz and Schreiber 1997], [Pikovsky et al. 2001], [Lehnertz et al. 2000], [Olbrich et al. 2011].

In review of these challenges, we would like to emphasize on four areas of research that mark out outstanding future potential, and are related to central keywords: Consciousness, Pathological Oscillations, Neuroprosthetics, and Neuroenhancement. In these four areas, computational models in a similar way connect natural sciences and neuroengineering research and pave a way towards medical applications.

Understanding consciousness, anaesthesia and sleep. Large-scale synchronized oscillations are easily observed experimentally and can be both of physiological and pathological character. The mammal sleep-wake cycle is a remarkably robust oscillation, in whose regulation again various neural oscillations are involved, including the cortical slow oscillation in the so-called delta band,

with frequencies around 1 Hz [Compte et al. 2003], [Ngo et al. 2010], [Mattia and Sanchez-Vives 2012]. Electrical stimulation of the brain at this slow wave frequency enhanced the slow oscillations themselves as well as increased their memory consolidation effect [Marshall et al. 2006]. The slow oscillations, comprised by the interplay between bursting activity and an activitydependent self-inhibition, exhibit a specific anticorrelation in the durations of Up and Down states; as Mattia and Sanchez-Vives show in comparison of ferret brain slice data, mean-field models, and simulations [Mattia and Sanchez-Vives 2012]. Mean-field models describe the gross activity of a neural subpopulation (at column level or below) but keep track of main types of neurons and their simplified connectivity. They are of great advantage in describing the consciousness transi-

tions of sleep and general anaesthesia [Hutt et al. 2012, Steyn-Ross et al. 2012]. As modeled by [Hutt et al. 2012]. GABAergic tonic inhibition influences the brain's arousal system during general anaesthesia including a loss of consciousness. Including the effect of gap junctions on the dynamics elucidated the influence on the propensity of generalized seizures [Stevn-Ross et al. 2012].

Medical treatment: pathological oscillations. – It is only a recent development that medicine has spotted the essential importance of dynamical phenomena for understanding and treatment of certain diseases. These are primarily those where oscillations themselves comprise the disease as in movement disorders, namely essential tremor and Parkinson's disease, for which electrical stimulation methods co-developed with theoretical work and computer simulations have found their way into clinical practice [Tass et al. 2006]. But even when there is no observable mechanical or electrical oscillation, as in the cortical spreading depression which is comprised by slow $(10^2-10^3 \text{s timescale}) \text{ Ca}^{2+}$ waves, control methods may become means of treatment [Dahlem et al. 2008]. In epilepsy, and in more severe cases of mood disorders, deep brain stimulation is applied [Abelson et al. 2005] and sophisticated technical

implementations as radio stimulation are developed [Delgado et al. 1968]. Also, mechanical damages to neural pathways can result in pathological dynamical phenomena, as in the case of spinal cord injury which leads to a hyperexcitability of motorneurons. The modeling approach by [Venugopal et al. 2012] goes down to the role of each ion channels and aims at a reduction of spasticity. Finding suitable parameter ranges for electrical stimulation is annother important issue.

[Krishnamurthi et al. 2012] investigate by measuring velocity reduction how an optimal amplitude of DBS can be found for Parkinson's disease. Considering a RempeJens Christian Claussen, Ulrich G. Hofmann

Terman based computational model of basal ganglia, in [Njap et al. 2012] it is demonstrated that a high concentration of the inhibiting neurotransmitter GABA together with electrical stimulation reestablishes faithful thalamocortical relaying. A more general question is addressed in [Schütt and Claussen 2012] by investigating low- to high-frequency stimulation in an Izhikevich-type cortical network model, with the observation that in a frequency range around 100Hz, a dynamical desynchronization is observed.

Neuroprosthetics and Brain-Computer Interfaces. -The consequent continuation of few-electrode stimulations and recordings are Human-Machine interfaces (HMIs) [Birbaumer 2006], the most apparent applications where theoretical brain science and engineering meet. In the loss of direct control through the natural pathways, prosthetic devices have to be controlled through an HMI which comprises a "thought-control" [Hochberg et al. 2006, Pfurtscheller et al. 2003]. The consequent continuation of that idea - and most immediate demonstration of an HMI at work is that a blind patient can use the HMI for reading [Zrenner et al. 2011], cochlea implants improve hearing [Edgerton et al. 1982]. and patients with locked-in syndrome can use an EEGbased BCI for expressing words [Wolpaw et al. 2002]. Overall, prosthetic applications require a sound modeling approach and understanding of the neural processing, and, if possible, also coding, to design an effective information interfacing with the brain. The visual system is a part of the brain where experimental research and detailed modeling have made large progress. Here, [Norheim et al. 2012] and [Einevoll and Plesser 2012] investigate both a minimal and a feedback - extended model for temporal processing in the lateral geniculate nucleus (LGN). The final step of brain-machine interfacing goes towards dissolution of the border between computer and brain itself: In their beautiful experimental setup, [Perez-Marcos et al. 2012] demonstrate virtual hand illusions and – at least for a part of the body – question the conscious awareness of our Self, and thereby connect two quite different fields.

Understanding and influencing neural plasticity. – Neural plasticity, the basis of all learning, can not only be influenced pharmacologically, but also by various means of electrical stimulation with remarkable effects on cognitive learning and consolidation

[Marshall et al. 2006]. Stimulation, as well as learning, can effect on both the dynamics [Mattia and Sanchez-Vives 2012], [Schütt and Claussen 2012] and on the plasticity [Clopath 2012], [Vogt and Hofmann 2012]. [Vogt and Hofmann 2012] demonstrate modulatory effects of dopamine based on an underlying STDP learning. Modulatory or multi-input based learning mechanisms are good candidates to explain memory consolidation. In this direction, [Clopath 2012] compares two recently proposed mechanisms, namely tag-trigger consolidation, and a metastate tagging model, and provides a comprehensive comparison of both concepts. Finally, [Weigenand et al. 2011], [Weigenand et al. 2012] DOI: 10.1063/1.2937120 investigate the phase-dependence of stimulation of cortical slow waves. It would be highly interesting, based on improved understanding of neural coding and plasticity mechanisms, to design specific stimulation protocols that selectively strengthen desired acquired memories.

Outlook. While it is tempting to go beyond the purpose of medical treatment by "enhancing the brain" [Farah et al. 2004], a deeper understanding of neural plasticity mechanisms by theoretical and computational models also is expected to offer pathways to the treatment of various neural disabilities and disorders, be them memory-related like Alzheimer, or mood-related disorders as depressive disorders. How emotion modulates learning, and how emotional processes dynamically regulate psychological states is an emerging field [Figueroa Helland et al. 2008, Huber et al. 1999] and can be expected to become the 'neuroengineering" extension of of computational modeling for the treatment of brain disorders.

Acknowledgments. - The collection of articles on these topics in the current and the subsequent issue of Cognitive Neurodynamics emerged from the 1^{st} Baltic Autumn School "Applied Computational Neuroscience: Sleep, Neuroengineering and Dynamics", hosted by the University of Lübeck with generous support from the Deutsche Forschungsgemeinschaft (DFG). We thank all invitees and participants for their contributions for this Special Issue as well as for their enthusiastic and stimulating discussions during the workshop.

References

- Abelson et al. 2005. Abelson JL, Curtis GC, Sagher O, Albucher RC, Harrigan M, Taylor SF, Martis B, Giordani B.: Deep brain stimulation for refractory obsessivecompulsive disorder. Biol Psychiatry 57, 510-6 (2005).
- Bi and Poo 2001. G. Bi, M. Poo, Synaptic modification by correlated activity: Hebb's postulate revisited. Ann. Rev. Neurosci. 24, 139-166 (2001)
- Birbaumer 2006. Birbaumer, N.: Brain-Computer-Interface Research: Coming of Age. Clin Neurophysiol. 117, pp. 479-483 (2006).
- Clopath 2012. Claudia Clopath, Synaptic consolidation: an approach to long-term learning, Cognitive Neurodynamics 6 (3), (2012) 10.1007/s11571-011-9177-6

- Compte et al. 2003. Compte A, Sanchez-Vives MV, Mc-Cormick DA, Wang X (2003) Cellular and network mechanisms of slow oscillatory activity (<1 hz) and wave propagations in a cortical network model. J Neurophysiol 89: 2707 - 2725.
- Dahlem et al. 2008. Markus A. Dahlem, Felix M. Schneider, and Eckehard Schöll, Failure of feedback as a putative common mechanism of spreading depolarizations in migraine and stroke. Chaos 18 026110 (2008).
- - Delgado et al. 1968. Delgado JM, Mark V, Sweet W, Ervin F, Weiss G, Bach YR, and Hagiwara R: Intracerebral radio stimulation and recording in completely free patients. J Nerv Ment Dis 147, 329-340 (1968).
 - Edgerton et al. 1982. Edgerton, B.J.; House, W.F.; Hitselberger, W. Hearing by cochlear nucleus stimulation in humans. Ann Otol Rhinol Laryngol Suppl 91 (2 Pt3), 117-124 (1982).
 - Einevoll and Plesser 2012. Gaute T. Einevoll and Hans E. Plesser, Extended difference-of-Gaussians model incorporating cortical feedback for relay cells in the lateral geniculate nucleus of cat, Cognitive Neurodynamics 6 (4), (2012)

DOI: 10.1007/s11571-011-9183-8

- Farah et al. 2004. Farah MJ, Illes J, Cook-Deegan R, Gardner H, Kandel E, King P, Parens E, Sahakian B, and Wolpe PR: Neurocognitive enhancement: what can we do and what should we do? Nat Rev Neurosci 5 2004, 421-425.
- Figueroa Helland et al. 2008. V. Carolina Figueroa Helland, Svetlana Postnova, Udo Schwarz, Jürgen Kurths, Bernd Kundermann, Ulrich Hemmeter, and Hans A. Braun, Comparison of Different Methods for the Evaluation of Treatment Effects from the Sleep EEG of Patients with Major Depression, J Biol Phys. 34, 393-404 (2008). DOI: 10.1007/s10867-008-9096-x
- Hochberg et al. 2006. Hochberg, L.R. et al.: Neuronal ensemble control of prosthetic devices by a human with tetraplegia. Nature 442, 164-171 (2006).
- Huber et al. 1999. Martin T. Huber, Hans A. Braun, Jürgen C. Krieg Consequences of deterministic and random dynamics for the course of affective disorders, Biol. Psychiatry 46,
 - 10.1016/S0006-3223(98)00311-4
- Hutt et al. 2012. Axel Hutt, The population firing rate in the presence of GABAergic tonic inhibition in single neurons and application to general anaesthesia, Cognitive Neurodynamics 6 (3), (2012) 10.1007/s11571-011-9182-9
- Kantz and Schreiber 1997. Holger Kantz and T. Schreiber, Nonlinear Time Series Analysis, Cambridge University Press (1997)
- Krishnamurthi et al. 2012. Naravanan Krishnamurthi, Stefani Mulligan, Padma Mahant, Johan Samanta and James J. Abbas, Deep brain stimulation amplitude alters posture shift velocity in Parkinsons disease, Cognitive Neurodynamics 6 (4), (2012) DOI: 10.1007/s11571-012-9201-5
- Lehnertz et al. 2000. K Lehnertz, C.E. Elger, J. Arnhold, P. Grassberger (Eds.), Chaos in Brain?, World Scientific (2000)
- Marshall et al. 2006. Lisa Marshall, Halla Helgadottir, Matthias Molle, Jan Born Boosting cillations during sleep potentiates memory, Nature 444, 610-613 (2006).
- Mattia and Sanchez-Vives 2012. Maurizio Mattia and Maria V. Sanchez-Vives, Exploring the spectrum of dynamical regimes and timescales in spontaneous cortical activity,

Cognitive Neurodynamics 6 (3), (2012)

DOI: 10.1007/s11571-011-9179-4

- Ngo et al. 2010. Ngo HV, Köhler J, Mayer J, Claussen JC, Schuster HG Triggering up states in all-to-all coupled neurons. epl Europhysics Letters 89, 68002 (2010). DOI: 10.1209/0295-5075/89/68002
- Njap et al. 2012. Felix Njap and Jens Christian Claussen and Andreas Moser and Ulrich G. Hofmann, Modeling Effect of GABAergic Current in Basal Ganglia Computational Model, Cognitive Neurodynamics 6 (4), (2012) DOI: 10.1007/s11571-012-9203-3
- Norheim et al. 2012. Eivind S. Norheim, John Wyller, Eilen Nordlie and Gaute T. Einevoll, A minimal mechanistic model for temporal signal processing in the lateral geniculate nucleus, Cognitive Neurodynamics 6 (3), (2012) DOI: 10.1007/s11571-012-9198-9
- Olbrich et al. 2011. Eckehard Olbrich and Jens Christian Claussen and Peter Achermann, The multiple time scales of sleep dynamics as a challenge for modeling the sleeping brain, Phil. Trans. R. Soc. A 369, 3884-3901 (2011) DOI: 10.1098/rsta.2011.0082
- Perez-Marcos et al. 2012. Daniel Perez-Marcos, Maria V. Sanchez-Vives and Mel Slater, Is my hand connected to my body? The impact of body continuity and arm alignment on the virtual hand illusion, Cognitive Neurodynamics 6 (4), (2012)
 - DOI: 10.1007/s11571-011-9178-5
- Pfurtscheller et al. 2003. Pfurtscheller, G., Mueller, G. R., Pfurtscheller, J., Gerner, H. J., and Rupp, R.: 'Thought'-Control of Functional Electrical Stimulation to Restore Hand Grasp in a Patient with Tetraplegia. Neurosci. Lett. 351, 33-36 (2003).
- Pikovsky et al. 2001. Arkady Pikovsky, Michael Rosenblum, Jürgen Kurths (Eds.), Synchronization: A Universal Concept in Nonlinear Sciences, Cambridge University Press (2001)
- Schütt and Claussen 2012. Markus Schütt and Jens Christian Claussen, Desynchronizing effect of high-frequency stimulation in a generic cortical network model, Cognitive Neurodynamics 6 (4), (2012) DOI: 10.1007/s11571-012-9199-8
- Siegelbaum and Kandel 1991. Steven A. Siegelbaum, Eric R. Kandel Learning-related synaptic plasticity: LTP and LTD, Current Opinion in Neurobiology 1, 113-120 (1991)
- Singer 1999. Wolf Singer. Neuronal Synchrony: A versatile Code for the Definition of Relations? Neuron 24, 49-65 (1999)
- Steyn-Ross et al. 2012. Moira L. Steyn-Ross, D. Alistair Steyn-Ross and Jamie W. Sleigh, Gap junctions modulate seizures in a mean-field model of general anesthesia for the cortex, Cognitive Neurodynamics 6 (3), (2012) DOI: 10.1007/s11571-012-9194-0
- Tass et al. 2006. Peter A Tass, Christian Hauptmann, Oleksandr V Popovych, Development of therapeutic brain stimulation techniques with methods from nonlinear dynamics and statistical physics, International Journal Of Bifurcation And Chaos 16, 1889-1911 (2006)
- Venugopal et al. 2012. Ranu Sharmila Venugopal, Thomas M. Hamm and Ranu Jung, Differential contributions of somatic and dendritic calcium-dependent potassium currents to the control of motoneuron excitability following spinal cord injury, Cognitive Neurodynamics 6 (3), (2012)

DOI: 10.1007/s11571-012-9191-3

Vogt and Hofmann 2012. Simon M. Vogt and Ulrich G. Hofmann Neuromodulation of STDP through short-term changes in firing causality, Cognitive Neurodynamics 6 (4), (2012)

DOI: 10.1007/s11571-012-9202-4

- Weigenand et al. 2011. Arne Weigenand, Thomas Martinetz, and Jens Christian Claussen, Switching of cortical Up and Down states: reproduction of the Shu-Hasenstaub-McCormick experiment from a conductance-based model BMC Neuroscience 12, S1, P39 (2011) DOI: 10.1186/1471-2202-12-S1-P39
- Weigenand et al. 2012. Arne Weigenand, Thomas Martinetz, and Jens Christian Claussen, Phase response of the cortical slow oscillation, Cognitive Neurodynamics 6(4), 367-375 (2012)

DOI: 10.1007/s11571-012-9207-z

- Wolpaw et al. 2002. Wolpaw, J. R., Birbaumer, N., McFarland, D. J., Pfurtscheller, G., and Vaughan, T. M.: Brain-Computer Interfaces for Communication and Control. Clin. Neurophysiol. 113, 767-791 (2002).
- Zrenner et al. 2011. Zrenner E, Bartz-Schmidt KU, Benav H, Besch D, Bruckmann A, Gabel VP, Gekeler F, Greppmaier U, Harscher A, Kibbel S, Koch J, Kusnyerik A, Peters T, Stingl K, Sachs H, Stett A, Szurman P, Wilhelm B, Wilke R. Subretinal electronic chips allow blind patients to read letters and combine them to words. Proc Biol Sci. 278, 1489-97 (2011).