

# Analysis of Compatible Discrete Operator Schemes for the Stokes Equations on Polyhedral Meshes

Jerome Bonelle

EDF R&D  
6, quai Watier, BP 49  
78401 Chatou cedex  
jerome.bonelle@edf.fr

Alexandre Ern

Université Paris-Est, CERMICS  
Ecole des Ponts ParisTech  
77455 Marne la Vallée Cedex 2, France  
ern@cermics.enpc.fr

July 24, 2018

## Abstract

Compatible Discrete Operator schemes preserve basic properties of the continuous model at the discrete level. They combine discrete differential operators that discretize exactly topological laws and discrete Hodge operators that approximate constitutive relations. We devise and analyze two families of such schemes for the Stokes equations in curl formulation, with the pressure degrees of freedom located at either mesh vertices or cells. The schemes ensure local mass and momentum conservation. We prove discrete stability by establishing novel discrete Poincaré inequalities. Using commutators related to the consistency error, we derive error estimates with first-order convergence rates for smooth solutions. We analyze two strategies for discretizing the external load, so as to deliver tight error estimates when the external load has a large irrotational or divergence-free part. Finally, numerical results are presented on three-dimensional polyhedral meshes. The detailed material is available from <http://hal.archives-ouvertes.fr/hal-00939164>

## 1 Introduction

Compatible Discrete Operator (CDO) schemes belong to the broad class of compatible or mimetic schemes, which preserve basic properties of the continuous model at the discrete level; see [2, 3, 7, 11, 12, 13, 15, 22, 27, 29, 32, 34] and references therein. Following the seminal ideas of [33] and [9], the degrees of freedom (DoFs) are defined using de Rham maps, and their localization results from the physical nature of the fields. Moreover, a distinction is operated between topological laws (that are discretized exactly) and constitutive relations (that are approximated). CDO schemes are formulated using discrete differential operators for the topological laws and discrete Hodge operators for the constitutive relations. The discrete differential operators produce a cochain complex, they commute with the de Rham maps, and discrete adjunction properties hold between these operators. The discrete Hodge operator is the key operator in the CDO framework. The design of this operator is not unique, and each design leads to a specific scheme; see [31, 23, 8]. CDO schemes involve two meshes: a primal mesh (which is the only one seen by the end-user) and a dual mesh. The discrete Hodge operator links DoFs defined on the primal mesh to DoFs defined on the dual mesh. In [8], CDO schemes have been analyzed for elliptic problems on polyhedral meshes.

The Stokes equations model flows of incompressible and viscous fluids where the advective inertial forces are negligible with respect to the viscous forces. In this paper, we focus on the stationary Stokes equations posed on an open, bounded and connected domain  $\Omega \subset \mathbb{R}^3$  with boundary  $\partial\Omega$  and outward normal  $\underline{\nu}_{\partial\Omega}$ . Our starting point is to formulate the viscous stresses in the momentum balance using the curl operator. This way, all the terms in the Stokes equations can be interpreted using scalar-valued differential forms. We analyze two formulations. The first one, hereafter called 2-field curl formulation,

takes the form

$$\begin{cases} \underline{\mathbf{curl}}(\underline{\mathbf{curl}}(\underline{\mathbf{u}})) + \underline{\mathbf{grad}}(p) &= \underline{\mathbf{f}}, & \text{in } \Omega, \\ \operatorname{div}(\underline{\mathbf{u}}) &= 0, & \text{in } \Omega, \end{cases} \quad (1)$$

where  $p$  is the pressure,  $\underline{\mathbf{u}}$  the velocity and  $\underline{\mathbf{f}}$  the external load. Introducing the vorticity  $\underline{\omega} := \underline{\mathbf{curl}} \underline{\mathbf{u}}$ , the second formulation, hereafter called 3-field curl formulation (also called Velocity-Vorticity-Pressure formulation in the literature), takes the form

$$\begin{cases} -\underline{\omega} + \underline{\mathbf{curl}}(\underline{\mathbf{u}}) &= \underline{\mathbf{0}}, & \text{in } \Omega, \\ \underline{\mathbf{curl}}(\underline{\omega}) + \underline{\mathbf{grad}}(p) &= \underline{\mathbf{f}}, & \text{in } \Omega, \\ \operatorname{div}(\underline{\mathbf{u}}) &= 0, & \text{in } \Omega. \end{cases} \quad (2)$$

Essential and natural boundary conditions (BCs) can be considered for both formulations. The first set of BCs enforces the value of the normal component of the velocity  $\underline{\mathbf{u}} \cdot \underline{\nu}_{\partial\Omega}$  and that of the tangential components of the vorticity  $\underline{\omega} \times \underline{\nu}_{\partial\Omega}$  at the boundary. These BCs are natural for (1) and essential for (2). As the pressure is then determined up to an additive constant, the additional requirement of  $p$  having zero mean-value is typically added. The second set of BCs enforces the value of the tangential components of the velocity  $\underline{\mathbf{u}} \times \underline{\nu}_{\partial\Omega}$  and the value of the pressure at the boundary. These BCs are essential for (1) and natural for (2).

Mimetic or compatible schemes for the Stokes equations using either the 2-field or the 3-field curl formulation with the above BCs have already been investigated. Most of the work dedicated to the Stokes equations in curl formulations addresses the 3-field curl formulation (2). Based on the seminal work of [28], [18, 19] first analyzed this formulation in the context of finite elements (FE), and [6] in the context of spectral discretization. More recently, [24] proposed a scheme based on the Mimetic Spectral Element method in 2D and 3D for general quadrilateral/hexahedral meshes, and [14] designed a Discrete Duality Finite Volume (DDFV) scheme for general 2D meshes. Concerning the 2-field curl formulation, only the papers of [10] and of [1] have addressed this formulation on simplicial meshes; see also [30] and [20] for staggered schemes on triangular Delaunay meshes.

In the present work, we devise and analyze CDO schemes for the Stokes problem in both 2-field and 3-field curl formulations. The CDO schemes involve two Hodge operators, one linking the velocity (seen as a circulation) to the mass flux and the other linking the vorticity to the viscous stress. One key feature of the present schemes is that they ensure local mass and momentum conservation on polyhedral meshes. We prove discrete stability by establishing novel discrete Poincaré inequalities in the CDO framework. The present schemes do not need any stabilization. Moreover, using commutators related to the consistency error as in [9, 23, 8], we derive *a priori* error estimates and establish first-order error estimates for smooth solutions. In addition, we show how the present CDO framework can deal with the practically important issue of discretizing the external load, so as to deliver tight error estimates when the external load is expected to have a large irrotational or divergence-free part (see [26] for a related work).

The more classical formulation of the Stokes equations uses the vector Laplacian of the velocity in the momentum balance. Various schemes were proposed to discretize this formulation on polygonal or polyhedral meshes, including Mixed Finite Volumes by [17], Mimetic Finite Differences (MFD) by [4, 5], DDFV by [25], and an extension of the Crouzeix–Raviart finite element by [16]. [21] investigated a scheme for triangular meshes within the Finite Element Exterior Calculus (FEEC) framework. Discretizing the vector Laplacian with CDO schemes is the subject of ongoing work.

This paper is organized as follows. In Section 2, we briefly recall the CDO framework. Then, we investigate CDO schemes for the Stokes equations in the curl formulations (1) and (2). Since the pressure is seen as a potential, its DoFs are located at primal or dual mesh vertices. The former case, treated in Section 3, hinges on the 2-field curl formulation (1) leading to vertex-based pressure schemes. The latter case, treated in Section 4, hinges on the 3-field curl formulation (2) leading to cell-based pressure schemes (since primal cells are in one-to-one correspondence with dual mesh vertices). The vertex-based pressure schemes are, to our knowledge, the first of this class on polyhedral meshes. The cell-based pressure schemes share common features with the recent Mimetic Spectral Element schemes of [24];

the present schemes can be deployed on polyhedral meshes and offer two strategies for discretizing the external load. Finally, we present numerical results in Section 5. For simplicity, we often assume in what follows that  $\Omega$  is simply connected and that its boundary  $\partial\Omega$  is connected. Whenever needed, we refer to this assumption as  $(\mathbf{H}_\Omega)$ .

---

The detailed material is available from <http://hal.archives-ouvertes.fr/hal-00939164>

---

## 2 Conclusions

In this work, we have analyzed CDO schemes for the Stokes equations on three-dimensional polyhedral meshes. The distinction between primal and dual meshes enabled us to devise vertex-based and cell-based pressure schemes. Vertex-based pressure schemes lead to an algebraic system of size  $(\#V + \#E)$  with two unknowns, the pressure located at primal vertices and the velocity at primal edges. Cell-based pressure schemes lead to a system of size  $(\#E + \#F + \#C)$  with three unknowns, the pressure located at primal cells, the mass flux at primal faces, and the viscous stress circulation at primal edges. For both schemes, two discrete Hodge operators, related to the mass density and the viscosity, are used; as for elliptic problems, these operators must satisfy a stability and a consistency property. Both schemes conserve mass and momentum, the vertex-based ones at dual cells and dual faces, respectively, and the cell-based ones at primal cells and dual edges, respectively. Finally, both schemes can be deployed with two possible load discretizations, so as to handle a large irrotational or divergence-free part of the load. Finally, various tracks are worth pursuing in future work, including the use of hybridization techniques for saddle-point problems and the treatment of essential (and more general) BCs as well as complex topologies for the domain  $\Omega$ .

## References

- [1] H. Abboud, F. E. Chami, and T. Sayah. A priori and a posteriori estimates for three-dimensional Stokes equations with nonstandard boundary conditions. *Numer. Methods Partial Differential Equations*, 28(4):1178–1193, 2012.
- [2] B. Andreianov, M. Bendahmane, F. Hubert, and S. Krell. On 3D DDFV discretization of gradient and divergence operators. I. meshing, operators and discrete duality. *IMA Journal of Numerical Analysis*, 32(4):1574–1603, 2012.
- [3] D. N. Arnold, R. S. Falk, and R. Winther. Finite Element Exterior Calculus: from Hodge Theory to Numerical Stability. *Bull. Amer. Math. Soc.*, 47:281–354, 2010. DOI: 10.1090/S0273-0979-10-01278-4.
- [4] L. Beirão da Veiga, V. Gyrya, K. Lipnikov, and G. Manzini. Mimetic Finite Difference Method for the Stokes Problem on Polygonal Meshes. *Journal of Computational Physics*, 228(19):7215–7232, 2009.
- [5] L. Beirão da Veiga, K. Lipnikov, and G. Manzini. Error analysis for mimetic discretization of the steady Stokes problem on polyhedral meshes. *SIAM J. Numer. Anal.*, 48(4):1419–1443, 2010.

- [6] C. Bernardi and N. Chorfi. Spectral discretization of the vorticity, velocity, and pressure formulation of the Stokes problem. *SIAM J. Numer. Anal.*, 44:826–850, 2006.
- [7] P. Bochev and J. M. Hyman. Principles of mimetic discretizations of differential operators. In D. Arnold, P. Bochev, R. Lehoucq, R. A. Nicolaides, and M. Shashkov, editors, *Compatible Spatial Discretization*, volume 142 of *The IMA Volumes in mathematics and its applications*, pages 89–120. Springer, 2005.
- [8] J. Bonelle and A. Ern. Analysis of Compatible Discrete Operator Schemes for Elliptic Problems on Polyhedral Meshes. *ESAIM: Mathematical Modelling and Numerical Analysis*, 48(2):553–581, 2014. DOI:10.1051/m2an/2013104.
- [9] A. Bossavit. Computational electromagnetism and geometry. *J. Japan Soc. Appl. Electromagn. & Mech.*, 7-8:150–9 (no 1), 294–301 (no 2), 401–8 (no 3), 102–9 (no 4), 203–9 (no 5), 372–7 (no 6), 1999-2000.
- [10] J. H. Bramble and P. Lee. On Variational Formulations for the Stokes Equations with Non-Standard Boundary Conditions. *M2AN Math. Model. Numer. Anal.*, 28(7):903–919, 1994.
- [11] F. Brezzi, A. Buffa, and K. Lipnikov. Mimetic Finite Difference for elliptic problem. *Mathematical Modelling and Numerical Analysis*, 43:277–295, 2009.
- [12] F. Brezzi, K. Lipnikov, and M. Shashkov. Convergence of the Mimetic Finite Difference method for diffusion problems on polyhedral meshes. *SIAM J. Numer. Anal.*, 43(5):1872–1896, 2005.
- [13] M. Clemens and T. Weiland. Discrete Electromagnetism with the Finite Integration Technique. *Progress In Electromagnetics Research*, 32:65–87, 2001.
- [14] S. Delcourte and P. Omnes. A Discrete Duality Finite Volume discretization of the vorticity-velocity-pressure formulation of the 2D Stokes problem on almost arbitrary two-dimensional grids. <http://hal-cea.archives-ouvertes.fr/cea-00772972>, 2013.
- [15] M. Desbrun, A. N. Hirani, M. Leok, and J. E. Marsden. Discrete Exterior Calculus. <http://arxiv.org/abs/math/0508341>, 2005.
- [16] D. A. Di Pietro and S. Lemaire. An extension of the Crouzeix-Raviart space to general meshes with application to quasi-incompressible linear elasticity and Stokes flow. *Math. Comp.*, 2014. Accepted for publication.
- [17] J. Droniou and R. Eymard. A mixed finite volume scheme for anisotropic diffusion problems on any grid. *Numer. Math.*, 105(1):35–71, 2006.
- [18] F. Dubois. Une formulation tourbillon-vitesse-presion pour le problème de Stokes. *Comptes Rendus de l'Académie des Sciences*, 314:277–280, 1992.
- [19] F. Dubois. Vorticity-velocity-pressure formulation for the Stokes problem. *Mathematical Methods in the Applied Sciences*, 25(13):1091–1119, 2002.
- [20] R. Eymard, J. Fuhrmann, and A. Linke. MAC schemes on triangular delaunay meshes. Technical Report 1654, WIAS, 2011.
- [21] R. Falk and M. Neilan. Stokes Complexes and the Construction of Stable Finite Elements with Pointwise Mass Conservation. *SIAM J. Numer. Anal.*, 51(2):1308–1326, 2013.
- [22] M. Gerritsma. An introduction to a compatible spectral discretization method. *Mechanics of Advanced Materials and Structures*, 19(1-3):48–67, 2012.
- [23] R. Hiptmair. Discrete Hodge operators: An algebraic perspective. *Progress In Electromagnetics Research*, 32:247–269, 2001.

- [24] J. Kreeft and M. Gerritsma. Mixed mimetic spectral element method for Stokes flow: A pointwise divergence-free solution. *Journal of Computational Physics*, 240:284–309, 2013.
- [25] S. Krell and G. Manzini. The Discrete Duality Finite Volume method for Stokes equations on three-dimensional polyhedral meshes. *SIAM Journal on Numerical Analysis*, 50, 2012.
- [26] Alexander Linke. On the role of the Helmholtz decomposition in mixed methods for incompressible flows and a new variational crime. *Comput. Methods Appl. Mech. Engrg.*, 268:782–800, 2014.
- [27] C. Mattiussi. The finite volume, finite element, and finite difference methods as numerical methods for physical field problems. *Advances in Imaging and electron physics*, 113:1–146, 2000.
- [28] J. C. Nédélec. Éléments Finis Mixtes Incompressibles pour l'Équation de Stokes dans  $\mathbb{R}^3$ . *Numer. Math.*, 39:97–112, 1982.
- [29] J. B. Perot. Discrete Conservation Properties of Unstructured Mesh Schemes. *Annual Review of Fluid Mechanics*, 43(1):299–318, 2011.
- [30] J. B. Perot and R. Nallapati. A moving unstructured staggered mesh method for the simulation of incompressible free-surface flows. *Journal of Computational Physics*, 184(1):192–214, 2003.
- [31] T. Tarhasaari, L. Kettunen, and A. Bossavit. Some realizations of a discrete Hodge operator: A reinterpretation of finite element techniques. *IEEE Transactions on magnetics*, 35(3):1494–1497, 1999.
- [32] F.L. Teixeira. Differential Forms in Lattice Field Theories: An Overview. *ISRN Mathematical Physics*, 2013:16p, 2013.
- [33] E. Tonti. *On the formal structure of physical theories*. Istituto di matematica, Politecnico, Milano, 1975.
- [34] E. Tonti. Finite formulation of the electromagnetic field. *Progress In Electromagnetics Research (PIER)*, 32:1–44, 2001.