# On the collapse of trial solutions for a damped-driven non-linear Schrödinger equation

## Sigurd Assing

Department of Statistics University of Warwick e-mail: s.assing@warwick.ac.uk

#### **Astrid Hilbert**

Department of Mathematics Linnéuniversitetet Vaxjo e-mail: astrid.hilbert@lnu.se

**Abstract:** We consider the focusing 2D non-linear Schrödinger equation, perturbed by a damping term, and driven by multiplicative noise. We show that a physically motivated trial solution does not collapse for any admissible initial condition although the exponent of the non-linearity is critical. Our method is based on the construction of a global solution to a singular stochastic Hamiltonian system used to connect trial solution and Schrödinger equation.

MSC 2010 subject classifications: Primary 60H10,74H35; secondary 74J30.

**Keywords and phrases:** Schrödinger equation, singular Hamiltonian system.

## 1. Motivation

Consider the formal equation,

$$\mathbf{i}\,\partial_t \psi + \Delta_u \psi + |\psi|^2 \psi - \Lambda\,\partial_t (|\psi|^2) \psi + \sigma(u, t) \psi = 0, \tag{1.1}$$

with Cauchy data at t=0, where  $[\sigma(u,t),(u,t)\in\mathbb{R}^2\times(0,\infty)]$  is radially symmetric centred Gaussian noise with covariance

$$\langle \sigma(u,t)\sigma(u',t')\rangle = \frac{D_r}{|u|} \delta_0(|u|-|u'|)\delta_0(t-t').$$

This equation was derived in [3] as the isotropic continuum approximation of a model for two-dimensional damped-driven exciton-phonon systems.

Note that (1.1), as derived in Section II of [3], is actually driven by coloured multiplicative noise. But, in Section III of [3], the authors say they would rather approximate the driving noise by space-time white noise which they had justified in [2]. Finally, in order to allow radially symmetric (i.e. isotropic) solutions, they

simplified space-time white noise to radially symmetric Gaussian noise as used in the formulation of (1.1)—the reader is referred to [2] for the definition of a parameter  $D_{white}$  which can be used to choose a physically meaningful value for  $D_r$ . The definition, in physical terms, of the positive damping parameter  $\Lambda$  can be found in [3], too.

In the case of  $\Lambda = D_r = 0$ , equation (1.1) is identical to the classical focusing (power) non-linear Schrödinger equation, and the power two in the non-linearity  $|\psi|^2\psi$  is known to be the smallest power-like non-linearity for which blow-up occurs in space dimension d=2.

For example, the wave function,

$$\psi(u,t) = |t-1|^{-\frac{d}{2}} Q(\frac{u}{t-1}) e^{i\frac{|u|^2}{4(t-1)} - i/(t-1)}, \tag{1.2}$$

where Q is the unique positive radially symmetric solution of

$$\Delta Q - Q + Q^{1 + \frac{4}{d}} = 0$$
 in  $\mathbb{R}^d$ , with  $Q(u) \to 0$  as  $|u| \to \infty$ ,

can be viewed as a solution of

$$\mathbf{i}\,\partial_t \psi + \Delta_u \psi + |\psi|^{\frac{4}{d}} \psi = 0, \quad \text{in } \mathbb{R}^d \times (0, \infty),$$

with Cauchy data.

$$\psi(u,0) = Q(u) e^{-i(\frac{|u|^2}{4}-1)}, \text{ at } t = 0,$$

which blows up at time t = 1. Note that Q is also called *ground state*. Now observe that

$$\int_{\mathbb{R}^d} |\psi(u,t)|^2 du = \int_{\mathbb{R}^d} Q(u)^2 du, \text{ for all } t \in [0,1),$$

in the above example. Hence, all  $L^2$ -mass is accumulated into blow-up, and, by the shape of Q, this  $L^2$ -mass is concentrated at u=0 at time t=1.

The above described phenomenon, also called the  $L^2$ -concentration phenomenon, is well-known for  $L^2$ -critical Schrödinger equations—the reader is referred to [8],[9],[10] for general results.

Now, in [3], the authors have asked if this phenomenon was possible for solutions of their model for damped-driven exciton-phonon coupled systems, as the balanced energy input could prevent solutions from blow-up.

Of course, (1.1) is hard to solve, and being a formal equation only, its rigorous meaning would need further discussion in the first place. This difficulty was by-passed in [3]. Instead, the authors introduced the following family of wave functions,

$$\psi(u,t) \stackrel{\text{def}}{=} \frac{\|\psi(\cdot,0)\|_{L^2}}{\sqrt{c_f^{1,2,0}}} \times \frac{1}{x(t)} f(\frac{|u|}{x(t)}) e^{\mathbf{i} \frac{\dot{x}(t)|u|^2}{4x(t)}}, \tag{1.3}$$

<sup>&</sup>lt;sup>1</sup> See Remark 1.1 for the definition of  $c_f^{m,n,p}$ .

parametrised by  $\|\psi(\cdot,0)\|_{L^2}$ , a smooth function  $f:\mathbb{R}\to(0,\infty)$  which is rapidly decreasing, and an unknown stochastic process  $x=[x(t),\,t\geq0]$  which plays the role of the width of the corresponding non-linear wave.

Note that f, in contrast to Q used in (1.2), does not have to satisfy any equation. Nevertheless, similar to (1.2), wave functions of this type would blow up, if x starting from a positive value hits zero in finite time, the initial  $L^2$ - mass being preserved in the process. Due to the nature of a blow-up with vanishing width of the wave function, we also call such a behaviour *collapse*.

So, the question asked can be scaled down to the following problems:

- a) choose x in a way such that the trial solution (1.3) has something to do with the primary equation (1.1);
- b) study whether x chosen this way reaches zero in finite time or not.

To shed some light on a), restricting ourselves to isotropic solutions, we can rewrite (1.1) as

$$\mathbf{i}\,\partial_t\psi + \partial_r^2\psi + \frac{1}{r}\partial_r\psi + |\psi|^2\psi - \Lambda\,\partial_t(|\psi|^2)\psi + \sigma(r,t)\psi = 0,$$

reducing the problem to one space dimension with radial coordinate r = |u|. Since  $\sigma$  has negative Hölder-regularity, non-trivial solutions of this equation are not smooth in r, but the trial solution (1.3) is. Therefore, (1.3) can at most be an approximate solution having some features of a true solution.

The features chosen in [3] involve the virial coefficient

$$\mathfrak{v}(t) = \int_{\mathbb{R}^2} |u|^2 |\psi(u,t)|^2 du = 2\pi \int_0^\infty r^3 |\psi(r,t)|^2 dr$$

which, when  $\psi$  is assumed to solve (1.1), would formally satisfy

$$\ddot{\mathfrak{v}} = 16\mathscr{H} + 8\pi \int_0^\infty r^2 |\psi|^2 \,\partial_r [-\Lambda \partial_t (|\psi|^2) + \sigma] \,\mathrm{d}r \tag{1.4}$$

where

$$\mathscr{H}(t) = \int_{\mathbb{R}^2} \left[ \frac{1}{2} |\nabla_u \psi(u, t)|^2 - \frac{1}{4} |\psi(u, t)|^4 \right] du$$

is the Hamiltonian of the focusing cubic non-linear Schrödinger equation.

Plugging  $\psi$  given by (1.3) into (1.4), and then performing the integration against r, transforms (1.4) into an equation for x, only. The authors of [3] now argue that this equation can be studied by the simpler equation,

$$\ddot{x} = \frac{\delta}{x^3} + \left[\sqrt{2D}\frac{\dot{W}}{x^2} - \frac{\gamma \dot{x}}{x^4}\right],\tag{1.5}$$

where  $W = [W(t), t \ge 0]$  stands for a standard one-dimensional Wiener process.

Remark 1.1. As stated in [3],

$$\delta = \frac{4}{c_f^{3,2,0}} \left( c_f^{1,0,2} - \frac{\|\psi(\cdot,0)\|_{L^2}^2 c_f^{1,4,0}}{2c_f^{1,2,0}} \right),\,$$

and

$$\gamma \, = \, 8 \Lambda \| \psi(\cdot,0) \|_{L^2}^2 \, c_f^{3,2,2} / (c_f^{1,2,0} c_f^{3,2,0}), \quad D \, = \, 32 \pi^2 D_r \, c_f^{3,2,2} / (c_f^{3,2,0})^2,$$

using

$$c_f^{m,n,p} \stackrel{\text{def}}{=} 2\pi \int_0^\infty r^m [f(r)]^n [f'(r)]^p dr.$$

The methodology for solving the first problem, a), as developed in [3]), can therefore be described as follows: for fixed  $\|\psi(\cdot,0)\|_{L^2}$  and f, fit the trial solution (1.3) to (1.4) to obtain equation (1.5) for the unknown x. This way, equation (1.1) is more or less replaced by (1.4) subject to a structure condition—the specific form of  $\psi$  given by (1.3)—which may put such a  $\psi$  in close vicinity of a true solution to (1.1), in particular for well-chosen f.

However, no rigorous analysis has been done to support such a quality of the trial solution (1.3). First, one would have to make rigorous sense of all formal calculations used to motivate both equations (1.4) & (1.5), and, second, one would have to study how close the trial solution (1.3) and true solutions to (1.1) really are, and in what sense.

These open but very interesting problems are beyond the present paper and left for future research. We should nevertheless mention that numerical experiments reported in physics journals confirm a close match of trial solutions and true solutions on short time intervals. So, studying the long-time behaviour of solutions to (1.5), and in particular answering the second problem, b), seems to be a natural next step in the analysis of the original problem. For example, once the answer to b) is known, blow-up of true solutions could be decided by merely comparing functionals of trial solutions and true solutions.

In this paper, we therefore study the physically motivated equation (1.5) and solve the second problem, b), for all parameters of interest. Apart from the above application, these results are interesting in themselves since equation (1.5) describes the dynamics of a perturbed stochastically driven Hamiltonian system with a singular potential (cf. [4, 5] and Remark 2.1 below).

## Remark 1.2.

- (i) Although the trial solution (1.3) has infinite degrees of freedom, studying it's blow-up through (1.5) means that only the interplay of five one-dimensional parameters has to be considered:  $x(0), \dot{x}(0), \delta, \gamma, D$ . Here, assuming  $\Lambda, D_r > 0$ , one also has  $\gamma, D > 0$  by Remark 1.1, and since the trial solution has a physical meaning, x(0) > 0 should hold, too.
- (ii) Note that the parameter  $\delta$  can have both signs depending on the relationship between  $\|\psi(\cdot,0)\|_{L^2}^2$  and integrals of f. Of course, compared with negative  $\delta$ , the width x of the wave function is less likely to reach zero in finite time when  $\delta$  is positive, and hence one might expect a phase transition in the behaviour of solutions to (1.5) depending on  $\delta$ . However, we are going to prove that, for any choice of  $\delta$ , the width x never collapses to zero in finite time.

#### 2. Results

Motivated by (1.5), we study the degenerated Itô diffusion equation,

$$dx = y dt,$$

$$dy = \frac{a}{x^{\alpha}} dt + \left[\frac{\sqrt{2T\gamma}}{x^{\beta}} dW - \frac{\gamma}{x^{2\beta}} y dt\right],$$
(2.1)

where W denotes a one-dimensional Wiener process.

Our goal is to construct a global solution  $[x(t), y(t), t \geq 0]$ , for any initial condition  $(x(0), y(0)) \in \mathbb{H} \stackrel{\text{def}}{=} (0, \infty) \times \mathbb{R}$ , if both  $\gamma, T > 0$ , but  $a \in \mathbb{R}$  is a real parameter with no restrictions on its sign.

We also assume  $\beta > 1/2$  for reasons explained in Section 3 below.

Furthermore, let  $\alpha > 1$ , though this will be relaxed for  $a \geq 0$ , later.

First, note that the infinitesimal operator associated with this equation can formally be written as

$$\mathcal{L} = \frac{\partial H}{\partial y} \partial_x - \frac{\partial H}{\partial x} \partial_y + \left[ T \frac{\gamma}{x^{2\beta}} \partial_y^2 - \frac{\gamma}{x^{2\beta}} y \, \partial_y \right], \tag{2.2}$$

using the Hamiltonian,

$$H(x,y) = U(x) + \frac{y^2}{2}$$
, with potential  $U(x) = \frac{a}{\alpha - 1} x^{1-\alpha}$ ,

so that (2.1) can be interpreted as a damped-driven Hamiltonian system, being forced by noise on an x-depending scale which exactly balances the x-depending dissipation placing the stochastic system at temperature T.

The difficulty we are facing is twofold: first, our potential U has a singularity at zero, and, second, intensity of noise =  $\sqrt{2T \times friction} = \sqrt{2T \times \gamma/x^{2\beta}}$  not only depends on x, but has a singularity at zero, too.

Nevertheless, because noise and dissipation are balanced,

$$\varrho_{\star}(x,y) = e^{-H(x,y)/T}$$
 satisfies  $\mathcal{L}^{\star}\varrho_{\star} = 0$  on  $\mathbb{H}$ ,

where  $\mathcal{L}^{\star}$  stands for the (formal) adjoint of  $\mathcal{L}$ . However, in our case,

$$\int_{0}^{\infty} \int_{\mathbb{R}} e^{-H(x,y)/T} \, \mathrm{d}y \, \mathrm{d}x = +\infty, \tag{2.3}$$

which means that this density cannot be normalised to become the density of the system's canonical invariant probability measure.

### Remark 2.1.

(i) There is recent work in [4] on stochastic dynamical systems associated with operators of type (2.2) where singular potentials of the form,

$$U(x) = a_1 x^{\alpha_1} + a_2 x^{-\alpha_2},$$

are considered assuming  $a_1, a_2 > 0$ ,  $\alpha_1 > 2$ , but where intensity of noise and friction are constant, i.e.  $\beta = 0$ . Here, the integral (2.3) converges, and the canonical invariant measure occurs to be the system's unique invariant measure. More importantly, the authors describe a general method of how to construct a Lyapunov function which gives control over the system's trajectories near the boundary of  $\mathbb{H}$ . Note that their parameter  $a_2$ —the analogue of our parameter a—is assumed to be positive, which makes their potential repulsive at zero. If our potential is repulsive at zero, i.e. a > 0, we are going to show that  $\lim_{t\to\infty} x(t) = +\infty$ , and hence a Lyapunov function cannot exist.

- (ii) The driven Rayleigh-Plesset equation considered in [5] has features being more similar to our equation (2.1). Only looking at the degenerate-diffusion-case (cf. Section 5 in [5]), their potential has no a₁-term either, and the leading singularity is of the form bx<sup>-3k</sup>, for some k≥ 1, but again b>0, so that the singular potential is repulsive at zero, as in [4]. Intensity of noise and friction, though, depend on both x and y, but in an unbalanced way. The special form of the unbalanced noise and dissipation terms together with the repulsive potential make it possible to find a Lyapunov function, and thus an invariant measure exists.
- (iii) It should be mentioned that the methods used in both papers [4] and [5] will not work when the potential is not repulsive at zero.

All in all, results on stochastic dynamical systems as singular as ours were only obtained when the system admitted a structure inducing contraction properties like existence of a Lyapunov function—see [4] for further references. As our system (2.1) does not have such a structure, a different technique is needed, even for showing existence of global solutions.

Let us briefly discuss the deterministic case, i.e.  $\gamma = 0$ . Here,  $\mathcal{L}H = 0$  on the upper half-plane  $\mathbb{H}$ , and thus, if a > 0, then  $f_L(x,y) = H(x,y) + x^2/2$  is a non-negative function on  $\mathbb{H}$ , satisfying

$$\lim_{(x,y)\to\partial\mathbb{H}} f_L(x,y) = +\infty \quad \text{and} \quad \mathcal{L}f_L \le const (1+f_L).$$

Since the coefficients of equation (2.1) are locally Lipschitz on  $\mathbb{H}$ , by standard arguments, for any  $(x(0), y(0)) \in \mathbb{H}$ , there is a global solution  $[x(t), y(t), t \geq 0]$  whose first component never reaches zero in finite time.

However, if a < 0, then there is no function  $f_L$  as above. Worse, the x-component of any solution in  $\mathbb{H}$  eventually collapses to zero in finite time, and hence there is no global solution.

So, the question is: can adding noise but also dissipation, both balancing each other as in equation (2.1), prevent collapse of solutions in  $\mathbb{H}$ , even for a < 0? The answer is YES for certain choices of the exponents  $\alpha, \beta$  depending on whether a > 0, a = 0, or a < 0.

The proof of our existence result is based on the following lemma which is the key-result of this paper. Lemma 2.2. Assume  $\gamma, T > 0$ , and

$$\beta > 1/2, \ \alpha \ge \beta \ge \alpha/2 \quad if \quad a > 0,$$

$$\beta > 1/2 \quad if \quad a = 0,$$

$$\alpha > 1, \ \alpha \ge \beta \ge (\alpha + 1)/2 \quad if \quad a < 0.$$

$$(2.4)$$

Consider the product

$$\Omega_{+} = \Omega_{-} \otimes C([0,\infty)), \quad \mathcal{F}_{+} = \mathcal{F}_{-} \otimes \mathcal{B}(C([0,\infty))), \quad \mathbf{Q} = \mathbf{P}_{-} \otimes \mathbf{P}_{W}$$

of a probability space  $(\Omega_-, \mathcal{F}_-, \mathbf{P}_-)$  and the standard Wiener space  $(C([0, \infty)), \mathcal{B}(C([0,\infty))), \mathbf{P}_W)$ , and let  $(x_-, y_-)$  be a given pair of random variables on  $(\Omega_-, \mathcal{F}_-, \mathbf{P}_-)$  satisfying  $\mathbf{P}_-(\{x_- > 0\}) = 1$ . Extend all random variables on either  $(\Omega_-, \mathcal{F}_-, \mathbf{P}_-)$  or  $(C([0,\infty)), \mathcal{B}(C([0,\infty))), \mathbf{P}_W)$  to  $(\Omega_+, \mathcal{F}_+, \mathbf{Q})$  in the canonical way without changing their notation.

Then there exists a filtration  $\mathbb{F}_+ = [\mathcal{F}_t^+, t \geq 0]$  of sub- $\sigma$ -algebras of  $\mathcal{F}_+$ , a probability measure  $\mathbf{P}_+$  on  $(\Omega_+, \mathcal{F}_{\infty}^+)$ , a pair  $[x_+(t), y_+(t), t \geq 0]$  of continuous  $\mathbb{F}_+$ -adapted processes, and an  $\mathbb{F}_+$ -Wiener-process  $W_+$ , such that

$$\mathbf{P}_{+} = \mathbf{Q} \ on \ \mathcal{F}_{0}^{+},$$

and

$$\begin{array}{rcl} x_{+}(t) &>& 0, \\ x_{+}(t) &=& x_{-} \,+\, \int_{0}^{t} y_{+}(s) \,\mathrm{d} s, \\ y_{+}(t) &=& y_{-} \,+\, a \int_{0}^{t} \frac{\mathrm{d} s}{x_{+}(s)^{\alpha}} + \sqrt{2T\gamma} \int_{0}^{t} \frac{\mathrm{d} W_{+}(s)}{x_{+}(s)^{\beta}} - \gamma \int_{0}^{t} \frac{y_{+}(s) \,\mathrm{d} s}{x_{+}(s)^{2\beta}}, \end{array}$$

for all  $t \in [0,1]$ ,  $\mathbf{P}_+$ -a.s. Note that  $\mathcal{F}_{\infty}^+$  can be strictly smaller than  $\mathcal{F}_+$ .

The above result can be used to prove existence of a global weak solution in  $\mathbb{H}$  which, by standard arguments, turns into a unique strong solution, as the equation's coefficients are locally Lipschitz on  $\mathbb{H}$ .

**Theorem 2.3.** Assume  $\gamma, T > 0$ , and (2.4). Then, equation (2.1) has a unique global strong solution in  $\mathbb{H}$ .

In the proof of the above theorem, the global solution is constructed by patching together solutions on finite intervals as constructed in the proof of Lemma 2.2. However, when a=0, the proof of Lemma 2.2 can be used to obtain a global solution without patching, and we are going to show that the x-component of this global solution is transient. Using a comparison argument, we can also verify this transient behaviour in the case of a>0, leading to

**Proposition 2.4.** Assume  $\gamma, T > 0$ , and (2.4). If  $a \ge 0$ , then the x-component of any global solution to (2.1) satisfies  $\lim_{t\to\infty} x(t) = +\infty$ , a.s.

Of course, if a process  $[x(t), y(t), t \ge 0]$  taking values in  $\mathbb{H}$ , when started at any initial condition, almost surely satisfies  $\lim_{t\to\infty} x(t) = +\infty$ , then its

dynamics would not allow for an invariant measure. As the Lyapunov functions constructed in [4, 5] guarantee the existence of an invariant measure, we also have the following

Corollary 2.5. Recall  $\mathcal{L}$  given by (2.2) with  $\gamma, T > 0$ , and assume both  $\beta > 1/2$  as well as  $\alpha \geq \beta \geq \alpha/2$ . If the potential U is repulsive at zero, i.e. a > 0, then  $\mathcal{L}$  does not admit Lyapunov functions of the types used in [4, 5].

However, if a<0 and  $\alpha>1$ , Theorem 2.3 gives a condition on the relationship between  $\alpha$  and  $\beta$  ensuring existence of a global solution, regardless the values taken by  $\gamma, T>0$ . But, we cannot tell whether this global solution  $[x(t),y(t),\,t\geq0]$  satisfies  $\lim_{t\to\infty}x(t)=+\infty$ , or  $\lim_{t\to\infty}x(t)=0$ , or whether x is recurrent—all three scenarios might be possible, depending on the choice of  $\gamma,T>0$ , which remains an open problem. So far we only know from (4.17) in the proof of Proposition 2.4 that

$$\lim_{t\to\infty} x(t) = +\infty$$
, on  $\{\int_0^\infty x(t)^{-2\beta} dt < \infty\}$ , a.s.,

but this set might have measure zero in cases where a < 0.

Finally, we return to our motivating example, (1.5), where  $\alpha = 3$  and  $\beta = 2$ , so that  $\beta = (\alpha + 1)/2$ , and hence Theorem 2.3 implies that (1.5) always has a positive global solution, regardless the values taken by  $\delta \in \mathbb{R}$  and  $\gamma, D > 0$ .

Corollary 2.6. If x solves (1.5), then the non-linear wave  $\psi$  given by (1.3) would never collapse.

#### 3. Discussion of Conditions

In this section we relate crucial steps in our proofs to the conditions they rely upon, which sheds some light on how essential these conditions actually are.

First, the base step of our construction consists in analysing functionals of the solution to equation (4.2), for a given Ornstein-Uhlenbeck process  $[\hat{y}(t), t \geq 0]$ .

If  $\beta \in [0, 1/2]$  then Lemma 4.2 would be wrong because, almost surely, the solution to (4.2) would be a continuous function on the compact interval  $[0, \tau]$ . We do not see an easy fix for our proofs without a valid Lemma 4.2, and that is the main reason why we want  $\beta$  to be greater than 1/2 throughout this paper.

Second, when  $a \neq 0$ , the base step of our construction is followed by a Girsanov transform, and we have to check Novikov's condition. Checking this condition is based on Hölder's inequality which can only be applied if  $\alpha$  and  $\beta$  are in the relation  $\alpha \geq \beta \geq \alpha/2$ .

The third and last crucial step of our construction is the proof of Lemma 4.5. It turns out that, if a>0, the conditions  $\beta>1/2$  and  $\alpha\geq\beta\geq\alpha/2$  assumed in the first and second step, respectively, are sufficient for this proof. But, if a<0, two extra conditions,  $\alpha>1$  and  $\beta\geq(\alpha+1)/2$ , are required. Note that these extra conditions imply  $\beta>1/2$  needed in the first step.

Furthermore, compared with  $\alpha > 1/2$ , which is the consequence of condition  $\beta > 1/2$ ,  $\alpha \ge \beta \ge \alpha/2$  from case  $\alpha > 0$ , the extra condition  $\alpha > 1$  is in a way

counter intuitive for negative a. Indeed, for negative a, a greater power of  $\alpha$  in (2.1) should push the trajectories of the x-component further to zero, once x is close to collapse. This push is obviously compensated by stronger fluctuations of the damped noise caused by the other extra condition  $\beta \geq (\alpha + 1)/2$ . The message of the proof of Lemma 4.5 seems to be that global solutions to (2.1) can only exist if  $\alpha$  and  $\beta$  are in the right ratio.

We finally discuss the most interesting case  $\delta < 0$  of our main application, equation (1.5), where  $\alpha = 3$  and  $\beta = 2$ , so that  $\beta = (\alpha + 1)/2$ , which is at the 'edge' of the condition ensuring global existence. This could mean that, when  $\delta < 0$ , solutions to (1.5) are 'just' global in the sense that  $\lim_{t\to\infty} x(t) = 0$ , almost surely or with positive probability, depending on the choice of  $\gamma, D > 0$ . This behaviour, which the authors called 'pseudo-collapse', has been conjectured and supported by numerical experiments in [3].

However, when  $\delta \geq 0$ , we know from Proposition 2.4 that solutions are even transient and cannot pseudo-collapse, and hence  $\delta \geq 0$  would be a condition for non-pseudo-collapse of the corresponding non-linear wave given by (1.3). Recalling the definition of  $\delta$  in Remark 1.1, this condition would read

$$\|\psi(\cdot,0)\|_{L^2}^2 \le 2c_f^{1,0,2} c_f^{1,2,0} / c_f^{1,4,0},$$

which compares the  $L^2$ -norm of the wave's initial condition with a product of integrals of f and f'.

A similar but structurally easier condition is well-known for the classical focusing non-linear Schrödinger equation: Weinstein's criterion, [11, Thm. A], says that solutions would never blow up if

$$\|\psi(\cdot,0)\|_{L^2} < \|Q\|_{L^2},$$

where Q is the ground state used for (1.2). Here, Q satisfies an equation, while f used for (1.3) does not. Therefore, the condition for non-pseudo-collapse could be relaxed to

$$\|\psi(\cdot,0)\|_{L^2}^2 \le \sup_f 2c_f^{1,0,2} c_f^{1,2,0}/c_f^{1,4,0},$$

where the supremum is taken over all smooth functions  $f: \mathbb{R} \to (0, \infty)$  which are rapidly decreasing.

#### 4. Proofs

Proof of Lemma 2.2. First, observe that  $\beta > 1/2$  in all three cases of (2.4). Let  $[B_t, t \geq 0]$  be the coordinate process on  $(C([0, \infty)), \mathcal{B}(C([0, \infty))), \mathbf{P}_W)$ , and define the filtration

$$\mathbb{G}_{+} = [\mathcal{G}_{t}^{+}, t \geq 0]$$
 by  $\mathcal{G}_{t}^{+} = \sigma(x_{-}, y_{-}) \vee \sigma(\{B_{s} : s \leq t\}).$ 

Note that  $[B_t, t \geq 0]$  is a  $\mathbb{G}_+$ -Wiener-process on  $(\Omega_+, \mathcal{F}_+, \mathbf{Q})$ .

Of course,

$$\hat{y}(t) \stackrel{\text{def}}{=} e^{-\gamma t} y_- + \sqrt{2T\gamma} \int_0^t e^{-\gamma(t-s)} dB_s, \quad t \ge 0,$$

satisfies

$$\hat{y}(t) = y_{-} + \sqrt{2T\gamma} B_{t} - \gamma \int_{0}^{t} \hat{y}(s) ds,$$
 (4.1)

for all  $t \geq 0$ , **Q**-a.s.

Next, for  $\beta > 1/2$ ,

$$\hat{x}(t) \stackrel{\text{def}}{=} \left[ x_{-}^{1-2\beta} - (2\beta - 1) \int_{0}^{t} \hat{y}(s) \, \mathrm{d}s \right]^{\frac{-1}{2\beta - 1}}$$

solves

$$\hat{x}(t) = x_{-} + \int_{0}^{t} \hat{x}(s)^{2\beta} \hat{y}(s) ds, \text{ for } t < \tau,$$
 (4.2)

where

$$\tau \stackrel{\text{def}}{=} \inf\{s \ge 0 : \int_0^s \hat{y}(r) \, \mathrm{d}r = \frac{x_-^{1-2\beta}}{2\beta - 1}\}.$$

**Lemma 4.1.** The  $\mathbb{G}_+$ -stopping time  $\tau$  satisfies  $\mathbf{Q}(\{\tau < \infty\}) = 1$ .

*Proof.* Rewrite (4.1) to obtain

$$\gamma \int_0^s \hat{y}(r) dr = y_- + \sqrt{2T\gamma} B_s - \hat{y}(s), \quad s \ge 0, \ \mathbf{Q}\text{-a.s.},$$

where

$$\sqrt{2T\gamma} B_s - \hat{y}(s) = -e^{-\gamma s} y_- + \sqrt{2T\gamma} \int_0^s [1 - e^{-\gamma(s-r)}] dB_r$$

by definition of  $\hat{y}(s)$ . Thus, since  $\lim_{s\to\infty} e^{-\gamma s} y_- = 0$ , the process  $s\mapsto \int_0^s \hat{y}(r)\,\mathrm{d}r$  is almost surely going to hit  $x_-^{1-2\beta}/(2\beta-1)$  in finite time, if the process  $s\mapsto \int_0^s [1-e^{-\gamma(s-r)}]\,\mathrm{d}B_r$  is recurrent.

To show recurrence of this process, we use the representation,

$$\int_0^s [1 - e^{-\gamma(s-r)}] dB_r = B_s - e^{-\gamma s} \tilde{B}(\frac{e^{2\gamma s} - 1}{2\gamma}), \quad s \ge 0, \ \mathbf{Q}\text{-a.s.},$$

where  $[\tilde{B}(t), t \geq 0]$  is another Wiener process on  $(\Omega_+, \mathcal{F}_+, \mathbf{Q})$ —see Thm.II.7.2 in [6]. Taking into account the law of iterated logarithm (cf. Thm.2.9.23 in [7]), i.e., for any one-dimensional standard Wiener process  $[W_t, t \geq 0]$ ,

$$\limsup_{t \to \infty} \frac{W_t}{\sqrt{2t \log \log t}} = 1, \text{ a.s., } \liminf_{t \to \infty} \frac{W_t}{\sqrt{2t \log \log t}} = -1, \text{ a.s.};$$

we can then conclude that the process

$$s \mapsto B_s - e^{-\gamma s} \tilde{B}(\frac{e^{2\gamma s} - 1}{2\gamma})$$

is recurrent if

$$\sqrt{2s\log\log s} \, - e^{-\gamma s} \sqrt{\frac{e^{2\gamma s} - 1}{\gamma} \log\log\frac{e^{2\gamma s} - 1}{2\gamma}}$$

diverges, when s goes to infinity, which is true.

Now introduce  $[T_t, t \geq 0]$  defined by

$$T_t \stackrel{\text{def}}{=} \begin{cases} \int_0^t \hat{x}(s)^{2\beta} \, \mathrm{d}s & : \quad t < \tau, \\ +\infty & : \quad t \ge \tau, \end{cases}$$

which gives an increasing right-continuous  $\mathbb{G}_+$ -adapted process.

**Lemma 4.2.** For 
$$\beta > 1/2$$
,  $\mathbf{Q}(\{\lim_{t \uparrow \tau} T_t = +\infty\}) = 1$ .

*Proof.* There exists  $\Omega_0 \in \mathcal{F}_+$  such that  $\mathbf{Q}(\Omega_0) = 1$  and both  $\hat{y}(s, \omega)$  is continuous in s as well as  $\tau(\omega) < \infty$  for all  $\omega \in \Omega_0$ . Choose  $\omega \in \Omega_0$  and assume that

$$\lim_{t \uparrow \tau(\omega)} T_t(\omega) = c < +\infty.$$

Then, for  $t < \tau(\omega)$ , it follows from the mean value theorem that

$$c - T_t(\omega) = (\tau(\omega) - t) \left[ x_-^{1-2\beta} - (2\beta - 1) \int_0^{\tilde{t}} \hat{y}(s, \omega) \, ds \right]^{\frac{-2\beta}{2\beta - 1}}$$

where  $\tilde{t} \in (t, \tau(\omega))$ . Also, by definition of  $\tau$  and continuity of  $\hat{y}(s, \omega)$  in s,

$$x_{-}^{1-2\beta} - (2\beta - 1) \int_{0}^{\tilde{t}} \hat{y}(s,\omega) \, \mathrm{d}s = (2\beta - 1)(\tau(\omega) - \tilde{t}) \, \hat{y}(\tilde{\tilde{t}},\omega)$$

for some  $\tilde{\tilde{t}} \in (\tilde{t}, \tau(\omega))$ , again applying the mean value theorem. Thus,

$$c - T_t(\omega) = O\left((\tau(\omega) - t)^{\frac{-1}{2\beta - 1}}\right), \quad t \uparrow \tau(\omega),$$

which means that  $c - T_t(\omega)$  should blow up when t goes to  $\tau(\omega)$ , since  $\beta > 1/2$ . But, such a blow-up would contradict our assumption of  $c < \infty$ , proving  $\lim_{t \uparrow \tau(\omega)} T_t(\omega) = +\infty$ , for all  $\omega \in \Omega_0$ .

As  $[T_t, t \geq 0]$  is continuous and strictly increasing on  $[0, \tau)$ , Lemma 4.2 implies that the right-inverse  $A = [A_t, t \geq 0]$  defined by

$$A_t = \inf\{s > 0 : T_s > t\}$$

is a continuous strictly increasing family of  $\mathbb{G}_+$ -stopping times satisfying

$$A_t < \tau$$
,  $t \ge 0$ , **Q**-a.s.

As a consequence, the time-changed processes  $x_+(t) = \hat{x}(A_t)$ ,  $y_+(t) = \hat{y}(A_t)$  are well-defined for all  $t \geq 0$  on a set in  $\mathcal{F}_+$  of **Q**-measure one. Furthermore, by time-change, equation (4.2) yields

$$x_{+}(t) = x_{-} + \int_{0}^{t} y_{+}(s) \, ds, \quad t \ge 0, \, \mathbf{Q}\text{-a.s.}$$
 (4.3)

**Remark 4.3.** It follows immediately from the construction of  $\hat{x}$  on  $[0, \tau)$  that  $x_{+}(t) > 0$ , for all  $t \geq 0$ , **Q**-a.s.

So, the first two statements of the lemma would be true under the measure  $\mathbf{Q}$ , and, if a=0, by time-change, equation (4.1) yields (4.9), for all  $t \geq 0$ ,  $\mathbf{Q}$ -a.s., in a straight forward way. Note that  $\beta > 1/2$  has been the only assumption we made, so far, proving the lemma in the case a=0.

In what follows, assume  $a \neq 0$ .

Then, the equation for  $y_+(t)$  requires a measure  $\mathbf{P}_+$  different to  $\mathbf{Q}$ . The next step is to construct this measure.

Introduce the process,  $\rho = [\rho(t), t \ge 0]$ , given by

$$\rho(t) \stackrel{\text{def}}{=} \exp\{\frac{a}{\sqrt{2T\gamma}} \int_0^t \mathbf{1}_{[0,A_1]}(s) \hat{x}(s)^{2\beta-\alpha} dB_s - \frac{a^2}{4T\gamma} \int_0^t \mathbf{1}_{[0,A_1]}(s) \hat{x}(s)^{4\beta-2\alpha} ds\},$$

which is well-defined since the stochastic integrand is a caglad  $\mathbb{G}_+$ -adapted process. Since  $\alpha \geq \beta \geq \alpha/2$ , by Hölder's inequality,

$$\int_0^t \mathbf{1}_{[0,A_1]}(s)\hat{x}(s)^{4\beta-2\alpha} \, \mathrm{d}s \leq t^{\frac{\alpha-\beta}{\beta}} \cdot \left(\int_0^{A_1} \hat{x}(s)^{2\beta} \, \mathrm{d}s\right)^{\frac{2\beta-\alpha}{\beta}} = t^{\frac{\alpha-\beta}{\beta}} \cdot \left(T_{A_1}\right)^{\frac{2\beta-\alpha}{\beta}},$$

where  $\beta > 1/2$  yields  $T_{A_1} = 1$ , **Q**-a.s., by Lemma 4.2, and hence

$$\int \exp\{\frac{a^2}{4T\gamma} \int_0^t \mathbf{1}_{[0,A_1]}(s)\hat{x}(s)^{4\beta-2\alpha} \, ds\} \, d\mathbf{Q} < \infty,$$

for all  $t \geq 0$ , so that  $\rho$  is a  $\mathbb{G}_+$ -martingale by Novikov's condition.

Copying the proof of Corollary 3.5.2 in [7], one can construct a probability measure  $P_+$  on  $\mathcal{G}_{\infty}^+$  such that

$$\hat{W}_{+}(t) \stackrel{\text{def}}{=} B_{t} - a \int_{0}^{t} \mathbf{1}_{[0,A_{1}]}(s)\hat{x}(s)^{2\beta-\alpha} \,\mathrm{d}s / \sqrt{2T\gamma}, \quad t \geq 0,$$

is a  $\mathbb{G}_+$ -Wiener-process.

# Remark 4.4.

(i) The measure constructed in the original proof of Corollary 3.5.2 in [7] would be defined on  $\sigma(\{B_s:s\geq 0\})$  but our integrand  $\mathbf{1}_{[0,A_1]}\hat{x}(\cdot)$  is not  $\sigma(\{B_s:s\geq 0\})$ -measurable. However, the proof still works when using  $\mathcal{G}_{\pm}^+$  instead. It is not needed that  $\mathbb{G}_+$  satisfies the usual conditions.

(ii) The  $\sigma$ -algebra  $\mathcal{G}_{\infty}^+$  may be smaller than  $\mathcal{F}_+$ .

The measure  $\mathbf{P}_{+}$  does not have to be absolutely continuous w.r.t.  $\mathbf{Q}$ , but it is on every  $\mathcal{G}_{t}^{+}$ , where  $\mathbf{P}_{+} = \rho(t) \cdot \mathbf{Q}$ . This allows to carry over some  $\mathbf{Q}$ -a.s. events to  $\mathbf{P}_{+}$ -a.s. events by approximation with monotone sequences of events, for example,

$$\mathbf{P}_{+}(\{\hat{y}(\cdot) \text{ continuous on } [0,\infty)\}) = 1. \tag{4.4}$$

As a consequence,

$$\left| \int_0^t \hat{y}(s) \, \mathrm{d}s \right| < \infty$$
, for all  $t \ge 0$ ,  $\mathbf{P}_+$ -a.s.,

which yields

$$\hat{x}(t) > 0$$
, for all  $t < \tau$ ,  $\mathbf{P}_{+}$ -a.s.

However, the results of both lemmas, 4.1 and 4.2, might not remain true under the new measure  $\mathbf{P}_{\!+}$ .

Indeed, though the definition of  $\hat{y}$  is still the same under  $\mathbf{P}_+$ , the process B is now a Wiener process with drift, and hence the recurrence of the stochastic integral process used to prove Lemma 4.1 might fail to hold. Thus, we have to take into account a positive  $\mathbf{P}_+$ -probability of the event  $\{\tau=\infty\}$ , and on this event the proof of Lemma 4.2 does not work. The next lemma gives conditions on the parameters  $\alpha, \beta$  ensuring that  $T_{\infty}$  cannot be finite, on  $\{\tau=\infty\}$ ,  $\mathbf{P}_+$ -a.s., and this property turns out to be crucial for the rest of the proof.

**Lemma 4.5.** Assume 
$$a \neq 0$$
 and (2.4). Then,  $P_{+}(\{\lim_{t \uparrow \tau} T_{t} = +\infty\}) = 1$ .

*Proof.* First, recall (4.4) and note that, by the same principle, (4.1) is also true, for all  $t \geq 0$ ,  $\mathbf{P}_+$ -a.s. Then, choose  $\Omega_0$  such that  $\mathbf{P}_+(\Omega_0) = 1$  and, on  $\Omega_0$ :  $\hat{y}(\cdot)$  is continuous, (4.1) is satisfied for all  $t \geq 0$ , and

$$\limsup_{t \to \infty} \frac{\hat{W}_{+}(t)}{\sqrt{2t \log \log t}} = 1, \quad \liminf_{t \to \infty} \frac{\hat{W}_{+}(t)}{\sqrt{2t \log \log t}} = -1.$$

Here we used both  $\beta > 1/2$  and  $\alpha \ge \beta \ge \alpha/2$  for validation of Novikov's condition to make sure that  $\hat{W}_+$  as defined above Remark 4.4 is a Wiener process.

Now, choose  $\omega \in \Omega_0$ . To simplify notation, for the rest of this proof, consider all random variables being evaluated at the chosen  $\omega$  without emphasising.

If  $\tau < \infty$ , as  $\beta > 1/2$ ,  $\lim_{t \uparrow \tau} T_t = +\infty$  can be shown following the arguments used in the proof of Lemma 4.2.

If  $\tau = \infty$ , using the definitions of  $\hat{y}$  and  $\hat{W}_{+}$ , a rewrite of equation (4.1) yields

$$\gamma z_t = y_- - e^{-\gamma t} y_- + a \int_0^t [1 - e^{-\gamma (t-s)}] \, \hat{x}(s)^{2\beta - \alpha} \, \mathrm{d}s$$
$$+ \sqrt{2T\gamma} \int_0^t [1 - e^{-\gamma (t-s)}] \, \mathrm{d}\hat{W}_+(s), \tag{4.5}$$

for all  $t \geq 0$ , where

$$z_t \stackrel{\text{def}}{=} \int_0^t \hat{y}(s) \, \mathrm{d}s, \quad t \ge 0.$$

Since  $\tau = \infty$ , and since  $t \mapsto z_t$  cannot explode in finite time by our choice of  $\Omega_0$ ,  $\hat{x}(\cdot)$  is a positive function on the entire domain  $[0, \infty)$ . Thus, simple differentiation reveals that the first integral in (4.5) is a monotonously increasing function in t, and hence the behaviour of the function  $t \mapsto z_t$  will be different depending on whether this monotone function dominates the stochastic integral or not, when t goes to infinity.

However, by our choice of  $\Omega_0$ , using the same arguments as in the proof of Lemma 4.1,

$$\limsup_{t \to \infty} \frac{\int_0^t [1 - e^{-\gamma(t-s)}] \, \mathrm{d}\hat{W}_+(s)}{\sqrt{2t \log \log t}} = 1, \quad \liminf_{t \to \infty} \frac{\int_0^t [1 - e^{-\gamma(t-s)}] \, \mathrm{d}\hat{W}_+(s)}{\sqrt{2t \log \log t}} = -1.$$

So, if a > 0, then the first integral in (4.5) adds to the upward-fluctuations of the stochastic integral leading to a finite value of  $\tau$ , and hence the case a > 0 cannot occur, once  $\tau = \infty$ . Therefore, the case a > 0 was covered above, only assuming  $\beta > 1/2$  and  $\alpha \ge \beta \ge \alpha/2$ .

If  $\tau = \infty$  and a < 0, we are going to show that, if  $\alpha > 1$  and  $\beta > \alpha/2$ , then  $T_{\infty} < +\infty$  would imply  $\beta < (\alpha + 1)/2$ , proving the lemma in this case, too.

For the rest of this proof, assume  $\tau = \infty$ , a < 0,  $\alpha > 1$ ,  $\beta > \alpha/2$ ,  $T_{\infty} < +\infty$ .

To begin with, we are going to verify that the first integral in (4.5) would always dominate the stochastic integral, pushing all fluctuations of  $z_t$  down to  $\lim_{t\to\infty} z_t = -\infty$ , eventually.

Indeed, if a < 0, when using the long-time behaviour of the stochastic integral in (4.5), we can deduce that, for some large enough  $t_0$ , there exists  $b_0 > 0$  such that, for all  $t \ge t_0$ ,

$$\gamma z_t \ge y_- - e^{-\gamma t} y_- - |a| \int_0^t \hat{x}(s)^{2\beta - \alpha} ds - b_0 \sqrt{t} \sqrt{\log \log t}.$$

Of course,  $y_- - e^{-\gamma t}y_-$  is bigger than some negative number, for all  $t \ge 0$ , and this negative number becomes even smaller when subtracting  $|a| \int_0^{t_0} \hat{x}(s)^{2\beta-\alpha} ds$ . Therefore, for all  $t \ge t_0$ , the above inequality can be written as follows,

$$z_t \ge -a_0 - \frac{b_0}{\gamma} \sqrt{t_0} \sqrt{\log \log t_0} - \frac{|a|}{\gamma} \int_{t_0}^t [c_1 - c_2 z_s]^{-\kappa} \, \mathrm{d}s - \frac{b_0}{\gamma} \int_{t_0}^t f(s) \, \mathrm{d}s,$$

writing f(s) for  $\frac{d}{ds}(\sqrt{t}\sqrt{\log\log t})$ , and substituting the definition of  $\hat{x}$ , so that:

$$c_1 = x_-^{1-2\beta}, \quad c_2 = 2\beta - 1, \quad \kappa = \frac{2\beta - \alpha}{2\beta - 1}.$$

Now, consider the ordinary differential equation (ODE),

$$\frac{\mathrm{d}}{\mathrm{d}t}\tilde{z} = -\frac{|a|}{\gamma}[c_1 - c_2\tilde{z}]^{-\kappa} - \frac{b_0}{\gamma}f(t).$$

If this equation, when started at  $t_0$  from  $-a_0 - \frac{b_0}{\gamma} \sqrt{t_0} \sqrt{\log \log t_0}$ , has a unique global solution, then, by standard comparison arguments,

$$z_t \geq \tilde{z}_t, \quad t \geq t_0,$$

and thus.

$$c_1 - c_2 z_t \le p_t \stackrel{\text{def}}{=} c_1 - c_2 \tilde{z}_t, \quad t \ge t_0,$$
 (4.6)

if  $[p_t, t \ge t_0]$  was the unique global solution of

$$p_t = c_1 + c_2 a_0 + c_2 \frac{|a|}{\gamma} \int_{t_0}^t p_s^{-\kappa} \, \mathrm{d}s + c_2 \frac{b_0}{\gamma} \sqrt{t} \sqrt{\log \log t} \,. \tag{4.7}$$

Yet, since  $c_1 + c_2 a_0 > 0$  and  $t_0$  was chosen large enough, this ODE (written in integral form) has local solutions, these local solutions are unique on their domain of definition (since the equation's coefficients are locally Lipschitz), and any local solution is monotonously increasing.

So, on its domain of definition, any local solution satisfies

$$p_t \ge c_2 \frac{b_0}{\gamma} \sqrt{t} \sqrt{\log \log t},$$

and hence, since  $\kappa > 0$ ,

$$\int_{t_0}^t p_s^{-\kappa} \, \mathrm{d}s \le \left(\frac{\gamma}{c_2 b_0}\right)^{\kappa} \int_{t_0}^t s^{-\kappa/2} \, \mathrm{d}s,$$

which means that  $p_t = c_1 - c_2 \tilde{z}_t$ ,  $t \ge t_0$ , is indeed the unique global solution of equation (4.7), because blow-up cannot occur in finite time.

Next, since  $\kappa \neq 2$ , the above inequality asserts

$$\int_{t_0}^t p_s^{-\kappa} \, \mathrm{d}s \le \left(\frac{\gamma}{c_2 b_0}\right)^{\kappa} \cdot t^{-\frac{\kappa}{2}+1}, \quad t \ge t_0,$$

which we apply to estimate the right-hand side of (4.7). Here, since  $\kappa < 1$ , the product  $\sqrt{t}\sqrt{\log\log t}$  is dominated by  $t^{-\frac{\kappa}{2}+1}$ , when t goes to infinity, and therefore (4.7) yields

$$p_t \le c_0 t^{-\frac{\kappa}{2}+1}, \quad t \ge t_0,$$

for some sufficiently large constant  $c_0 > 0$ .

Using this bound, (4.6), the definition of  $\hat{x}$ , and  $\kappa > 0$ , we obtain that

$$\frac{1}{(c_0 t^{-\frac{\kappa}{2}+1})^{\kappa}} \le \hat{x}(t)^{2\beta-\alpha}, \quad t \ge t_0,$$

and hence the first integral in (4.5) is bounded below by

$$c_0^{-\kappa} \int_{t_0}^t [1 - e^{-\gamma(t-s)}] s^{-\kappa(1-\kappa/2)} ds, \quad t \ge t_0.$$

Since  $\kappa(1 - \kappa/2) > 0$ , by l'Hospital,

$$\lim_{t \to \infty} \int_{t_0}^t e^{-\gamma(t-s)} \, s^{-\kappa(1-\kappa/2)} \, \mathrm{d}s \, = \, 0,$$

and since  $\kappa(1-\kappa/2) < 1/2$ ,

$$\lim_{t\to\infty} \frac{\int_{t_0}^t s^{-\kappa(1-\kappa/2)} \, \mathrm{d}s}{\sqrt{2t\log\log t}} \, = \, +\infty,$$

finally proving our claim that the first integral in (4.5) would dominate the stochastic integral, for any a < 0.

As a consequence, for any a < 0, we can now conclude that

$$\lim_{t \to \infty} z_t = -\lim_{t \to \infty} \int_0^t \left[ 1 - e^{-\gamma(t-s)} \right] \hat{x}(s)^{2\beta - \alpha} \, \mathrm{d}s = -\infty,$$

and thus, by l'Hospital,

$$\lim_{t \to \infty} \int_0^t e^{-\gamma(t-s)} \,\hat{x}(s)^{2\beta - \alpha} \, \mathrm{d}s = \lim_{t \to \infty} \int_0^t e^{-\gamma(t-s)} \, [c_1 - c_2 z_s]^{-\kappa} \, \mathrm{d}s = 0.$$

Therefore, the long-time behaviour of the right-hand side of (4.5) is fully determined by the long-time behaviour of the function,

$$t \mapsto \int_0^t \hat{x}(s)^{2\beta - \alpha} \, \mathrm{d}s = \int_0^t [c_1 - c_2 z_s]^{-\kappa} \, \mathrm{d}s,$$

and, choosing  $t_0$  large enough, we can conclude that

$$\frac{-|a|-\varepsilon}{\gamma} \int_0^t [c_1 - c_2 z_s]^{-\kappa} \, \mathrm{d}s \le z_t \le \frac{-|a|+\varepsilon}{\gamma} \int_0^t [c_1 - c_2 z_s]^{-\kappa} \, \mathrm{d}s, \quad t \ge t_0,$$

for some  $\varepsilon > 0$ , such that  $-|a| + \varepsilon$  is still negative, leading to

$$c_{-} t^{\frac{1}{\kappa+1}} \le [c_{1} - c_{2} z_{t}] \le c_{+} t^{\frac{1}{\kappa+1}}, \quad t \ge t_{0},$$

by standard comparison arguments, where  $c_+ > c_- > 0$ , of course.

Using the definition of  $\hat{x}$ , the above sandwich-bound translates into

$$\frac{1}{c_{+}} t^{-\frac{2\beta}{(\kappa+1)(2\beta-1)}} \le \hat{x}^{2\beta}(t) \le \frac{1}{c_{-}} t^{-\frac{2\beta}{(\kappa+1)(2\beta-1)}}, \quad t \ge t_{0},$$

which means that, if  $T_{\infty} < +\infty$ , the exponent  $2\beta/(\kappa+1)/(2\beta-1)$  would have to be bigger than one, i.e.  $\beta < (\alpha+1)/2$ .

We continue with the proof of Lemma 2.2.

All in all, if  $a \neq 0$  and (2.4), then there exists a measure  $\mathbf{P}_+$  on  $\mathcal{G}_{\infty}^+$  such that equation (4.3) and Remark 4.3 remain true, when the measure  $\mathbf{Q}$  is replaced by  $\mathbf{P}_+$ . Furthermore, equation (4.1) can be written as

$$\hat{y}(t) = y_{-} + a \int_{0}^{t} \mathbf{1}_{[0,A_{1}]}(s) \,\hat{x}(s)^{2\beta - \alpha} \,ds + \sqrt{2T\gamma} \,\hat{W}_{+}(t) - \gamma \int_{0}^{t} \hat{y}(s) \,ds,$$

for all  $t \geq 0$ , **P**<sub>+</sub>-a.s., which gives

$$y_{+}(t) = y_{-} + a \int_{0}^{t} \frac{\mathbf{1}_{[0,A_{1}]}(A_{s})}{x_{+}(s)^{\alpha}} \, \mathrm{d}s + \sqrt{2T\gamma} \, \hat{W}_{+}(A_{t}) - \gamma \int_{0}^{t} \frac{y_{+}(s)}{x_{+}(s)^{2\beta}} \, \mathrm{d}s \quad (4.8)$$

since

$$A_t = \int_0^{A_t} \hat{x}(s)^{-2\beta} dT_s = \int_0^t x_+(s)^{-2\beta} ds,$$

for all  $t \geq 0$ ,  $\mathbf{P}_{+}$ -a.s.

Now, let  $\mathbb{F}_+$  be the time-changed filtration given by  $\mathcal{F}_t^+ \stackrel{\text{def}}{=} \mathcal{G}_{A_t}^+$ ,  $t \geq 0$ , so that

$$\mathbf{P}_{+} = \mathbf{Q}$$
 on  $\mathcal{F}_{0}^{+} = \mathcal{G}_{0}^{+}$ 

because  $\rho(0) = 1$ . Also, note that  $[x_+(t), y_+(t), t \ge 0]$  are  $\mathbb{F}_+$ -adapted processes which are both  $\mathbf{P}_+$ -a.s. continuous. Of course, when switching to the filtration  $\mathbb{F}_+$ , the measure  $\mathbf{P}_+$  can be restricted to  $\mathcal{F}_+^+$  which might be smaller than  $\mathcal{G}_+^+$ .

Next, the continuous local  $\mathbb{F}_+$ -martingale  $M_+(t) \stackrel{\text{def}}{=} \hat{W}_+(A_t)$ ,  $t \geq 0$ , has quadratic variation  $\langle M_+ \rangle = A$ . Since,  $\mathbf{P}_+$ -a.s., this quadratic variation takes the form,  $\int_0^t x_+(s)^{-2\beta} \, \mathrm{d}s$ ,  $t \geq 0$ , where the integrand  $x_+(s)^{-2\beta}$ ,  $s \geq 0$ , is positive and continuous, Theorem II.7.1 in [6] implies that there is an  $\mathbb{F}_+$ -Wiener-process  $W_+$  on  $(\Omega_+, \mathcal{F}_+^+, \mathbf{P}_+)$  such that

$$M_{+}(t) = \hat{W}_{+}(A_{t}) = \int_{0}^{t} \frac{dW_{+}(s)}{x_{+}(s)^{\beta}}, \quad t \ge 0, \, \mathbf{P}_{+}\text{-a.s.}$$

Hence, (4.8) translates into

$$y_{+}(t) = y_{-} + a \int_{0}^{t} \frac{\mathbf{1}_{[0,1]}(s)}{x_{+}(s)^{\alpha}} \, \mathrm{d}s + \sqrt{2T\gamma} \int_{0}^{t} \frac{\mathrm{d}W_{+}(s)}{x_{+}(s)^{\beta}} - \gamma \int_{0}^{t} \frac{y_{+}(s)}{x_{+}(s)^{2\beta}} \, \mathrm{d}s, \tag{4.9}$$

for all  $t \geq 0$ ,  $P_+$ -a.s., finally proving the lemma.

*Proof of Theorem 2.3.* As explained in Section 2 in the paragraph above the theorem, it suffices to show existence of a global weak solution.

Choose an arbitrary initial condition  $(x(0), y(0)) \in \mathbb{H}$ , set  $\Omega_{-} = \mathbb{R}^{2}$ ,  $\mathcal{F}_{-} = \mathcal{B}(\mathbb{R}^{2})$ , and denote by  $\mathbf{P}_{-}$  the Dirac measure at the point (x(0), y(0)). Let  $(x_{-}, y_{-})$  be the random variable on  $(\Omega_{-}, \mathcal{F}_{-}, \mathbf{P}_{-})$  induced by the identity on  $\mathbb{R}^{2}$ . Observe that  $\mathbf{P}_{-}(\{x_{-} > 0\}) = 1$  is an immediate consequence of x(0) > 0.

Hence, there is a tupel  $(\Omega_+, \mathcal{F}_{\infty}^+, \mathbb{F}_+, \mathbf{P}_+, [x_+(t), y_+(t), W_+(t), t \geq 0])$  the components of which satisfy the properties stated in the conclusion of Lemma

2.2. Moreover, using  $dx_+(t) = y_+(t) dt$  and (4.9) when multiplying  $x_+(t)^{\beta}$  by  $y_+(t)$ , we obtain that

$$\sqrt{2T\gamma} W_{+}(t) = x_{+}(t)^{\beta} y_{+}(t) - x_{-}^{\beta} y_{-} - \beta \int_{0}^{t} x_{+}(s)^{\beta-1} y_{+}(s)^{2} ds 
- a \int_{0}^{t} \frac{\mathbf{1}_{[0,1]} ds}{x_{+}(s)^{\alpha-\beta}} + \gamma \int_{0}^{t} \frac{y_{+}(s) ds}{x_{+}(s)^{\beta}}, \tag{4.10}$$

for all  $t \geq 0$ ,  $\mathbf{P}_+$ -a.s., and hence  $W_+$  can be considered a Wiener process with respect to the filtration  $[\sigma(\{(x_+(s),y_+(s)):s\leq t\}),\,t\geq 0]$ . Note that the proof of Lemma 2.2 makes clear that no extra sets of measure zero have to be added to this filtration.

The next step is to construct, by induction, a sequence  $(\Omega_n, \mathcal{F}_n, \mathbb{F}_n, \mathbf{P}_n, [x_n(t), y_n(t), W_n(t), t \ge 0]), n = 1, 2, \ldots$ , such that

$$x_n(t) > 0,$$
  
 $x_n(t) = x(0) + \int_0^t y_n(s) \, ds,$   
 $y_n(t) = y(0) + a \int_0^t \frac{ds}{x_n(s)^{\alpha}} + \sqrt{2T\gamma} \int_0^t \frac{dW_n(s)}{x_n(s)^{\beta}} - \gamma \int_0^t \frac{y_n(s) \, ds}{x_n(s)^{2\beta}},$ 

for all  $t \in [0, n]$ ,  $\mathbf{P}_n$ -a.s., where  $[x_n(t), y_n(t), t \ge 0]$  are continuous processes,  $\mathbb{F}_n$  stands for the filtration  $\mathcal{F}_t^n = \sigma(\{(x_n(s), y_n(s)) : s \le t\}), t \ge 0$ , and  $[W_n(t), t \ge 0]$  is an  $\mathbb{F}_n$ -Wiener process.

Observe that the tupel  $(\Omega_+, \mathcal{F}_{\infty}^+, \mathbb{F}_1, \mathbf{P}_+, [x_+(t), y_+(t), W_+(t), t \geq 0])$  found in the first part of the proof plays the role of the initial case n = 1, of course.

So, fix  $n \geq 2$ , and suppose that  $(\Omega_{n-1}, \mathcal{F}_{n-1}, \mathbb{F}_{n-1}, \mathbf{P}_{n-1}, [x_{n-1}(t), y_{n-1}(t), W_{n-1}(t), t \geq 0])$  has already been constructed.

Reset  $\Omega_{-} = \Omega_{n-1}$ ,  $\mathcal{F}_{-} = \mathcal{F}_{n-1}$ ,  $\mathbf{P}_{-} = \mathbf{P}_{n-1}$ , and choose  $x_{-} = x_{n-1}(n-1)$ ,  $y_{-} = y_{n-1}(n-1)$ . Then, again by Lemma 2.2, there is a corresponding tupel  $(\Omega_{+}, \mathcal{F}_{\infty}^{+}, \mathbb{F}_{+}, \mathbf{P}_{+}, [x_{+}(t), y_{+}(t), W_{+}(t), t \geq 0])$  which we now denote by  $(\Omega_{n}, \mathcal{F}_{n}, \mathbb{F}_{+}, \mathbf{P}_{n}, [x_{+}(t), y_{+}(t), W_{+}(t), t \geq 0])$ .

**Remark 4.6.** In Lemma 2.2, the filtration  $\mathbb{F}_+$  was given by

$$\mathcal{F}_t^+ = \mathcal{G}_{A_t}^+, \text{ using } \mathcal{G}_t^+ = \sigma(x_-, y_-) \vee \sigma(\{B_s : s \leq t\}),$$

for the purpose of Remark 4.4. But, when applying Lemma 2.2 in the context of the present proof, we are going to work with

$$\mathcal{G}_t^+ = \sigma(\{(x_{n-1}(s), y_{n-1}(s)) : s < n-1\}) \vee \sigma(\{B_s : s < t\})$$

instead, without violating the truth of Remark 4.4.

Recall that  $[x_{n-1}(t), y_{n-1}(t), W_{n-1}(t), t \ge 0]$  are extended to  $(\Omega_n, \mathcal{F}_n, \mathbf{P}_n)$  in the canonical way without changing their notation. Define, for  $t \ge 0$ ,

$$x_n(t) = x_{n-1}(t \wedge (n-1)) + x_+((t-n+1) \vee 0) - x_{n-1}(n-1),$$
  
$$y_n(t) = y_{n-1}(t \wedge (n-1)) + y_+((t-n+1) \vee 0) - y_{n-1}(n-1),$$

$$W_n(t) = W_{n-1}(t \wedge (n-1)) + W_+((t-n+1) \vee 0),$$

and build a filtration  $\hat{\mathbb{F}}_n$  from both  $\mathbb{F}_{n-1}$  and  $\mathbb{F}_+$  by

$$\hat{\mathcal{F}}_t^n \stackrel{\text{def}}{=} \left\{ \begin{array}{ll} \mathcal{F}_t^{n-1} & : & t < n-1, \\ \\ \mathcal{F}_{t-n+1}^+ & : & t \ge n-1. \end{array} \right.$$

All in all, because of

$$\mathbf{P}_n = \mathbf{P}_+ = \mathbf{Q} = \mathbf{P}_- \otimes \mathbf{P}_W = \mathbf{P}_{n-1} \otimes \mathbf{P}_W$$
 on  $\mathcal{F}_0^+$ ,

and because  $\mathcal{F}_t^{n-1} \subseteq \mathcal{F}_0^+$  (see Remark 4.6), the processes  $[x_n(t), y_n(t), W_n(t), t \geq 0]$  would have all properties needed for the induction step, except that  $W_n$  is a Wiener process with respect to the filtration  $\hat{\mathbb{F}}_n$  which is possibly bigger than  $\mathbb{F}_n$ .

However, as a consequence of (4.9), we also have

$$y_n(t) = y(0) + a \int_0^t \frac{\mathbf{1}_{[0,n]} \, \mathrm{d}s}{x_n(s)^{\alpha}} + \sqrt{2T\gamma} \int_0^t \frac{\mathrm{d}W_n(s)}{x_n(s)^{\beta}} - \gamma \int_0^t \frac{y_n(s) \, \mathrm{d}s}{x_n(s)^{2\beta}},$$

for all  $t \geq 0$ ,  $\mathbf{P}_n$ -a.s., leading to the nth-step analogue of (4.10), i.e.,

$$\sqrt{2T\gamma} W_n(t) = x_n(t)^{\beta} y_n(t) - x(0)^{\beta} y(0) - \beta \int_0^t x_n(s)^{\beta - 1} y_n(s)^2 ds$$
$$- a \int_0^t \frac{\mathbf{1}_{[0,n]} ds}{x_n(s)^{\alpha - \beta}} + \gamma \int_0^t \frac{y_n(s) ds}{x_n(s)^{\beta}}, \tag{4.11}$$

for all  $t \geq 0$ ,  $\mathbf{P}_n$ -a.s., so that  $W_n$  is  $\mathbb{F}_n$ -adapted, and therefore it must be an  $\mathbb{F}_n$ -Wiener process, too.

The next step of the proof consists in constructing a measure on  $((\mathbb{R}^2)^{[0,\infty)}, \mathcal{B}((\mathbb{R}^2)^{[0,\infty)}))$  whose finite-dimensional distributions are induced by the laws of the two-dimensional processes  $[x_n(t), y_n(t), t \geq 0], n = 1, 2, \ldots$ , in the following way.

For an arbitrary finite sequence of non-negative mutually different numbers  $\mathbf{t} = (t_1, \dots, t_k)$ , define

$$\mathbf{P_t}(\Gamma) = \mathbf{P_n}(\overbrace{\{(x_n(t_1), y_n(t_1), \dots, x_n(t_k), y_n(t_k)) \in \Gamma\}}^{cylinder\ set}), \quad \Gamma \in \mathcal{B}(\mathbb{R}^{2k}),$$

where n is the smallest integer such that  $n \ge \max\{t_1, \ldots, t_k\}$ . Then,  $\{\mathbf{P_t}\}$  is a consistent family of finite-dimensional distributions in the sense of Kolmogorov. Hence, there is a probability measure  $\mathbf{P}$  on  $((\mathbb{R}^2)^{[0,\infty)}, \mathcal{B}((\mathbb{R}^2)^{[0,\infty)}))$  satisfying

$$\mathbf{P_t}(\Gamma) = \mathbf{P}(\{(x(t_1), y(t_1), \dots, x(t_k), y(t_k)) \in \Gamma\}), \quad \Gamma \in \mathcal{B}(\mathbb{R}^{2k}),$$

where  $[x(t), y(t), t \geq 0]$  denotes the coordinate process on  $\Omega = (\mathbb{R}^2)^{[0,\infty)}$ . Let  $\mathcal{F}$  be the completion of  $\mathcal{B}((\mathbb{R}^2)^{[0,\infty)})$  with respect to  $\mathbf{P}$ , and let  $\mathbb{F}$  be the filtration obtained by  $\mathbf{P}$ -augmentation of  $\sigma(\{x(s), y(s) : s \leq t\})$ ,  $t \geq 0$ .

Note that, almost surely,  $[x(t), y(t), t \ge 0]$  is a pair of continuous  $\mathbb{F}$ -adapted processes. This is seen by two simple arguments. First, since<sup>2</sup>

$$\mathbf{P}_n(\{t \mapsto (x_n(t), y_n(t)) \text{ continuous on } [0, n]\}) = 1,$$

each of the events  $\{t \mapsto (x(t), y(t)) \text{ continuous on } [0, n]\}$  is in  $\mathcal{F}$ ,  $n = 1, 2, \ldots$ , and second,

$$\mathbf{P}(\{t \mapsto (x(t), y(t)) \text{ continuous on } [0, n]\})$$

$$= \mathbf{P}_n(\{t \mapsto (x_n(t), y_n(t)) \text{ continuous on } [0, n]\}),$$

for each n > 1.

In what follows,  $[x(t), y(t), t \ge 0]$  always stands for a fixed continuous version indistinguishable of the coordinate process.

Of course, in a similar way, one shows that x(t) > 0,  $t \ge 0$ , P-a.s., as well as

$$x(t) = x(0) + \int_0^t y(s) ds, \quad t \ge 0,$$
 **P**-a.s. (4.12)

Now, introduce

$$W(t) \stackrel{\text{def}}{=} \frac{x(t)^{\beta}y(t) - x(0)^{\beta}y(0)}{\sqrt{2T\gamma}} - \beta \int_{0}^{t} \frac{x(s)^{\beta - 1}y(s)^{2}}{\sqrt{2T\gamma}} ds$$
$$- a \int_{0}^{t} \frac{ds}{\sqrt{2T\gamma}x(s)^{\alpha - \beta}} + \gamma \int_{0}^{t} \frac{y(s) ds}{\sqrt{2T\gamma}x(s)^{\beta}}, \quad t \ge 0,$$

and observe that this process is a Wiener process because, by (4.11), it satisfies

$$\mathbf{P}(\{(W(t_1),\ldots,W(t_k))\in\Gamma\}) = \mathbf{P}_n(\{(W_n(t_1),\ldots,W_n(t_k))\in\Gamma\}),$$

for every  $\mathbf{t} = (t_1, \dots, t_k)$ ,  $n \ge \max\{t_1, \dots, t_k\}$  and  $\Gamma \in \mathcal{B}(\mathbb{R}^k)$ , and because the continuity of  $[x(t), y(t), t \ge 0]$  makes it a continuous process. Furthermore, as  $W_n$  can be considered a Wiener process with respect to the filtration obtained by  $\mathbf{P}_n$ -augmentation of  $\sigma(\{(x_n(s), y_n(s)) : s \le t\})$ ,  $t \ge 0$ , one can also consider W to be an  $\mathbb{F}$ -Wiener-process.

Finally, again using the corresponding property on  $(\Omega_n, \mathcal{F}_n, \mathbf{P}_n)$ , each of the processes  $[y(t \wedge n), t \geq 0]$  is an  $\mathbb{F}$ -semimartingale,  $n = 1, 2, \ldots$ , and hence  $[y(t), t \geq 0]$  is one, too. As a consequence, by partial integration,

$$\sqrt{2T\gamma}W(t) = \int_0^t x(s)^\beta \,\mathrm{d}y(s) - a \int_0^t \frac{\mathrm{d}s}{x(s)^{\alpha-\beta}} + \gamma \int_0^t \frac{y(s) \,\mathrm{d}s}{x(s)^\beta}, \quad t \ge 0, \, \mathbf{P}\text{-a.s.},$$

follows from (4.12) and the definition of W(t). Thus, using the above right-hand side when calculating  $\int_0^t x(s)^{-\beta} dW(s)$  for any  $t \geq 0$ , eventually proves the theorem.

<sup>&</sup>lt;sup>2</sup>As this probability can be approximated by probabilities of cylinder sets.

Proof of Proposition 2.4. To begin with, assume  $a \in \mathbb{R}$ , and let  $[x(t), y(t), t \geq 0]$  be the solution of (2.1), started at  $(x(0), y(0)) \in \mathbb{H}$ , and driven by a Wiener process  $[W(t), t \geq 0]$ , given on a probability space  $(\Omega, \mathcal{F}, \mathbf{P})$ , i.e.,

$$x(t) = x(0) + \int_0^t y(s) \, ds,$$
  
$$y(t) = y(0) + a \int_0^t \frac{ds}{x(s)^{\alpha}} + \sqrt{2T\gamma} \int_0^t \frac{dW(s)}{x(s)^{\beta}} - \gamma \int_0^t \frac{y(s)}{x(s)^{2\beta}} \, ds,$$

for all t > 0, **P**-a.s.

Since the stochastic integral in the above equation is well-defined for all  $t \geq 0$ , its quadratic variation,  $\tilde{A}_t = \int_0^t x(s)^{-2\beta} ds$ , is well-defined for all  $t \geq 0$ , too, and

$$\tilde{T}_t = \inf\{s \ge 0 : \tilde{A}_s > t\}, \quad t \ge 0,$$

can be used as a time-change. Furthermore, since x(t) > 0, for all  $t \ge 0$ , **P**-a.s., this time-change is strictly increasing and continuous on  $[0, \tilde{A}_{\infty})$ , **P**-a.s., and

$$\mathbf{P}(\{\lim_{t \uparrow \tilde{A}_{\infty}} \tilde{T}_t = +\infty\}) = 1. \tag{4.13}$$

So, on the one hand, the time-changed processes,  $t \mapsto \tilde{x}(t) = x(\tilde{T}_t)$  and  $t \mapsto \tilde{y}(t) = y(\tilde{T}_t)$ , are almost surely continuous processes on  $[0, \tilde{A}_{\infty})$ . On the other hand, since

$$\tilde{T}_t = \int_0^t \tilde{x}(s)^{2\beta} \, \mathrm{d}s, \quad t < \tilde{A}_\infty, \text{ P-a.s.},$$

property (4.13) implies that

$$\limsup_{t \uparrow \tilde{A}_{\infty}} \tilde{x}(t) = +\infty, \quad \text{on } \{\tilde{A}_{\infty} < \infty\}, \quad \textbf{P-a.s.}, \tag{4.14}$$

and we are going to show next that, on  $\{\tilde{A}_{\infty} < \infty\}$ ,  $\liminf_{t \uparrow \tilde{A}_{\infty}} \tilde{x}(t)$  cannot be finite with positive probability, either, even if the parameter a is negative.

Applying the time-change to the above equation yields

$$\tilde{x}(t) = x(0) + \int_0^t \tilde{x}(s)^{2\beta} \, \tilde{y}(s) \, ds, 
\tilde{y}(t) = y(0) + a \int_0^t \tilde{x}(s)^{2\beta - \alpha} \, ds + \sqrt{2T\gamma} \, \tilde{W}(t) - \gamma \int_0^t \tilde{y}(s) \, ds, \tag{4.15}$$

for all  $t < \tilde{A}_{\infty}$ , **P**-a.s., where  $[\tilde{W}(t), t \ge 0]$  is another Wiener process on a possibly enlarged<sup>3</sup> probability space (cf. Theorem II.7.2' in [6]).

Of course, by the continuity properties of the time-changed processes,

$$\tilde{x}(t) = \left[ x(0)^{1-2\beta} - (2\beta - 1) \int_0^t \tilde{y}(s) \, ds \right]^{\frac{-1}{2\beta - 1}},$$

<sup>&</sup>lt;sup>3</sup>By standard convention, the enlarged space is denoted by  $(\Omega, \mathcal{F}, \mathbf{P})$ , too.

for at least all  $t < \tilde{A}_{\infty},$  **P**-a.s., where

$$\tilde{y}(s) = e^{-\gamma s} y(0) + \sqrt{2T\gamma} \int_0^s e^{-\gamma(s-r)} d\tilde{W}(r) + a \int_0^s e^{-\gamma(s-r)} \tilde{x}(r)^{2\beta-\alpha} dr,$$

for all  $s < \tilde{A}_{\infty}$ , **P**-a.s.

The question is now whether different sequences of time points,  $(t_n)_{n=1}^{\infty}$ , converging to  $\tilde{A}_{\infty} < \infty$ , can lead to different limits of  $\tilde{x}(t_n)$ ,  $n \to \infty$ , which can only happen if  $\int_0^{t_n} \tilde{y}(s) \, \mathrm{d}s$  has different limits for different sequences of time points, which can only happen if  $\tilde{y}(t_n)$  has different limits for different sequences of time points.

The key to the answer of this question is writing

$$\tilde{y}(s)$$
 as  $\hat{y}(s) + ae^{-\gamma s}R_s$ , (4.16)

where  $[\hat{y}(s), s < \tilde{A}_{\infty}]$  is indistinguishable of an Ornstein-Uhlenbeck process restricted to  $[0, \tilde{A}_{\infty})$ , and

$$R_s = \int_0^s e^{\gamma r} \, \tilde{x}(r)^{2\beta - \alpha} \, \mathrm{d}r, \quad s < \tilde{A}_{\infty}.$$

Note that  $s \mapsto R_s$  is almost surely a continuous monotone function on  $[0, \tilde{A}_{\infty})$ , so that  $ae^{-\gamma t_n}R_{t_n}$  can only have one limit for any sequence of time points converging to  $\tilde{A}_{\infty}$ , on  $\{\tilde{A}_{\infty} < \infty\}$ , **P**-a.s., and the same applies to  $\hat{y}(t_n)$ , as this function can almost surely be extended to a continuous function on  $[0, \infty)$ .

All in all,  $\liminf_{t\uparrow \tilde{A}_{\infty}} \tilde{x}(t)$  has indeed to coincide with  $\limsup_{t\uparrow \tilde{A}_{\infty}} \tilde{x}(t)$ , on  $\{\tilde{A}_{\infty} < \infty\}$ , **P**-a.s., proving

$$\lim_{t\to\infty} x(t) = +\infty$$
, on  $\{\tilde{A}_{\infty} < \infty\}$ , **P**-a.s., (4.17)

by (4.14), for any parameter  $a \in \mathbb{R}$ , because  $x(t) = \tilde{x}(\tilde{A}_t), t \geq 0$ , **P**-a.s.

In the second part of the proof, we will show that the event  $\{\tilde{A}_{\infty} < \infty\}$  has probability one, for any  $a \geq 0$ , eventually proving the proposition.

First, the Ornstein-Uhlenbeck process used in (4.16) and the process  $[\hat{y}(t), t \ge 0]$  used in the proof of Lemma 2.2 have the same law, when started at y(0), justifying the same notation. Therefore,

$$\hat{x}(t) = \left[ x(0)^{1-2\beta} - (2\beta - 1) \int_0^t \hat{y}(s) \, ds \right]^{\frac{-1}{2\beta - 1}}, \ t < \tau,$$

$$\tau = \inf\{ s \ge 0 : \int_0^s \hat{y}(r) \, dr = \frac{x(0)^{1-2\beta}}{2\beta - 1} \},$$

and the corresponding objects given in the proof of Lemma 2.2 have the same law, too, when using x(0) instead of  $x_{-}$ , so that

$$\mathbf{P}(\{\tau < \infty\} \cap \{\lim_{t \uparrow \tau} \hat{x}(t) = +\infty\}) = 1 \tag{4.18}$$

can easily be derived from Lemma 4.1. Furthermore,

$$\hat{x}(t) = x(0) + \int_0^t \hat{x}(s)^{2\beta} \, \hat{y}(s) \, ds, 
\hat{y}(t) = y(0) + \sqrt{2T\gamma} \, \tilde{W}(t) - \gamma \int_0^t \hat{y}(s) \, ds,$$
(4.19)

for all  $t < \tau$ , **P**-a.s.

The next step is to show that, if  $a \geq 0$ , then  $(\tilde{x}(t), \tilde{y}(t))$  satisfying (4.15) dominates  $(\hat{x}(t), \hat{y}(t))$  satisfying (4.19), for all  $t < \tau \land \hat{A}_{\infty}$ . In fact, since the drift coefficient of equation (4.19), i.e.  $(\hat{x}, \hat{y}) \mapsto (\hat{x}^{2\beta} \hat{y}, -\gamma \hat{y})$ , is quasi-monotonously increasing (cf. Def.3.1 in [1]), and since solutions to (4.19) are pathwise unique up to  $\tau \land \hat{A}_{\infty}$ , and since the difference of the drift coefficients of (4.15) and (4.19) is a vector field on  $\mathbb{H}$  with non-negative components, it follows from Prop.3.3 in [1] that

$$\tilde{x}(t) > \hat{x}(t), \quad \tilde{y}(t) > \hat{y}(t), \quad t < \tau \wedge \tilde{A}_{\infty}, \text{ P-a.s.}$$

**Remark 4.7.** The results in [1] were obtained for coefficients defined on  $\mathbb{R}^d$ , but it is an easy exercise to show their validity for coefficients defined on domains like our half-plane,  $\mathbb{H}$ .

Now, if  $\tilde{A}_{\infty}$  was bigger than  $\tau$  with positive probability, then the process  $t \mapsto \tilde{x}(t)$ , being continuous on  $[0, \tilde{A}_{\infty})$ , would blow up before  $\tilde{A}_{\infty}$ , by (4.18), which is a contradiction. Thus,  $\tilde{A}_{\infty}$  is almost surely bounded by  $\tau$  from above leading to  $\mathbf{P}(\{\tilde{A}_{\infty} < \infty\}) = 1$ , again by (4.18).

#### References

- Assing, S., Manthey, R.: The behavior of solutions of stochastic differential inequalities. *Probab. Theory Relat. Fields*, 103, (1995), 493–514. MR-1360202
- [2] Bang, O., Christiansen, P.L., Gaididei, Y.B., If, F., Rasmussen, K.O.: Temperature effects in a nonlinear model of monolayer Scheibe aggregates. *Physical Review E*, **49**, (1994), no. 5, 4627–4635.
- [3] Christiansen, P.L., Gaididei, Y.B., Johansson, M., Rasmussen, K.O., Yakimenko, I.I.: Collapse of solitary excitations in the nonlinear Schrödinger equation with nonlinear damping and white noise. *Physical Review E*, **54**, (1996), no. 1, 924–930.
- [4] Cooke, B., Herzog, D.P., Mattingly, J.C., McKinley, S.A., Schmidler, S.C.: Geometric Ergodicity of Tow-Dimensional Hamiltonian Systems with a LENARD-JONES-LIKE Repulsive Potential. ArXiv e-prints, arxiv:1104.3842v2 (2016). 39pp.
- [5] Funaki, T., Ohnawa, M., Suzuki, Y., Yokoyama, S.: Existence and uniqueness of solutions to stochastic Reyleigh-Plesset equations. J. Math. Anal. Appl., 425, (2015), 20–32. MR-3299647
- [6] Ikeda, N. and Watanabe, S.: Stochastic Differential Equations and Diffusion Processes. Second edition. North-Holland mathematical library 24. North-Holland Publishing, Amsterdam, 1989. xiv+555 pp. MR-1011252
- [7] Karatzas, I. and Shreve, E.: Brownian Motion and Stochastic Calculus. Second edition. Graduate Texts in Mathematics 113. Springer Verlag, New York, 1991. xxiv+470 pp. MR-1121940
- [8] Merle, F.: Construction of solutions with exactly k blow-up points for the Schrödinger equation with critical nonlinearity. *Comm. Math. Phys.* **129**, (1990), no. 2, 223–240. MR-1048692
- [9] Merle, F., Tsutsumi, Y.:  $L^2$ -concentration of blow up solutions for the non-linear Schrödinger equation with critical power nonlinearity. *J. Diff. Eq.* **84**, (1990), 205–214. MR-1047566
- [10] Nawa, H.: Asymptotic and limiting profiles of blowup solutions of the non-linear Schrödinger Equation with critical power. Comm. Pure Appl. Math. 52, (1999), no. 2, 193–270. MR-1653454
- [11] Weinstein, M.I.: Nonlinear Schrödinger equations and sharp interpolation estimates. Comm. Math. Phys. 87, (1983), 567–576. MR-0691044