

## A NOTE ON HARMONIC FUNCTIONS ON SURFACES

JEAN C. CORTISSOZ

ABSTRACT. We review and give elementary proofs of Liouville type properties of harmonic and subharmonic functions in the plane endowed with a complete Riemannian metric, and prove a gap theorem for the possible growth of harmonic functions when this metric has nonnegative Gaussian curvature.

## 1. INTRODUCTION.

Given a function  $u : \Omega \rightarrow \mathbb{R}$ , where  $\Omega$  is an open subset of  $\mathbb{R}^2$ , the Laplacian is defined, in rectangular coordinates, as

$$\Delta u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2},$$

and we say that  $u$  is harmonic (resp. subharmonic) if  $\Delta u = 0$  ( $\Delta u \geq 0$ ). The classical Liouville's Theorem in  $\mathbb{R}^2$  states that a bounded harmonic function is constant (For a beautiful proof of this fact we recommend [9]). A stronger version says that if  $u$  is a subharmonic function bounded above then it must be constant; we refer to this property of the plane as parabolicity, and, no doubt, it is an amazing fact that being a solution to a partial differential identity or a partial differential inequality may determine the growth properties of a function.

Over the years, the analysis studied on  $\mathbb{R}^n$  has been carried over to Riemannian manifolds, a realm where a differential and an inner product structure coexist. In particular, a Laplacian operator acting over functions can be defined, and hence it is a framework in which the concept of harmonic, subharmonic and superharmonic function have a natural extension. So it is also natural to ask which properties of harmonic functions, such as Liouville's Theorem, or parabolicity are preserved in a Riemannian manifold.

In this paper we are interested in Liouville type theorems and gap theorems on surfaces with a pole (i.e., surfaces where polar coordinates can be defined). By a Liouville type property we mean a theorem that states that if a harmonic function is conveniently bounded then it must be constant, and by a gap theorem we mean a theorem that imposes restriction on how fast a harmonic function must grow so that it does not belong to a class of strictly lower growth.

Before starting to throw definitions formulas and theorems at the reader, let us mention some interesting results related to the work we will present in this paper. Regarding Liouville type theorems, of great importance are the results of Ahlfors and Milnor, which in the case of surfaces endowed with a rotationally symmetric metric relates an intrinsic quantity, the curvature, to the behavior of subharmonic functions, and to be more precise to the parabolicity of the surface. Green and Wu ([6]) extended the Ahlfors-Milnor theorem to the case of surfaces with

a pole. We will give a relatively simple proof of part of the Ahlfors-Milnor-Greene-Wu parabolicity criterion, in which our main tool will be the Strong Maximum Principle.

Regarding gap properties for harmonic functions, on the classical side, that is, in the complex plane, it is well known that if the rate of growth of a harmonic function is bounded by a power of the distance to a fixed point, then it must be a polynomial. This is a consequence of the analyticity of harmonic functions in  $\mathbb{R}^2$  and of the Cauchy estimates. A most recent result has been proved by Ni and Tam in [4]: Here the authors show how fast, in a Kähler manifold of positive bisectional curvature, a superlinear harmonic function must grow. As a treat for the reader, we improve upon Ni and Tam's result in the case of a surface with a pole: This is the only new result in this paper (at least to the best of our knowledge).

The main ideas of our proofs are contained in, and someone could even say they are transplanted, via a classical comparison theorem due to Sturm, from the beautiful book "Maximum Principles in Differential Equations" [8], which we highly recommend. We also hope that this note serves as an introduction, assuming as little as possible prerequisites from the reader, to the study of harmonic functions in Riemannian geometry.

## 2. PRELIMINARIES.

For the convenience of the reader, let us give a quick review of a few concepts in Riemannian geometry that we shall be using in what follows. We will consider  $\mathbb{R}^2$  and fix polar coordinates  $(r, \theta)$  with respect to the origin, and we will endow it with a family of inner products of the form

$$g = dr^2 + (f(r, \theta))^2 d\theta^2,$$

where  $f : (0, \infty) \times [0, 2\pi] \rightarrow (0, \infty)$  is a smooth function such that

$$f(r, 0) = f(r, 2\pi), \quad \lim_{r \rightarrow 0^+} f(r, \theta) = 0 \quad \text{and} \quad \lim_{r \rightarrow 0^+} f'(r, \theta) = 1,$$

where we have used (and will use in what follows)  $f'$  to denote differentiation with respect to  $r$ .

For those not familiar with Riemannian manifolds,  $g$  represents a way of measuring vectors, and it is called a Riemannian metric. It defines an inner product for vectors based at the point  $(r, \theta)$ , and represented in the basis

$$\frac{\partial}{\partial r} = \frac{x}{\sqrt{x^2 + y^2}} \mathbf{i} + \frac{y}{\sqrt{x^2 + y^2}} \mathbf{j}, \quad \frac{\partial}{\partial \theta} = y \mathbf{i} - x \mathbf{j}$$

in the following way. If we have

$$\mathbf{v}_j = a_j \frac{\partial}{\partial r} + b_j \frac{\partial}{\partial \theta},$$

based at the point  $(r, \theta)$  (the point  $(r \cos \theta, r \sin \theta)$  in rectangular coordinates), then

$$g(\mathbf{v}_1, \mathbf{v}_2) = a_1 a_2 + b_1 b_2 (f(r, \theta))^2.$$

Notice that with the choice  $f(r, \theta) = r$  we obtain the usual inner product of vectors in the plane. The pair  $(M, g)$  is called a Riemannian surface (as opposed to a Riemann surface), and, as discovered by Gauss, Riemannian surfaces have an

important intrinsic estimate: the curvature. From the expression for a Riemannian metric given above, the curvature can be computed as

$$K_g(r, \theta) = -\frac{f''(r, \theta)}{f(r, \theta)}.$$

In a Riemannian surface the gradient of a function  $u : M \rightarrow \mathbb{R}$  can be defined, due to the fact that given a nondegenerate scalar product a metric dual of the derivative of a function can be defined. In our case then, the gradient of  $u$  can be computed as

$$\nabla_g u = \frac{\partial u}{\partial r} \frac{\partial}{\partial r} + \frac{1}{f^2} \frac{\partial u}{\partial \theta} \frac{\partial}{\partial \theta}.$$

We can also define a Laplacian, which is the operator of our utmost interest:

$$\Delta_g = \frac{\partial^2}{\partial r^2} + \frac{f'}{f} \frac{\partial}{\partial r} + \frac{1}{f} \frac{\partial^2}{\partial \theta^2} - \frac{f_\theta}{f^3} \frac{\partial}{\partial \theta},$$

and here  $f_\theta$  denotes  $\frac{\partial f}{\partial \theta}$ .

Given a Laplacian, we can define a  $C^2$  function  $u$  as harmonic if  $\Delta_g u = 0$ , subharmonic if  $\Delta_g u \geq 0$  and superharmonic if  $\Delta_g u \leq 0$ .

The term  $h_g := \frac{f'}{f}$  in the expression for the Laplacian is rather important for the following discussion. It gives the mean curvature of the circle of radius  $r$  with respect to the metric  $g_0$ . We will need to estimate this term, and the tool we will employ is the following comparison, due to Sturm, and which is known as the Laplacian Comparison Theorem among geometers.

**Theorem 1.** *Let  $h, f : [a, \infty) \rightarrow (0, \infty)$  be  $C^2$  functions. If  $\frac{f''}{f}(r) \leq \frac{h''}{h}(r)$ ,  $r > a$ , and  $\frac{f'}{f}(a) \leq \frac{h'}{h}(a)$  then  $\frac{f'}{f}(r) \leq \frac{h'}{h}(r)$  for  $r > a$ .*

The proof of this theorem is based upon the following observation:

$$\left(\frac{w'}{w}\right)' = -\left(\frac{w'}{w}\right)^2 + \frac{w''}{w}.$$

Notice then that the hypothesis imply that  $\left(\frac{f'}{f}(r)\right)' \leq \left(\frac{h'}{h}(r)\right)'$  whenever  $\frac{f''}{f}(r) = \frac{h''}{h}(r)$ .

Another important tool in the arguments that follow is the Maximum Principle.

**Theorem 2.** *Let  $\Omega$  be a bounded open subset of  $\mathbb{R}^2$ ,  $\mathbb{R}^2$  endowed with a Riemannian metric  $g$ , and let  $u : \overline{\Omega} \rightarrow \mathbb{R}$  be a  $C^2(\Omega) \cap C(\overline{\Omega})$  function. Then we have that:*

- (i) *if  $\Delta_g u \leq 0$  and  $u$  has a minimum in the interior of  $\Omega$  then  $u$  is constant.*
- (ii) *if  $\Delta_g u \geq 0$  and  $u$  has a maximum in the interior of  $\Omega$  then  $u$  is constant.*

This theorem follows from the Strong Maximum Principle for elliptic operators (See Section 3.2 in [5], in particular Theorem 3.5).

## 3. BOUNDED HARMONIC FUNCTIONS.

Before we state the main result of this section, we shall make an observation. Let  $z : [a, \infty) \rightarrow (0, \infty)$  be a  $C^2$  function. Direct calculation shows that the function

$$h(r) = \int_a^r \frac{1}{z(\sigma)} d\sigma$$

satisfies the identity

$$\frac{d^2 h}{dr^2} + \frac{z'(r)}{z(r)} \frac{dh}{dr} = 0,$$

i.e, it is harmonic with respect to the metric

$$g_0 = dr^2 + z(r)^2 d\theta^2,$$

defined on the complement of the open ball of radius  $a > 0$ .

Hence if  $g = dr^2 + (f(r, \theta))^2 d\theta^2$  is a metric on  $\mathbb{R}^2$  such that  $K_g(r, \theta) \geq -\frac{z''}{z}(r)$ , then  $\Delta_g h \geq 0$ . Indeed, notice that  $\lim_{r \rightarrow a^+} \frac{z'}{z}(r) = \infty$ , and hence for  $a_0$  close to  $a$ ,  $\frac{f'}{f}(a) \leq \frac{z'}{z}(a)$ , so the hypothesis of Sturm's comparison theorem hold, and we have that  $\frac{f'}{f}(r) \leq \frac{z'}{z}(r)$ , for  $r \geq a$ . Therefore,

$$\Delta_g h = \frac{d^2 h}{dr^2} + \frac{f'}{f} \frac{dh}{dr} = \frac{d^2 h}{dr^2} + \frac{f'}{f} \frac{1}{z} \leq \frac{d^2 h}{dr^2} + \frac{z'}{z} \frac{1}{z} = 0.$$

Given a continuous function  $v$  define

$$M(v; r) = \sup_{\theta \in [0, 2\pi)} |v(r, \theta)|.$$

We are ready to show the following result.

**Theorem 3.** *Let  $g = dr^2 + f(r, \theta)^2 d\theta^2$  be a metric on  $\mathbb{R}^2$ . Let  $z$  be as above, and assume that  $K_g(r, \theta) \geq -\frac{z''}{z}(r)$  for  $r$  large enough. Then, any subharmonic function  $u$  which satisfies*

$$\liminf_{r \rightarrow \infty} \frac{M(u; r)}{h(r)} = 0$$

*must be constant.*

*Proof.* In what follows we will denote by  $B_R$  the ball of radius  $R$  centered at  $(0, 0)$ . Fix  $R_1 > 0$ , fix  $\delta > 0$  also small, and define the function

$$w_{\delta, \eta} = u - \delta h(r) - M(u; R_1).$$

By hypothesis, we can take  $R_2$  larger than  $R_1$  and such that  $w_{\delta, \eta} \leq 0$  on both  $\partial B_{R_1}$  and  $\partial B_{R_2}$ . It is also clear from its definition that  $\Delta_g w_{\delta, \eta} \geq 0$ . From the Maximum Principle it follows that

$$w_{\delta, \eta} \leq 0$$

on the annulus of inner radius  $R_1$  and outer radius  $R_2$ . Since  $\delta > 0$  can be made arbitrarily small, we conclude that for all  $P$  in the annulus the estimate

$$|u(P)| \leq M(u; R_1)$$

holds, and hence that  $|u| \leq M_{R_1}$  on the ball of radius  $R_2$ , and from this via the Maximum Principle (as  $u$  attains its maximum at an interior point of  $B_{R_2}$ ) we

can conclude that  $u$  is constant in the ball of radius  $R_2$ . Since  $R_2$  can be taken arbitrarily large, the theorem follows.  $\square$

The proof given above is presented in [8] in the case of a flat metric (See Theorem 19 in Section 2 of [8] and the example thereafter), but its generalization to other metrics is straightforward. Besides, the previous theorem has the following interesting consequence:

**Corollary 1.** *Let  $u$  be a harmonic function on a surface such that for  $r \geq r_0 \geq 1$  its curvature function satisfies  $K_g \geq -\frac{1}{r^2 \log r}$ . If*

$$\liminf_{r \rightarrow \infty} \frac{M(u; r)}{\log \log r} = 0$$

*then  $u$  is constant. In particular, if  $u$  is bounded then it is constant.*

This corollary is due to Greene and Wu (Theorem D in [6]), and its proof is quite simple: take  $z(r) = r \log\left(\frac{r}{r_0}\right)$  for  $r \geq r_0$ . Notice that, since  $r_0 \geq 1$ , then

$$-\frac{z''}{z} = -\frac{1}{r^2 \log\left(\frac{r}{r_0}\right)} \leq -\frac{1}{r^2 \log r} \leq K_g.$$

On the other hand,  $\lim_{r \rightarrow r_0^+} \frac{z'}{z}(r) = +\infty$ , and hence by Sturm's comparison theorem,  $\Delta_g h \geq 0$ , with  $h$  as defined above; by Theorem 3 the result follows.

We want to point out that in higher dimensions, Liouville's Theorem has been extended by Yau, via his gradient estimate (see below), to manifolds of nonnegative Ricci curvature.

**3.1. On the Ahlfors-Milnor-Greene-Wu parabolicity criterion: an application of Hadamard's three circles theorem.** A surface  $M$  is called parabolic if any subharmonic function (i.e.  $\Delta u \geq 0$ ) bounded above is constant. Milnor in [7] showed that given a rotationally symmetric metric on  $\mathbb{R}^2$ , it is parabolic if for large enough  $r$  the curvature of the metric is larger than or equal to  $-\frac{1}{r^2 \log r}$ .

We use the Maximum Principle to give a proof of a generalization of Milnor's criterion (which is a theorem due to Greene and Wu). We assume that we have endowed  $\mathbb{R}^2$  with a metric of the form  $dr^2 + (f(r, \theta))^2 d\theta^2$  whose curvature is larger than or equal to  $-\frac{1}{r^2 \log r}$ .

First, we have the following version of Hadamard's three circles theorem. Let  $u$  be a subharmonic function, and let

$$M_r = \max_{\theta \in [0, 2\pi)} u(r, \theta).$$

Notice that if  $r_2 > r_1$  then  $M_{r_2} \geq M_{r_1}$ , by the Maximum Principle. So let  $r_1 < r < r_2$ , and define

$$\varphi(r) = \frac{1}{\log\left(\frac{\log r_2}{\log r_1}\right)} \left( M_{r_1} \log\left(\frac{\log r_2}{\log r}\right) + M_{r_2} \log\left(\frac{\log r}{\log r_1}\right) \right).$$

It is easy to check that  $\Delta\varphi \leq 0$ , and hence  $u - \varphi$  is subharmonic, and also  $u - \varphi \leq 0$  on both  $\partial B_1$  and  $\partial B_2$ . We can conclude via the Maximum Principle that

$$M_r \leq \varphi(r).$$

This last inequality is our version of Hadamard's three circles theorem. Now assume that  $u$  is bounded above. By taking  $r_2 \rightarrow \infty$ , we obtain the estimate

$$M_r \leq M_{r_1}.$$

But then, since  $r > r_1$ , and  $r$  is arbitrary by the Maximum Principle  $u$  must be constant. We can conclude that the surface is parabolic. Again, we must point out that our proof follows closely the arguments given in Chapter 2, Section 12 in [8].

#### 4. A GAP THEOREM FOR SURFACES WITH NONNEGATIVE GAUSSIAN CURVATURE.

It is an exercise in complex analysis to prove the following result. Given a holomorphic function  $f$  such that  $|f(z)| \leq C|z|^k + B$  then  $f$  is a polynomial of degree at most  $[k]$ . From this we can conclude that there are gaps between the possible growth that a holomorphic, and in consequence, harmonic functions can have. For instance, a harmonic function of subquadratic growth (i.e.,  $k < 2$ ) must be of at most linear growth (i.e.,  $k$  must be less or equal to 1).

In this section we prove a gap theorem on the possible growth of harmonic functions on a complete noncompact surface  $M$  of positive Gaussian curvature. To be more precise we will show the following gap theorem.

**Theorem 4.** *Let  $g = dr^2 + f(r, \theta)^2 d\theta^2$  be a metric on  $\mathbb{R}^2$  with nonnegative Gaussian curvature. Let  $u : M \rightarrow \mathbb{R}$  be a harmonic function. If for all  $\delta > 0$*

$$\liminf_{r \rightarrow \infty} \frac{M(u; r)}{r^{1+\delta}} = 0,$$

*then there is a constant  $C > 0$  such that  $|u(r, \theta)| \leq Cr$  (i.e., it must be of linear growth).*

Let us comment on the significance of Theorem 4 in view of what is known. Ni and Tam in [4] showed that on a Kähler manifold if a harmonic function satisfies that

$$\limsup_{r \rightarrow \infty} \frac{u(r, \theta)}{r^{1+\delta}} = 0$$

for all  $\delta > 0$  then it must be of linear growth. This estimate is sharp in the following sense: It is known that there are complete noncompact surfaces of nonnegative Gaussian curvature that support harmonic functions which grow like  $r^{1+\delta}$  for  $0 < \delta < 1$ . Theorem 4 can be compared to Ni and Tam's theorem since in dimension 2 every orientable Riemannian manifold is Kähler.

**4.1. Proof of Theorem 4.** First we must recall, without a proof (one of which is via the Maximum Principle), a classical estimate due to Yau (see [10]) for the gradient of a harmonic function. To simplify the notation, in what follows, we will suppress the subindex  $g$  indicating the dependence on the metric of the gradient and Laplace operator.

**Theorem 5.** *Let  $u : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ , be a positive harmonic function,  $\mathbb{R}^2$  being endowed with a Riemannian metric of positive curvature. Then the estimate*

$$|\nabla \log u| \leq \frac{C}{r}$$

holds on any ball of radius  $r$  contained in  $\Omega$ .

Before engaging in the proof of Theorem 4, we shall show a weaker result. To this end, let us introduce Bochner's identity

$$\frac{1}{2}\Delta |\nabla u|^2 = |Hess u|^2 + g(\nabla \Delta u, \nabla u) + K_g g(\nabla u, \nabla u).$$

We recommend the reader non familiar with this formula to prove it in the case of  $\mathbb{R}^n$ , where the term involving the curvature does not appear, and  $g$  is the usual inner product. Recall that in this case  $Hess u$ , the Hessian of  $u$ , is the matrix of second derivatives of  $u$ . The general case in a Riemannian manifold follows from the noncommutativity of the covariant derivatives, which is measured by the curvature (for a proof of this formula see Lemma 1.36 and Exercise 1.37 in [2], and beware that  $Hess u = \nabla \nabla u$ ).

Now observe that if  $u$  is harmonic and  $K_g \geq 0$ , then from Bochner's identity follows that  $|\nabla u|^2$  is subharmonic. Let us assume that

$$\liminf_{r \rightarrow \infty} \frac{M(u; r)}{r\sqrt{\log r}} = 0.$$

Then, by Yau's estimate,

$$\liminf_{r \rightarrow \infty} \frac{M(|\nabla u|^2; r)}{\log r} = 0$$

so  $|\nabla u|^2$  is constant by Theorem 3 (take  $z(r) = r$ , and the hypothesis hold since  $K_g \geq 0$ ), and hence  $u$  must be of linear growth.

However we can do much better: It is time to give a proof of Theorem 4. Following the work of Ni and Tam, instead of using Bochner's identity, we will make use of the following identity, which is valid for any harmonic function  $u$  defined on a surface:

$$(1) \quad \Delta \log(1 + |\nabla u|^2) = \frac{2|Hess u|^2 + 2K_g |\nabla u|^2 (1 + |\nabla u|^2)}{(1 + |\nabla u|^2)^2}.$$

Not being as well known as Bochner's, we shall give a proof of this identity in the last paragraphs of this paper, so let us then continue with the proof of Theorem 4.

Again, identity (1) implies in the case of nonnegative curvature that  $\log(1 + |\nabla u|^2)$  is subharmonic. Now use the hypothesis: If for  $\delta > 0$

$$\liminf_{r \rightarrow \infty} \frac{M(u; r)}{r^{1+\delta}} = 0,$$

then by Yau's gradient estimate there is a sequence  $r_{\delta, k} \rightarrow \infty$  such that

$$|\nabla u| \leq C r_{\delta, k}^\delta$$

and hence,

$$\log(1 + |\nabla u|^2) \leq C \log r_{\delta, k}^\delta.$$

Since  $\delta > 0$  can be as small as we wish, as a consequence we can deduce that

$$\liminf_{r \rightarrow \infty} \frac{M(\log(1 + |\nabla u|^2), r)}{\log r} = 0$$

and from Theorem 3 we can conclude that  $\log(1 + |\nabla u|^2)$  is constant, and hence that  $|\nabla u|$  is constant, i.e.,  $u$  is of linear growth.

**4.2. Proof of formula (1).** As is customary, we pick a local orthonormal frame  $e_1, e_2$  around the point where we will be performing our computations. A subindex  $i$  will denote covariant differentiation with respect to (or in the direction of, as you prefer)  $e_i$ . We shall enforce Einstein's summation condition, i.e., we add over repeated subindices. For instance, we have:

$$\Delta u = u_{jj} = u_{11} + u_{22}, \quad \text{and} \quad |Hess u|^2 = u_{ij}u_{ij} = u_{11}^2 + u_{12}^2 + u_{21}^2 + u_{22}^2.$$

We are ready to begin:

$$\begin{aligned} \left( \log(1 + |\nabla u|^2) \right)_{jj} &= \left( \frac{2}{1 + |\nabla u|^2} u_{ij}u_i \right)_j \\ &= - \frac{4}{(1 + |\nabla u|^2)^2} u_{ij}u_{kj}u_iu_k \\ &\quad + \frac{2}{1 + |\nabla u|^2} u_{ij}u_{ij} + \frac{2}{1 + |\nabla u|^2} u_{ijj}u_i. \end{aligned}$$

The term  $u_{ij}u_{kj}u_iu_k$  when written in expanded form is

$$u_{11}^2u_1^2 + 2u_{12}u_{11}u_1u_2 + u_{12}^2u_2^2 + u_{22}^2u_2^2 + 2u_{21}u_{22}u_1u_2 + u_{21}^2u_1^2,$$

which by using that  $u_{11} = -u_{22}$  and  $u_{21} = u_{12}$ , reduces to

$$u_{11}^2u_1^2 + u_{12}^2u_2^2 + u_{22}^2u_2^2 + u_{21}^2u_1^2.$$

Now we need to be a little bit careful. Notice that

$$u_{11}u_i = -u_{22}u_i,$$

so we get that

$$u_i^2 |\nabla u|^2 = u_{11}^2u_1^2 + u_{22}^2u_2^2,$$

and hence

$$(u_{11}^2 + u_{22}^2) |\nabla u|^2 = 2(u_{11}^2u_1^2 + u_{22}^2u_2^2),$$

which leads to

$$(1 + |\nabla u|^2) u_{ij}u_{ij} = 2(u_{11}^2u_1^2 + u_{22}^2u_2^2) + 2u_{12}^2u_1^2 + 2u_{12}^2u_2^2 + u_{ij}u_{ij},$$

from which we obtain

$$- \frac{4}{(1 + |\nabla u|^2)^2} u_{ij}u_{kj}u_iu_k + \frac{2}{1 + |\nabla u|^2} u_{ij}u_{ij} = \frac{2u_{ij}u_{ij}}{(1 + |\nabla u|^2)^2}.$$

It is from  $u_{ijj}u_i$  that we obtain the curvature term. Indeed, the Ricci identity (Lemma 1.36 in [2]) tells us that

$$\Delta u_j = (\Delta u)_j + R_{ij}u_i,$$

where  $R_{ij}$  is the Ricci tensor of the metric. In the case of a surface,  $R_{ij} = K_g g_{ij}$ ,  $K_g$  being the Gaussian curvature, and hence we have that

$$\frac{2}{1 + |\nabla u|^2} u_{ijj}u_i = \frac{2}{1 + |\nabla u|^2} R_{ki}u_ku_i = \frac{2K_g |\nabla u|^2}{1 + |\nabla u|^2},$$

and formula (1) follows.



## REFERENCES

- [1] Cheeger, J.; Colding, T. H.; Minicozzi, W. P., II. Linear growth harmonic functions on complete manifolds with nonnegative Ricci curvature. *Geom. Funct. Anal.* 5 (1995), no. 6, 948–954.
- [2] Chow, Bennett; Lu, Peng; Ni, Lei Hamilton’s Ricci flow. *Graduate Studies in Mathematics*, 77. American Mathematical Society, Providence, RI; Science Press, New York, 2006.
- [3] Li, Peter; Tam, Luen-Fai. Linear growth harmonic functions on a complete manifold. *J. Differential Geom.* 29 (1989), no. 2, 421–425.
- [4] Ni, Lei; Tam, Luen-Fai. Plurisubharmonic functions and the structure of complete Kähler manifolds with nonnegative curvature. *J. Differential Geom.* 64 (2003), no. 3, 457–524.
- [5] Gilbarg, David; Trudinger, Neil S. *Elliptic partial differential equations of second order.* Reprint of the 1998 edition. *Classics in Mathematics.* Springer-Verlag, Berlin, 2001. xiv+517 pp.
- [6] Greene, R. E.; Wu, H. *Function theory on manifolds which possess a pole.* *Lecture Notes in Mathematics*, 699. Springer, Berlin, 1979. ii+215 pp.
- [7] Milnor, John. On deciding whether a surface is parabolic or hyperbolic. *Amer. Math. Monthly* 84 (1977), no. 1, 43–46.
- [8] Protter, Murray H.; Weinberger, Hans F. *Maximum principles in differential equations.* Prentice-Hall, Inc., Englewood Cliffs, N.J. 1967 x+261 pp.
- [9] Nelson, Edward. A proof of Liouville’s theorem. *Proc. Amer. Math. Soc.* 12 (1961), 995.
- [10] Yau, S.-T. Harmonic functions on complete Riemannian manifolds. *Comm. Pure Appl. Math.* 28 (1975), 201–228.