# INEQUALITIES FOR TRACE ON $\tau$ -MEASURABLE OPERATORS

MOHAMMAD SAL MOSLEHIAN<sup>1</sup> AND GHADIR SADEGHI<sup>2</sup>

ABSTRACT. Let  $\mathfrak{M}$  be a semifinite von Neumann algebra on a Hilbert space equipped with a faithful normal semifinite trace  $\tau$ . A closed densely defined operator x affiliated with  $\mathfrak{M}$ is called  $\tau$ -measurable if there exists a number  $\lambda \geq 0$  such that  $\tau \left( e^{|x|}(\lambda, \infty) \right) < \infty$ . A number of useful inequalities, which are known for the trace on Hilbert space operators, are extended to trace on  $\tau$ -measurable operators. In particular, these inequalities imply Clarkson inequalities for *n*-tuples of  $\tau$ -measurable operators. A general parallelogram law for  $\tau$ -measurable operators are given as well.

### 1. INTRODUCTION AND PRELIMINARIES

Let  $\mathfrak{M}$  be a semifinite von Neumann algebra on a Hilbert space  $\mathfrak{H}$ , with unit element 1, equipped with a faithful normal semifinite trace  $\tau$ . For standard facts concerning von Neumann algebras, we refer the reader to [2]. A closed densely defined linear operator  $x : \mathcal{D}(x) \subseteq \mathfrak{H} \to \mathfrak{H}$  is called affiliated with  $\mathfrak{M}$  if ux = xu for all unitaries u in the commutant  $\mathfrak{M}'$  of  $\mathfrak{M}$ . If x is in the algebra  $\mathcal{B}(\mathfrak{H})$  of all bounded linear operators on the Hilbert space  $\mathfrak{H}$ , then x is affiliated with  $\mathfrak{M}$  if and only if  $x \in \mathfrak{M}$ . A closed densely defined operator xaffiliated with  $\mathfrak{M}$  is called  $\tau$ -measurable if there exists a number  $\lambda \geq 0$  such that

$$\tau\left(e^{|x|}(\lambda,\infty)\right) < \infty$$

where  $|x| = (x^*x)^{1/2}$  and  $e^a$  denotes the spectral measure of the self-adjoint operator a which is a  $\sigma$ -additive (w.r.t. the strong operator topology) from the Borel  $\sigma$ -algebra of  $\mathbb{R}$  into the orthogonal projections. The collection of all  $\tau$ -measurable operators is denoted by  $\widetilde{\mathfrak{M}}$ . With the sum and product defined as the respective closure of the algebraic sum and product, it is well known that  $\widetilde{\mathfrak{M}}$  is a \*-algebra with respect to the operations of strong sum and strong product and taking adjoint; see [11]. For  $x \in \widetilde{\mathfrak{M}}$ , the generalized singular value function

<sup>2010</sup> Mathematics Subject Classification. Primary 46L51; Secondary 47A30.

Key words and phrases. Semifinite von Neumann algebra,  $\tau$ -measurable operator, trace, Clarkson inequality.

 $\mu(x): [0,\infty] \to [0,\infty]$  is defined by

$$\mu_t(x) = \inf \left\{ \lambda \ge 0 : \quad \tau \left( e^{|x|}(\lambda, \infty) \right) \le t \right\}, \quad t \ge 0.$$

The generalized singular value function  $\mu(x)$  is decreasing right-continuous. Moreover,

$$\mu(uxv) \le \|u\| \|v\| \mu(x)$$

for all  $u, v \in \mathfrak{M}$  and  $x \in \widetilde{\mathfrak{M}}$ . Moreover,

$$\mu(f(x)) = f(\mu(x))$$

whenever  $0 \leq x \in \widetilde{\mathfrak{M}}$  and f is an increasing continuous function on  $[0, \infty)$  satisfying f(0) = 0. The space  $\widetilde{\mathfrak{M}}$  is a partially ordered vector space under the ordering  $x \geq 0$  defined by  $\langle x\xi,\xi \rangle \geq 0, \xi \in \mathcal{D}(x)$ . The trace  $\tau$  on  $\mathfrak{M}^+$  extends uniquely to an additive, positively homogeneous, unitarily invariant and normal functional  $\widetilde{\tau} : \widetilde{\mathfrak{M}} \to [0, \infty]$ , which is given by  $\widetilde{\tau}(x) = \int_0^\infty \mu_t(x) dt, x \in \mathfrak{M}^+$  [4]. This extension is also denoted by  $\tau$ . Further,

$$\tau(f(x)) = \int_0^\infty f(\mu_t(x))dt \tag{1.1}$$

whenever  $0 \leq x \in \widetilde{\mathfrak{M}}$  and f is non-negative Borel function which is bounded on a neighborhood of 0 and satisfies f(0) = 0. For  $0 , <math>L^p(\mathfrak{M}, \tau)$  is defined as the set of all densely defined closed operators x affiliated with  $\mathfrak{M}$  such that

$$||x||_p = \tau (|x|^p)^{\frac{1}{p}} = \left(\int_0^\infty \mu_t(x)^p dt\right)^{\frac{1}{p}} < \infty.$$

For further details and proofs, we refer the reader to [15, 5, 3, 12, 13]. Important special cases of these noncommutative spaces are usual  $L^p$ -spaces and the Schatten *p*-classes  $C_p$ .

The classical Clarkson inequalities, for schatten *p*-norms of Hilbert space operators, assert that if  $A, B \in \mathcal{B}(\mathfrak{H})$ , then

$$2(\|A\|_{p}^{p} + \|B\|_{p}^{p}) \le \|A + B\|_{p}^{p} + \|A - B\|_{p}^{p} \le 2^{p-1}(\|A\|_{p}^{p} + \|B\|_{p}^{p})$$
(1.2)

for  $2 \leq p < \infty$ , and

$$2^{p-1}(\|A\|_p^p + \|B\|_p^p) \le \|A + B\|_p^p + \|A - B\|_p^p \le 2(\|A\|_p^p + \|B\|_p^p)$$
(1.3)

for 0 (see [6]). Bahatia and Kittaneh [1] generalized the Clarkson inequalities (1.2)and (1.3) for*n* $-tuples operators as follows. If <math>A_0, \ldots, A_{n-1} \in \mathcal{B}(\mathfrak{H})$  and  $\omega_0, \ldots, \omega_{n-1}$  are *n* roots of unity with  $\omega_j = e^{2\pi i j/n}$ ,  $0 \leq j \leq n-1$ , then

$$n\sum_{j=0}^{n-1} \|A_j\|_p^p \le \sum_{k=0}^{n-1} \left\| \sum_{j=0}^{n-1} \omega_j^k A_j \right\|_p^p \le n^{p-1} \sum_{j=0}^{n-1} \|A_j\|_p^p$$
(1.4)

for  $2 \leq p \leq \infty$ , and

$$n^{p-1} \sum_{j=0}^{n-1} \|A_j\|_p^p \le \sum_{k=0}^{n-1} \left\| \sum_{j=0}^{n-1} \omega_j^k A_j \right\|_p^p \le n \sum_{j=0}^{n-1} \|A_j\|_p^p$$
(1.5)

for 0 . Related Clarkson inequalities for*n*-tuples of operators have been recentlygiven by Kissin [8] and Hirzalleh and Kitaneh [7]. These inequalities have been found tobe very powerful tools in operator theory and in mathematical physics (see, e.g., [14]). In $[5], Fack and Kosaki proved Clarkson inequalities for measurable operators in Haagerup <math>L^p$ spaces. Let x be a  $\tau$ -measurable operator in  $L^p(\mathfrak{M}, \tau)$ . Since  $||x||_p^p = \tau(|x|^p)$ , for 0 ,our generalization of the inequalities (1.2) and (1.3) will be clear if we rewrite them as

$$2[\tau(|x|^p) + \tau(|y|^p)] \le \tau(|x+y|^p) + \tau(|x-y|^p) \le 2^{p-1}[\tau(|x|^p) + \tau(|y|^p)]$$
(1.6)

for  $2 \leq p < \infty$ , and

$$2^{p-1}[\tau(|x|^p) + \tau(|y|^p)] \le \tau(|x+y|^p) + \tau(|x-y|^p) \le 2[\tau(|x|^p) + \tau(|y|^p)]$$
(1.7)

for 0 .

There are several extensions of the classical parallelogram law in the literature. Generalizations of the parallelogram law for the Schatten p-norms have been given in the form of the celebrated Clarkson inequalities; see [7, 9, 10] and references cited therein.

A number of useful inequalities relating the traces of operators on a Hilbert space are known when the trace is defined in the usual way. In this paper, we present some inequalities for trace on  $\tau$ -measurable operators. These inequalities to trace generalized Clarkson inequalities for *n*-tuples of  $\tau$ -measurable operators. In fact, we give natural generalizations of inequalities (1.6) and (1.7) for *n*-tuples of  $\tau$ -measurable operators and show that the power functions  $\varphi(t) = t^p$  can be replaced by more general classes of functions. A general parallelogram law for  $\tau$ -measurable operators are given as well.

### 2. Trace inequalities for *n*-tuple of $\tau$ -measurable operators

In this section, we give new trace inequalities for *n*-tuple of  $\tau$ -measurable operators that considerably generalize inequalities (1.6) and (1.7). Before we give the main theorems of this section, we need the following lemma, which clearly is an extension of [5, proposition 4.6]) for *n*-tuples of  $\tau$ -measurable operators.

**Lemma 2.1.** Let  $x_0, \ldots, x_{n-1}$  be positive  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real numbers such that  $\sum_{j=0}^{n-1} \alpha_j = 1$ . Then, for every continuous increasing function f on  $\mathbb{R}_+$  with f(0) = 0,

$$\tau\left(f\left(\sum_{j=0}^{n-1}\alpha_j x_j\right)\right) \le \sum_{j=0}^{n-1}\alpha_j \tau(f(x_j)) \quad when \ f \ is \ convex.$$

$$(2.1)$$

$$\sum_{j=0}^{n-1} \alpha_j \tau(f(x_j)) \le \tau \left( f\left(\sum_{j=0}^{n-1} \alpha_j x_j\right) \right) \quad when \ f \ is \ concave.$$
(2.2)

$$\sum_{j=0}^{n-1} \tau\left(f\left(x_{j}\right)\right) \leq \tau\left(f\left(\sum_{j=0}^{n-1} x_{j}\right)\right) \quad when \ f \ is \ convex.$$

$$(2.3)$$

$$\tau\left(f\left(\sum_{j=0}^{n-1} x_j\right)\right) \le \sum_{j=0}^{n-1} \tau\left(f\left(x_j\right)\right) \quad when \ f \ is \ concave.$$

$$(2.4)$$

Now we are in a position to present our main result, providing a generalization of inequality (1.6) for *n*-tuples of  $\tau$ -measurable operators.

**Theorem 2.2.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real numbers such that  $\sum_{j=0}^{n-1} \alpha_j = 1$ . Let  $\varphi$  be a continuous increasing function on  $\mathbb{R}_+$  with  $\varphi(0) = 0$  such that  $\psi(t) = \varphi(\sqrt{t})$  is convex. Then

$$\tau\left(\varphi\left(\left|\sum_{j=0}^{n-1}\alpha_j x_j\right|\right)\right) + \sum_{0 \le j < k \le n-1} \tau\left(\varphi\left(\sqrt{\alpha_j \alpha_k} \left|x_j - x_k\right|\right)\right) \le \sum_{j=0}^{n-1} \alpha_j \tau(\varphi(\left|x_j\right|)).$$
(2.5)

*Proof.* The following useful identity can be easily verified,

$$\left|\sum_{j=0}^{n-1} \alpha_j x_j\right|^2 + \sum_{0 \le j < k \le n-1} \alpha_j \alpha_k |x_j - x_k|^2 = \sum_{j=0}^{n-1} \alpha_j |x_j|^2.$$
(2.6)

Now, by using identity (2.6) and inequalities (2.1), (2.3) we get

$$\sum_{j=0}^{n-1} \alpha_j \tau(\varphi(|x_j|)) = \sum_{j=0}^{n-1} \alpha_j \tau(\psi(|x_j|^2))$$

$$\geq \tau \left( \psi \left( \sum_{j=0}^{n-1} \alpha_j |x_j|^2 \right) \right)$$

$$= \tau \left( \psi \left( \left| \sum_{j=0}^{n-1} \alpha_j x_j \right|^2 + \sum_{0 \le j < k \le n-1} \alpha_j \alpha_k |x_j - x_k|^2 \right) \right)$$

$$\geq \tau \left( \psi \left( \left| \sum_{j=0}^{n-1} \alpha_j x_j \right|^2 \right) \right) + \sum_{0 \le j < k \le n-1} \tau \left( \psi \left( \alpha_j \alpha_k |x_j - x_k|^2 \right) \right)$$

$$= \tau \left( \varphi \left( \left| \sum_{j=0}^{n-1} \alpha_j x_j \right|^2 \right) \right) + \sum_{0 \le j < k \le n-1} \tau \left( \varphi \left( \sqrt{\alpha_j \alpha_k} |x_j - x_k|^2 \right) \right)$$

Based on Lemma 2.1 and inequalities (2.2) and (2.4), one can employ an argument similar to that used in the proof of Theorem 2.2 to prove the following related result.

**Theorem 2.3.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real numbers such that  $\sum_{j=0}^{n-1} \alpha_j = 1$ . Let  $\varphi$  be a continuous increasing function on  $\mathbb{R}_+$  with  $\varphi(0) = 0$  such that  $\psi(t) = \varphi(\sqrt{t})$  is concave. Then

$$\sum_{j=0}^{n-1} \alpha_j \tau(\varphi(|x_j|)) \le \tau \left( \varphi\left( \left| \sum_{j=0}^{n-1} \alpha_j x_j \right| \right) \right) + \sum_{0 \le j < k \le n-1} \tau \left( \varphi\left( \sqrt{\alpha_j \alpha_k} \left| x_j - x_k \right| \right) \right).$$
(2.7)

Applying Theorem 2.2 and Theorem 2.3 for the functions  $\varphi(t) = t^p$   $(2 \le p < \infty)$  and  $\varphi(t) = t^p$  (0 , respectively, we obtain natural generalization of Clarkson inequalities for*n* $-tuples of <math>\tau$ -measurable operators.

**Corollary 2.4.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real numbers such that  $\sum_{j=0}^{n-1} \alpha_j = 1$ . Then

$$\left\|\sum_{j=0}^{n-1} \alpha_j x_j\right\|_p^p + \sum_{0 \le j < k \le n-1} (\alpha_j \alpha_k)^{\frac{p}{2}} \|x_j - x_k\|_p^p \le \sum_{j=0}^{n-1} \alpha_j \|x_j\|_p^p$$

for  $2 \leq p < \infty$ , and

$$\sum_{j=0}^{n-1} \alpha_j \|x_j\|_p^p \le \left\|\sum_{j=0}^{n-1} \alpha_j x_j\right\|_p^p + \sum_{0 \le j < k \le n-1} (\alpha_j \alpha_k)^{\frac{p}{2}} \|x_j - x_k\|_p^p$$

for 0 . In particular

$$\left\|\sum_{j=0}^{n-1} x_j\right\|_p^p + \sum_{0 \le j < k \le n-1} \|x_j - x_k\|_p^p \le n^{p-1} \sum_{j=0}^{n-1} \|x_j\|_p^p$$

for  $2 \leq p < \infty$ , and

$$n^{p-1} \sum_{j=0}^{n-1} \|x_j\|_p^p \le \left\| \sum_{j=0}^{n-1} x_j \right\|_p^p + \sum_{0 \le j < k \le n-1} \|x_j - x_k\|_p^p$$

for 0 .

Let  $\varphi(t) = e^{t^2} - 1$ . Then  $\varphi$  is a continuous increasing function on  $\mathbb{R}_+$ ,  $\varphi(0) = 0$  and  $\psi(t) = \varphi(\sqrt{t}) = e^t - 1$  is convex. Applying Theorem 2.2 to this spacial function, we get the next result.

**Corollary 2.5.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real numbers such that  $\sum_{j=0}^{n-1} \alpha_j = 1$ . Then

$$\tau \left( e^{\left| \sum_{j=0}^{n-1} \alpha_j x_j \right|^2} - \mathbf{1} \right) + \sum_{0 \le j < k \le n-1} \tau \left( e^{\alpha_j \alpha_k |x_j - x_k|^2} - \mathbf{1} \right) \le \sum_{j=0}^{n-1} \alpha_j \tau \left( e^{|x_j|^2} - \mathbf{1} \right).$$

Now let  $\varphi(t) = \log(t+1)$ . Then  $\psi(t) = \varphi(\sqrt{t}) = \log(\sqrt{t}+1)$  is concave. Applying Theorem 2.3 to this function, we obtain the following corollary.

**Corollary 2.6.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real numbers such that  $\sum_{j=0}^{n-1} \alpha_j = 1$ . Then

$$\sum_{j=0}^{n-1} \alpha_j \tau \left( \log(|x_j|+1) \right) \le \tau \left( \log\left( \left| \sum_{j=0}^{n-1} \alpha_j x_j \right| + 1 \right) \right) + \sum_{0 \le j < k \le n-1} \tau \left( \log\left( \sqrt{\alpha_j \alpha_k} |x_j - x_k| + 1 \right) \right)$$

6

# 3. Refinement of trace inequalities

In this section, by using some trace inequalities for *n*-tuples of  $\tau$ -measurable operators, we obtain a refinement of the Clarkson inequalities, which was established by Bahatia and Kittaneh in [1]. To this end, we need the following useful identity

$$\frac{1}{n} \sum_{k=0}^{n-1} \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right|^2 = \sum_{j=0}^{n-1} |x_j|^2, \tag{3.1}$$

where  $\omega_0, \ldots, \omega_{n-1}$  are the *n* roots of unity with  $\omega_j = e^{2\pi i j/n}, 0 \le j \le n-1$  (see [1]).

**Theorem 3.1.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators and let  $\varphi$  be a continuous increasing function on  $\mathbb{R}_+$  with  $\varphi(0) = 0$  such that  $\psi(t) = \varphi(\sqrt{t})$  is convex. Then

$$\sum_{k=0}^{n-1} \tau \left( \varphi \left( \frac{1}{\sqrt{n}} \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right| \right) \right) \le \tau \left( \varphi \left( \left( \sum_{j=0}^{n-1} |x_j|^2 \right)^{\frac{1}{2}} \right) \right) \right)$$

$$\le \frac{1}{n} \sum_{k=0}^{n-1} \tau \left( \varphi \left( \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right| \right) \right).$$
(3.2)

*Proof.* Using inequalities (2.1), (2.3) and identity (3.1), we reach the first inequality in (3.2).

$$\sum_{k=0}^{n-1} \tau \left( \varphi \left( \frac{1}{\sqrt{n}} \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right| \right) \right) = \sum_{k=0}^{n-1} \tau \left( \psi \left( \frac{1}{n} \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right|^2 \right) \right) \right)$$
$$\leq \tau \left( \psi \left( \frac{1}{n} \sum_{k=0}^{n-1} \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right|^2 \right) \right)$$
$$= \tau \left( \psi \left( \sum_{j=0}^{n-1} |x_j|^2 \right) \right)$$
$$= \tau \left( \varphi \left( \left( \sum_{j=0}^{n-1} |x_j|^2 \right)^{\frac{1}{2}} \right) \right),$$

as desired.

Now using Lemma 2.1, inequality (2.1) and identity (3.1) we get the second inequality in (3.2).

$$\tau \left( \varphi \left( \left( \sum_{j=0}^{n-1} |x_j|^2 \right)^{\frac{1}{2}} \right) \right) = \tau \left( \varphi \left( \left( \frac{1}{n} \sum_{k=0}^{n-1} \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right|^2 \right)^{\frac{1}{2}} \right) \right)$$
$$= \tau \left( \psi \left( \frac{1}{n} \sum_{k=0}^{n-1} \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right|^2 \right) \right)$$
$$\leq \frac{1}{n} \sum_{k=0}^{n-1} \tau \left( \psi \left( \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right|^2 \right) \right)$$
$$= \frac{1}{n} \sum_{k=0}^{n-1} \tau \left( \varphi \left( \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right|^2 \right) \right).$$

Based on inequalities (2.2), (2.4) and identity (3.1), one can employ an argument similar to that used in the proof of Theorem 3.1 to get the following related result.

**Theorem 3.2.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators and let  $\varphi$  be a continuous increasing function on  $\mathbb{R}_+$  with  $\varphi(0) = 0$  such that  $\psi(t) = \varphi(\sqrt{t})$  is concave. Then

$$\frac{1}{n}\sum_{k=0}^{n-1}\tau\left(\varphi\left(\left|\sum_{j=0}^{n-1}\omega_j^k x_j\right|\right)\right) \le \tau\left(\varphi\left(\left(\sum_{j=0}^{n-1}|x_j|^2\right)^{\frac{1}{2}}\right)\right) \le \sum_{k=0}^{n-1}\tau\left(\varphi\left(\frac{1}{\sqrt{n}}\left|\sum_{j=0}^{n-1}\omega_j^k x_j\right|\right)\right).$$

Applying Theorem 3.1 and Theorem 3.2 for the functions  $\varphi(t) = t^p$   $(2 \le p < \infty)$  and  $\varphi(t) = t^p$  (0 , respectively, we obtain refinements of the Clarkson inequalities for*n* $-tuples of <math>\tau$ -measurable operators.

**Corollary 3.3.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators. Then

$$n^{-\frac{p}{2}} \sum_{k=0}^{n-1} \left\| \sum_{j=0}^{n-1} \omega_j^k x_j \right\|_p^p \le \left\| \left( \sum_{j=0}^{n-1} |x_j|^2 \right)^{\frac{1}{2}} \right\|_p^p \le \frac{1}{n} \sum_{k=0}^{n-1} \left\| \sum_{j=0}^{n-1} \omega_j^k x_j \right\|_p^p$$

for  $2 \leq p < \infty$ , and

$$\frac{1}{n}\sum_{k=0}^{n-1}\left\|\sum_{j=0}^{n-1}\omega_j^k x_j\right\|_p^p \le \left\|\left(\sum_{j=0}^{n-1}|x_j|^2\right)^{\frac{1}{2}}\right\|_p^p \le n^{-\frac{p}{2}}\sum_{k=0}^{n-1}\left\|\sum_{j=0}^{n-1}\omega_j^k x_j\right\|_p^p$$

for 0 .

Applying Theorem 3.1 and Theorem 3.2 to special functions  $\varphi(t) = e^{t^2} - 1$  and  $\varphi(t) = \log(t+1)$  respectively, we obtain the following results.

**Corollary 3.4.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators. Then

$$\sum_{k=0}^{n-1} \tau \left( e^{\frac{1}{n} \left| \sum_{j=0}^{n-1} \omega_j^k x_j \right|^2} - \mathbf{1} \right) \le \tau \left( e^{\sum_{j=0}^{n-1} |x_j|^2} - \mathbf{1} \right) \le \frac{1}{n} \sum_{k=0}^{n-1} \tau \left( e^{\left| \sum_{j=0}^{n-1} \omega_j^k x_j \right|^2} - \mathbf{1} \right).$$

**Corollary 3.5.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators. Then

$$\frac{1}{n}\sum_{k=0}^{n-1}\tau\left(\log\left(\left|\sum_{j=0}^{n-1}\omega_{j}^{k}x_{j}\right|+1\right)\right) \leq \tau\left(\log\left(\left(\sum_{j=0}^{n-1}|x_{j}|\right)^{\frac{1}{2}}+1\right)\right)$$
$$\leq \frac{1}{n}\sum_{k=0}^{n-1}\tau\left(\log\left(\frac{1}{n}\left|\sum_{j=0}^{n-1}\omega_{j}^{k}x_{j}\right|+1\right)\right).$$

#### 4. A general parallelogram law for $\tau$ -measurable operators

The parallelogram law for *n*-tuples of operators has been recently investigated by the second author [9]. In this section, by using some trace inequalities for *n*-tuples of  $\tau$ -measurable operators, we obtain an extension of the main result of [9] for *n*-tuples of  $\tau$ -measurable operators. Let  $x_0, \ldots, x_{n-1}, y_1, \ldots, y_n$  be  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real numbers. Then the following identity can be given similar to [9, Corollary 2.3],

$$\sum_{0 \le i < j \le n-1} \left| \sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} x_j \right|^2 + \sum_{0 \le i < j \le n-1} \left| \sqrt{\frac{\alpha_i}{\alpha_j}} y_i - \sqrt{\frac{\alpha_j}{\alpha_i}} y_j \right|^2$$

$$= \sum_{i,j=0}^{n-1} \left| \sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} y_j \right|^2 - \left| \sum_{i=0}^{n-1} (x_i - y_i) \right|^2.$$
(4.1)

If we set  $y_1 = \ldots = y_n = 0$  in (4.1) and  $\sum_{j=0}^{n-1} \frac{1}{\alpha_j} = 1$ , then the following identity can be given

$$\sum_{0 \le i < j \le n-1} \left| \sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} x_j \right|^2 = \sum_{i=0}^{n-1} \alpha_i |x_j|^2 - \left| \sum_{i=0}^{n-1} x_i \right|^2.$$
(4.2)

**Theorem 4.1.** Let  $x_0, \ldots, x_{n-1}, y_1, \ldots, y_n$  be  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real number such that  $\sum_{i,j=0}^{n-1} \frac{1}{\sqrt{\alpha_i \alpha_j}} = 1$ . Let  $\varphi$  be a continuous increasing function

on  $\mathbb{R}_+$  with  $\varphi(0) = 0$  such that  $\psi(t) = \varphi(\sqrt{t})$  is convex. Then

$$\sum_{\substack{i,j=0\\ j\leq n-1}}^{n-1} \frac{1}{\sqrt{\alpha_i \alpha_j}} \tau(\varphi(|\alpha_i x_i - \alpha_j y_j|)) \geq$$

$$\sum_{\substack{0\leq i< j\leq n-1}} \tau\left(\varphi\left(\left|\sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} x_j\right|\right)\right) + \sum_{\substack{0\leq i< j\leq n-1}} \tau\left(\varphi\left(\left|\sqrt{\frac{\alpha_i}{\alpha_j}} y_i - \sqrt{\frac{\alpha_j}{\alpha_i}} y_j\right|\right)\right) + \tau\left(\varphi\left(\left|\sum_{i=0}^{n-1} (x_i - y_i)\right|\right)\right).$$

$$(4.3)$$

If  $\psi(t)$  is concave, then the reverse of inequality (4.3) holds.

Proof.

$$\begin{split} \sum_{i,j=0}^{n-1} \frac{1}{\sqrt{\alpha_i \alpha_j}} \tau(\varphi(|\alpha_i x_i - \alpha_j y_j|)) \\ &= \sum_{i,j=0}^{n-1} \frac{1}{\sqrt{\alpha_i \alpha_j}} \tau\left(\psi\left(|\alpha_i x_i - \alpha_j y_j|^2\right)\right) \\ &\geq \tau\left(\psi\left(\sum_{i,j=0}^{n-1} \frac{1}{\sqrt{\alpha_i \alpha_j}} |\alpha_i x_i - \alpha_j y_j|^2\right)\right) \qquad (by (2.1)) \\ &= \tau\left(\psi\left(\sum_{0 \le i < j \le n-1} \left|\sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} x_j\right|^2 + \sum_{0 \le i < j \le n-1} \left|\sqrt{\frac{\alpha_i}{\alpha_j}} y_i - \sqrt{\frac{\alpha_j}{\alpha_i}} y_j\right|^2\right) + \left|\sum_{i=0}^n (x_i - y_i)\right|^2\right) \\ &\qquad (by (4.1)) \\ &\geq \sum_{i,j=0}^{n-1} \tau\left(\psi\left(\left|\sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} x_j\right|^2\right)\right) + \sum_{i,j=0}^{n-1} \tau\left(\psi\left(\left|\sqrt{\frac{\alpha_i}{\alpha_j}} y_i - \sqrt{\frac{\alpha_j}{\alpha_i}} y_j\right|^2\right)\right) \\ &+ \tau\left(\psi\left(\left|\sum_{i=0}^n (x_i - y_i)\right|^2\right)\right) \qquad (by (2.3)) \\ &= \sum_{i,j=0}^{n-1} \tau\left(\varphi\left(\left|\sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} x_j\right|\right)\right) + \sum_{i,j=0}^{n-1} \tau\left(\varphi\left(\left|\sqrt{\frac{\alpha_i}{\alpha_j}} y_i - \sqrt{\frac{\alpha_j}{\alpha_i}} y_j\right|\right)\right) \\ &+ \tau\left(\varphi\left(\left|\sum_{i=0}^n (x_i - y_i)\right|\right)\right) \end{matrix}$$

10

**Corollary 4.2.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real numbers such that  $\sum_{j=0}^{n-1} \frac{1}{\alpha_j} = 1$ . Let  $\varphi$  be a continuous increasing function on  $\mathbb{R}_+$  with  $\varphi(0) = 0$  such that  $\psi(t) = \varphi(\sqrt{t})$  is convex. Then

$$\sum_{j=0}^{n-1} \frac{1}{\alpha_j} \tau(\varphi(|\alpha_i x_i|)) \ge \sum_{0 \le i < j \le n-1} \tau\left(\varphi\left(\left|\sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} x_j\right|\right)\right) + \tau\left(\varphi\left(\left|\sum_{i=0}^{n-1} x_i\right|\right)\right). \quad (4.4)$$

If  $\psi(t)$  is concave, then the reverse of inequality 4.4 holds.

*Proof.* It is sufficient that put  $y_1 = \ldots = y_n = 0$  in Theorem 4.1.

Applying Corollary 4.2 for the function  $\varphi(t) = t^p (2 \le p < \infty)$  and  $\varphi(t) = t^p (0 respectively, we get the following corollary.$ 

**Corollary 4.3.** Let  $x_0, \ldots, x_{n-1}$  be  $\tau$ -measurable operators and  $\alpha_0, \ldots, \alpha_{n-1}$  be positive real numbers such that  $\sum_{j=0}^{n-1} \frac{1}{\alpha_j} = 1$ . Then

$$\sum_{i=0}^{n-1} \alpha_i^{p-1} \|x_i\|^p \ge \sum_{0 \le i < j \le n-1} \left\| \sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} x_j \right\|_p^p + \left\| \sum_{i=0}^{n-1} x_i \right\|_p^p$$

for  $2 \leq p < \infty$ , and

$$\sum_{0 \le i < j \le n-1} \left\| \sqrt{\frac{\alpha_i}{\alpha_j}} x_i - \sqrt{\frac{\alpha_j}{\alpha_i}} x_j \right\|_p^p + \left\| \sum_{i=0}^{n-1} x_i \right\|_p^p \ge \sum_{i=0}^{n-1} \alpha_i^{p-1} \|x_i\|^p$$

for 0 .

Acknowledgement. The first author would like to thank Professor Qing-Wen Wang for his great hospitality during his visit of Shanghai University in 2013.

#### References

- R. Bahatia and F. Kittaneh, *Clarkson inequalities with several operators*, Bull. London Math. Soc. 36: 820–832, 2004.
- [2] J. Dixmier, Von Neumann Algebras, North-Holland Mathematical Library, Vol. 27, North-Holland, Amesterdam, 1981.
- [3] P. G. Dodds, T. K. Dodds and B. de Pagter, Noncommutative Banach function spaces, Math. Z. 201: 47–57, 1989.
- [4] P. G. Dodds, T.K. Dodds and B. de Pagter, *Noncommutative Köthe duality*, Trans. Amer. Math. Soc. 339: 717–750,1993.

- [5] Th. Fack and H. Kosaki, Generalized s-numbers of  $\tau$ -measurable operators, Pacific J. Math. 123: 269–300, 1986.
- [6] O. Hirzallah and F. Kittaneh, Noncommutative Clarkson inequalities for unitarily invariant norms, Pacific J. Math. 202: 363–369, 2002.
- [7] O. Hirzallah and F. Kittaneh, Noncommutative Clarkson inequalities for n-tuples of operators, Integral equations Operator theory 60: 369–379, 2008.
- [8] E. Kissin, On Clarkson-McCarthy inequalities for n-tuples of operators, Proc. Amer. Math. Soc. 135: 2483–2495, 2007.
- M. S. Moslehian, An operator extension of the parallelogram law and related norm inequalities, Math. Inequal. Appl. 14(2): 717–725, 2011.
- [10] M. S. Moslehian, M. Tominaga and K. S. Saito, Schatten p-norm inequalities related to an extended operator parallelogram law, Linear Algebra Appl. 435(4): 823–829, 2011.
- [11] E. Nelson, Note on noncommutative integration, J. Funct. Anal. 15: 103–116, 1974.
- [12] B. de Pagter, Noncommutative Banach function spaces, Positivity, 197-227, Trends Math., Birkhüser, Basel, 2007.
- [13] Gh. Sadeghi, Non-commutative Orlicz spaces associated to a modular on τ-measurable operators, J. Math. Anal. Appl. 395(2): 705-715, 2012.
- [14] B. Simon, Trace Ideals and Their Applications, Cambridge University press, Cambridge, 1979.
- [15] F. J. Yeadon, Noncommutative L<sup>p</sup>-spaces, Proc. Camb. Phil. Soc. 77: 91–102, 1975

<sup>1</sup> Department of Pure Mathematics, Center of Excellence in Analysis on Algebraic Structures (CEAAS), Ferdowsi University of Mashhad, P. O. Box 1159, Mashhad 91775, Iran

 $E\text{-}mail\ address: \texttt{moslehianQum.ac.ir, moslehianQmember.ams.org}$ 

<sup>2</sup> DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES, HAKIM SABZEVARI UNIVERSITY, P.O. Box 397, SABZEVAR, IRAN.

*E-mail address*: ghadir540gmail.com, g.sadeghi@hsu.ac.ir