Evidence of the Big Fix

Yuta Hamada, Hikaru Kawai, and Kiyoharu Kawana

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

October 2, 2018

Abstract

We give an evidence of the Big Fix. The theory of wormholes and multiverse suggests that the parameters of the Standard Model are fixed in such a way that the total entropy at the late stage of the universe is maximized, which we call the maximum entropy principle. In this paper, we discuss how it can be confirmed by the experimental data, and we show that it is indeed true for the Higgs vacuum expectation value v_h . We assume that the baryon number is produced by the sphaleron process, and that the current quark masses, the gauge couplings and the Higgs self coupling are fixed when we vary v_h . It turns out that the existence of the atomic nuclei plays a crucial role to maximize the entropy. This is reminiscent of the anthropic principle, however it is required by the fundamental low in our case.

Keywards: Quantum gravity, maximum entropy principle, sphaleron, baryon decay, multiverse, wormhole.

PACS numbers: 11-10.-z, 04.60.-m,

^{*}E-mail: hamada@gauge.scphys.kyoto-u.ac.jp

[†]E-mail: hkawai@gauge.scphys.kyoto-u.ac.jp

[‡]E-mail: kiyokawa@gauge.scphys.kyoto-u.ac.jp

1 Introduction

There are many open questions that the Standard Model can not answer. The Cosmological Constant Problem is the biggest one. The naive expectation value of the vacuum energy is given by

$$\rho \simeq M_{pl}^4 \simeq 10^{72} \text{GeV}^4,\tag{1}$$

if we assume that the cutoff scale exists around the Planck scale. However, from cosmological observations, we know that the vacuum energy density of our universe is

$$\rho_{\Lambda} \simeq 10^{-48} \text{GeV}^4. \tag{2}$$

The above mismatch between Eq.(1) and Eq.(2) is the Cosmological Constant Problem. There are other important problems of fine tuning in the Standard Model(SM): the Higgs mass, the Yukawa couplings, the QCD θ parameter \cdots and so on. The ordinary approach to these problems is to expect the emergence of a new physics at some high energy scale. For example, the supersymmetry is one of the candidates to answer the Higgs mass hierarchy. However, the recent observations of the Higgs mass [16, 17], and the related analyses [18] show the possibility that that there is no new physics between the weak and the Planck scale¹. If it is the case, we must answer the above problems by the Planck scale physics in the framework of the quantum gravity or string theory².

One possibility along this direction is to consider the dynamics of wormholes and multiverse [1]. Recently, there has been a development [2, 3, 4] in which the wave function of the multiverse is constructed based on the Lorentzian path integral. As a result, assuming that we live in the S^3 universe, it is found that the parameters of the SM are fixed in such a way that the total energy of the universe at the late stage E_{tot} is maximized, see Appendix B for the details. It is called The Big Fix. If the universe at the late stage is radiation(matter) dominated, it means that the total radiation(matter) is maximized, which we call the maximum entropy(matter) principle³. In order to test this principle, it is sufficient to examine how E_{tot} varies when we change each of the SM parameters from the experimental values. If we find that E_{tot} decreases under such change, the principle is verified. It seems that this program can be accomplished only if we know the whole history of the universe. However, for some of the SM parameters, we know how they participate the history of the universe.

In particular, we can check the maximum entropy principle for the Higgs vacuum expectation value v_h . In this paper, we show that this principle is actually satisfied when v_h is around $v_h = 246 \text{GeV}$. Here we assume that the dark matter(DM) decays much earlier than baryons so that the universe at the late stage becomes dominated by the radiation S_{rad} produced by the decay of baryons.

¹The observed Higgs mass may indicate that Higgs is the inflaton [26].

²The other possibilities are to introduce new physical principles such as the multiple point principle [19, 20, 21], the classical conformality [22], hidden duality [23] and the asymptotic safety [24]. There is also an interesting argument [25] which might be related to the maximum entropy principle.

³We regard the total radiation as the total entropy.

Note that S_{rad} is defined by

$$S_{rad} = a^4 \times \rho_{rad}(a) \tag{3}$$

where $\rho_{rad}(a)$ is the energy density of the radiation and a is the radius of the universe.

The overview of how S_{rad} is produced in the universe is as follows:

- Stage 1: The baryon number N_B is produced by the sphaleron process if we assume the standard leptogenesis scenario.
- Stage 2: The ratio of neutrons to all nucleons X_n is fixed by the following two processes:
 - A) The weak interaction is frozen out.
 - B) Some of the neutrons are converted to protons through the beta decay before the Big Bang Nucleosynthesis (BBN).
- Stage 3: S_{rad} is created by the decay of baryons. There are two contributions; hydrogens (free protons) and atomic nuclei.

The quantities that appear in the above stages can be expressed as a function of v_h , y_u and y_d . In fact, at the stage 1, using the recent numerical analysis of the sphaleron process[27], we can determine N_B as a function of v_h if we fix the gauge couplings, the Higgs self coupling, and the top quark mass. At the stage 2, X_n is determined by the Fermi constant, the proton neutron mass difference and the electron mass. Therefore, it can be expressed as an explicit function of v_h and the difference of the Yukawa couplings $y_- = y_d - y_u$. At the stage 3, for the case of the proton decay, S_{rad} can be expressed in terms of the mass and the life time of protons; m_p , τ_p . On the other hands, for the case of the decay of the helium nuclei, the situation is rather complicated. We must consider the effect that the pion produced by the decay of a nucleon loses its energy through a collision with the other nucleons. Then, in addition to the mass and life time of helium nuclei, we need a quantity that indicates the average radiation energy created by the decay of a helium nucleus, which we express as $(1-2\epsilon)m_p$. In principle, these five quantities

$$m_p$$
 , τ_p , m_{He} , τ_{He} , ϵ (4)

can be calculated by the QCD once the Λ_{QCD} and the current quark masses

$$m_{u,d} = y_{u,d} \cdot \frac{v_h}{\sqrt{2}} \tag{5}$$

are given⁴. However, since it is technically difficult, we try several phenomenological values for them. Using the above results, we can show that S_{rad} in deed takes the maximum around the experimental values of v_h .

⁴One need not worry about the GUT scale dependence because it appears as an overall factor.

This paper is organized as follows. In Section2, we discuss the history of the baryons, and give general differential equations that explain the time evolution of the radiation. In Section3, the qualitative and numerical results are shown. In Section5, we give summary and discussion. In Appendix A, we present the WKB solution of the wave function of the universe. In Appendix B, the probabilistic interpretation of the wave function of the universe is discussed. In Appendix C, we study the sphaleron process. In Appendix D, we estimate τ_{He} and ϵ . In Appendix E, we consider the scenario in which the universe at the late stage becomes dominated by the DM.

2 History of Baryons

We first mention the possible scenarios of the universe at the late stage. Since we do not know the fate of the DM, there are typically three possibilities⁵:

- (I): The DM decays earlier than the baryons, and the universe becomes dominated by the radiation after the baryon decay.
- (II): The baryons decay earlier than the DM, and the universe becomes dominated by the radiation after the dark matter decay.
- (III): The baryons decay, but the DM does not, then the universe is dominated by DM.

In this paper, we consider the scenario (I), and study how the baryons of the universe are produced at the early universe, and decay at the late stage of the universe. We give our argument along the stages indicated in the introduction. The consideration about the scenario (III) is presented in Appendix E.

Stage 1: Baryogenesis by the Sphaleron Process

In this subsection, we study how the baryon number is produced and how it depends on the SM parameters. We assume that it is produced by the sphaleron process. If the decoupling temperature of that process is comparable with the heavy quark masses, we must consider the decrease of the baryon number caused by them. The detailed calculations are presented in Appendix C.

Using the recent numerical results [27], we can obtain the decoupling temperature of the sphaleron process

$$T_{sph} \simeq \frac{7}{8} T_c = 140 \text{GeV} \times \frac{v_h}{246 \text{GeV}}$$
 (6)

⁵We assume that the Cosmological Constant at the late stage of the universe is fixed to the critical value Λ_{cri} so that it balances with the curvature. This is one of the predictions of the maximum entropy principle. See Appendix B for the details.




Figure 1: The baryon number as a function of v_h when the current quark masses, the gauge couplings and the Higgs self coupling are fixed. We can see that $\frac{N_B}{N_B-N_L}$ decreases around $v_h = 150 \text{GeV}$ corresponding to the decrease of v_h .

where T_c is the critical temperature of the phase transition. Here we fix the gauge couplings and the Higgs self coupling to the observed values, and regard v_h as a variable. Furthermore, we can find the temperature dependence of the Higgs expectation value from [27],

$$v_h(T) = v_h \sqrt{\frac{T_c - T}{T_c}}. (7)$$

This is consistent with the behavior of the Landau theory. Since the particle masses are proportional to $v_h(t)$, the suppression factors caused by the quarks and W boson at T_{sph} are given by

$$\frac{m_i}{T_{sph}} = \frac{m_{0i}\sqrt{\frac{T_c - T_{sph}}{T_c}}}{T_{sph}} = \frac{12}{7\sqrt{8}} \times \frac{m_{0i}}{v_h} = \frac{3}{7} \times \frac{y_i v_h}{v_h},\tag{8}$$

$$\frac{m_W}{T_{sph}} = \frac{m_{0W}\sqrt{\frac{T_c - T_{sph}}{T_c}}}{T_{sph}} = \frac{12}{7\sqrt{8}} \times \frac{m_{0W}}{v_h} = \frac{6}{7\sqrt{8}} \times \frac{g_2 v_h}{v_h}$$
(9)

where m_{0i} and m_{0W} are the physical masses and g_2 is the gauge coupling. When v_h becomes comparable with m_i , Eq.(8) starts to have an effect, and the baryon number decreases. Since T_{sph} is of order 100GeV, we need not to consider the light quarks. Namely, only the top quark and W boson are crucial to find the maximum of the entropy. In this approximation, we can obtain the baryon number of our universe (see Eq.(125) in Appendix C);

$$N_B = \frac{N_B}{N_{B-L}} N_{B-L} = \frac{2 (4 + g_b(m_W)) \cdot (5 + g_f(m_t))}{24 (5 + g_f(m_t)) + g_b(m_W) (37 + 2g_f(m_t))} N_{B-L}$$
(10)

where $g_f(m_t)$ and $g_b(m_W)$ are given by

$$g_f(m_t) := \frac{3}{2\pi^2} \int_0^\infty dx \frac{x^2}{\cosh^2\left(\frac{\sqrt{x^2 + (m_t/T)^2}}{2}\right)},$$

$$g_b(m_W) := \frac{3}{2\pi^2} \int_0^\infty dx \frac{x^2}{\sinh^2\left(\frac{\sqrt{x^2 + (m_W/T)^2}}{2}\right)}.$$
(11)

If we fix the Yukawa couplings and the gauge couplings, there is no v_h dependence. Therefore, in this case, the baryon number does not depend on v_h . On the other hands, if we fix the current quark masses, Eq.(8) depends on v_h . As we will discuss in the following sections, it is appropriate to fix the current quark masses, the gauge couplings and the Higgs self coupling when we vary v_h . We plot Eq.(10) as a function of v_h in Fig.1. Especially, since the coupling constants are fixed, the suppression factor of the W boson does not depend on v_h . One can see that N_B starts to decrease around $v_h \simeq 150 \text{GeV}$ when we make v_h small. This is, of course, the effect of the top quark mass. In the history of the universe, one might think that the top quark hardly plays an important role. However, the above result suggests that it has an effect in determining the baryon number. We will use Eq.(10) to calculate the radiation.

Stage 2: Neutron to Baryon Ratio X_n

In this section, we discuss how X_n is determined at the early universe. Because the total number of helium nuclei is given by

$$N_{\rm He} = N_B \frac{X_n}{2},\tag{12}$$

the determination of X_n corresponds to that of N_{He} . We denote the photon temperature by T. For T>>1MeV, neutrons and protons are in thermal equilibrium through the six processes:

$$n + \nu \leftrightarrow p + e^- \quad n + e^+ \leftrightarrow p + \bar{\nu} \quad n \leftrightarrow p + e^- + \bar{\nu},$$
 (13)

and X_n is given by

$$X_n = \frac{1}{1 + \exp(\frac{Q}{T})}. (14)$$

where $Q := m_n - m_p$ is the mass difference between a proton and a neutron. However, at a temperature between 1MeV and 0.1MeV, the reaction rates of the above processes except for the beta decay become smaller than the expansion rate of the universe. This decoupling temperature T_{dec} is determined by

$$\Gamma(p \to n) = H := \frac{\dot{a}}{a} \tag{15}$$




Figure 2: The value of Xn as a function of v_h when the current quark masses are fixed. We can see that X_n depends strongly on v_h . For example, if $v_h \simeq 100 \text{GeV}$, X_n is almost zero, and there are no atomic nuclei.

where $\Gamma(p \to n)$ is the proton to neutron reaction rate and H is the Hubble expansion rate. For T >> 1MeV, $\Gamma(p \to n)$ is given by [28]

$$\Gamma(p \to n) = 0.400 \text{sec}^{-1} \times \left(\frac{T}{1 \text{MeV}}\right)^5 \times \left(\frac{246 \text{GeV}}{v_h}\right)^4, \tag{16}$$

and the expansion rate at the time t is

$$H = \frac{1}{2t} = \frac{1}{2} \sqrt{\frac{2\pi^2 \mathcal{N}}{45M_{pl}^2}} T^2, \tag{17}$$

where \mathcal{N} represents the degree of freedoms. By solving Eq.(15), we can determine T_{dec} as a function of v_h ⁶.

Below T_{dec} , X_n decreases through the beta decay process, and we obtain

$$X_n(t) = \exp(-\tau_n^{-1}(t - t_{dec})) \times X_n(t_{dec}) = \exp(-\tau_n^{-1}(t - t_{dec})) \times \frac{1}{1 + \exp(\frac{Q}{T_{dec}})}$$
(18)

where t_{dec} is the decoupling time, and τ_n is the neutron life time⁷

$$\tau_n^{-1} = 885^{-1} \text{sec}^{-1} \times \left(\frac{246 \text{GeV}}{v_h}\right)^4.$$
 (19)

⁶Note that, since T_{dec} is an increasing function of v_h , if T_{dec} becomes comparable with the electron mass, we must consider the effect of the electron-positron annihilation.

⁷The overall coefficient depends on the current quark masses and the electron mass, which are not important because we fix them in this paper.

When T reaches the BBN temperature $T_{BBN}^{8} (\simeq 0.1 \text{MeV})$, neutrons are convert to helium nuclei rapidly. Thus, X_n is fixed to the value given by

$$X_n = \exp(-\tau_n^{-1}(t_{\text{BBN}} - t_{dec})) \times \frac{1}{1 + \exp(\frac{Q}{T_{dec}})},$$
 (20)

where t_{BBN} is the BBN time. Using Eq.(17), we can convert t_{dec} and t_{BBN} to T_{dec} and T_{BBN} . Fig.2 shows X_n as a function of v_h when we fix the current quark masses. One can see that X_n depends strongly on v_h .

Stage 3: Baryon Decay and Production of Radiation

As is discussed in the introduction, the radiation due to the baryon decay comes from not only free protons but also the nucleons in helium nuclei. Before considering the realistic case, we first study the simplified case [3] where the baryons are all protons. In this case, S_{rad} can be estimated as follows. If we simplify the situation so that protons decay simultaneously, from the energy conservation, we have

$$\frac{N_B m_p}{a(\tau_p)^3} = \frac{S_{rad}}{a(\tau_p)^4} \tag{21}$$

where τ_p is the proton life time. From Eq.(21), we have

$$S_{rad} = N_B m_p a(\tau_p). (22)$$

Then we use the Friedman equation to express $a(\tau_p)$ in terms of τ_p ;

$$\frac{1}{\tau_p^2} \simeq \frac{1}{M_{nl}^2} \frac{N_B m_p}{a(\tau_p)^3},\tag{23}$$

and we obtain

$$S_{rad} = constant \times (\frac{1}{M_{pl}^2})^{\frac{1}{3}} \times (N_B m_p)^{\frac{4}{3}} \tau_p^{\frac{2}{3}}.$$
 (24)

This is the qualitative expression of the radiation without atomic nuclei. In [3], it is discussed that the values of the current quark masses m_u, m_d may be determined by requiring that Eq.(24) should be maximized.

Let's start to consider the effect of atomic nuclei. We denote the number of the protons at the time t by $N_p(t)$, and that of the helium nuclei by $N_{He}(t)$. They decrease as follows ⁹:

$$\frac{dN_p(t)}{dt} = -\tau_p^{-1} \cdot N_p(t) + 3\tau_{He}^{-1} \cdot N_{He}(t)$$
 (25)

 $^{^8}T_{BBN}$ depends on the current quark masses and v_h . However, since its dependences are rather weak, we regard it as a constant in the following discussion.

⁹The neutrons produced by the decay of helium nuclei are converted to protons through the beta decay immediately compared to the cosmological time.

$$\frac{dN_{He}(t)}{dt} = -\tau_{He}^{-1} \cdot N_{He}(t). \tag{26}$$

The Friedman equation determines the evolution of the scale factor;

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{1}{3M_{pl}^2} \cdot \left(\frac{M(t)}{a^3} + \frac{S_{rad}(t)}{a^4} - \frac{M_{pl}^2}{a^2} + M_{pl}^2 \Lambda\right)$$
(27)

where

$$M(t) = m_p N_p(t) + m_{He} N_{He}(t)$$
 (28)

is the amount of the matter, and m_{He} is the mass of a helium nucleus. In this paper, we express it as

$$m_{He} = 2m_p + 2m_n - \Delta \tag{29}$$

where $\Delta = 28 \text{MeV}$ is the binding energy. Since what we need is the total radiation S_{rad} , we can neglect the curvature term and the cosmological constant term in Eq.(27). The increase rate of the radiation is given by

$$\frac{dS_{rad}(t)}{dt} = a(t)m_p \times \left(\tau_p^{-1} \cdot N_p(t) + (1 - 2\epsilon) \cdot \tau_{He}^{-1} N_{He}(t)\right). \tag{30}$$

Here we have assumed that while each proton creates the radiation m_p when it decays, each helium nucleus creates the radiation $(1-2\epsilon)m_p$. The initial values of $N_p(t)$, $N_{He}(t)$ are given by

$$N_p(0) = N_B(1 - 2X_n), (31)$$

$$N_{He}(0) = N_B \frac{X_n}{2},\tag{32}$$

where we use Eqs.(10)(20) as X_n and N_B . Using the above equations, we can calculate S_{rad} if we give the five quantities

$$m_p$$
 , τ_p , m_{He} , τ_{He} , ϵ . (33)

In principle, these quantities are determined by the QCD. In particular, if we fix the current quark masses, they do not depend on v_h . Therefore, the v_h dependence appears only in N_B and X_n .

3 The Big Fix of v_h

In this section, we first give qualitative arguments and then give numerical results. As discussed before, we study whether the radiation of the universe becomes maximum around the experimental value $v_h=246{\rm GeV}$.




Figure 3: The radiation produced by the baryons as a function of X_n . We can see that S_{rad} depends linearly on X_n and that its amount decreases if ϵ becomes large.

3.1 The Qualitative Argument

Although we have given the differential equations to calculate S_{rad} , the v_h dependence is almost the same as that of the simplified case(24). The problem is how atomic nuclei change it. To see this, we first need to know the X_n dependence of the radiation, and this can be checked by solving the differential equations numerically. See Fig.3. One can see that the radiation depends on X_n almost linearly. Thus, we can write S_{rad} as

$$S_{rad} = constant \times \left(\frac{1}{M_{pl}^2}\right)^{\frac{1}{3}} \times (N_B m_p)^{\frac{4}{3}} \tau_p^{\frac{2}{3}} \times \left(1 - c(\epsilon, \frac{\tau_{He}}{\tau_p}, \frac{m_{He}}{m_p}) X_n\right), \tag{34}$$

where the coefficient c is a function of ϵ , $\frac{\tau_{He}}{\tau_p}$ and $\frac{m_{He}}{m_p}$ because we can rewrite the differential equations in terms of them if we rescale a and t as

$$a(t) \to a'(t) := \frac{a(t)}{m_p^{\frac{1}{3}}} , \quad t \to t' := \frac{t}{\tau_p}.$$
 (35)

c represents how much the radiation is reduced by the existence of neutrons. The important fact is that, since the five parameters $m_p, m_{He}, \tau_p, \tau_{He}, \epsilon$ should be determined by the nuclear physics, we can treat c as a constant when we fix Λ_{QCD} and the current quark masses. This is the reason why we fix the current quark masses in the next subsection. In this case, the v_h dependence of S_{rad} is given by

$$S_{rad} = constant \times \underbrace{N_B^{\frac{4}{3}} \times (1 - c \cdot X_n)}_{v_h \text{ dependence when } m_q = \text{fixed}}$$
 (36)

In the previous section, we have studied the behavior of X_n and N_B as a function of v_h . Thus, one can easily understand why Eq.(36) has a maximum around $v_h = 246 \text{GeV}$ as follows. In the region of small $v_h < 200 \text{GeV}$, since X_n is approximately zero as is shown in Fig.2, we can write

$$S_{rad} \propto N_B^{\frac{4}{3}},\tag{37}$$

where N_B is an increasing function of v_h as is shown in Fig.1. Thus, S_{rad} is a increasing function of v_h in this region. On the other hands, in the region of $v_h >> 200 \text{GeV}$, since X_n increases and N_B is almost constant, the radiation decreases due to the effect of $c \cdot X_n$.

3.2 The Numerical Result

As discussed in the previous subsection, the reason why we fix the current quark masses is that we need not to consider complicated effects of nuclear physics. Fig.4 shows the numerical results at a sufficiently late time $(t=20\tau_p)$ by using the differential equations in Section 2. One can see that S_{rad} has a global maximum around $v_h \simeq 200 \text{GeV}$ if the life time of a helium nucleus τ_{He} is not too big. Since τ_{He} is expected to be within (see Appendix D)

$$\frac{1}{4}\tau_p < \tau_{He} < \frac{1}{4} \times \frac{7}{6}\tau_p,\tag{38}$$

we can conclude that the radiation of our universe has a global maximum very generally. We can also see how precisely the approximate formula

$$S_{rad} \propto N_B^{\frac{4}{3}} \times (1 - c \cdot X_n) \tag{39}$$

reproduces the above numerical result. For example, Fig.5 shows the case of the following parameters:

$$\epsilon = \frac{1}{32} , \frac{\tau_{He}}{\tau_p} = \frac{1}{4} \times \frac{49}{48} , \frac{m_{He}}{m_p} = 4 - \frac{28}{938}.$$
(40)

One can see that the numerical result (the green line) of the differential equations is almost completely reproduced by Eq.(39) if we choose the best fit value c = 0.010 (the black line). Note that the coefficient c is around

$$c \simeq \mathcal{O}(0.01). \tag{41}$$

The problem is how such a small value is obtained. By solving the differential equations for various initial conditions and parameters, we can read the behavior of c:

$$c\left(\epsilon, \frac{\tau_{He}}{\tau_p}, \frac{m_{He}}{m_p}\right) = 0.545 \times \epsilon - 1.829 \times \left(\frac{\tau_{He}}{\tau_p} - \frac{1}{4}\right) + 0.118 \times \left(4 - \frac{m_{He}}{m_p}\right) + \cdots$$
 (42)

One can see that while ϵ and the binding energy $4 - \frac{m_{He}}{m_p}$ have effects to decease the radiation, $\frac{\tau_{He}}{\tau_p} - \frac{1}{4}$ has an increasing effect. Thus, it is not so unnatural for c to become $\mathcal{O}(0.01)$. Actually, we can check that c stays of order 0.01 when we change ϵ , $\frac{\tau_{He}}{\tau_p}$ and $\frac{m_{He}}{m_p}$ within the physically reasonable region. Therefore, we can conclude that the QCD naturally gives the above small value, and the maximum entropy principle is justified for v_h .




Figure 4: The radiation as a function of v_h when the gauge couplings, the Higgs self coupling and the current quark masses are fixed. The upper(lower) graph shows the case $\epsilon = \frac{1}{32}(\frac{1}{64})$ for various values of $\frac{\tau_{He}}{\tau_p}$. We can see that S_{rad} has a global maximum around $v_h = 246 \,\mathrm{GeV}$ without the fine tunings of τ_{He} and ϵ .




Figure 5: The fitting result when $\epsilon = \frac{1}{32}, \frac{\tau_{He}}{\tau_p} = \frac{1}{4} \times \frac{49}{48}$ and $\frac{m_{He}}{m_p} = 4 - \frac{28}{938}$. The green line shows the result of the differential equations, and the black line shows the fit by Eq.(39) with the best fit value c = 0.010. They overlap almost completely.

4 Summary and Discussion




Figure 6: The radiation of the universe as a function of the sum of m_u and m_d . Here, we have used the approximate formulas which are given by Eq.(43)(44), and fixed v_h and $y_d - y_u$ at the observed values.

In this paper, we have shown that the radiation at the late stage of the universe becomes maximum around $v_h = 246 \text{GeV}$ assuming that the baryon number is produced by the sphaleron process and that the current quark masses m_i , the gauge copings and the Higgs self coupling are fixed. Since the radiation of the universe can be regarded as the entropy, this conclusion is the maximum entropy principle. The reason why we have fixed m_i is that we can examine the v_h dependence without going through the complicated nuclear physics. Actually the maximum of S_{rad} is determined by the behavior of N_B and X_n as functions of v_h . We have seen that the existence of atomic nuclei plays a very important role to maximize S_{rad} . In other words, the maximum entropy principle forces the existence of atomic nuclei, and as a consequence, the Higgs expectation value is fixed to around 246GeV. This is reminiscent of the anthropic principle [29, 30, 31] in which the existence of human beings requires atomic nuclei. In our case, however, they are naturally required by the fundamental low.

Although what we actually want to do is to regard S_{rad} as a function of the SM parameters, and confirm the maximum entropy principle by changing them independently, in this paper, we have considered only v_h so that we need not to consider the effects of nuclear physics. However, by using the approximate formulas such as

$$m_p = \alpha \Lambda_{QCD} + \beta (2m_u + m_d) \quad , \quad m_\pi^2 = \gamma \Lambda_{QCD} \frac{m_u + m_d}{2}, \tag{43}$$

$$\Gamma_p = \tau_p^{-1} \propto g^2 \frac{m_p^5}{M_X^4} \times \left(1 - \frac{m_\pi^2}{m_p^2}\right)^2,$$
(44)

it may be possible to determine also y_u and y_d . For example, Fig.6 shows the numerical result of S_{rad} as a function of $m_+ := m_u + m_d = (y_u + y_d) \cdot v_h / \sqrt{2}$ when the difference of the Yukawa couplings $y_- = y_d - y_u$ and v_h are fixed at the observed values. Here, we have used the ad-hoc values

$$\Lambda_{QCD} = 300 \text{MeV} , \quad \alpha = 3.1 , \quad \beta = 1.2 , \quad \frac{\tau_{He}}{\tau_p} = \frac{1}{4} , \quad \frac{m_{He}}{m_p} = 4 , \quad \epsilon = \frac{1}{32}, \quad (45)$$

and experimental value¹⁰

$$m_{\pi|\text{phys}} = 135 \text{MeV}.$$
 (46)

One can see that S_{rad} has a maximum around the experimental values. Furthermore, since S_{rad} is an increasing function of y_{-} if we use the above equations (see the qualitative equation (36)), we may conclude that the Yukawa couplings are determined by the maximum entropy principle at

$$(y_u, y_d) \simeq (0, 4 \times 10^{-5}).$$
 (47)

Making the above argument more precisely is the future work. Although the understanding of the maximum entropy principle is in a primitive level at present, it is very interesting to study whether the radiation of the universe at the late stage is really decreased when we change each of the SM parameters independently.

¹⁰This can be used to determine $\gamma \times \Lambda_{QCD}$.

Acknowledgement

We thank Takahiro Kubota and Hiroshi Itoyama for valuable discussions. The work of Y. H. is supported by a Grant-in-Aid for Japan Society for the Promotion of Science(JSPS) Fellows No.25·1107.

Appendix A WKB Solution

Eigenstates of the wave function of the universe are given by

$$\hat{\mathcal{H}} \cdot \phi_E(a) = \left(-\frac{\hat{p}_a^2}{2M_{pl}^2 a} + \frac{a^3 \rho(a)}{6} \right) \cdot \phi_E(a) = E\phi_E(a) \tag{48}$$

$$\leftrightarrow \left(-\frac{1}{2}\hat{p}_a^2 - V(a)\right)\phi_E(a) = M_{pl}^2 a E \phi_E(a) \tag{49}$$

$$V(a) := -\frac{M_{pl}^2 a^4 \rho(a)}{3} = -\frac{M_{pl}^2}{3} \times a^4 \times \left(\frac{M}{a^3} + \frac{R}{a^4} - \frac{M_{pl}^2}{a^2} + M_{pl}^2 \Lambda\right)$$
 (50)

where $p_a = -M_{pl}^2 a\dot{a}$. According to the ordinary quantum mechanics, the WKB wave function is given by

$$\phi_E(a) \propto \exp\left(\frac{i}{\hbar}S_0 + iS_1 + o(\hbar)\right).$$
 (51)

We have to pay attention to the order of a and \hat{p}_a because the Hamiltonian has the kinetic term that depends on the scale factor a. The correct order is given by

$$\frac{p_a^2}{a} = -\hbar^2 \frac{d}{da} a \frac{d}{da} = -\hbar^2 \left(\frac{1}{a} \frac{d^2}{da^2} - \frac{1}{a^2} \frac{d}{da} \right). \tag{52}$$

If we assume this order, the WKB wave function (51) satisfies the following Schrödinger equation:

$$-\frac{1}{2} \left(\frac{d(S_0 + \hbar S_1)}{da} \right)^2 + \frac{i\hbar}{2} \frac{d^2(S_0 + \hbar S_1)}{da^2} - \frac{i\hbar}{2a} \frac{dS_0}{da} = M_{pl}^2 a E + V(a), \tag{53}$$

which leads to

$$\mathcal{O}(\hbar^0) : -\left(\frac{dS_0}{da}\right)^2 = 2\left(M_{pl}^2 a E + V(a)\right) := -p_{cl}^2,\tag{54}$$

$$\mathcal{O}(\hbar) : -2\frac{dS_0}{da} \cdot \frac{dS_1}{da} + i\left(\frac{d^2S_0}{da^2} - \frac{1}{2}\frac{dS_0}{da}\right) = 0.$$
 (55)

Solutions to this equations are

$$S_0 = \int^a da' p_{cl} \quad , \quad S_1 = \frac{i}{2} (\log p_{cl} - \log a) .$$
 (56)

Then, except for the normalization factor, Eq.(51) becomes

$$\phi_E \propto \exp\left(\frac{i}{\hbar}S_0 + iS_1 + o(\hbar)\right) = \frac{\sqrt{a}}{\sqrt{p_{cl}}} \exp\left(\frac{i}{\hbar}\int^a da' p_{cl}\right).$$
 (57)

Note that the factor \sqrt{a} is different from the ordinary WKB solution. For large a, the classical momentum becomes

$$p_{cl} = \sqrt{-2\left(V(a) + M_{pl}^2 a E\right)} \simeq \sqrt{-2M_{pl}^2 \left(-\frac{M_{pl}^2 \Lambda a^4}{3} + a E\right)} \simeq \sqrt{\frac{2\Lambda}{3}} \cdot (aM_{pl})^2 - \sqrt{\frac{3}{2\Lambda}} \cdot \frac{E}{a},$$

$$(58)$$

and the corresponding WKB solution (57) is

$$\phi_E \simeq \left(\frac{3}{2\Lambda a^2}\right)^{\frac{1}{4}} \times \exp\left[\frac{i}{\hbar} \left(\sqrt{\frac{2\Lambda}{3}} \frac{M_{pl}^2 a^3}{3} - \sqrt{\frac{3}{2\Lambda}} E \log a\right)\right].$$
 (59)

Finally we can obtain the correct inner product between these eigenstates:

$$(\phi_E, \phi_{E'}) \propto \sqrt{\frac{3}{2\Lambda}} \times \int da \frac{\exp\left[\frac{i}{\hbar}\sqrt{\frac{3}{2\Lambda}}(E - E')\log a\right]}{a}$$
$$= \hbar\delta\left(E - E'\right). \tag{60}$$

From Eq.(60), we can see the normalization factor as $\hbar^{\frac{1}{2}}$. Thus, the correct WKB solution is

$$\phi_E = \sqrt{\frac{a}{\hbar p_{cl}}} \times \exp\left(\frac{i}{\hbar} \int^a da' p_{cl}\right). \tag{61}$$

Appendix B Probabilistic Interpretation of the Wave Function of the Multiverse

In this Appendix, we review the previous works [2, 3] that interpret the squared absolute value of the wave function of the multiverse as the probability distribution. See also [5, 6, 7, 8, 9, 10, 11, 12, 13, 14].

Appendix B.1 Multiverse Path Integral

Coleman's first idea is that, by taking the wormhole configurations into account, we can obtain the multiverse path integral. We start with the Euclidean Einstein Gravity,

$$Z = \sum_{all\mathcal{M}} \int_{\mathcal{M}} \mathcal{D}g \mathcal{D}\phi \exp(-S_E^G - S_E^M)$$
 (62)

$$S_E^G = -\frac{1}{16\pi G} \int d^4x \sqrt{g(x)} (R + 2\Lambda_o)$$
(63)

$$S_E^M = \int d^4x \sqrt{g(x)} \mathcal{L}(\phi, \partial \phi), \tag{64}$$

where \mathcal{M} represents a four dimensional manifold, Λ_o is the bare cosmological constant and ϕ is matter field. Classical solution of the Euclidean Gravity is called the wormhole and Coleman discussed the dilute gas approximation of the wormhole configurations, and obtained

$$Z = \sum_{n=0}^{\infty} \int_{\mathcal{M}_n} \mathcal{D}g \mathcal{D}\phi \exp\{\sum_{i,j} c_{ij} \int dx^4 dy^4 \sqrt{g(x)} \sqrt{g(y)} \mathcal{O}(x)_i \mathcal{O}(y)_j \exp(-2S_W)\} \times \exp(-S_E^G - S_E^M).$$
(65)

where S_W is the classical action of a wormhole and \mathcal{M}_n represents a manifold such that there are n universes, which are disconnected each other. In fact, this result takes into account only the wormhole configurations having two legs. Including all the different types of wormholes, we obtain the following multi local effective action¹¹:

$$Z = \sum_{n=0}^{\infty} \int_{\mathcal{M}_n} \mathcal{D}g \mathcal{D}\phi \exp(-S_E^G - S_E^M - \sum_{k=2}^{\infty} Z_k)$$
 (66)

where Z_k comes from wormholes having k legs;

$$Z_k := \sum_{i_1, i_2 \cdots, i_k} c_{i_1 i_2 \cdots i_k} \prod_{j=1}^k \int d^4 x_j \sqrt{g(x_j)} \mathcal{O}^{i_j}(x_j) \exp(-S_W) := \sum_{i_1, i_2 \cdots, i_k} c'_{i_1 i_2 \cdots i_k} S_{i_1} S_{i_2} \cdots S_{i_k}.$$

$$(67)$$

Here,

$$S_i := \int dx^4 \sqrt{g(x)} \mathcal{O}^i(x) \tag{68}$$

is the ordinary local action. We use the Lorentzian counterpart of Eq.(67) by the Wick rotation:

$$\sum_{n=0}^{\infty} \int_{\mathcal{M}_n} \mathcal{D}g \mathcal{D}\phi \exp(iS_E^G + iS_E^M + i\sum_{k=2}^{\infty} Z_k) := \sum_{n=0}^{\infty} \int_{\mathcal{M}_n} \mathcal{D}g \mathcal{D}\phi \exp(iS_{eff})$$
 (69)

$$S_{eff} = S_E^G + S_E^M + \sum_{ij} c_{ij} S_i S_j + \sum_{ijk} c_{ijk} S_i S_j S_k + \cdots$$
 (70)

By making the Fourier transform, we can write Eq.(69) as the usual path integral form:

$$Z = \sum_{K} \int d\overrightarrow{\lambda} \omega(K, \lambda_{1}, \lambda_{2}, \cdots) \sum_{n=0}^{\infty} \int_{\mathcal{M}_{n}} \mathcal{D}g \mathcal{D}\phi \exp(i \sum_{i} \lambda_{i} S_{i})$$

$$:= \sum_{K} \int d\overrightarrow{\lambda} \omega(K, \lambda_{1}, \lambda_{2}, \cdots) \sum_{n=0}^{\infty} \frac{1}{n!} (Z_{universe}^{(\lambda)})^{n}.$$
(71)

¹¹We can also obtain the multi local action in matrix model. See [15].

Here,

$$Z_{universe}^{(\lambda)} := \int \mathcal{D}g\mathcal{D}\phi \, \exp(i\sum_{i}\lambda_{i}S_{i})$$
 (72)

is the ordinary path integral of the universe, and $\omega(K, \{\lambda_i\})$ is the Fourier coefficient and K represents the topology of a single universe. One can see that $\{\lambda_i\}$ serve as coupling constants of a single universe. Eq.(71) indicates that they are variables (not the constants). In the following discussion, we assume the two conditions besides the Wick rotation:

1. We consider the homogeneous, isotropic universe with $S^3(K=1)$ topology:

$$d^2s = -N(t)d^2t + a^2(t)\left(d\mathbf{x}^2 + \frac{(\mathbf{x} \cdot d\mathbf{x})^2}{1 - \mathbf{x}^2}\right). \tag{73}$$

2. Matter and radiation fields are included as the potential of the scale factor a(t), namely the Hamiltonian is given by

$$\mathcal{H}(\lambda) = -\frac{1}{2M_{pl}^2 a} p_a^2 + a^3 \rho(a). \tag{74}$$

where $\rho(a)$ is the energy density of a single universe.

Based on this assumptions, Eq. (72) becomes

$$Z_{universe}^{(\lambda)}(a_f, a_i) = \int \mathcal{D}p_a \int_{t=0, a(0)=a_i}^{t=1, a(1)=a_f} \mathcal{D}a\mathcal{D}N \exp\{i \int_0^1 dt (p_a \dot{a} - N\mathcal{H}(\lambda))\}.$$
 (75)

where we have written the boundary condition explicitly. By choosing the gauge such that N(t) = T(constant) and rescaling t as $t \to t \times T$, Eq.(75) becomes

$$Z_{universe}^{(\lambda)}(a_f, a_i) = \int_{-\infty}^{\infty} dT \int_{t=0, a(0)=a_i}^{t=T, a(T)=a_f} \mathcal{D}a \exp\{i \int_0^T dt (p_a \dot{a} - \mathcal{H}(\lambda))\}$$

$$= \int_{\infty}^{\infty} dT \langle a_f | \exp(-i\hat{\mathcal{H}}(\lambda)) | a_i \rangle$$

$$= \langle a_f | \delta(\hat{\mathcal{H}}(\lambda)) | a_i \rangle.$$
(76)

If the cosmological constant $\Lambda(\{\lambda_i\})$ is positive, the eigenvalues of $\mathcal{H}(\lambda)$ are continuous, so we can choose the complete set as follows:

$$\langle \phi_E | \phi_{E'} \rangle = \delta(E - E'),$$
 (77)

$$\hat{\mathcal{H}}(\lambda)|\phi_E\rangle = E|\phi_{E=0}\rangle,\tag{78}$$

$$1 = \int_{-\infty}^{\infty} |E\rangle\langle E|. \tag{79}$$

Inserting Eq.(79) to Eq.(76), we obtain

$$= \langle a_f | \delta(\hat{\mathcal{H}}(\lambda)) \int_{-\infty}^{\infty} |E\rangle \langle E| |a_i\rangle$$

= $\phi_{E=0}(a_f) \phi_{E=0}^*(a_i)$. (80)

On the other hands, when $\Lambda(\{\lambda_i\})$ is negative, the complete set becomes

$$\langle \phi_E | \phi_{E'} \rangle = \delta_{E E'}, \tag{81}$$

$$\hat{\mathcal{H}}(\lambda)|\phi_E\rangle = E|\phi_{E=0}\rangle,\tag{82}$$

$$1 = \sum_{E} |E\rangle\langle E|,. \tag{83}$$

because the eigenvalues of $\hat{\mathcal{H}}(\lambda)$ are discrete. As well as the $\Lambda(\{\lambda_i\}) > 0$ case, the path integral given by Eq.(76) becomes the same expression as Eq.(80).

Appendix B.2 Multiverse Wave Function and Probabilistic Interpretation

In this subsection, we construct the multiverse wave function. Because we have to treat the coupling constants $\{\lambda_i\}$ of the universe as variables, the quantum state of the n universes is given by

$$|\Psi_n, \{\lambda_i\}\rangle = \frac{1}{\sqrt{n!}} \mu_K^n |\phi_{universe}\rangle \otimes \cdots \otimes |\phi_{universe}\rangle \otimes \omega(\lambda) |\{\lambda_i\}\rangle, \tag{84}$$

where $|\phi_{universe}\rangle$ is the state of a single universe, and μ_K is the probability amplitude of a universe emerging from nothing. Here we have divided $|\Psi_n, \{\lambda_i\}\rangle$ by the factor $\sqrt{n!}$ because we can not distinguish each universe. We can now express the wave function of the n universes as

$$\Psi(a_1, a_2, \cdots, a_n, \{\lambda_i\}) = \frac{1}{\sqrt{n!}} \mu_K^n \cdot \omega(\lambda) \times \phi_{universe}(a_1, \{\lambda_i\}) \cdot \phi_{universe}(a_2, \{\lambda_i\}) \cdots \phi_{universe}(a_n, \{\lambda_i\})$$
(85)

where

$$\phi_{universe}(a, \{\lambda_i\}) := \langle a | \phi_{universe} \rangle = \int da' \ Z_{universe}^{(\lambda)}(a, a') \langle a' | \phi_{universe} \rangle \tag{86}$$

is the wave function of a single universe. We interpret Eq. (85) as follows:

Probabilistic Interpretation of Wave Function of Multiverse

We interpret that

$$|\Psi(a_1, a_2, \cdots, a_n, \{\lambda_i\})|^2$$
 (87)

represents the probability that n universes having coupling constants $\{\lambda_i\}$ and scale factors (a_1, a_2, \dots, a_n) are observed.

We can now obtain the probability distribution $P(\{\lambda_i\})$ by tracing out the number of universes and the scale factors a_i :

$$P(\{\lambda_i\}) = \sum_{n=0}^{\infty} \int \cdots \int \prod_{k=1}^{n} da_k |\Psi(a_1, a_2, \cdots, a_n, \{\lambda_i\})|^2$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} |\mu_K|^{2n} \cdot |\omega(\lambda)|^2 \cdot \prod_{k=1}^{n} \left(\int da_k |\phi_{universe}(a_k, \{\lambda_i\})|^2 \right)$$

$$= |\omega(\lambda)|^2 \times \exp\left(|\mu_K|^2 \cdot \int da |\phi_{universe}(a, \{\lambda_i\})|^2 \right). \tag{88}$$

To analyze Eq.(88), we must determine the wave function of the single universe $\phi_{universe}(a, \{\lambda_i\})$. In this paper, we assume that a universe emerges from nothing with a small size ϵ :

$$|\phi_{universe}\rangle = |\epsilon\rangle.$$
 (89)

Generally speaking, there is no restriction on $|\phi_{universe}\rangle$. For the present, Eq.(89) is our proposal for the quantum state of the universe.

By this assumption, the wave function of the universe (86) becomes

$$\phi_{universe}(a, \{\lambda_i\}) = \int da' Z_{universe}^{(\lambda)}(a, a') \langle a' | \epsilon \rangle$$

$$= Z_{universe}^{(\lambda)}(a, \epsilon) = \phi_{E=0}(a) \phi_{E=0}^*(\epsilon), \tag{90}$$

where we have used Eq.(80). Here, note that $\phi_{E=0}(a)$ depends on $\{\lambda_i\}$. Now we obtain the explicit expression of $P(\{\lambda_i\})$:

$$P(\{\lambda_i\}) = |\omega(\lambda)|^2 \times \exp\left(|\mu_K \phi_{E=0}(\epsilon)|^2 \cdot \int da |\phi_{E=0}(a)|^2\right). \tag{91}$$

The intuitive understanding of this equation is as follows. By using Eqs. (76)(80) and regularizing the time integral by the cut off T_{Max} , the integral included in Eq. (91) becomes

$$|\phi_{E=0}(\epsilon)|^2 \int da |\phi_{E=0}(a)|^2 = \int da Z_{universe}^{(\lambda)*}(a,\epsilon) \times Z_{universe}^{(\lambda)}(a,\epsilon)$$

$$= \int da \langle \epsilon | \int_{-T_{Max}}^{T_{Max}} dT e^{-i\mathcal{H}(\lambda)} |a\rangle \langle a|E=0\rangle \times \langle \epsilon|E=0\rangle$$

$$= \int_{-T_{Max}}^{T_{Max}} dT \langle \epsilon|E=0\rangle \langle \epsilon|E=0\rangle$$

$$= 2T_{Max} \times |\phi_{E=0}(\epsilon)|^2. \tag{92}$$

Thus, Eq.(91) has a strong peak where the life time of the universe becomes maximum. In conclusion, the dynamics of the multiverse and wormholes naturally fixes the low energy parameters $\{\lambda_i\}$ in such a way that the life time of the universe becomes maximum.




Figure 7: The potentials of the S^3 universe for the various values of the Cosmological Constant Λ . If $\Lambda = \Lambda_{cr}$, the universe spends infinite times to grow up to the size a_* .

Appendix B.3 The Maximum Entropy Principle

In this section, we show how the maximum entropy principle is deduced for the S^3 universe by using Eq.(91). Eq.(91) has the following $\{\lambda_i\}$ dependences:

$$\omega(\lambda) , |\phi_{E=0}(\epsilon)|^2 , \int da |\phi_{E=0}|^2.$$
 (93)

Among them, $\omega(\lambda)$ and $|\phi_{E=0}(\epsilon)|^2$ are not so important because they do not have a sharp peak. On the other hands, as we will see, the integral

$$\int da |\phi_{E=0}(a)|^2 \tag{94}$$

depends on $\{\lambda_i\}$ most strongly. The hamiltonian of the S^3 universe is given by (see Appendix A)

$$\mathcal{H}(\lambda) = -\frac{1}{2}p_a^2 - V(a),\tag{95}$$

where

$$V(a) := -\frac{M_{pl}^2 a^4 \rho(a)}{3} = -\frac{M_{pl}^2}{3} \times a^4 \times \left(\frac{M}{a^3} + \frac{R}{a^4} - \frac{M_{pl}^2}{a^2} + M_{pl}^2 \Lambda\right)$$
(96)

is the potential of the universe. As discussed in [2][3], there is the critical surface of codimension one on which we have

$$\Lambda_{cr}(\{\lambda_i\}_{cr}) \simeq \begin{cases}
\frac{M_{pl}^2}{R(\lambda_{cr})} & \text{(for radiation dominated universe)} \\
\frac{M_{pl}^4}{M(\lambda_{cr})^2} & \text{(for matter dominated universe)}
\end{cases}$$
(97)

such that the maximum value of the potential $V(a^*)$ becomes 0 (see Fig.7). In this case, the integral (94) becomes

$$\int da |\phi_{E=0}(a)|^2 \simeq \begin{cases} \frac{R^{\frac{1}{2}}(\lambda_{cr})}{M_{pl}} \log(\frac{R^{\frac{1}{2}}(\lambda_{cr})}{M_{pl}\Delta a_{WKB}}) + \frac{R^{\frac{1}{2}}(\lambda_{cr})}{M_{pl}} \log(\frac{a_{IR}}{a^*}) & \text{(for radiation dominated universe)} \\ \frac{M(\lambda_{cr})}{M_{pl}^2} \log(\frac{M(\lambda_{cr})}{\Delta a_{WKB}}) + \frac{M(\lambda_{cr})}{M_{pl}^2} \log(\frac{a_{IR}}{a^*}) & \text{(for matter dominated universe)} \end{cases}$$

$$(98)$$

where we have introduced the maximum radius of the universe a_{IR} , and Δa_{WKB} represents the length of the violation the WKB approximation, namely, we can not use the WKB approximation in the $|a - a^*| < \Delta a_{WKB}$ region. In (98), the first term comes from the neighborhood of the critical point a^* , and the second term comes form the IR region. In this approximation, the both results of (98) have a peak at the point on the critical surface where

 $R(\lambda_i)$ $(M(\lambda_i))$ becomes maximum for the radiation (matter) dominated universe.

Therefore, the probability density (91) has a strong peak on the critical surface where R(M) is maximized, which we call the maximum entropy (matter) principle.

Appendix C Sphaleron Process and its v_h Dependence

In this appendix, we use the convention in [33]. In the SM, the baryon and lepton number are not conserved as a consequence of the anomaly. This is related to the topological charge of the SU(2) gauge field,

$$B(t_f) - B(t_i) = N_f \{ N_{cs}(t_f) - N_{cs}(t_i) \},$$
(99)

where

$$N_{cs}(t) = \frac{g^2}{32\pi^2} \int d^3x \epsilon^{ijk} \text{Tr}[A_i \partial_j A_k + \frac{2}{3} ig A_i A_j A_k]$$
 (100)

is the Cern-Simon number, and N_f is the number of the generations. The transition rate at zero temperature is given by the instanton action,

$$\Gamma \simeq e^{-S_{\text{Ins}}} = e^{-\frac{4\pi}{g^2}}$$

$$= \mathcal{O}(10^{-165}). \tag{101}$$

Since this rate is very small, the violation of B (or L) does not occur at zero temperature. However, in a thermal bath, one can make transition between the gauge vacua, through thermal fluctuations. This transition rate is determined by the sphaleron configuration, a classical solution of the field theory [34]. The transition rate of this process is [33]

$$\Gamma_{sph} \simeq \alpha_W^4 \times T \times e^{-\frac{E_{sph}}{T}}$$
 (102)

where $\alpha_W = g_2^2/4\pi$ and E_{sph} is the sphaleron energy. If the expansion rate of the universe $\frac{\dot{a}}{a} = H$ becomes lager than Γ_{aph} , the sphaleron process does not occur, and the total baryon number is fixed. The decoupling temperature is determined by equating H and Γ_{sph} ;

$$H \simeq \frac{T_{dec}^2}{M_{vl}} \simeq \alpha_W^4 \times T_{dec} \times e^{-\frac{E_{sph}}{T_{dec}}}.$$
 (103)

Note that we have used $H \simeq \frac{T^2}{M_{pl}}$ because this process happens in the radiation dominated era. The important fact is that, if T_{dec} is comparable with the mass of the heavy particle, its abundance is suppressed by the factor $e^{-\frac{m}{T_{dec}}}$. To analyze the decoupling temperature, we will use the recent numerical result [27] in which the gauge couplings and the Higgs self couplings are fixed to the experimental value. One can obtain the sphaleron energy from the Fig.2 presented in [27] as

$$\log\left(\frac{\Gamma_{sph}}{T}\right) = \log\left(\alpha_W^4\right) - \frac{E_{sph}(T)}{T}$$

$$\simeq \log(10^{-6}) + 123 \times \frac{T - T_c}{T},$$
(104)

which leads to

$$\Gamma_{sph} \simeq T \times e^{-13.8 + 123 \frac{T - T_c}{T}},\tag{105}$$

where $T_c = 160 \text{GeV}$ is the critical temperature of the phase transition. By replacing T_c to

$$T_c = 160 \text{GeV} \times \frac{v_h}{246 \text{GeV}},\tag{106}$$

we obtain the v_h dependence of these quantities when the gauge couplings and the Higgs self coupling are fixed to the experimental values. Then, solving Eq.(103), we obtain the decoupling temperature

$$T_{dec} \simeq \frac{7}{8} T_c = \frac{7}{8} \times 160 \text{GeV} \times \frac{v_h}{246 \text{GeV}} = 140 \text{GeV} \times \frac{v_h}{246 \text{GeV}}.$$
 (107)

The number density of the particle species i is given by

$$n_{i} = \frac{g_{i}}{(2\pi)^{3}} \int \frac{dp^{3}}{\exp\left(\frac{\sqrt{p^{2}+m_{i}^{2}}-\mu_{i}}{T}\right) \pm 1}$$

$$= 4\pi g_{i} \left(\frac{T}{2\pi}\right)^{3} \int_{0}^{\infty} dx \frac{x^{2}}{\exp\left(\sqrt{x^{2}+\left(\frac{m_{i}}{T}\right)^{2}}-\frac{\mu_{i}}{T}\right) \pm 1}$$
(108)

where g_i represents the degree of freedom, and the sign \pm is - for bosons and + for fermions. We can obtain the antiparticle density \bar{n}_i by replacing μ_i with $-\mu_i$ in Eq.(108). Then, the difference between the number density of particle and antiparticle is

$$n_{i} - \bar{n}_{i} = 4\pi g_{i} \left(\frac{T}{2\pi}\right)^{3} \int_{0}^{\infty} dx \ x^{2} \times \left(\frac{1}{\exp\left(\sqrt{x^{2} + (\frac{m_{i}}{T})^{2}} - \frac{\mu_{i}}{T}\right) \pm 1} - \frac{1}{\exp\left(\sqrt{x^{2} + (\frac{m_{i}}{T})^{2}} + \frac{\mu_{i}}{T}\right) \pm 1}\right)$$

$$\simeq 8\pi g_{i} \mu_{i} \frac{T^{2}}{(2\pi)^{3}} \int_{0}^{\infty} dx \ x^{2} \times \frac{\exp\sqrt{x^{2} + (m_{i}/T)^{2}}}{\left(\exp\sqrt{x^{2} + (m_{i}/T)^{2}} \pm 1\right)^{2}}$$

$$= 2\pi g_{i} \mu_{i} \frac{T^{2}}{(2\pi)^{3}} \int_{0}^{\infty} dx \ x^{2} \begin{cases} \frac{1}{\cosh^{2}(\sqrt{x^{2} + (m_{i}/T)^{2}})} & \text{(for fermion)} \\ \frac{1}{\sinh^{2}(\sqrt{x^{2} + (m_{i}/T)^{2}})} & \text{(for boson)} \end{cases}$$

$$:= g_{i} \mu_{i} \frac{T^{2}}{6} \begin{cases} g_{f}(m_{i}) & \text{(for fermion)}, \\ g_{b}(m_{i}) & \text{(for boson)}, \end{cases}$$

$$(109)$$

where

$$g_f(m_i) := \frac{3}{2\pi^2} \int_0^\infty dx x^2 \frac{1}{\cosh^2(\frac{\sqrt{x^2 + (m_i/T)^2}}{2})},$$

$$g_b(m_i) := \frac{3}{2\pi^2} \int_0^\infty dx x^2 \frac{1}{\sinh^2(\frac{\sqrt{x^2 + (m_i/T)^2}}{2})}.$$
(110)

Here we can take $\frac{T^2}{6}$ as a unit:

$$n_i - \bar{n}_i = g_i \mu_i \begin{cases} g_f(m_i) & \text{(for fermion)} \\ g_b(m_i) & \text{(for boson).} \end{cases}$$
(111)

where $g_f(0) = 1, g_b(0) = 2$. Note that the above density difference depends linearly on the chemical potential μ_i . We denote the chemical potentials for the conserved quantities as μ_a . Since the chemical potentials $\{\mu_i\}$ are conserved in thermal equilibrium, they can be written as linear combinations of the conserved quantum numbers;

$$\mu_i = \sum_a q_{ia} \mu_a. \tag{112}$$

In the broken phase, since the conserved quantum numbers are B-L and the electromagnetic charge Q, we can write the chemical potentials μ_i as follows:

$$\mu_u = \frac{1}{3}\mu_{B-L} + \frac{2}{3}\mu_Q \tag{113}$$

$$\mu_d = \frac{1}{3}\mu_{B-L} - \frac{1}{3}\mu_Q \tag{114}$$

$$\mu_e = -\mu_{B-L} - \mu_Q \tag{115}$$

$$\mu_{\nu_e} = -\mu_{B-L} \tag{116}$$

$$\mu_W = -\mu_Q. \tag{117}$$

We can eliminate μ_Q by using the fact that Q is zero:

$$Q = \frac{2}{3} \sum_{i:\text{u sector}} g_i g_f(m_i) \left(\frac{1}{3} \mu_{B-L} + \frac{2}{3} \mu_Q \right) - \frac{1}{3} \sum_{i:\text{d sector}} g_i g_f(m_i) \left(\frac{1}{3} \mu_{B-L} - \frac{1}{3} \mu_Q \right) - \sum_{i:\text{u sector}} g_i g_f(m_i) (-\mu_{B-L} - \mu_Q) + 3g_b(m_W) \mu_Q = 0.$$
(118)

This leads to

$$\mu_Q = -\mu_{B-L} \frac{4\sum_{i:u} g_f(m_i) - 2\sum_{i:d} g_f(m_i) + 6\sum_{i:e} g_f(m_i)}{8\sum_{i:u} g_f(m_i) + 2\sum_{i:d} g_f(m_i) + 6\sum_{i:e} g_f(m_i) + 9g_b(m_W)}.$$
 (119)

As a result, the baryon and lepton number densities are given by

$$B := n_{B} - \bar{n}_{B} = 6 \cdot \frac{1}{3} \left(\frac{1}{3} \mu_{B-L} + \frac{2}{3} \mu_{Q} \right) \cdot \sum_{i:\text{u sector}} g_{f}(m_{i}) + 6 \cdot \frac{1}{3} \left(\frac{1}{3} \mu_{B-L} - \frac{1}{3} \mu_{Q} \right) \cdot \sum_{i:\text{d sector}} g_{f}(m_{i})$$

$$= \frac{2\mu_{B-L}}{3} \left[\frac{6 \sum_{i:\text{d}} g_{f}(m_{i}) - 6 \sum_{i:\text{e}} g_{f}(m_{i}) + 9g_{b}(m_{W})}{8 \sum_{i:\text{u}} g_{f}(m_{i}) + 2 \sum_{i:\text{d}} g_{f}(m_{i}) + 6 \sum_{i:\text{e}} g_{f}(m_{i}) + 9g_{b}(m_{W})} \sum_{i:\text{u}} g_{f}(m_{i}) + \frac{12 \sum_{i:\text{u}} g_{f}(m_{i}) + 12 \sum_{i:\text{e}} g_{f}(m_{i}) + 9g_{b}(m_{W})}{8 \sum_{i:\text{u}} g_{f}(m_{i}) + 2 \sum_{i:\text{d}} g_{f}(m_{i}) + 6 \sum_{i:\text{e}} g_{f}(m_{i}) + 9g_{b}(m_{W})} \sum_{i:\text{d}} g_{f}(m_{i}) \right]$$

$$L := n_{L} - \bar{n}_{L} = 2 \cdot (-\mu_{B-L} - \mu_{Q}) \sum_{i:e} g_{f}(m_{i}) - 3\mu_{B-L}$$

$$= \mu_{B-L} \{ -2 \frac{4\{\sum_{i:u} g_{f}(m_{i}) + \sum_{i:d} g_{f}(m_{i})\} + 9g_{b}(m_{W})}{8\sum_{i:u} g_{f}(m_{i}) + 2\sum_{i:d} g_{f}(m_{i}) + 6\sum_{i:e} g_{f}(m_{i}) + 9g_{b}(m_{W})} \sum_{i:e} g_{f}(m_{i}) - 3 \}$$

$$(121)$$

where we have assumed that neutrinos are massless particles. Once N_{B-L} is given, the baryon number is given by

$$N_B = \frac{B}{B - L} N_{B - L}. \tag{122}$$

Since the general formula is very complicated, we do not write it here.

What we have to consider is whether the SM particles are relativistic or not at the sphaleron decoupling temperature. If a particle is non-relativistic, the integral (110) becomes very small, so it does not contribute to the baryon number. We can also read the T dependence of the vacuum expectation value v_h from the numerical result [27]:

$$v_h(T) = v_h \sqrt{\frac{T_c - T}{T_c}}. (123)$$

Therefore, at the decoupling temperature, the quark mass is given by

$$m_i(T_{dec}) = \frac{y_i v_h}{\sqrt{2}} \sqrt{\frac{T_c - T_{dec}}{T_c}} = m_{i0} \sqrt{\frac{T_c - T_{dec}}{T_c}}$$
 (124)

where m_{i0} is the current quark mass, and T_{dec} is given by Eq.(107). Since the suppression factors of the light particles start to have an effect at a very low temperature, it is almost valid to assume that all the particles except for the top quark and the W boson are massless. In this approximation, Eq.(122) becomes

$$N_B = \frac{2\{4 + g_b(m_W)\}\{5 + g_f(m_t)\}}{24\{5 + g_f(m_t)\} + g_b(m_W)\{37 + 2g_f(m_t)\}} N_{B-L}$$
(125)

This is the baryon number of the universe produced by the sphaleron process. We will use Eq.(125) in Section2 to estimate the radiation of the universe.

Appendix D Rough Estimations of τ_{He} and ϵ




Figure 8: The distribution of the remaining nucleons around the pion produced by the decay of a nucleon.

If we can apply the proton decay formula (44) to a nucleon in a helium nucleus, its life time is given by

$$\tau_{\rm N} \propto g^{-2} \frac{M_X^4}{(m_{p,n} - \delta)^5} (1 - \frac{m_\pi^2}{(m_{p,n} - \delta)^2})^{-2}$$

$$\simeq \tau_p (1 + 5 \frac{\delta}{m_{p,n}}) \tag{126}$$

where δ represents the fraction of the binding energy, and is expected to be

$$0 < \delta < \Delta. \tag{127}$$

Here $\Delta = 28 \text{MeV}$ is the binding energy of a helium nucleus. The life time of a helium nucleus is given by

$$\tau_{He} = \frac{\tau_N}{4} = \tau_p \frac{1 + 5\frac{\delta}{m_p}}{4},\tag{128}$$

because it has four nucleons. As a result, τ_{He} is in the region

$$\frac{1}{4}\tau_p < \tau_{He} < \frac{7}{24}\tau_p. \tag{129}$$

In this paper, we sweep τ_{He} in this region.

Another important fact about the atomic nuclei is that the pion produced by the decay of a nucleon in a helium nucleus may be scattered by the remaining nucleons, and loses its energy. See Fig.8. We express this effect by the dimensionless parameter ϵ . Namely, ϵ represents the amount of the decrease of the radiation for each neutron due to the pion scattering. In principle, ϵ can be determined by the QCD, but here we give a rough estimate. The cross section of the pion-nucleon scattering is given by [32]

$$\sigma \simeq 3.0 \text{fm}^2, \tag{130}$$

and the radius of nucleon is

$$r_N = 0.86 \text{fm}.$$
 (131)

We assume that the pion produced by the decay of a nucleon in a helium nucleus is surrounded by the other nucleons with a distance

$$R \simeq 2r_N,\tag{132}$$

Then, the scattering probability is estimated as

$$P_{\text{sca}} = \frac{3\sigma}{4\pi R^2}$$

$$\simeq \frac{1}{4}.$$
(133)

When a free proton decays, it produces the radiation energy m_p , on the other hands, when a nucleon in a helium nucleus decays, the radiation is reduced to $\frac{1}{2}m_p$ if the pion loses the energy completely by the scattering. Therefore, the radiation energy is maximally reduced by $\frac{1}{4}m_p$ per neutrons. This means the maximum value of ϵ is

$$\epsilon_{Max} = \frac{1}{4} \times P_{\text{sca}} = \frac{1}{16}.\tag{134}$$

The real value of ϵ is, of course, lower than Eq.(134), since the pion does not necessary lose all of its energy by the scattering. Fig.9 shows the kinematical situation. Since this is




Figure 9:

Figure 10: The kinematic which shows that a pion is scattered by a nucleon and loses its energy partly.

easily determined by solving the energy and momentum conservations, we do not follow the calculation here. When the pion is scattered to the angle $\theta = \pi$, the energy loss becomes maximum,

$$\frac{E_2}{E_0}(\theta = \pi) \simeq \frac{1}{2}.\tag{135}$$

Therefore, the real value of ϵ should be smaller than

$$\frac{1}{16} \times \frac{1}{2} = \frac{1}{32}.\tag{136}$$

To make more quantitative argument, we need to know how the nucleons are distributed in a helium nucleus.

Appendix E What if DM Dominates at the Late Stage of the Universe?

In this Appendix, we propose a scenario in which the DM physics is related to the Big Fix of v_h . As a simple extension of the SM which includes DM, we consider the Higgs portal scalar DM [35, 36]. This DM only interacts with the Higgs field and its stability is guaranteed by Z_2 symmetry. The Lagrangian is as follows:

$$\mathcal{L} = \mathcal{L}_{SM} + \frac{1}{2} (\partial_{\mu} S)^2 - \frac{1}{2} m_S^2 S^2 - \frac{\rho}{4!} S^4 - \frac{\kappa}{2} S^2 H^{\dagger} H, \tag{137}$$

where S is the DM field.

Let us assume that, at the late stage of the universe, the energy of the universe is dominated by the DM. If this is the case, various parameters are fixed so that the energy of the DM, $M_{\rm DM} = m_{\rm DM} n_{\rm DM} a_{\rm end}^3$, is maximized. This quantity is determined when the DM decouples from the thermal bath. Following the argument of Lee and Weinberg [37], $M_{\rm DM}$ is




Figure 11: The annihilation cross section of Higgs portal DM. The parameters are set to be $\kappa = 0.5, m_h = 126 \,\text{GeV}.$

roughly proportional to

$$M_{\rm DM} \propto \frac{1}{\langle \sigma v \rangle}$$
 (138)

Here σ and v denote the annihilation cross section and the relative velocity, respectively. $\langle ... \rangle$ means the thermal average. The cross section of annihilation processes of the scalar singlet DM can be found in [36]:¹²

$$\begin{split} \langle \sigma v \rangle &= \frac{\kappa^2}{8\pi m_S^2 \left[\left(4 - \frac{m_h^2}{m_S^2} \right)^2 + \frac{m_h^2}{m_S^2} \frac{\Gamma_h^2}{m_S^2} \right]} \\ &= \begin{cases} 6 \frac{m_b^2}{m_S^2} \left(1 - \frac{m_b^2}{m_S^2} \right)^{3/2} & b\bar{b} \end{cases} \\ &= \begin{cases} \delta_{W(Z)} \left(\frac{m_{W(Z)}^2}{m_S^2} \right)^2 \left(2 + \left(1 - 2 \frac{m_S^2}{m_{W(Z)}^2} \right)^2 \right) \sqrt{1 - \frac{m_{W(Z)}^2}{m_S^2}} & WW(ZZ) \\ 2 \left(\frac{\kappa}{\lambda} \frac{1 - \frac{1}{4} \frac{m_h^2}{m_S^2}}{1 - 2 \frac{m_S^2}{m_h^2}} + 1 + \frac{1}{2} \frac{m_h^2}{m_S^2} \right)^2 \sqrt{1 - \frac{m_h^2}{m_S^2}} & hh \end{cases} \end{split}$$

¹²This expression does not include the contributions from 4-body decay through virtual W and Z. Although such contributions become important around $m_S = m_{Z,W}$, this does not change our conclusion, and we ignore this effect.

where $\delta_W = 1, \delta_Z = 1/2$ and $\Gamma_h = 4.07 \text{MeV}$. We plot $\langle \sigma v \rangle^{-1}$ as a function of m_{DM} in Fig.11. It is found that there are three local minima at $m_{\text{DM}} \simeq 80, 90$, and 126 GeV. The global minimum corresponds to runaway m_{DM} . These local minima come from the threshold of W, Z boson and Higgs boson. This observation means that if m_{DM} is equal to m_W, m_Z or m_h , then M_{DM} is locally maximized. However, singlet scalar with $m_{\text{DM}} \simeq 80, 90 \text{GeV}$ has already excluded by direct detection [36]. Therefore, only the possibility is that the mass of the Higgs and that of the DM are the same, $m_h = m_{\text{DM}} \simeq 126 \text{GeV}$. This possibility is testable by the direct detection experiments [38].

References

- [1] S. R. Coleman, Nucl. Phys. B **310** (1988) 643.
- [2] H. Kawai and T. Okada, Int. J. Mod. Phys. A 26 (2011) 3107 [arXiv:1104.1764 [hep-th]].
- [3] H. Kawai and T. Okada, Prog. Theor. Phys. 127 (2012) 689 [arXiv:1110.2303 [hep-th]].
- [4] H. Kawai, Int. J. Mod. Phys. A 28, 1340001 (2013).
- [5] I. R. Klebanov, L. Susskind and T. Banks, Nucl. Phys. B **317**, 665 (1989).
- [6] J. Polchinski, Phys. Lett. B **219**, 251 (1989).
- [7] S. B. Giddings and A. Strominger, Nucl. Phys. B **321**, 481 (1989).
- [8] H. B. Nielsen and M. Ninomiya, Phys. Rev. Lett. 62, 1429 (1989).
- [9] B. Grinstein and M. B. Wise, Phys. Lett. B **212**, 407 (1988).
- [10] J. Preskill, S. P. Trivedi and M. B. Wise, Phys. Lett. B 223, 26 (1989).
- [11] V. A. Rubakov, Phys. Lett. B **214**, 503 (1988).
- [12] A. Strominger, Nucl. Phys. B **319**, 722 (1989).
- [13] W. Fischler, I. R. Klebanov, J. Polchinski and L. Susskind, Nucl. Phys. B 327, 157 (1989).
- [14] J. M. Cline, Phys. Lett. B **224**, 53 (1989).
- [15] Y. Asano, H. Kawai and A. Tsuchiya, Int. J. Mod. Phys. A 27, 1250089 (2012) [arXiv:1205.1468 [hep-th]].
- [16] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716** (2012) 30 [arXiv:1207.7235 [hep-ex]].
- [17] G. Aad et al. [ATLAS Collaboration], Phys. Lett. B 716 (2012) 1 [arXiv:1207.7214 [hep-ex]].
- [18] M. Holthausen, K. S. Lim and M. Lindner, JHEP 1202 (2012) 037 [arXiv:1112.2415 [hep-ph]]; F. Bezrukov, M. Y. .Kalmykov, B. A. Kniehl and M. Shaposhnikov, JHEP 1210 (2012) 140 [arXiv:1205.2893 [hep-ph]]; G. Degrassi, S. Di Vita, J. Elias-Miro,

- J. R. Espinosa, G. F. Giudice, G. Isidori and A. Strumia, JHEP **1208** (2012) 098 [arXiv:1205.6497 [hep-ph]]; S. Alekhin, A. Djouadi and S. Moch, Phys. Lett. B **716** (2012) 214 [arXiv:1207.0980 [hep-ph]]; I. Masina, Phys. Rev. D **87** (2013) 5, 053001 [arXiv:1209.0393 [hep-ph]]; Y. Hamada, H. Kawai and K. -y. Oda, Phys. Rev. D **87** (2013) 5, 053009 [arXiv:1210.2538 [hep-ph]]; arXiv:1305.7055 [hep-ph]; F. Jegerlehner; arXiv:1304.7813 [hep-ph]; arXiv:1305.6652 [hep-ph]; D. Buttazzo, G. Degrassi, P. P. Giardino, G. F. Giudice, F. Sala, A. Salvio and A. Strumia, JHEP **1312** (2013) 089 [arXiv:1307.3536].
- [19] C. D. Froggatt and H. B. Nielsen, Phys. Lett. B 368, 96 (1996) [hep-ph/9511371].
- [20] C. D. Froggatt, H. B. Nielsen and Y. Takanishi, Phys. Rev. D 64, 113014 (2001) [hep-ph/0104161].
- [21] H. B. Nielsen, arXiv:1212.5716 [hep-ph].
- [22] K. A. Meissner and H. Nicolai, Phys. Lett. B 648, 312 (2007) [hep-th/0612165];
 K. A. Meissner and H. Nicolai, Phys. Lett. B 660, 260 (2008) [arXiv:0710.2840 [hep-th]];
 S. Iso, N. Okada and Y. Orikasa, Phys. Lett. B 676, 81 (2009) [arXiv:0902.4050 [hep-ph]];
 S. Iso, N. Okada and Y. Orikasa, Phys. Rev. D 80, 115007 (2009) [arXiv:0909.0128 [hep-ph]];
 H. Aoki and S. Iso, Phys. Rev. D 86, 013001 (2012) [arXiv:1201.0857 [hep-ph]];
 S. Iso and Y. Orikasa, PTEP 2013, 023B08 (2013) [arXiv:1210.2848 [hep-ph]];
 M. Hashimoto, S. Iso and Y. Orikasa, Phys. Rev. D 89, 016019 (2014) [arXiv:1310.4304 [hep-ph]];
 M. Hashimoto, S. Iso and Y. Orikasa, Phys. Rev. D 89, 056010 (2014) [arXiv:1401.5944 [hep-ph]];
 R. Foot, A. Kobakhidze, K. L. McDonald and R. R. Volkas, Phys. Rev. D 77, 035006 (2008) [arXiv:0709.2750 [hep-ph]].
- [23] Y. Kawamura, PTEP 2013, no. 11, 113B04 (2013) [arXiv:1308.5069 [hep-ph]];
 Y. Kawamura, arXiv:1311.2365 [hep-ph].
- [24] M. Shaposhnikov and C. Wetterich, Phys. Lett. B 683, 196 (2010) [arXiv:0912.0208 [hep-th]].
- [25] A. Alves, E. R. Barreto and A. G. Dias, arXiv:1312.5333 [hep-ph].
- [26] F. L. Bezrukov and M. Shaposhnikov, Phys. Lett. B 659 (2008) 703 [arXiv:0710.3755 [hep-th]]; K. Allison, JHEP 1402 (2014) 040 [arXiv:1306.6931 [hep-ph]]; Y. Hamada, H. Kawai and K. -y. Oda, PTEP 2014 (2014) 023B02 [arXiv:1308.6651 [hep-ph]]; J. L. Cook, L. M. Krauss, A. J. Long and S. Sabharwal, arXiv:1403.4971 [astro-ph.CO]; Y. Hamada, H. Kawai, K. -y. Oda and S. C. Park, arXiv:1403.5043 [hep-ph]; F. Bezrukov and M. Shaposhnikov, arXiv:1403.6078 [hep-ph].
- [27] M. D'Onofrio, K. Rummukainen and A. Tranberg, PoS LATTICE 2012, 055 (2012) [arXiv:1212.3206].
- [28] S. Weinberg, Cosmology, first edition, Oxford 2008.
- [29] H. Martel, P. R. Shapiro and S. Weinberg, Astrophys. J. 492, 29 (1998) [astro-ph/9701099].

- [30] V. Agrawal, S. M. Barr, J. F. Donoghue and D. Seckel, Phys. Rev. D 57, 5480 (1998) [hep-ph/9707380].
- [31] V. Agrawal, S. M. Barr, J. F. Donoghue and D. Seckel, Phys. Rev. Lett. 80, 1822 (1998) [hep-ph/9801253].
- [32] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).
- [33] W. Buchmuller, R. D. Peccei and T. Yanagida, Ann. Rev. Nucl. Part. Sci. **55**, 311 (2005) [hep-ph/0502169].
- [34] F. R. Klinkhamer and N. S. Manton, Phys. Rev. D 30, 2212 (1984).
- [35] V. Silveira and A. Zee, Phys. Lett. B 161 (1985) 136; J. McDonald, Phys. Rev. D 50 (1994) 3637 [hep-ph/0702143 [HEP-PH]]; C. P. Burgess, M. Pospelov and T. ter Veldhuis, Nucl. Phys. B 619 (2001) 709 [hep-ph/0011335]; H. Davoudiasl, R. Kitano, T. Li and H. Murayama, Phys. Lett. B 609 (2005) 117 [hep-ph/0405097]; B. Patt and F. Wilczek, hep-ph/0605188; N. Haba, K. Kaneta and R. Takahashi, JHEP 1404 (2014) 029 [arXiv:1312.2089 [hep-ph]]; N. Haba and R. Takahashi, arXiv:1404.4737 [hep-ph]; A. Eichhorn and M. M. Scherer, arXiv:1404.5962 [hep-ph]; Y. Hamada, H. Kawai and K. -y. Oda, arXiv:1404.6141 [hep-ph].
- [36] J. M. Cline and K. Kainulainen, JCAP 1301 (2013) 012 [arXiv:1210.4196 [hep-ph]];
 J. M. Cline, K. Kainulainen, P. Scott and C. Weniger, Phys. Rev. D 88 (2013) 055025 [arXiv:1306.4710 [hep-ph]].
- [37] B. W. Lee and S. Weinberg, Phys. Rev. Lett. **39** (1977) 165.
- [38] E. Aprile [XENON1T Collaboration], arXiv:1206.6288 [astro-ph.IM]; D. S. Akerib *et al.* [LUX Collaboration], arXiv:1310.8214 [astro-ph.CO].