CANONICAL CARTAN CONNECTION FOR 4-DIMENSIONAL CR-MANIFOLDS BELONGING TO GENERAL CLASS ||

SAMUEL POCCHIOLA

ABSTRACT

We study the equivalence problem for 4-dimensional CR-manifolds of CR-dimension 1 and codimension 2 which have been referred to as belonging to general class II in [9], and which are also known as Engel CR-manifolds. We construct a canonical Cartan connection on such CR-manifolds through Cartan equivalence's method, thus providing an alternative approach to the results contained in [1]. In particular, we give the explicit expression of 4 biholomorphic invariants, the annulation of which is a necessary and sufficient condition for an Engel manifold to be locally biholomorphic to Beloshapka's cubic in \mathbb{C}^3 .

1. INTRODUCTION

As highlighted by Henri Poincaré [14] in 1907, the (local) biholomorphic equivalence problem between two submanifolds M and M' of \mathbb{C}^N is to determine whether or not there exists a (local) biholomorphism ϕ of \mathbb{C}^N such that $\phi(M) = M'$. Elie Cartan [2, 3] solved this problem for hypersurfaces $M^3 \subset \mathbb{C}^2$ in 1932, as he constructed a "hyperspherical connection" on such hypersurfaces by using the powerful technique which is now referred to as Cartan's equivalence method.

Given a manifold M and some geometric data specified on M, which usually appears as a G-structure on M (i.e. a reduction of the bundle of coframes of M), Cartan's equivalence method seeks to provide a principal bundle P on M together with a coframe ω of 1-forms on P which is adapted to the geometric structure of M in the following sense: an isomorphism between two such geometric structures M and M' lifts to a unique isomorphism between P and P' which sends ω on ω' . The equivalence problem between M and M' is thus reduced to an equivalence problem between $\{e\}$ -structures, which is well understood [10, 15].

We recall that a CR-manifold M is a real manifold endowed with a subbundle L of $\mathbb{C} \otimes TM$ of even rank 2n such that

(1) $L \cap \overline{L} = \{0\}$

(2) L is formally integrable, i.e. $[L, L] \subset L$.

The integer n is the CR-dimension of M and $k = \dim M - 2n$ is the codimension of M. In a recent attempt [9] to solve the equivalence problem for CR-manifolds up to dimension 5, it has been shown that one can restrict the study to six different general classes of CR-manifolds of dimension ≤ 5 , which have been referred to as general classes I, II, III₁, III₂, IV₁ and IV₂. The aim of this paper is to provide a solution to the equivalence problem for CR-manifolds which belong to general class II, that is the CR-manifolds of dimension 4 and of CR-dimension 1 whose CR-bundle L satisfy the additional non-degeneracy condition:

$$\mathbb{C} \otimes TM = L + \overline{L} + [L, \overline{L}] + [L, [L, \overline{L}]],$$

meaning that $\mathbb{C} \otimes TM$ is spanned by L, \overline{L} and their Lie brackets up to order 3.

This problem has already been solved by Beloshapka, Ezhov and Schmalz in [1], where the CR-manifolds we study are called Engel manifolds. The present paper provides thus an alternative solution to the results contained in [1]. The main result is the following:

Theorem 1. Let M be a CR-manifold belonging to general class II. There exists a 5-dimensional subbundle P of the bundle of coframes $\mathbb{C} \otimes F(M)$ of M and a coframe $\omega := (\Lambda, \sigma, \rho, \zeta, \overline{\zeta})$ on P such that any CRdiffeomorphism h of M lifts to a bundle isomorphism h^* of P which satisfy $h^*(\omega) = \omega$. Moreover the structure equations of ω on P are of the form:

$$\begin{split} d\sigma &= 3\Lambda \wedge \sigma + \rho \wedge \zeta + \rho \wedge \overline{\zeta}, \\ d\rho &= 2\Lambda \wedge \rho + i\,\zeta \wedge \overline{\zeta} \\ d\zeta &= \Lambda \wedge \zeta + \Im_1\,\sigma \wedge \rho + \Im_2\,\sigma \wedge \zeta + \Im_3\,\sigma \wedge \overline{\zeta} + \Im_4\,\rho \wedge \zeta + \Im_5\,\rho \wedge \overline{\zeta}, \\ d\overline{\zeta} &= \Lambda \wedge \overline{\zeta} + \overline{\jmath_1}\,\sigma \wedge \rho + \overline{\jmath_3}\,\sigma \wedge \zeta + \overline{\jmath_2}\,\sigma \wedge \overline{\zeta} + \overline{\jmath_5}\,\rho \wedge \zeta + \overline{\jmath_4}\,\rho \wedge \overline{\zeta}, \\ d\Lambda &= \frac{i}{2}\Im_1\,\sigma \wedge \overline{\zeta} - \frac{i}{2}\,\overline{\jmath_1}\,\sigma \wedge \zeta - \frac{1}{3}\,(\Im_2 + \overline{\jmath_3})\,\rho \wedge \zeta - \frac{1}{3}\,(\overline{\jmath_2} + \Im_3)\,\rho \wedge \zeta \\ &+ \Im_0\,\sigma \wedge \zeta \,, \end{split}$$

where \mathfrak{I}_0 , \mathfrak{I}_1 , \mathfrak{I}_2 , \mathfrak{I}_3 , \mathfrak{I}_4 , \mathfrak{I}_5 , are functions on P.

An example of CR-manifold belonging to general class II is provided by Beloshapka's cubic $B \subset \mathbb{C}^3$, which is defined by the equations:

B:

$$w_1 = \overline{w_1} + 2 i z \overline{z},$$

$$w_2 = \overline{w_2} + 2 i z \overline{z} (z + \overline{z}).$$

Cartan's equivalence method has been applied to Beloshapka's cubic in [12] where it has been shown that the coframe $(\Lambda, \sigma, \rho, \zeta, \overline{\zeta})$ of theorem 1 satisfy

the simplified structure equations:

$$d\sigma = 3 \Lambda \wedge \sigma + \rho \wedge \zeta + \rho \wedge \zeta ,$$

$$d\rho = 2 \Lambda \wedge \rho + i \zeta \wedge \overline{\zeta} ,$$

$$d\zeta = \Lambda \wedge \zeta ,$$

$$d\overline{\zeta} = \Lambda \wedge \overline{\zeta} ,$$

$$d\Lambda = 0,$$

corresponding to the case where the biholomorphic invariants \mathfrak{I}_i vanish identically. From this result together with theorem 1, we deduce the existence of a Cartan connection on CR-manifolds belonging to general class II in section 4.

We start in section 2 with the construction of a canonical G-structure P^1 on M, (e.g. a subbundle of the bundle of coframes of M), which encodes the equivalence problem for M under CR-automorphisms in the following sense: a diffeomorphism

$$h: M \longrightarrow M$$

is a CR-automorphism of M if and only if

$$h^*: P^1 \longrightarrow P^1$$

is a G-structure isomorphism of P^1 . We refer to [9, 6, 7] for details on the results summarized in this section and to [15] for an introduction to G-structures. Section 3 is devoted to reduce successively P^1 to three subbundles:

$$P^4 \subset P^3 \subset P^2 \subset P^1,$$

which are still adapted to the biholomorphic equivalence problem for M. We use Cartan equivalence method, for which we refer to [10]. Eventually a Cartan connection is constructed on P^4 in section 4.

2. INITIAL G-STRUCTURE

Let M be a 4-dimensional CR-manifold belonging to general class II and \mathscr{L} be a local generator of the CR-bundle L of M. As M belongs to general class II, the two vector fields \mathscr{T}, \mathscr{S} , defined by:

$$\begin{aligned} \mathcal{T} &:= i \left[\mathcal{L}, \overline{\mathcal{L}} \right], \\ \mathcal{S} &:= \left[\mathcal{L}, \mathcal{T} \right], \end{aligned}$$

are such that:

$$4 = \operatorname{rank}_{\mathbb{C}} \left(\mathscr{L}, \overline{\mathscr{L}}, \mathscr{T}, \mathscr{S} \right),$$

namely

$$(\mathscr{L}, \overline{\mathscr{L}}, \mathscr{T}, \mathscr{S})$$
 is a frame on M .

As a result there exist two functions A and B such that:

$$\overline{\mathscr{S}} = A \cdot \mathscr{T} + B \cdot \mathscr{S}.$$

From the fact that $\overline{\overline{\mathscr{S}}} = \mathscr{S}$, the functions A and B satisfy the relations:

(1)
$$BB = 1,$$
$$\overline{A} + \overline{B}A = 0.$$

There also exist two functions P, Q such that:

$$[\mathscr{L},\mathscr{S}] = P \cdot \mathscr{T} + Q \cdot \mathscr{S}.$$

The conjugate of P and Q, \overline{P} and \overline{Q} , are given by the relations: (2)

$$\overline{Q} = \mathscr{L}(B) + BQ + 2A + \frac{\mathscr{L}(B)}{B},$$

$$\overline{P} = B\mathscr{L}(A) - A\mathscr{L}(B) - BAQ - A^2 - A\frac{\overline{\mathscr{L}}(B)}{B} + \overline{\mathscr{L}}(A) + B^2P.$$

The four functions A, B, P, Q appear to be fundamental as all other Lie brackets between the vector fields $\mathscr{L}, \overline{\mathscr{L}}, \mathscr{T}$ and \mathscr{S} are expressed in terms of these five functions and their $\{\mathscr{L}, \overline{\mathscr{L}}\}$ -derivatives ([7]).

In the case of an embedded CR-manifold $M \subset \mathbb{C}^3$, we can give an explicit formula for the fundamental vector field \mathscr{L} , and hence for the functions A, B, P, Q, in terms of a graphing function of M. We refer to [8] for details on this question. Let us just mention that the submanifold $M \subset \mathbb{C}^3$ is represented in local coordinates:

$$(z, w_1, w_2) := (x + iy, u_1 + iv_1, u_2 + iv_2)$$

as a graph:

$$v_1 = \phi_1(x, y, u_1, u_2)$$

$$v_2 = \phi_2(x, y, u_1, u_2).$$

There exists then a unique local generator \mathscr{L} of $T^{1,0}M$ of the form:

$$\mathscr{L} = \frac{\partial}{\partial z} + A^1 \frac{\partial}{\partial u_1} + A^2 \frac{\partial}{\partial u_2}$$

having conjugate:

$$\overline{\mathscr{L}} = \frac{\partial}{\partial \overline{z}} + \overline{A^1} \frac{\partial}{\partial u_1} + \overline{A^2} \frac{\partial}{\partial u_2}$$

which is a generator of $T^{0,1}M$, where the functions A^1 and A^2 are given by the determinants:

$$A^{1} := \frac{\begin{vmatrix} -\phi_{1,z} & \phi_{1,u_{2}} \\ -\phi_{2,z} & i + \phi_{2,u_{2}} \end{vmatrix}}{\begin{vmatrix} i + \phi_{1,u_{1}} & \phi_{1,u_{2}} \\ \phi_{2,u_{1}} & i + \phi_{2,u_{2}} \end{vmatrix}}, \qquad A^{2} := \frac{\begin{vmatrix} i + \phi_{1,u_{1}} & -\phi_{1,z} \\ \phi_{2,u_{1}} & -\phi_{2,z} \end{vmatrix}}{\begin{vmatrix} i + \phi_{1,u_{1}} & \phi_{1,u_{2}} \\ \phi_{2,u_{1}} & i + \phi_{2,u_{2}} \end{vmatrix}}.$$

Returning to the general case of abstract CR-manifolds, let us introduce the coframe

$$\omega_0 := \left(\sigma_0, \rho_0, \zeta_0, \overline{\zeta}_0\right),\,$$

as the dual coframe of $(\mathscr{S}, \mathscr{T}, \mathscr{L}, \overline{\mathscr{L}})$. We have [7]:

Lemma 1. The structure equations enjoyed by ω_0 are of the form:

$$d\sigma_{0} = H \sigma_{0} \wedge \rho_{0} + F \sigma_{0} \wedge \overline{\zeta}_{0} + Q \sigma_{0} \wedge \zeta_{0} + B \rho_{0} \wedge \overline{\zeta}_{0} + \rho_{0} \wedge \zeta_{0},$$

$$d\rho_{0} = G \sigma_{0} \wedge \rho_{0} + E \sigma_{0} \wedge \overline{\zeta}_{0} + P \sigma_{0} \wedge \zeta_{0} + A \rho_{0} \wedge \overline{\zeta}_{0} + i \zeta_{0} \wedge \overline{\zeta}_{0},$$

$$d\zeta_{0} = 0,$$

$$d\overline{\zeta}_{0} = 0,$$

where the four functions:

can be expressed in terms of the four fundamental functions:

and their $\{\mathscr{L}, \overline{\mathscr{L}}\}$ -derivatives as: $E := \mathscr{L}(A) + B P,$ $F := \mathscr{L}(B) + B Q + A,$ $G := i \mathscr{L}(\mathscr{L}(A)) + i P \mathscr{L}(B) - i \mathscr{L}(P) - i Q \mathscr{L}(A) + i P \mathscr{L}(B) + i B \mathscr{L}(P),$ $H := i \mathscr{L}(\mathscr{L}(B)) + i Q \mathscr{L}(B) + i B \mathscr{L}(Q) + 2i \mathscr{L}(A) - i \mathscr{L}(Q).$

Let $h: M \longrightarrow M$ be a CR-automorphism of M. As we have

$$h_*\left(L\right) = L,$$

there exists a non-vanishing complex-valued function a on M such that:

$$h_*(\mathscr{L}) = \mathsf{a}\mathscr{L}.$$

From the definition of \mathscr{T}, \mathscr{S} , and the invariance

$$h_*([X,Y]) = [h_*(X), h_*(Y)]$$

for any vector fields X, Y on M, we easily get the existence of four functions

$$\mathsf{b},\mathsf{c},\mathsf{d},\mathsf{e}:M\longrightarrow\mathbb{C},$$

such that:

$$h_* \begin{pmatrix} \mathscr{L} \\ \overline{\mathscr{L}} \\ \mathscr{T} \\ \mathscr{T} \\ \mathscr{S} \end{pmatrix} = \begin{pmatrix} \mathsf{a} & 0 & 0 & 0 \\ 0 & \overline{\mathsf{a}} & 0 & 0 \\ \mathsf{b} & \overline{\mathsf{b}} & \mathsf{a}\overline{\mathsf{a}} & 0 \\ \mathsf{e} & \mathsf{d} & \mathsf{c} & \mathsf{a}^2\overline{\mathsf{a}} \end{pmatrix} \cdot \begin{pmatrix} \mathscr{L} \\ \overline{\mathscr{L}} \\ \mathscr{T} \\ \mathscr{T} \\ \mathscr{S} \end{pmatrix}.$$

This is summarized in the following lemma [6]:

Lemma 2. Let $h : M \longrightarrow M$ a CR-automorphism of M and let G_1 be the subgroup of $GL_4(\mathbb{C})$

$$G_1 := \left\{ \begin{pmatrix} \mathsf{a}^2\overline{\mathsf{a}} & 0 & 0 & 0\\ \mathsf{c} & \mathsf{a}\overline{\mathsf{a}} & 0 & 0\\ \mathsf{d} & \mathsf{b} & \mathsf{a} & 0\\ \mathsf{e} & \overline{\mathsf{b}} & 0 & \overline{\mathsf{a}} \end{pmatrix}, \, \mathsf{a} \in \mathbb{C} \setminus \{0\}, \, \mathsf{b}, \mathsf{c}, \mathsf{d}, \mathsf{e} \in \mathbb{C} \right\}.$$

Then the pullback ω of ω_0 by h, $\omega := h^* \omega_0$, satisfies:

$$\omega = g \cdot \omega_0,$$

where g is smooth (locally defined) function $M \xrightarrow{g} G_1$.

This motivates the introduction of the subbundle P^1 of the bundle of coframes on M constituted by the coframes ω of the form

$$\omega := g \cdot \omega_0, \qquad g \in G_1$$

The next section is devoted to reduce successively P^1 to three subbundles:

$$P^4 \subset P^3 \subset P^2 \subset P^1$$

which are adapted to the biholomorphic equivalence problem for M.

3. Reductions of P^1

The coframe ω_0 gives a natural (local) trivialisation $P^1 \xrightarrow{tr} M \times G_1$ from which we may consider any differential form on M (resp. G_1) as a differential form on P^1 through the pullback by the first (resp. the second) component of tr. With this identification, the structure equations of P^1 are naturally obtained by the formula:

(3)
$$d\omega = dg \cdot g^{-1} \wedge \omega + g \cdot d\omega_0.$$

The term $g \cdot d\omega_0$ contains the so-called torsion coefficients of P^1 . A 1-form $\tilde{\alpha}$ on P^1 is called a modified Maurer-Cartan form if its restriction to any fiber of P^1 is a Maurer-Cartan form of G_1 , or equivalently, if it is of the form:

$$\widetilde{\alpha} := \alpha - x_{\sigma} \, \sigma - x_{\rho} \, \rho - x_{\zeta} \, \zeta - x_{\overline{\zeta}} \, \overline{\zeta},$$

where x_{σ} , x_{ρ} , x_{ζ} , $x_{\overline{\zeta}}$, are arbitrary complex-valued functions on M and where α is a Maurer-Cartan form of G_1 .

A basis for the Maurer-Cartan forms of G_1 is given by the following 1-forms:

$$\begin{split} \alpha^{1} &:= \frac{d\mathbf{a}}{\mathbf{a}}, \\ \alpha^{2} &:= -\frac{\mathbf{b}d\mathbf{a}}{\mathbf{a}^{2}\overline{\mathbf{a}}} + \frac{d\mathbf{b}}{\mathbf{a}\overline{\mathbf{a}}}, \\ \alpha^{3} &:= -\frac{\mathbf{c}d\mathbf{a}}{\overline{\mathbf{a}}\,\mathbf{a}^{3}} - \frac{\mathbf{c}d\overline{\mathbf{a}}}{\overline{\mathbf{a}}^{2}\mathbf{a}^{2}} + \frac{d\mathbf{c}}{\mathbf{a}^{2}\overline{\mathbf{a}}}, \\ \alpha^{4} &= -\frac{(\mathbf{d}\mathbf{a}\overline{\mathbf{a}} - \mathbf{b}\mathbf{c})\,d\mathbf{a}}{\mathbf{a}^{4}\overline{\mathbf{a}}^{2}} - \frac{\mathbf{c}d\mathbf{b}}{\mathbf{a}^{3}\overline{\mathbf{a}}^{2}} + \frac{d\mathbf{d}}{\mathbf{a}^{2}\overline{\mathbf{a}}}, \\ \alpha^{5} &= -\frac{(\mathbf{e}\mathbf{a}\overline{\mathbf{a}} - \overline{\mathbf{b}}\,\mathbf{c})\,d\overline{\mathbf{a}}}{\mathbf{a}^{3}\overline{\mathbf{a}}^{3}} - \frac{\mathbf{c}d\overline{\mathbf{b}}}{\mathbf{a}^{3}\overline{\mathbf{a}}^{2}} + \frac{d\mathbf{e}}{\mathbf{a}^{2}\overline{\mathbf{a}}}, \end{split}$$

together with their conjugate.

We derive the structure equations of P^1 from the relations (3), from which we extract the expression of $d\sigma$:

$$\begin{split} d\sigma &= 2 \; \alpha^1 \wedge \sigma + \overline{\alpha^1} \wedge \sigma \\ &+ T^{\sigma}_{\sigma\rho} \; \sigma \wedge \rho - T^{\sigma}_{\sigma\zeta} \; \sigma \wedge \zeta - T^{\sigma}_{\sigma\overline{\zeta}} \; \sigma \wedge \overline{\zeta} + \rho \wedge \zeta + \frac{\mathsf{a}}{\mathsf{a}} \; B \; \rho \wedge \overline{\zeta} \; , \end{split}$$

or equivalently:

$$d\sigma = 2 \ \widetilde{\alpha}^1 \wedge \sigma + \overline{\widetilde{\alpha}^1} \wedge \sigma + \rho \wedge \zeta + \frac{\mathsf{a}}{\mathsf{a}} B \ \rho \wedge \overline{\zeta} \ ,$$

for a modified Maurer-Cartan form $\tilde{\alpha}^1$. The coefficient

$$\frac{a}{\overline{a}}B,$$

which can not be absorbed for any choice of the modified Maurer-Cartan form $\tilde{\alpha}^1$, is referred to as an essential torsion coefficient. From standard results on Cartan theory (see [10, 15]), a diffeomorphism of M is an isomorphism of the G_1 -structure P^1 if and only if it is an isomorphism of the reduced bundle $P^2 \subset P^1$ consisting of those coframes ω on M such that

$$\frac{\mathsf{a}}{\mathsf{a}}B = 1$$

This is equivalent to the normalization:

$$\overline{\mathbf{a}} = \mathbf{a}B.$$

A coframe $\omega \in P^2$ is related to the coframe ω_0 by the relations:

$$\begin{cases} \boldsymbol{\sigma} = \mathbf{a}^3 \, B \, \sigma_0 \\ \boldsymbol{\rho} = \mathbf{c} \, \sigma_0 + \mathbf{a}^2 \, B \, \rho_0 \\ \boldsymbol{\zeta} = \mathbf{d} \, \sigma_0 + \mathbf{b} \, \rho_0 + \mathbf{a} \, \zeta_0 \\ \overline{\boldsymbol{\zeta}} = \mathbf{e} \, \sigma_0 + \overline{\mathbf{b}} \, \rho_0 + \mathbf{a} \, B \, \overline{\boldsymbol{\zeta}}_0, \end{cases}$$

which are equivalent to:

$$\begin{cases} \sigma = \mathbf{a'}^3 \, \sigma_1 \\ \rho = \mathbf{c'} \, \sigma_1 + \mathbf{a'}^2 \, \rho_1 \\ \zeta = \mathbf{d'} \, \sigma_1 + \mathbf{b} \, \rho_1 + \mathbf{a'} \, \zeta_1 \\ \overline{\zeta} = \mathbf{e'} \, \sigma_1 + \overline{\mathbf{b}} \, \rho_1 + \mathbf{a'} \, \overline{\zeta}_1, \end{cases}$$

where:

$$\sigma_1 := \frac{\sigma_0}{B^{\frac{1}{2}}}, \qquad \rho_1 := \rho_0, \qquad \zeta_1 := \frac{\zeta_0}{B^{\frac{1}{2}}},$$

and

$$x' := x \cdot B^{\frac{1}{2}},$$
 for $x = \mathsf{a}, \mathsf{c}, \mathsf{d}, \mathsf{e}$

We notice that a' is a real parameter, and that σ_1 is a real 1-form. Let ω_1 be the coframe $\omega_1 := (\sigma_1, \rho_1, \zeta_1, \overline{\zeta_1})$, and G_2 be the subgroup of G_1 :

$$G_2 := \left\{ \begin{pmatrix} \mathsf{a}^3 & 0 & 0 & 0 \\ \mathsf{c} & \mathsf{a}^2 & 0 & 0 \\ \mathsf{d} & \mathsf{b} & \mathsf{a} & 0 \\ \mathsf{e} & \overline{\mathsf{b}} & 0 & \mathsf{a} \end{pmatrix}, \, \mathsf{a} \in \mathbb{R} \setminus \{0\}, \, \mathsf{b}, \mathsf{c}, \mathsf{d}, \mathsf{e} \in \mathbb{C} \right\}.$$

A coframe ω on M belongs to P^2 if and only if there is a local function $g: M \xrightarrow{g} G_2$ such that $\omega = g \cdot \omega_1$, namely P^2 is a G_2 structure on M. The Maurer-Cartan forms of G_2 are given by:

$$\begin{split} \beta^1 &:= \frac{d\mathbf{a}}{\mathbf{a}}, \\ \beta^2 &:= -\frac{bd\mathbf{a}}{\mathbf{a}^3} + \frac{d\mathbf{b}}{\mathbf{a}^2}, \\ \beta^3 &:= -2\frac{\mathbf{c}d\mathbf{a}}{\mathbf{a}^4} + \frac{d\mathbf{c}}{\mathbf{a}^3}, \\ \beta^4 &= -\frac{(\mathbf{d}\mathbf{a}^2 - \mathbf{b}\mathbf{c})\,d\mathbf{a}}{\mathbf{a}^6} - \frac{\mathbf{c}d\mathbf{b}}{\mathbf{a}^5} + \frac{d\mathbf{d}}{\mathbf{a}^3}, \\ \beta^5 &= -\frac{\left(\mathbf{e}\mathbf{a}^2 - \overline{\mathbf{b}}\,\mathbf{c}\right)\,d\mathbf{a}}{\mathbf{a}^6} - \frac{\mathbf{c}d\overline{\mathbf{b}}}{\mathbf{a}^5} + \frac{d\mathbf{e}}{\mathbf{a}^3}, \end{split}$$

together with $\overline{\beta^2}$, $\overline{\beta^3}$, $\overline{\beta^4}$, $\overline{\beta^5}$. Using formula (3), we get the structure equations of P^2 :

$$d\sigma = 3 \beta^{1} \wedge \sigma + U^{\sigma}_{\sigma\rho} \sigma \wedge \rho + U^{\sigma}_{\sigma\zeta} \sigma \wedge \zeta + U^{\sigma}_{\sigma\overline{\zeta}} \sigma \wedge \overline{\zeta} + \rho \wedge \zeta + \rho \wedge \overline{\zeta}$$

$$\begin{split} d\rho &= 2\beta^1 \wedge \rho + \beta^3 \wedge \sigma \\ &+ U^{\rho}_{\sigma\rho} \, \sigma \wedge \rho + U^{\rho}_{\sigma\zeta} \, \sigma \wedge \zeta + U^{\rho}_{\sigma\overline{\zeta}} \, \sigma \wedge \overline{\zeta} \\ &+ U^{\rho}_{\rho\zeta} \, \rho \wedge \zeta + U^{\rho}_{\rho\overline{\zeta}} \, \rho \wedge \overline{\zeta} + i \, \zeta \wedge \overline{\zeta}, \end{split}$$

$$\begin{split} d\zeta &= \beta^1 \wedge \zeta + \beta^2 \wedge \rho + \beta^4 \wedge \sigma \\ &+ U^{\zeta}_{\sigma\rho} \, \sigma \wedge \rho + U^{\zeta}_{\sigma\zeta} \, \sigma \wedge \zeta + U^{\zeta}_{\sigma\overline{\zeta}} \, \sigma \wedge \overline{\zeta} + U^{\zeta}_{\rho\zeta} \, \rho \wedge \zeta \\ &+ U^{\zeta}_{\rho\overline{\zeta}} \, \rho \wedge \overline{\zeta} + U^{\zeta}_{\zeta\overline{\zeta}} \, \zeta \wedge \overline{\zeta}. \end{split}$$

Introducing the modified Maurer-Cartan forms:

$$\widetilde{\beta}^i = \beta^i - y_\sigma \, \sigma - y^i_\rho \, \rho - y^i_\zeta \, \zeta \, - y^i_{\overline{\zeta}} \, \overline{\zeta},$$

the structure equations rewrite:

$$d\sigma = 3 \ \widetilde{\beta}^{1} \wedge \sigma + \left(U^{\sigma}_{\sigma\rho} - 3 y^{1}_{\rho} \right) \ \sigma \wedge \rho + \left(U^{\sigma}_{\sigma\zeta} - 3 y^{1}_{\zeta} \right) \ \sigma \wedge \zeta + \left(U^{\sigma}_{\sigma\overline{\zeta}} - 3 y^{1}_{\overline{\zeta}} \right) \ \sigma \wedge \overline{\zeta} + \rho \wedge \zeta + \rho \wedge \overline{\zeta}$$

$$\begin{split} d\rho &= 2\widetilde{\beta}^1 \wedge \rho + \widetilde{\beta}^3 \wedge \sigma \\ &+ \left(U^{\rho}_{\sigma\rho} + 2\,y^1_{\sigma} - y^3_{\rho} \right) \,\, \sigma \wedge \rho + \left(U^{\rho}_{\sigma\zeta} - y^3_{\zeta} \right) \,\, \sigma \wedge \zeta \\ &+ \left(U^{\rho}_{\sigma\overline{\zeta}} - y^3_{\overline{\zeta}} \right) \,\, \sigma \wedge \overline{\zeta} + \left(U^{\rho}_{\rho\zeta} - 2\,y^1_{\zeta} \right) \,\, \rho \wedge \zeta \\ &+ \left(U^{\rho}_{\rho\overline{\zeta}} - 2\,y^1_{\overline{\zeta}} \right) \,\, \rho \wedge \overline{\zeta} + i \,\, \zeta \wedge \overline{\zeta} \,\,, \end{split}$$

$$\begin{split} d\zeta &= \widetilde{\beta^1} \wedge \zeta + \widetilde{\beta^2} \wedge \rho + \widetilde{\beta^4} \wedge \sigma \\ &+ \left(U_{\sigma\rho}^{\zeta} + y_{\sigma}^2 - y_{\rho}^4 \right) \ \sigma \wedge \rho + \left(U_{\sigma\zeta}^{\zeta} + y_{\sigma}^1 - y_{\zeta}^4 \right) \ \sigma \wedge \zeta \\ &+ \left(U_{\sigma\overline{\zeta}}^{\zeta} - y_{\overline{\zeta}}^4 \right) \ \sigma \wedge \overline{\zeta} + \left(U_{\rho\zeta}^{\zeta} + y_{\rho}^1 - y_{\zeta}^2 \right) \ \rho \wedge \zeta \\ &+ \left(U_{\rho\overline{\zeta}}^{\zeta} - y_{\overline{\zeta}}^2 \right) \ \rho \wedge \overline{\zeta} + \left(U_{\zeta\overline{\zeta}}^{\zeta} - y_{\overline{\zeta}}^1 \right) \ \zeta \wedge \overline{\zeta} \ , \end{split}$$

which leads to the following absorbtion equations:

$$\begin{array}{ll} 3 \ y_{\rho}^{1} = U_{\sigma\rho}^{\sigma}, & 3 \ y_{\zeta}^{1} = U_{\sigma\zeta}^{\sigma}, & 3 \ y_{\overline{\zeta}}^{1} = U_{\sigma\zeta}^{\sigma}, \\ -2 \ y_{\sigma}^{1} + y_{\rho}^{3} = U_{\sigma\rho}^{\rho}, & y_{\zeta}^{3} = U_{\sigma\zeta}^{\rho}, & y_{\overline{\zeta}}^{3} = U_{\sigma\overline{\zeta}}^{\rho}, \\ 2 \ y_{\zeta}^{1} = U_{\rho\zeta}^{\rho}, & 2 \ y_{\overline{\zeta}}^{1} = U_{\rho\overline{\zeta}}^{\rho}, & -y_{\sigma}^{2} + y_{\rho}^{4} = U_{\sigma\rho}^{\zeta}, \\ -y_{\sigma}^{1} + y_{\zeta}^{4} = U_{\sigma\zeta}^{\zeta}, & y_{\overline{\zeta}}^{4} = U_{\sigma\overline{\zeta}}^{\zeta}, & -y_{\rho}^{1} + y_{\zeta}^{2} = U_{\rho\zeta}^{\zeta}, \\ y_{\overline{\zeta}}^{2} = U_{\rho\overline{\zeta}}^{\zeta}, & y_{\overline{\zeta}}^{1} = U_{\zeta\overline{\zeta}}^{\zeta}. \end{array}$$

Eliminating $y_{\overline{\zeta}}^1$ among these equations leads to:

$$U^{\zeta}_{\zeta\overline{\zeta}} = \frac{1}{2} U^{\rho}_{\rho\overline{\zeta}} = \frac{1}{3} U^{\sigma}_{\sigma\overline{\zeta}},$$

from which we deduce the following normalizations:

$$c = a^2 C_0$$

and

$$\mathsf{b} = \mathsf{a} \, \mathbf{B}_0,$$

where:

$$\mathbf{C}_{0} := \left(\frac{1}{2}\frac{\mathscr{L}(B)}{B^{\frac{1}{2}}} + \frac{1}{2}QB^{\frac{1}{2}}\right),\,$$

and

$$\mathbf{B}_{0} := \left(\frac{i}{3} \, \overline{\frac{\mathscr{L}(B)}{B^{\frac{3}{2}}}} - \frac{i}{3} \, \frac{A}{B^{\frac{1}{2}}} - \frac{i}{6} \, B^{\frac{1}{2}}Q - \frac{i}{6} \, \frac{\mathscr{L}(B)}{B^{\frac{1}{2}}}\right).$$

We introduce the coframe $\omega_2 := \left(\sigma_2, \rho_2, \zeta_2, \overline{\zeta}_2\right)$ on M, defined by:

$$\begin{cases} \sigma_2 := \sigma_1, \\ \rho_2 := \rho_1 + \mathbf{C}_0 \, \sigma_1, \\ \zeta_2 := \zeta_1 + \mathbf{B}_0 \, \rho_1, \end{cases}$$

and the 3-dimensional subgroup $G_3 \subset G_2$:

$$G_3 := \left\{ \begin{pmatrix} \mathsf{a}^3 & 0 & 0 & 0 \\ 0 & \mathsf{a}^2 & 0 & 0 \\ \frac{\mathsf{d}}{\mathsf{d}} & 0 & \mathsf{a} & 0 \\ \frac{\mathsf{d}}{\mathsf{d}} & 0 & 0 & \mathsf{a} \end{pmatrix}, \, \mathsf{a} \in \mathbb{R} \setminus \{0\}, \, \mathsf{d} \in \mathbb{C} \right\}$$

The normalizations:

$$\mathsf{b} := \mathsf{a} \, \mathbf{B}_0, \qquad \qquad \mathsf{c} := \mathsf{a}^2 \, \mathbf{C}_0,$$

amount to consider the subbundle $P^3 \subset P^2$ consisting of those coframes ω of the form

$$\omega := g \cdot \omega_2, \quad \text{where } g \text{ is a function } g : M \xrightarrow{g} G_3.$$

A basis of the Maurer Cartan forms of G_3 is given by:

$$\gamma^1 := \frac{d\mathbf{a}}{\mathbf{a}}, \qquad \gamma^2 := -\frac{dd\mathbf{a}}{\mathbf{a}^4} + \frac{d\mathbf{d}}{\mathbf{a}^3}, \qquad \overline{\gamma}_2.$$

The structure equations of P^3 are:

$$\begin{split} d\sigma &= 3\,\gamma^1 \wedge \sigma \\ &+ V^{\sigma}_{\sigma\rho}\,\sigma \wedge \rho + V^{\sigma}_{\sigma\zeta}\,\sigma \wedge \zeta + V^{\sigma}_{\sigma\overline{\zeta}}\,\sigma \wedge \overline{\zeta} + \rho \wedge \zeta + \rho \wedge \overline{\zeta}, \\ d\rho &= 2\gamma^1 \wedge \rho + V^{\rho}_{\sigma\rho}\,\sigma \wedge \rho + V^{\rho}_{\sigma\zeta}\,\sigma \wedge \zeta + V^{\rho}_{\sigma\overline{\zeta}}\,\sigma \wedge \overline{\zeta} \\ &+ V^{\rho}_{\rho\zeta}\,\rho \wedge \zeta + V^{\rho}_{\rho\overline{\zeta}}\,\rho \wedge \overline{\zeta} + i\,\zeta \wedge \overline{\zeta}, \end{split}$$

$$\begin{split} d\zeta &= \gamma^1 \wedge \zeta + \gamma^2 \wedge \sigma \\ &+ V^{\zeta}_{\sigma\rho} \, \sigma \wedge \rho + V^{\zeta}_{\sigma\zeta} \, \sigma \wedge \zeta + V^{\zeta}_{\sigma\overline{\zeta}} \, \sigma \wedge \overline{\zeta} + V^{\zeta}_{\rho\zeta} \, \rho \wedge \zeta \\ &+ V^{\zeta}_{\rho\overline{\zeta}} \, \rho \wedge \overline{\zeta} + V^{\zeta}_{\zeta\overline{\zeta}} \, \zeta \wedge \overline{\zeta}. \end{split}$$

$$\begin{split} d\overline{\zeta} &= \gamma^1 \wedge \zeta + \gamma^3 \wedge \sigma \\ &+ V^{\overline{\zeta}}_{\sigma\rho} \, \sigma \wedge \rho + V^{\overline{\zeta}}_{\sigma\zeta} \, \sigma \wedge \zeta + V^{\overline{\zeta}}_{\sigma\overline{\zeta}} \, \sigma \wedge \overline{\zeta} + V^{\overline{\zeta}}_{\rho\zeta} \, \rho \wedge \zeta \\ &+ V^{\overline{\zeta}}_{\rho\overline{\zeta}} \, \rho \wedge \overline{\zeta} + V^{\overline{\zeta}}_{\zeta\overline{\zeta}} \, \zeta \wedge \overline{\zeta}. \end{split}$$

It is straightforward to notice that $V^{\rho}_{\sigma\zeta}$ and $V^{\rho}_{\sigma\overline{\zeta}}$ are two essential torsion coefficients. The first one leads to the normalization:

$$\overline{\mathsf{d}}=\mathsf{a}\,\overline{\mathbf{D}}_0,$$

with

$$\overline{\mathbf{D}}_0 := \frac{i}{2} \frac{\mathscr{L}(B)^2}{B} + \frac{i}{3} Q \mathscr{L}(B) - \frac{i}{2} \mathscr{L}(\mathscr{L}(B)) - \frac{i}{2} B \mathscr{L}(Q) + \frac{i}{2} A \frac{\mathscr{L}(B)}{B} + \frac{i}{6} A Q + i B P,$$

while the second essential torsion coefficient gives the normalization:

$$d = a D_0$$

with:

$$\mathbf{D}_{0} := -\frac{2i}{3} \mathscr{L}(B) Q - \frac{i}{6} \frac{\mathscr{L}(B) A}{B} - \frac{i}{6} A Q + \frac{i}{6} \overline{\mathscr{L}(B)} Q - \frac{i}{3} \frac{\mathscr{L}(B)^{2}}{B} - \frac{i}{3} B Q^{2} - i \mathscr{L}(A) - \frac{i}{3} \overline{\mathscr{L}(B)} \mathscr{L}(B) + \frac{i}{2} \overline{\mathscr{L}(B)} + \frac{i}{2} \overline{\mathscr{L}(Q)} - i B P.$$

The coherency of the above formulae can be checked using the relations (1) and (2).

Let G_4 be the 1-dimensional Lie subgroup of G_3 whose elements g are of the form:

$$g := \begin{pmatrix} \mathsf{a}^3 & 0 & 0 & 0\\ 0 & \mathsf{a}^2 & 0 & 0\\ 0 & 0 & \mathsf{a} & 0\\ 0 & 0 & 0 & \mathsf{a} \end{pmatrix}, \qquad \mathsf{a} \in \mathbb{R} \setminus \{0\},$$

and let $\omega_3 := \left(\sigma_3, \rho_3, \zeta_3, \overline{\zeta}_3\right)$ be the coframe defined on M by:

$$\sigma_3 := \sigma_2, \qquad \qquad \rho_3 := \rho_2, \qquad \qquad \zeta_3 := \zeta_2 + \mathbf{D}_0 \, \sigma_2.$$

The normalization of d is equivalent to the reduction of P^3 to a subbundle P^4 consisting of those coframes ω on M such that:

$$\omega := g \cdot \omega_3$$
, where g is a function $g : M \xrightarrow{g} G_4$.

The Maurer-Cartan forms of G_4 are spanned by:

$$\alpha := \frac{d\mathsf{a}}{\mathsf{a}}.$$

Proceeding as in the previous steps, we compute the structure equations of P^4 :

$$\begin{split} d\sigma &= 3 \frac{d\mathbf{a}}{\mathbf{a}} \wedge \sigma \\ &+ W^{\sigma}_{\sigma\rho} \, \sigma \wedge \rho + W^{\sigma}_{\sigma\zeta} \, \sigma \wedge \zeta + W^{\sigma}_{\sigma\overline{\zeta}} \, \sigma \wedge \overline{\zeta} + \rho \wedge \zeta + \rho \wedge \overline{\zeta}, \\ d\rho &= 2 \frac{d\mathbf{a}}{\mathbf{a}} \wedge \rho + W^{\rho}_{\sigma\rho} \, \sigma \wedge \rho + W^{\rho}_{\rho\zeta} \, \rho \wedge \zeta + W^{\rho}_{\rho\overline{\zeta}} \, \rho \wedge \overline{\zeta} + i \, \zeta \wedge \overline{\zeta}, \\ d\zeta &= \frac{d\mathbf{a}}{\mathbf{a}} \wedge \zeta \\ &+ W^{\zeta}_{\sigma\rho} \, \sigma \wedge \rho + W^{\zeta}_{\sigma\zeta} \, \sigma \wedge \zeta + W^{\zeta}_{\sigma\overline{\zeta}} \, \sigma \wedge \overline{\zeta} + W^{\zeta}_{\rho\zeta} \, \rho \wedge \zeta \\ &+ W^{\zeta}_{\rho\overline{\zeta}} \, \rho \wedge \overline{\zeta} + W^{\zeta}_{\zeta\overline{\zeta}} \, \zeta \wedge \overline{\zeta}, \end{split}$$

$$\begin{split} d\overline{\zeta} &= \frac{d\mathbf{a}}{\mathbf{a}} \wedge \overline{\zeta} \\ &+ W^{\overline{\zeta}}_{\sigma\rho} \, \sigma \wedge \rho + W^{\overline{\zeta}}_{\sigma\zeta} \, \sigma \wedge \zeta + W^{\overline{\zeta}}_{\sigma\overline{\zeta}} \, \sigma \wedge \overline{\zeta} + W^{\overline{\zeta}}_{\rho\zeta} \, \rho \wedge \zeta \\ &+ W^{\overline{\zeta}}_{\rho\overline{\zeta}} \, \rho \wedge \overline{\zeta} + W^{\overline{\zeta}}_{\zeta\overline{\zeta}} \, \zeta \wedge \overline{\zeta}. \end{split}$$

Introducing the modified Maurer-Cartan form Λ :

$$\Lambda := \frac{d\mathbf{a}}{\mathbf{a}} + \frac{W^{\rho}_{\sigma\rho}}{2}\rho - \frac{W^{\sigma}_{\sigma\rho}}{3}\sigma - \frac{W^{\sigma}_{\sigma\rho}}{3}\zeta - \frac{W^{\sigma}_{\sigma\overline{\zeta}}}{3}\overline{\zeta},$$

these equations rewrite in the absorbed form as: (4)

$$\begin{split} d\sigma &= 3\Lambda \wedge \sigma + \rho \wedge \zeta + \rho \wedge \overline{\zeta}, \\ d\rho &= 2\Lambda \wedge \rho + i\,\zeta \wedge \overline{\zeta}, \\ d\zeta &= \Lambda \wedge \zeta + \frac{I_1}{a^4}\,\sigma \wedge \rho + \frac{I_2}{a^3}\,\sigma \wedge \zeta + \frac{I_3}{a^3}\,\sigma \wedge \overline{\zeta} + \frac{I_4}{a^2}\,\rho \wedge \zeta + \frac{I_5}{a^2}\,\rho \wedge \overline{\zeta}, \\ d\overline{\zeta} &= \Lambda \wedge \overline{\zeta} + \frac{\overline{I_1}}{a^4}\,\sigma \wedge \rho + \frac{\overline{I_3}}{a^3}\,\sigma \wedge \zeta + \frac{\overline{I_2}}{a^3}\,\sigma \wedge \overline{\zeta} + \frac{\overline{I_5}}{a^2}\,\rho \wedge \zeta + \frac{\overline{I_4}}{a^2}\,\rho \wedge \overline{\zeta}, \end{split}$$

where the invariants I_i , $i = 2 \dots 5$, are given by:

$$\begin{split} \overline{I_2} &= \frac{i}{8} \frac{\mathcal{QL}(B)^2}{B^{\frac{1}{2}}} - \frac{i}{8} B^{\frac{1}{2}} \mathcal{L}(B) Q^2 - \frac{3i}{4} \frac{\mathcal{L}(\mathcal{L}(B))\mathcal{L}(B)}{B^{\frac{1}{2}}} + \frac{i}{4} B^{\frac{1}{2}} \mathcal{L}(B) \mathcal{L}(Q) \\ &- \frac{i}{2} B^{\frac{1}{2}} P \mathcal{L}(B) - \frac{i}{4} B^{\frac{1}{2}} Q \mathcal{L}(\mathcal{L}(B)) - \frac{i}{4} B^{\frac{1}{2}} Q \mathcal{L}(\mathcal{L}(B)) \\ &- \frac{3i}{4} B^{\frac{3}{2}} Q \mathcal{L}(Q) + \frac{i}{2} B^{\frac{3}{2}} P Q + \frac{3i}{8} \frac{\mathcal{L}(B)^3}{B^{\frac{3}{2}}} + \frac{i}{8} B^{\frac{3}{2}} Q^3 \\ &+ \frac{i}{2} B^{\frac{3}{2}} \mathcal{L}(\mathcal{L}(Q)) + \frac{i}{2} B^{\frac{1}{2}} \mathcal{L}(\mathcal{L}(\mathcal{L}(B))) - i B^{\frac{3}{2}} \mathcal{L}(P) \end{split}$$

$$\begin{split} I_{3} &= -\mathbf{D}_{0}\mathbf{C}_{0} + \frac{\mathscr{L}(B)}{B^{\frac{1}{2}}} \mathbf{D}_{0} + B^{\frac{1}{2}}Q\mathbf{D}_{0} + \frac{A}{B^{\frac{1}{2}}}\mathbf{D}_{0} - \frac{\overline{\mathscr{L}}(\mathbf{D}_{0})}{B^{\frac{1}{2}}} - i\,\mathbf{B}_{0}\mathbf{D}_{0} + i\,\mathbf{B}_{0}^{2}\mathbf{C}_{0} \\ &- \frac{A}{B^{\frac{1}{2}}}\mathbf{B}_{0}\mathbf{C}_{0} + \mathbf{B}_{0}\,\mathscr{L}(A) + BP\,\mathbf{B}_{0} + \frac{\overline{\mathscr{L}}(\mathbf{B}_{0})}{B^{\frac{1}{2}}}\,\mathbf{C}_{0} + \frac{1}{2}\,\frac{\overline{\mathscr{L}}(B)}{B^{\frac{3}{2}}}\,\mathbf{B}_{0}\mathbf{C}_{0} \\ &\overline{I_{4}} = \frac{3}{4}\,i\,\frac{\mathscr{L}(B)^{2}}{B} + \frac{1}{6}\,i\,\mathscr{L}(B)\,Q + \frac{11}{36}\,i\,B\,Q^{2} - i\,\mathscr{L}(\mathscr{L}(B)) - \frac{2}{3}\,i\,B\,\mathscr{L}(Q) + i\,BP, \end{split}$$

$$\overline{I_5} = \frac{i}{3} \mathscr{L}(A) + \frac{i}{3} \overline{\mathscr{L}}(Q) - i \frac{\overline{\mathscr{L}}(\mathscr{L}(B))}{B} + \frac{5}{12} i \frac{\mathscr{L}(B)^2}{B} - \frac{i}{3} B\mathscr{L}(Q) + \frac{11}{36} i B Q^2 + i B P + \frac{2}{3} i \frac{\mathscr{L}(\overline{\mathscr{L}}(B))}{B} - \frac{i}{3} \mathscr{L}(\mathscr{L}(B)) + \frac{i}{3} \frac{A \mathscr{L}(B)}{B} + \frac{7}{18} i \mathscr{L}(B) Q - \frac{i}{9} \frac{\overline{\mathscr{L}}(B) Q}{B} + \frac{i}{9} A Q,$$

and I_1 is given by:

$$I_1 = \frac{2i}{3} (I_3)_{\zeta} - \frac{2i}{3} (I_2)_{\overline{\zeta}}$$

The exterior derivative of Λ can be determined by taking the exterior derivative of the four equations (4), which leads to the so-called Bianchi-Cartan's identities. For example, taking the exterior derivative of the first

equation of (4), one gets:

$$0 = \left[3 d\Lambda + \left(\frac{I_2}{\mathsf{a}^3} + \frac{\overline{I_3}}{\mathsf{a}^3}\right) \rho \wedge \zeta + \left(\frac{\overline{I_2}}{\mathsf{a}^3} + \frac{I_3}{\mathsf{a}^3}\right) \rho \wedge \zeta\right] \wedge \sigma_{\gamma}$$

while taking the exterior derivative of the second equation gives:

$$0 = \left[2 d\Lambda - i \frac{I_1}{\mathsf{a}^4} \sigma \wedge \overline{\zeta} + i \frac{\overline{I_1}}{\mathsf{a}^4} \sigma \wedge \zeta\right] \wedge \rho.$$

Eventually we get:

(5)

$$d\Lambda = \frac{i}{2} \frac{I_1}{\mathsf{a}^4} \, \sigma \wedge \overline{\zeta} - \frac{i}{2} \frac{\overline{I_1}}{\mathsf{a}^4} \, \sigma \wedge \zeta - \frac{1}{3} \left(\frac{I_2}{\mathsf{a}^3} + \frac{\overline{I_3}}{\mathsf{a}^3} \right) \, \rho \wedge \zeta - \frac{1}{3} \left(\frac{\overline{I_2}}{\mathsf{a}^3} + \frac{I_3}{\mathsf{a}^3} \right) \, \rho \wedge \zeta + \frac{I_0}{\mathsf{a}^4} \, \sigma \wedge \zeta \;,$$

where I_0 is given by:

$$I_0 := -\frac{1}{2\mathsf{a}^4} \left(I_1 \right)_{\zeta} - \frac{1}{2\mathsf{a}^4} \left(\overline{I_1} \right)_{\overline{\zeta}}.$$

4. CARTAN CONNECTION

We recall that the model for CR-manifolds belonging to general class II is Beloshapka's cubic $B \subset \mathbb{C}^3$, which is defined by the equations:

B:

$$w_1 = \overline{w_1} + 2 i z \overline{z},$$

$$w_2 = \overline{w_2} + 2 i z \overline{z} (z + \overline{z}).$$

Its Lie algebra of infinitesimal CR-automorphisms is given by the following theorem:

Theorem 2. [12]. Beloshapka's cubic,

B:

$$w_1 = \overline{w_1} + 2 i z \overline{z},$$

 $w_2 = \overline{w_2} + 2 i z \overline{z} (z + \overline{z}),$

has a 5-dimensional Lie algebra of CR-automorphisms $\operatorname{aut}_{CR}(B)$. A basis for the Maurer-Cartan forms of $\operatorname{aut}_{CR}(B)$ is provided by the 5 differential 1-forms σ , ρ , ζ , $\overline{\zeta}$, α , which satisfy the structure equations:

$$d\sigma = 3 \alpha \wedge \sigma + \rho \wedge \zeta + \rho \wedge \zeta ,$$

$$d\rho = 2 \alpha \wedge \rho + i \zeta \wedge \overline{\zeta} ,$$

$$d\zeta = \alpha \wedge \zeta ,$$

$$d\overline{\zeta} = \alpha \wedge \overline{\zeta} ,$$

$$d\alpha = 0.$$

SAMUEL POCCHIOLA

Let us write \mathfrak{g} instead of $\operatorname{aut}_{CR}(B)$ for the Lie algebra of inifinitesimal automorphisms of Beloshapka's cubic, and let $(e_{\alpha}, e_{\sigma}, e_{\rho}, e_{\zeta}, e_{\overline{\zeta}})$ be the dual basis of the basis of Maurer-Cartan 1-forms: $(\alpha, \sigma, \rho, \zeta, \overline{\zeta})$ of \mathfrak{g} . From the above structure equations, the Lie brackets structure of \mathfrak{g} is given by:

$$\begin{split} [e_{\alpha}, e_{\sigma}] &= -3 e_{\sigma}, & [e_{\alpha}, e_{\rho}] &= -2 e_{\rho}, & [e_{\alpha}, e_{\zeta}] &= -e_{\zeta}, \\ [e_{\alpha}, e_{\overline{\zeta}}] &= -e_{\overline{\zeta}}, & [e_{\rho}, e_{\zeta}] &= -e_{\sigma}, & [e_{\rho}, e_{\overline{\zeta}}] &= -e_{\sigma}, \\ [e_{\zeta}, e_{\overline{\zeta}}] &= -i e_{\rho}, \end{split}$$

the remaining brackets being equal to zero.

We refer to [5], p. 127-128, for the definition of a Cartan connection. Let $\mathfrak{g}_0 \subset \mathfrak{g}$ be the subalgebra spanned by e_α , \mathfrak{G} the connected, simply connected Lie group whose Lie algebra is \mathfrak{g} and \mathfrak{G}_0 the connected closed 1-dimensional subgroup of \mathfrak{G} generated by \mathfrak{g}_0 . We notice that $\mathfrak{G}_0 \cong G_4$, so that P^4 is a principal bundle over M with structure group \mathfrak{G}_0 , and that $\dim \mathfrak{G}/\mathfrak{G}_0 = \dim M = 4$.

Let $(\Lambda, \sigma, \rho, \zeta, \overline{\zeta})$ be the coframe of 1-forms on P^4 whose structure equation are given by (4) – (5) and ω the 1-form on P with values in \mathfrak{g} defined by:

$$\omega(X) := \Lambda(X) e_{\alpha} + \sigma(X) e_{\sigma} + \rho(X) e_{\rho} + \zeta(X) e_{\zeta}, +\overline{\zeta}(X) e_{\overline{\zeta}},$$

for $X \in T_p P^4$. We have:

Theorem 3. ω is a Cartan connection on P^4 .

Proof. We shall check that the following three conditions hold:

- (1) $\omega(e_{\alpha}^{*}) = e_{\alpha}$, where e_{α}^{*} is the vertical vector field on P^{4} generated by the action of e_{α} ,
- (2) $R_a^* \omega = \operatorname{Ad}(a^{-1}) \omega$ for every $a \in \mathfrak{G}_0$,
- (3) for each $p \in P^4$, ω_p is an isomorphism $T_p P^4 \xrightarrow{\omega_p} \mathfrak{g}$.

Condition (3) is trivially satisfied as $(\Lambda, \sigma, \rho, \zeta, \overline{\zeta})$ is a coframe on P^4 and thus defines a basis of $T_p^* P^4$ at each point p.

Condition (1) follows simply from the fact that Λ is a modified-Maurer Cartan form on P^4 :

$$\Lambda = \frac{d\mathbf{a}}{\mathbf{a}} + \frac{W^{\rho}_{\sigma\rho}}{2}\rho - \frac{W^{\sigma}_{\sigma\rho}}{3}\sigma - \frac{W^{\sigma}_{\sigma\rho}}{3}\zeta - \frac{W^{\sigma}_{\sigma\overline{\zeta}}}{3}\overline{\zeta},$$

so that

$$\omega(e_{\alpha}^*) = \Lambda(e_{\alpha^*}) = e_{\alpha},$$

as

$$\sigma(e_{\alpha}^{*}) = \rho(e_{\alpha}^{*}) = \zeta(e_{\alpha}^{*}) = \overline{\zeta}(e_{\alpha}^{*}) = 0, \qquad \frac{d\mathsf{a}}{\mathsf{a}}(e_{\alpha}^{*}) = 1,$$

since e^*_{α} is a vertical vector field on P^4 .

Condition (2) is equivalent to its infinitesimal counterpart:

$$\mathscr{L}_{e^*_{\alpha}} \omega = -\mathsf{ad}_{e_{\alpha}} \omega,$$

where $\mathscr{L}_{e_{\alpha}^{*}} \omega$ is the Lie derivative of ω by the vector field e_{α}^{*} and where $\mathrm{ad}_{e_{\alpha}}$ is the linear map $\mathfrak{g} \to \mathfrak{g}$ defined by: $\mathrm{ad}_{e_{\alpha}}(X) = [e_{\alpha}, X]$. We determine $\mathscr{L}_{e_{\alpha}^{*}} \omega$ with the help of Cartan's formula:

$$\mathscr{L}_{e^*_{\alpha}}\omega = e_{\alpha^*} \,\lrcorner\, d\omega + d\left(e^*_{\alpha} \,\lrcorner\, \omega\right),$$

with

$$d\left(e_{\alpha}^{*} \,\lrcorner\, \omega\right) = 0$$

from condition (1). The structure equations (4)–(5) give:

$$e_{\alpha^*} \,\lrcorner\, d\omega = \begin{pmatrix} 0\\ 3 \,\sigma\\ 2 \,\rho\\ \frac{\zeta}{\zeta} \end{pmatrix},$$

which is easily seen being equal to $-ad_{e_{\alpha}}\omega$ from the Lie bracket structure of g.

From theorem 3, the structure equations (4) and (5), and the fact that the invariants I_0 and I_1 are expressed in terms of I_2 , I_3 , I_4 , I_5 , we have:

Theorem 4. A CR-manifold M belonging to general class || is locally biholomorphic to Beloshapka's cubic $B \subset \mathbb{C}^3$ if and only if the condition

$$I_2 \equiv I_3 \equiv I_4 \equiv I_5 \equiv 0$$

holds locally on M.

SAMUEL POCCHIOLA

REFERENCES

- Beloshapka, V.K.; Ezhov, V.; Schmalz, G.: Canonical Cartan connection and holomorphic invariants on Engel CR manifolds, Russian J. Mathematical Physics 14 (2007), no. 2, 121–133.
- [2] Cartan, E.: Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes, Ann. Mat. Pura Appl. **11** (1933), no. 1, 17–90.
- [3] Cartan, E.: Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes II, Ann. Scuola. Norm. Sup. Pisa 1 (1932), no. 4, 333–354.
- [4] Ezhov, V.; McLaughlin, B.; Schmalz, G.: *From Cartan to Tanaka: Getting Real in the Complex World*, Notices of the AMS **58** (2011), no. 1.
- [5] Kobayashi, S.: Transformation groups in differential geometry, Ergebnisse der Mathematik und ihrer Grenzgebiete, 70, Springer-Verlag, New-York Heidelberg Berlin, 1972.
- [6] Merker, J.: Equivalences of 5-dimensional CR manifolds, III: Six models and (very) elementary normalizations, 54 pages, arxiv.org/abs/1311.7522.
- [7] Merker, J.: Equivalences of 5-dimensional CR-manifolds, IV: Six ambiguity matrix groups, (Initial G-structures), 34 pages, arxiv.org/abs/1312.1084
- [8] Merker, J.: Equivalences of 5-dimensional CR-manifolds V: Six initial frames and coframes; Explicitness obstacles, 63 pages, arxiv.org/abs/1312.3581
- [9] Merker, J.; Pocchiola, S.; Sabzevari, M.: Equivalences of 5-dimensional CRmanifolds, II: General classes I, II, III-1, III-2, IV-1, IV-2, arxiv.org/abs/1311.5669.
- [10] Olver, P.J.: *Equivalence, Invariance and Symmetries*. Cambridge University Press, Cambridge, 1995, xvi+525 pp.
- [11] Pocchiola, S.: Absolute parallelism for 2-nondegenerate real hypersurfaces $M^5 \subset \mathbb{C}^3$ of constant Levi rank 1, arxiv.org/abs/1312.6400, 56 pp.
- [12] Pocchiola, S.: Lie algebras of infinitesimal CR-automorphisms for the model manifolds of general classes II, III₂, IV₂, 44 pp.
- [13] Pocchiola, S.: Canonical Cartan connection for 5-dimensional CR-manifolds belonging to general class III₂, 19 pp.
- [14] Poincaré, H.: Les fonctions analytiques de deux variables et la représentation conforme, *Rend. Circ. Math. Palermo*, 23, (1907), 185-220.
- [15] Sternberg, S.: Lectures on Differential Geometry. Prentice-Hall mathematical series, Inc., Englewood Cliffs, N.J. 1964, xv+390 pp.

SAMUEL POCCHIOLA — DÉPARTEMENT DE MATHÉMATIQUES, BÂTIMENT 425, FACULTÉ DES SCIENCES D'ORSAY, UNIVERSITÉ PARIS-SUD, F-91405 ORSAY CEDEX, FRANCE

E-mail address: samuel.pocchiola@math.u-psud.fr