Analytic Hilbert fields and hermitian Hilbert bundles

Dat Tran

2014

1 Introduction

This paper was directly motivated by the paper [LS14] of Lempert and Szőke. In [LS14], the authors defined the notion of a smooth Hilbert field. Hilbert field was introduced by Godement in [God51]. It also appeared in [Dix69] with a continuous structure attached to it. The smooth Hilbert field can be thought of as a generalization of a smooth hermitian Hilbert bundle with a hermitian connection. Smooth Hilbert fields has a curvature, just like a Hilbert bundle endowed with a connection on it. In [LS14], Lempert and Szőke defined what it means to be an analytic Hilbert field and proved that any analytic Hilbert field with curvature zero induces a Hilbert bundle. They also provided an example of a Hilbert field with curvature zero, yet does not induce any Hilbert bundle. In this paper, we will give an example of an analytic Hilbert field which does not come from any Hilbert bundle. This example occured as a direct image of a bundle in the dissertation of the author, but in this paper, it will be presented in a more general form.

For the sake of completeness, we first provide the definitions of a Hilbert bundle, Hilbert field, smooth and analytic Hilbert field. Most of these definitions are also in [LS14].

Definition 1.1. A Banach manifold is a Hausdorf space M with an open cover \mathcal{U} such that for every $U \in \mathcal{U}$, we are given a Banach space B_U and a homeomorphism φ_U between U and an open subset V_U of B_U . These pairs (U, φ_U) are called coordinate charts and it is required that for every two charts $(U, \varphi_U), (U', \varphi_{U'})$, the composition $\varphi_{U'} \circ \varphi_U^{-1}$ should be smooth wherever it is defined.

Definition 1.2. Let M and N be two Banach manifolds. A map $f : M \to N$ is smooth if for any pair of charts (U, φ_U) , (V, ψ_V) on M, resp. N, the map $\psi_V \circ f \circ \varphi_U^{-1}$ is smooth where it is defined.

Definition 1.3. A smooth Hilbert bundle is a smooth map $\pi : H \to S$ of Banach manifolds, each fiber $\pi^{-1}s$ is endowed with the structure of a complex vector space; for each $s \in S$, there should exist an open neighborhood $U \subset S$, a complex Hilbert space X_U and a smooth map $F_U : \pi^{-1}U \to X_U$ whose restriction to each fiber $\pi^{-1}s$, $s \in U$ is linear and $\pi \times F :$ $\pi^{-1}U \to U \times X_U$ is diffeomorphic.

We use $\Gamma^{\infty}(S, H)$ (or $\Gamma^{\infty}(H)$ when there is no ambiguity about the base) to denote the set of smooth sections of the smooth Hilbert bundle $H \to S$.

Definition 1.4. A smooth Hermitian metric on a smooth Hilbert bundle $\pi : H \to S$ is a function $h : H \oplus H \to \mathbb{C}$ such that any local trivialization $F_U : \pi^{-1}U \to X_U$ as in the previous definition can be chosen so that $h(u, v) = \langle F(u), F(v) \rangle$ where u, v are in the same fiber and \langle , \rangle is the inner product of X_U . We call the pair $(H, h) \to S$ a hermitian Hilbert bundle.

Let S be a smooth manifold, we use VectS to denote the set of smooth complex valued vector fields on S. If $(H, h) \to S$ is a smooth hermitan Hilbert bundle and $\varphi, \psi \in \Gamma^{\infty}(H)$, then $h(\varphi, \psi) \in C^{\infty}(S)$.

Definition 1.5. A connection ∇ on a Hilbert bundle $H \to S$ associates with every $\xi \in \text{Vect}S$ a linear map $\nabla_{\xi} : \Gamma^{\infty}(H) \to \Gamma^{\infty}(H)$. It is required that for every local trivialization $F_U : \pi^{-1}U \to U \times X_U$, there be a smooth map $A : \mathbb{C} \otimes TU \to U \times \text{End}X_U$, mapping fiber to fiber, linear on each fiber $\mathbb{C} \otimes T_s U$ such that on U,

 $F_U(\nabla_{\xi}\varphi) = \xi F_U(\varphi) + A(\xi)F_U(\varphi)$

for every $\varphi \in \Gamma^{\infty}(S, H)$.

If $(H,h) \to S$ is a hermitian Hilbert bundle and ∇ is a connection on H, then ∇ is a hermitian connection if for every $\xi \in \text{Vect}S$, and for every smooth section φ, ψ of H, we have $\xi h(\varphi, \psi) = h(\nabla_{\xi}\varphi, \psi) + h(\varphi, \nabla_{\bar{\xi}}\psi)$.

Now, we go over the definitions of Hibert fields.

Definition 1.6. A Hilbert field is a map between sets $p : H \to S$ such that for all $s \in S$, there is a Hilbert space structure endowed on $H_s = p^{-1}s$.

Definition 1.7 (Smooth Hilbert field). Let S be a smooth manifold, and $H \to S$ a Hilbert field. A smooth structure on H is given by specifying a set Γ^{∞} of sections of H, closed under addition and under multiplication by elements of $C^{\infty}(S)$, and linear operators $\nabla_{\xi} : \Gamma^{\infty} \to$ Γ^{∞} for each $\xi \in \text{Vect}S$ such that for $\xi, \eta \in \text{Vect}S$, $f \in C^{\infty}(S)$, $\varphi, \psi \in \Gamma^{\infty}$, the following conditions are satisfied.

1. $\nabla_{\xi+\eta}\varphi = \nabla_{\xi}\varphi + \nabla_{\eta}\varphi; \ \nabla_{(f\xi)}\varphi = f\nabla_{\xi}\varphi; \ \nabla_{\xi}(f\varphi) = (\xi f)\varphi + f\nabla_{\xi}\varphi.$

2. $h(\varphi, \psi) \in C^{\infty}(S, \mathbb{C})$ and $\xi h(\varphi, \psi) = h(\nabla_{\xi} \varphi, \psi) + h(\varphi, \nabla_{\bar{\xi}} \psi)$.

3. The set $\{\varphi(s): \varphi \in \Gamma^{\infty}\}$ is dense in H_s for all $s \in S$.

Definition 1.8. Let $H \to S$ be a Hilbert field with a smooth structure $(\Gamma^{\infty}, \nabla)$, and $(\hat{H}, \hat{h}) \to S$ be a hermitian Hilbert bundle with a hermitian connection $\tilde{\nabla}$. If there is a fiber preserving map $F : H \to \tilde{H}$ such that $F|H_s$ are isometric for all $s, F\varphi \in \Gamma^{\infty}(S, \tilde{H}) \quad \forall \varphi \in \Gamma^{\infty}$, and $\tilde{\nabla}_{\xi}(F\varphi) = F(\nabla_{\xi}\varphi), \quad \forall \varphi \in \Gamma^{\infty}, \forall \xi \in \text{Vect } S, \text{ we then say that } H \to S \text{ induces from the hermitian Hilbert bundle } (\tilde{H}, \tilde{h}) \to S.$

Let $H \to S$ be a Hilbert field with smooth structure $(\Gamma^{\infty}, \nabla)$. For every $\xi, \eta \in \text{Vect}S$, $\varphi \in \Gamma^{\infty}$, we set

$$R(\xi,\eta)\varphi := \nabla_{\xi}\nabla_{\eta}\varphi - \nabla_{\eta}\nabla_{\xi}\varphi - \nabla_{[\xi,\eta]}\varphi.$$

and call the operator R the curvature of this smooth Hilbert field. It was proved in [LS14], that $R(\xi, \eta)\varphi(s)$ only depends on $\xi(s), \eta(s)$, and $\varphi(s)$. Therefore, for each ξ, η in $T_s S \otimes \mathbb{C}$, $R(\xi, \eta)$ is a densely defined operator on H_s , which we also denote $R(\xi, \eta)$.

If $(H, h) \to S$ is a hermitian Hilbert bundle, then the curvature of the bundle is defined exactly as the definition of the curvature of smooth Hilbert field above. Yet, it is known that for every $\xi, \eta \in \text{Vect}S$, the operator $R(\xi, \eta)$ is a bounded operator on each fiber. **Definition 1.9** (Analytic section of smooth Hilbert field with analytic base and analytic Hilbert field). Let S be a finite dimensional analytic manifold and $H \to S$ a smooth Hilbert field with smooth structure $(\Gamma^{\infty}, \nabla)$.

1. A section $\varphi \in \Gamma^{\infty}$ is said to be analytic if for any compact subset C of S and any finite subset Ξ of Vect S such that every $\xi \in \Xi$ is analytic in a neighborhood of C, there is an $\epsilon > 0$ such that

$$\sup \frac{\epsilon^m}{m!} h(\nabla_{\xi_m} \dots \nabla_{\xi_1} \varphi)^{1/2}(s) < \infty$$
(1)

where the sup is taken over $m = 0, 1, \ldots; \xi_j \in \Xi$; and $s \in C$.

- 2. The set of analytic sections of S is denoted Γ^{ω} .
- 3. If the set $\{\varphi(s) : \varphi \in \Gamma^{\omega}\}$ is dense in H_s for all $s \in S$, then this Hilbert field is said to be analytic.

The method of defining analytic sections like above can also be applied in defining realanalytic functions. For examples of this, please see [KP02].

2 Example of an anlytic Hilbert field not coming from any bundle

Let $S = \mathbb{C}$ and $H \to S$ be a Hilbert field such that every fiber H_s is infinite dimensional and separable. Define φ_j , $j = 0, 1, \ldots$ be sections of this Hilbert field such that for each s, $\{\varphi_i(s)\}$ form an orthonormal basis on H_s . Let

$$\Gamma^{\infty} = \left\{ \sum_{l=0}^{k} a_l(s) \varphi_l | k < \infty, a_l \in C^{\infty}(S) \right\}.$$

Since the set of linear combinations of an orthonormal basis will be dense in any Hilbert space, the set $\{\varphi(s)|\varphi\in\Gamma^{\infty}\}$ is dense in H_s .

Let k be a complex-valued, real-analytic function on S. Let $\xi = \frac{\partial}{\partial s}$ and define:

$$\nabla_{\xi}\varphi_j = (j+1)k\varphi_j$$
 and $\nabla_{\bar{\xi}}\varphi_j = -(j+1)\bar{k}\varphi_j$.

Since $\{\xi(s)\}$ span $T_s^{(1,0)}S$ for all s, we can extend by linearity and get $\nabla_{\eta} : \Gamma^{\infty} \to \Gamma^{\infty}$ as required by the definition of smooth structure for all $\eta \in \text{Vect}S$. So, $(\Gamma^{\infty}, \nabla)$ is a smooth structure of the Hilbert field $H \to S$.

Lemma 2.1. All the sections φ_j are analytic and consequently, the Hilbert field $H \rightarrow S$ constructed above is analytic. Moreover, if g is a real-valued, analytic, non-harmonic function, and $k = \frac{\partial g}{\partial s}$ then this Hilbert field does not induce any hermitian Hilbert bundle.

Proof. The goal here is to show that φ_j are analytic, i.e. for each fixed j, for every compact set $K \subset S$, there is a $\delta > 0$ such that

$$\sup \frac{\delta^m}{m!} h(\nabla_{\eta_1} \dots \nabla_{\eta_m} \varphi_j)^{1/2}(s) < \infty,$$

where the sup is taken over all $m = 0, 1, ..., \eta_1, ..., \eta_m \in \left\{\frac{\partial}{\partial s}, \frac{\partial}{\partial \bar{s}}\right\}$, and all $s \in K$. It was shown in Corollary 3.3.4 of [LS14] that we only need to prove this for $\eta_1, ..., \eta_m \in \left\{\frac{\partial}{\partial s}, \frac{\partial}{\partial \bar{s}}\right\}$, instead of for $\eta_1, ..., \eta_m$ in any possible predetermined finite subset of Vect^{ω} S.

Let f be a smooth function on S, for $\eta \in \left\{\frac{\partial}{\partial s}, \frac{\partial}{\partial \bar{s}}\right\}$, we have

$$\nabla_{\eta}(f\varphi_j) = (\eta f)\varphi_j + a(\eta)f\varphi_j$$

where

$$a\left(\frac{\partial}{\partial s}\right) = (j+1)k$$
 and $a\left(\frac{\partial}{\partial \bar{s}}\right) = -(j+1)\bar{k}$

Fix $K \subset S$, a nonnegative integer j, and an analytic function f defined in a neighborhood of K. Since f, g are all analytic in a neighborhood of K, there is an $\epsilon \in (0, 1)$ and an M > 1such that

$$\sup \frac{\epsilon^m}{m!} |\eta_1 \dots \eta_m h(s)| < M \tag{2}$$

where the sup is taken over all $m = 0, 1, ...; \eta_1, ..., \eta_m \in \left\{\frac{\partial}{\partial s}, \frac{\partial}{\partial \bar{s}}\right\};$ $h \in \{f, (j+1)k, -(j+1)\bar{k}\}; \text{ and all } s \in K.$ (This is a result in [KP02])

Before the next step of the proof, we want to introduce some notations.

- 1. If $I \subset \{1, \ldots, m\}$ for $m \ge 1$ and $I = \{i_1, \ldots, i_l\}$ where $l \le m$ and $i_1 > \ldots > i_l$, then $\eta_I = \eta_{i_1} \ldots \eta_{i_l}$. We also write $\eta_{\emptyset} = id$.
- 2. $a_i = a(\eta_i)$.
- 3. A collection of I_1, \ldots, I_l is a sub-splitting of $\{1, \ldots, m\}$ if $I_{\alpha} \cap I_{\beta} = \emptyset$ for $\alpha \neq \beta$ and $\bigcup_{\alpha} I_{\alpha} \subseteq \{1, \ldots, m\}$. I_{α} can be empty set.
- 4. A k-splitting of m consists of a sub-splitting I_1, \ldots, I_k and $i_1 > \ldots > i_{k-1}$ of $\{1, \ldots, m\}$ such that every element of I_{α} is larger than i_{α} for $\alpha < k$ and $\bigcup_{\alpha} I_{\alpha} \cup \{i_1, \ldots, i_{k-1}\} =$ $\{1, \ldots, m\}$. Every m has k-splittings for $k = 1, \ldots, m+1$. For m = 0, the empty set is the unique 1-splitting of m.

Let m be a non-negative integer. For k = 1, ..., m + 2, a k-splitting of (m + 1) is of the following two types.

1. A k-splitting $(I_1, \ldots, I_k; i_1, \ldots, i_{k-1})$ is of type 1 if $\bigcup_{\alpha} I_{\alpha} \subset \{1, \ldots, m\}$. This means $i_1 = m + 1$ due to $i_1 > \ldots > i_{k-1}$. Because any element of I_1 is larger than $i_1, I_1 = \emptyset$. Suppose k = 1, then $\bigcup_{\alpha} I_{\alpha} \cup \{i_1, \ldots, i_{k-1}\} = \emptyset \cup \emptyset \neq \{1, \ldots, m+1\}$

and we have a contradiction. So, no 1-splitting is of type 1. Hence, any k-splitting $(I_1, \ldots, I_k; i_1, \ldots, i_{k-1})$ of m+1 of type 1 corresponds to a (k-1)-splitting of m which is $(I_2, \ldots, I_k; i_2, \ldots, i_{k-1})$. Vice versa, for any (k-1)-splitting of m, we have a unique type 1 k-splitting of m+1 by adjoining $I_1 = \emptyset$ and $i_1 = m+1$ to it. So there is a 1-1 correspondence between type 1 k-splitting of m+1 and (k-1)-splitting of m.

2. Type 2 k-splittings are those that are not type 1. It means that if $(I_1, \ldots, I_k; i_1, \ldots, i_{k-1})$ is type 2, then $m + 1 \in \bigcup_{\alpha} I_{\alpha}$. And so, there must be an α such that I_{α} contains m + 1. If we replace this I_{α} by $\tilde{I}_{\alpha} = I_{\alpha} \setminus \{m + 1\}$ we get a k-splitting of m. Suppose k = m + 2, then by counting, $\{i_1, \ldots, i_{k-1}\} = \{1, \ldots, m + 1\}$ and so there cannot be any I_{α} that contain m + 1. Hence, there is no type 2 (m + 2)-splitting. From these facts, we can see that for each k in $\{1, \ldots, m + 1\}$ there is a 1 - k correspondence between the k-splitting of m and the k-splitting of m + 1 of type 2 as follow: for each k-splitting $(I_1, \ldots, I_k; i_1, \ldots, i_{k-1})$ of m, we acquire k different k-splittings of m + 1 by inserting m + 1 to the begining of each I_{α} .

Let
$$\eta_1, \ldots, \eta_m \in \left\{\frac{\partial}{\partial s}, \frac{\partial}{\partial \bar{s}}\right\}$$
. For any f as above, we define $T^m \in C^\infty(S)$. We define

$$T^m = \sum_{k=1}^{m+1} \sum_{k=1}^{m+1} (\eta_{I_1} a_{i_1}) \ldots (\eta_{I_{k-1}} a_{i_{k-1}}) \eta_{I_k} f$$

where the inner sum is taken over all distinct k-splittings,

$$S_1^m = \sum_{k=2}^{m+1} \sum (\eta_{I_1} a_{i_1}) \dots (\eta_{I_{k-1}} a_{i_{k-1}}) \eta_{I_k} f$$

where the inner sum is taken over all distinct type 1 k-splittings, and

$$S_2^m = \sum_{k=1}^m \sum (\eta_{I_1} a_{i_1}) \dots (\eta_{I_{k-1}} a_{i_{k-1}}) \eta_{I_k} f$$

where the inner sum is taken over all distinct type 2 k-splittings. Clearly,

$$T^m = S_1^m + S_2^m. (3)$$

We have

$$S_1^{m+1} = \sum_{k=2}^{m+2} \sum (\eta_{I_1} a_{i_1}) \dots (\eta_{I_{k-1}} a_{i_{k-1}}) \eta_{I_k} f = a_{m+1} T^m$$
(4)

as explained when we described the first collection. We also have

$$S_2^{m+1} = \sum_{k=1}^{m+1} \sum (\eta_{I_1} a_{i_1}) \dots (\eta_{I_{k-1}} a_{i_{k-1}}) \eta_{I_k} f$$
(5)

where we can group terms in the inner sum into distinct groups of k-splitting of m+1, each group consists of elements correspond to the same k-splitting of m. For each k-splitting

 $(I_1, \ldots, I_k; i_1, \ldots, i_{k-1})$ of m, $\eta_{m+1}[(\eta_{I_1}a_{i_1}) \ldots (\eta_{I_{k-1}}a_{i_{k-1}})\eta_{I_k}f]$ gives us the sum of all the k-splittings of m + 1 corresponds to it. So

$$S_2^{m+1} = \sum_{k=1}^{m+1} \sum (\eta_{I_1} a_{i_1}) \dots (\eta_{I_{k-1}} a_{i_{k-1}}) \eta_{I_k} f$$
(6)

$$=\sum_{k=1}^{m+1}\sum \eta_{m+1}[(\eta_{I_1}a_{i_1})\dots(\eta_{I_{k-1}}a_{i_{k-1}})\eta_{I_k}f]$$
(7)

$$=\eta_{m+1}\left[\sum_{k=1}^{m+1}\sum(\eta_{I_1}a_{i_1})\dots(\eta_{I_{k-1}}a_{i_{k-1}})\eta_{I_k}f\right]$$
(8)

$$=\eta_{m+1}T^m \tag{9}$$

where the inner sums of (3.5) and (3.6) are taken over all possible k-splitting of m.

Claim: $\nabla_{\eta_m} \dots \nabla_{\eta_1} (f\varphi_j) = T^m \varphi_j.$

We prove the above claim by induction.

For m = 1, $T^1 = \eta_1 f + a_1 f$ and $\nabla_{\eta_1} \varphi_j = (\eta_1 f) \varphi_j + (a_1 f) \varphi_j$. Clearly, the claim is true. Suppose the claim is true for m. We have

$$\nabla_{\eta_{m+1}} \dots \nabla_{\eta_1} (f\varphi_j) = \nabla_{\eta_{m+1}} (\nabla_{\eta_m} \dots \nabla_{\eta_1} (f\varphi_j)) = \nabla_{\eta_{m+1}} (T^m \varphi_j)$$

= $\eta_{m+1} (T^m) \varphi_j + a_{m+1} T^m \varphi_j = S_2^{m+1} \varphi_j + S_1^{m+1} \varphi_j = T^{m+1} \varphi_j$

Therefore, we have

$$\nabla_{\eta_m} \dots \nabla_{\eta_1} (f\varphi_j) = \left[\sum_{k=1}^{m+1} \sum (\eta_{I_1} a_{i_1}) \dots (\eta_{I_{k-1}} a_{i_{k-1}}) \eta_{I_k} f\right] \varphi_j,$$
(10)

where the inner sum is taken over all of the k-splittings of m.

Fix a k in $\{1, \ldots, m+1\}$ and pick any set of positive integers l_1, \ldots, l_k such that $l_1 + \ldots + l_k = m + 1$. We call a summand in (10) with $|I_1| = l_1 - 1, \ldots, |I_{k-1}| = l_{k-1} - 1, |I_k| = l_k$ a term of type (l_1, \ldots, l_k) . From equation (2), each of the terms of type (l_1, \ldots, l_k) is bounded above by

$$\frac{M^k}{\epsilon^{m+1-k}}(l_1-1)!\dots(l_k-1)!.$$

The number of terms of type (l_1, \ldots, l_k) is at most $\binom{m+1}{l_1; \ldots; l_k}$. So the sum of all the terms of type (l_1, \ldots, l_k) is bounded above by

$$\binom{m+1}{l_1;\ldots;l_k} \frac{M^k}{\epsilon^{m+1-k}} (l_1-1)!\ldots(l_k-1)!.$$

For each fixed k, the number of choices of (l_1, \ldots, l_k) is $\binom{m}{k-1}$. And hence we can bound

both sides of equation (10) for any $s \in K$ as below

$$h(\nabla_{\eta_1} \dots \nabla_{\eta_m} \varphi_j)^{1/2}(s)$$

$$= \sum_{k=1}^{m+1} \binom{m}{k-1} \binom{m+1}{l_1; \dots; l_k} \frac{M^k}{\epsilon^{m+1-k}} (l_1-1)! \dots (l_k-1)!$$

$$\leq \sum_{k=1}^{m+1} \binom{m}{k-1} (m+1)! \frac{M^k}{\epsilon^{m+1-k}}$$

$$\leq \sum_{k=0}^m \binom{m}{k} (m+1)! \frac{M^{k+1}}{\epsilon^{m-k}}$$

$$\leq \frac{(m+1)!M}{\epsilon^m} \sum_{k=0}^m \binom{m}{k} (M\epsilon)^k$$

$$\leq (m+1)! M(\frac{1+M\epsilon}{\epsilon})^m$$

Pick $\delta = \frac{\epsilon}{2(1+M\epsilon)}$, we get

$$\frac{\delta^m}{m!} h(\nabla_{\eta_1} \dots \nabla_{\eta_m} \varphi_j)^{1/2}(s)$$

$$\leq \left(\frac{\epsilon}{2(1+M\epsilon)}\right)^m \frac{1}{m!} (m+1)! M(\frac{1+M\epsilon}{\epsilon})^m$$

$$\leq (m+1) M\left(\frac{1}{2}\right)^m \longrightarrow 0 \text{ as } m \to \infty.$$

So, we conclude

$$\sup \frac{\delta^m}{m!} h(\nabla_{\eta_1} \dots \nabla_{\eta_m} \varphi_j)^{1/2}(s) < \infty,$$

where the sup is taken over all $m = 0, 1, ...; \eta_1, ..., \eta_m \in \left\{\frac{\partial}{\partial s}, \frac{\partial}{\partial \bar{s}}\right\}$; and all $s \in K$.

Taking $f \equiv 1$, this shows that φ_j is an analytic section of S for any fixed j. Since $\{\varphi_j(s)_{j=0}^\infty\}$ spans H_s for any s, we have $H \to S$ is an analythic Hilbert field.

Now, if g is real-analytic and non-harmonic, and $k = \frac{\partial g}{\partial s}$, then the curvature operator $R\left(\frac{\partial}{\partial s}, \frac{\partial}{\partial \bar{s}}\right)$ are unbounded operators at points where $\Delta g \neq 0$. This means that this Hilbert field does not induce any Hilbert bundle.

References

[Dix69] Jacques Dixmier, Les C*-algèbres et leurs représentations, Deuxième édition. Cahiers Scientifiques, Fasc. XXIX, Gauthier-Villars Éditeur, Paris, 1969. MR 0246136 (39 #7442)

- [God51] R. Godement, Sur la théorie des représentations unitaires, Ann. of Math. (2) 53 (1951), 68–124. MR 0038571 (12,421d)
- [KP02] Steven G. Krantz and Harold R. Parks, A primer of real analytic functions, second ed., Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks], Birkhäuser Boston, Inc., Boston, MA, 2002. MR 1916029 (2003f:26045)
- [LS14] László Lempert and Róbert Szőke, Direct Images, Fields of Hilbert Spaces, and Geometric Quantization, Comm. Math. Phys. 327 (2014), no. 1, 49–99. MR 3177932