

Kakutani type Borsuk–Ulam theorem

Oleg R. Musin

Abstract

In this paper we present a Kakutani type theorem that is equivalent to the Borsuk–Ulam theorem for manifolds.

For a set S denote by 2^S the set of all subsets of S , including the empty set and S itself. Let X and Y be topological spaces. Say that $F : X \rightarrow 2^Y$ is a *set-valued function on X with a closed graph* if $\text{graph}(F) = \{(x, y) \in X \times Y : y \in F(x)\}$ is a closed as a subset of $X \times Y$.

The Kakutani fixed point theorem [1] states:

Let S be a non-empty, compact and convex subset of \mathbb{R}^d . Let $F : S \rightarrow 2^S$ be a set-valued function on S with a closed graph and the property that $F(x)$ is non-empty and convex for all $x \in S$. Then there is $y \in S$ such that $y \in F(y)$. (The point y is called a fixed point.)

In our paper [2] we extended the Borsuk–Ulam theorem for manifolds.

Let M be a compact PL (piece-wise linear) d -dimensional manifold without boundary with a free simplicial involution $A : M \rightarrow M$, i. e. $A^2(x) = A(A(x)) = x$ and $A(x) \neq x$. We say that a pair (M, A) is a *BUT (Borsuk-Ulam Type) manifold* if for any continuous $g : M \rightarrow \mathbb{R}^d$ there is a point $x \in M$ such that $g(A(x)) = g(x)$. Equivalently, if a continuous map $f : M \rightarrow \mathbb{R}^d$ is antipodal, i. e. $f(A(x)) = -f(x)$, then the zeros set $Z_f := f^{-1}(0)$ is not empty.

In [2], we found several equivalent necessary and sufficient conditions for manifolds to be BUT. For instance, *M is a BUT manifold if and only if M admits an antipodal continuous transversal to zeros map $h : M \rightarrow \mathbb{R}^d$ with $|Z_h| = 2 \pmod{4}$.*

The class of BUT manifolds is sufficiently large. It is clear that (\mathbb{S}^d, A) with $A(x) = -x$ is a BUT-manifold. Suppose that M can be represented as a connected sum $N \# N$, where N is a closed PL manifold. Then M admits a free involution. Indeed, M can be “centrally symmetric” embedded to \mathbb{R}^k with some k and the antipodal symmetry $x \rightarrow -x$ in \mathbb{R}^k implies a free involution $A : M \rightarrow M$ [2, Corollary 1]. For instance, orientable two-dimensional manifolds M_g^2 with even genus g and non-orientable manifolds P_m^2 with even m , where m is the number of Möbius bands, are BUT-manifolds.

Theorem 1. *Let M be a compact PL BUT-manifold of dimension d with a free involution A . Let $F : M \rightarrow 2^{\mathbb{R}^d}$ be a set-valued function on M with a closed graph and the property that for all $x \in M$ the set $F(x)$ is non-empty and convex in \mathbb{R}^d and there is $y \in F(x)$ such that $(-y) \in F(A(x))$. Then there is $x_0 \in M$ such that $F(x_0)$ covers the origin $0 \in \mathbb{R}^d$.*

Proof. Note that any PL manifold admit a metric. For a triangulation T of M the norm of T , denoted by $|T|$, is the diameter of the largest simplex in T .

Let $\{T_i, i = 1, 2, \dots\}$ be a sequence of antipodal triangulations of M with $|T_i| \rightarrow 0$. Now we define an antipodal mapping $f_i : M \rightarrow \mathbb{R}^d$. Note that if we define f_i for all vertices $V(T_i)$, then f_i can be piece-wise linearly extended to all T_i , i. e. to $f_i : M \rightarrow \mathbb{R}^d$.

Let $x \in V(T_i)$. Then $A(x)$ is also a vertex of T_i . Let $y \in F(x)$ be a point in \mathbb{R}^d such that $(-y) \in F(A(x))$. Then for the pair $(x, A(x))$ set $f_i(x) := y$ and $f_i(A(x)) := -y$.

Now we have a continuous antipodal mapping $f_i : M \rightarrow \mathbb{R}^d$. By assumption, M is BUT, therefore, there is a point $x_i \in M$ such that $f_i(x_i) = 0$. Now, suppose that x_i lies in a d -simplex s_i of T_i with vertices v_0^i, \dots, v_d^i and let $y_k^i := f_i(v_k^i)$. (Note that $y_k^i \in F(v_k^i) \subset \mathbb{R}^d$.) We have $0 \in \text{conv}(y_0^i, \dots, y_d^i)$ in \mathbb{R}^d . Let t_0^i, \dots, t_d^i be the barycentric coordinates of 0 relative to the simplex in \mathbb{R}^d with vertices y_0^i, \dots, y_d^i . Then

$$\sum_{k=0}^d t_k^i y_k^i = 0.$$

Since M is compact the sequences $\{x_i\}_{i=1}^\infty, \{t_k^i\}_{i=1}^\infty, \{y_k^i\}_{i=1}^\infty, k = 0, 1, \dots, d$, may, after possibly renumbering them, be assumed to converge to points x_0, t_k and $y_k, k = 0, 1, \dots, d$ respectively.

Finally, since the set-valued function F has closed graph, $y_k \in F(x_0), k = 0, 1, \dots, d$. Since F takes convex values, $F(x_0)$ is convex and so $0 \in \mathbb{R}^d$, being a convex combination of the y_k is in $F(x_0)$. Thus x_0 is the required point in M . \square

Corollary 1. *Let M be a compact PL BUT-manifold of dimension d with a free involution A . Let $F : M \rightarrow 2^{\mathbb{R}^d}$ be a set-valued function on M with a closed graph. Then there is $x_0 \in M$ such that the sets $F(x_0)$ and $F(A(x_0))$ have a common intersection point.*

Proof. Let $\bar{F}(x), x \in M$, be the Minkowski difference of $F(x)$ and $F(A(x))$ in \mathbb{R}^d ,

$$\bar{F}(x) := F(x) - F(A(x)) = \{a - b | a \in F(x), b \in F(A(x))\}.$$

Then $\bar{F}(A(x)) = -\bar{F}(x)$. Thus, the theorem for \bar{F} yields this corollary. \square

References

- [1] S. Kakutani, A generalization of Brouwers fixed point theorem, *Duke Math. J.*, **8** (1941): 457–459.
- [2] O. R. Musin, Borsuk-Ulam type theorems for manifolds, *Proc. Amer. Math. Soc.* **140** (2012), 2551-2560.