

MOVABLE INTERSECTION AND BIGNESS CRITERION

JIAN XIAO

ABSTRACT. In this note, we give a Morse-type bigness criterion for the difference of two pseudo-effective $(1,1)$ -classes by using movable intersections. As an application, we give a Morse-type bigness criterion for the difference of two movable $(n-1, n-1)$ -classes.

1. INTRODUCTION

Let X be a smooth projective variety of dimension n , and let A, B be two nef line bundles, then we have the fundamental inequality

$$\text{vol}(A - B) \geq A^n - nA^{n-1} \cdot B,$$

which is first discovered as a consequence of Demailly's holomorphic Morse inequalities (see [Dem85], [Siu93], [Tra95]). Thus the above inequality is usually called algebraic Morse inequality for line bundles. Recall that the volume of a holomorphic line bundle L is defined as

$$\text{vol}(L) := \limsup_{k \rightarrow +\infty} \frac{n!}{k^n} h^0(X, \mathcal{O}(kL)).$$

And L is called a big line bundle if $\text{vol}(L) > 0$. In particular, the Morse-type inequality for A, B implies that $A - B$ must be a big line bundle if $A^n - nA^{n-1} \cdot B > 0$. This provides a very effective way to construct holomorphic sections; see [DMR10, Dem11] for related applications.

Assume that L is a holomorphic line bundle over a compact Kähler manifold X , then it is proved by [Bou02b, Theorem 1.2] that the volume of L can be characterized as the maximum of the Monge-Ampère mass of the positive curvature currents contained in the class $c_1(L)$. This naturally extends the volume function $\text{vol}(\cdot)$ to transcendental $(1,1)$ -classes over compact complex manifold (see [Bou02b, Definition 1.3] or [BDPP13, Definition 3.2]).

Recall that Demailly's conjecture on (weak) transcendental holomorphic Morse inequality over compact Kähler manifolds is the following statement.

Conjecture 1.1. (see [BDPP13, Conjecture 10.1])¹ Let X be a compact Kähler manifold of dimension n , and let $\alpha, \beta \in H^{1,1}(X, \mathbb{R})$ be two nef classes. Then we have

$$\text{vol}(\alpha - \beta) \geq \alpha^n - n\alpha^{n-1} \cdot \beta.$$

In particular, if $\alpha^n - n\alpha^{n-1} \cdot \beta > 0$ then there exists a Kähler current in the class $\alpha - \beta$.

Based on the method of [Chi16], in our previous work [Xia15], it was proved that if $\alpha^n - 4n\alpha^{n-1} \cdot \beta > 0$ then there exists a Kähler current in the class $\alpha - \beta$. Recently, by keeping the same method as in [Chi16, Xia15] and with the new technique introduced by [Pop16], D. Popovici proved that the constant $4n$ can be improved to be the natural and optimal constant n . Thus we have a Morse-type criterion for the difference of two transcendental nef classes – indeed, our results in this note depend mainly on this important improvement.

It is natural to ask whether the above Morse-type bigness criterion “ $\alpha^n - n\alpha^{n-1} \cdot \beta > 0 \Rightarrow \text{vol}(\alpha - \beta) > 0$ ” for nef classes can be generalized to pseudo-effective $(1,1)$ -classes. Towards this generalization, we need the movable intersection products (denoted by $\langle - \rangle$) of pseudo-effective $(1,1)$ -classes developed in [Bou02a, BDPP13]. Then our problem can be stated as following:

¹For projective manifolds, this conjecture has been confirmed by [WN16].

Problem 1.1. Let X be a compact Kähler manifold of dimension n , and let $\alpha, \beta \in H^{1,1}(X, \mathbb{R})$ be two pseudo-effective classes. Does $\text{vol}(\alpha) - n\langle \alpha^{n-1} \rangle \cdot \beta > 0$ imply that there exists a Kähler current in the class $\alpha - \beta$?

Unfortunately, a very simple example due to [Tra95] implies that the above generalization does not always hold.

Example 1.1. (see [Tra95, Example 3.8]) Let $\pi : X \rightarrow \mathbb{P}^2$ be the blow-up of \mathbb{P}^2 along a point p . Let $R = \pi^*H$, where H is the hyperplane line bundle on \mathbb{P}^2 . Let $E = \pi^{-1}(p)$ be the exceptional divisor. Then for every positive integral k , the space of global holomorphic sections of $k(R - 2E)$ is the space of homogeneous polynomials in three variables of degree at most k and vanishes of order $2k$ at p ; hence $k(R - 2E)$ does not have any global holomorphic sections. The space $H^0(X, \mathcal{O}(k(R - 2E))) = \{0\}$ implies that $R - 2E$ can not be big. However, we have $R^2 - R \cdot 2E > 0$ as $R^2 = 1$ and $R \cdot E = 0$.

As the first result of this note, we show that the answer to Problem 1.1 is YES if β is movable. Here β being movable means that the negative part of β vanishes in its divisorial Zariski decomposition (see [Bou04]). In particular, if $\beta = c_1(L)$ for some pseudo-effective line bundle, then β being movable is equivalent to that the base locus of $mL + A$ is of codimension at least two for a fixed ample line bundle A and for large m .

Theorem 1.1. Let X be a compact Kähler manifold of dimension n , and let $\alpha, \beta \in H^{1,1}(X, \mathbb{R})$ be two pseudo-effective classes with β movable. Then $\text{vol}(\alpha) - n\langle \alpha^{n-1} \rangle \cdot \beta > 0$ implies that there exists a Kähler current in the class $\alpha - \beta$.

Remark 1.1. In the case when $\beta = 0$, Theorem 1.1 is just [Bou02b, Theorem 4.7], and when α is also nef, it is [DP04, Theorem 0.5].

An ancillary goal of the note is to explain the fact that Demailly's conjecture on weak transcendental holomorphic Morse inequality over compact Kähler manifolds is equivalent to the \mathcal{C}^1 differentiability of the volume function for transcendental $(1, 1)$ -classes. Though not stated explicitly, this fact is essentially contained in [BFJ09, Section 3] and the key ingredients are also implicitly contained in [BDPP13].

Proposition 1.1. Let X be a compact Kähler manifold of dimension n . Then the following statements are equivalent:

- (1) Let $\alpha, \beta \in H^{1,1}(X, \mathbb{R})$ be two nef classes, then we have

$$\text{vol}(\alpha - \beta) \geq \alpha^n - n\alpha^{n-1} \cdot \beta.$$

- (2) Let $\alpha, \beta \in H^{1,1}(X, \mathbb{R})$ be two pseudo-effective classes with β movable, then

$$\text{vol}(\alpha - \beta) \geq \langle \alpha^n \rangle - n\langle \alpha^{n-1} \rangle \cdot \beta.$$

- (3) Let $\alpha, \gamma \in H^{1,1}(X, \mathbb{R})$ be two $(1, 1)$ -classes with α big, then we have

$$\left. \frac{d}{dt} \right|_{t=0} \text{vol}(\alpha + t\gamma) = n\langle \alpha^{n-1} \rangle \cdot \gamma.$$

As an application of Proposition 1.1 and the \mathcal{C}^1 differentiability of the volume function for line bundles (see [BFJ09, Theorem A]), the algebraic Morse inequality can be generalized as following. It generalizes the previous result [Tra11, Corollary 3.2].

Theorem 1.2. Let X be a smooth projective variety of dimension n , and let α, β be the first Chern classes of two pseudo-effective line bundles with β movable. Then we have

$$\text{vol}(\alpha - \beta) \geq \text{vol}(\alpha) - n\langle \alpha^{n-1} \rangle \cdot \beta.$$

Remark 1.2. In particular, if α is nef and β is movable then we have $\text{vol}(\alpha - \beta) \geq \alpha^n - n\alpha^{n-1} \cdot \beta$ which is just [Tra11, Corollary 3.2].

Finally, as an application of Theorem 1.1, we give a Morse-type bigness criterion for the difference of two movable $(n-1, n-1)$ -classes.

Theorem 1.3. Let X be a compact Kähler manifold of dimension n , and let $\alpha, \beta \in H^{1,1}(X, \mathbb{R})$ be two pseudo-effective classes. Then $\text{vol}(\alpha) - n\alpha \cdot \langle \beta^{n-1} \rangle > 0$ implies that there exists a strictly positive $(n-1, n-1)$ -current in the class $\langle \alpha^{n-1} \rangle - \langle \beta^{n-1} \rangle$.

2. TECHNICAL PRELIMINARIES

2.1. Resolution of singularities of positive currents. Let X be a compact complex manifold, and let T be a d -closed almost positive $(1, 1)$ -current on X , that is, there exists a smooth $(1, 1)$ -form γ such that $T \geq \gamma$. If $\gamma = 0$, then T is called a positive $(1, 1)$ -current and the class $\{T\}$ is called pseudo-effective; And if γ is a hermitian metric, then T is called a Kähler current and the class $\{T\}$ is called big; see [Dem12] for the basic theory of positive currents.

Demailly's regularization theorem (see [Dem92]) implies that we can always approximate the almost positive $(1, 1)$ -current T by a family of almost positive closed $(1, 1)$ -currents T_k with analytic singularities such that $T_k \geq \gamma - \varepsilon_k \omega$, where $\varepsilon_k \downarrow 0$ is a sequence of positive constants and ω is a fixed hermitian metric. In particular, when T is a Kähler current, it can be approximated by a family of Kähler currents with analytic singularities.

When T has analytic singularities along an analytic subvariety $V(\mathcal{I})$ where $\mathcal{I} \subset \mathcal{O}_X$ is a coherent ideal sheaf, by blowing up along $V(\mathcal{I})$ and then resolving the singularities, we get a modification $\mu: \tilde{X} \rightarrow X$ such that $\mu^*T = \tilde{\theta} + [D]$ where $\tilde{\theta}$ is an almost positive smooth $(1, 1)$ -form with $\tilde{\theta} \geq \mu^*\gamma$ and D is an effective \mathbb{R} -divisor; see e.g. [BDPP13, Theorem 3.1]. In particular, if T is positive, then $\tilde{\theta}$ is a smooth positive $(1, 1)$ -form. We call such a modification the log-resolution of singularities of T .

For almost positive $(1, 1)$ -current T , we can always decompose T with respect to the Lebesgue measure; see e.g. [Bou02b, Section 2.3]. We write $T = T_{ac} + T_{sg}$ where T_{ac} is the absolutely continuous part and T_{sg} is the singular part. The absolutely part T_{ac} can be seen as a form with L^1_{loc} coefficients, and the wedge product $T_{ac}^k(x)$ makes sense for almost every point x . We always have $T_{ac} \geq \gamma$ since γ is smooth. If T has analytic singularities along V , then $T_{ac} = \mathbf{1}_{X \setminus V} T$. However, in general T_{ac} is not closed even if T is closed (see [Bou02a]). We have the following proposition.

Proposition 2.1. Let T_1, \dots, T_k be k almost positive closed $(1, 1)$ -currents with analytic singularities on X and let ψ be a smooth $(n-k, n-k)$ -form. Let $\mu: \tilde{X} \rightarrow X$ be a simultaneous log-resolution with $\mu^*T_i = \tilde{\theta}_i + [D_i]$. Then

$$\int_X T_{1,ac} \wedge \dots \wedge T_{k,ac} \wedge \psi = \int_{\tilde{X}} \tilde{\theta}_1 \wedge \dots \wedge \tilde{\theta}_k \wedge \mu^*\psi.$$

Proof. This is obvious since μ is an isomorphism outside a proper analytic subvariety and $T_{1,ac} \wedge \dots \wedge T_{k,ac}$ puts no mass on such subset and $\tilde{\theta}_i$ is smooth on \tilde{X} . \square

2.2. Movable cohomology classes. We first briefly recall the definition of divisorial Zariski decomposition and the definition of movable $(1, 1)$ -class on compact complex manifold; see [Bou04], see also [Nak04] for the algebraic approach.

Let X be a compact complex manifold of dimension n and let α be a pseudo-effective $(1, 1)$ -class over X , then one can always associate an effective divisor $N(\alpha) := \sum \nu(\alpha, D)D$ to α where the sum ranges among all prime divisors on X . The class $\{N(\alpha)\}$ is called the negative part of α . And $Z(\alpha) = \alpha - \{N(\alpha)\}$ is called the positive part of α . The decomposition $\alpha = Z(\alpha) + \{N(\alpha)\}$ then is the divisorial Zariski decomposition of α .

Definition 2.1. Let X be a compact complex manifold of dimension n , and let α be a pseudo-effective $(1, 1)$ -class. Then α is called movable if $\alpha = Z(\alpha)$.

Proposition 2.2. (see [Bou04, Proposition 2.3]) Let α be a movable $(1, 1)$ -class and let ω be a Kähler class, then for any $\delta > 0$ there exist a modification $\mu : Y \rightarrow X$ and a Kähler class $\tilde{\omega}$ over Y such that $\alpha + \delta\omega = \mu_*\tilde{\omega}$.

Remark 2.1. In [Bou04], α is called modified nef if $\alpha = Z(\alpha)$ (see [Bou04, Definition 2.2 and Proposition 3.8]). Here we call it movable in order to keep the same notation as the algebraic geometry situation. Let L be a line bundle over a smooth projective variety and let $\alpha = c_1(L)$, then α is modified nef if and only if L is movable.

2.3. Movable intersections. We take the opportunity to point out the well known fact that the several definitions of movable intersections of pseudo-effective $(1, 1)$ -classes over compact Kähler manifold coincide; see [Bou02a, BDPP13, BEGZ10] for the analytic constructions over compact Kähler manifold. And it also coincides with the algebraic construction of [BFJ09] on smooth projective variety for specified degrees. We remark that it is helpful to know the definition of movable intersections of pseudo-effective $(1, 1)$ -classes can be interpreted in several equivalent ways.

Let $\alpha_1, \dots, \alpha_k \in H^{1,1}(X, \mathbb{R})$ be pseudo-effective classes on a compact Kähler manifold of dimension n . By the common basic property of these definitions of movable intersection products, we only need to verify the respectively defined positive (k, k) cohomology classes $\langle \alpha_1 \cdot \dots \cdot \alpha_k \rangle$ coincide when all the classes are big. Firstly, by the definition of Riemann-Zariski space, it is clear from [BDPP13, Theorem 3.5] and [BFJ09, Definition 2.5] that the two definitions of movable intersection products are the same when X is a smooth projective variety defined over \mathbb{C} and all the classes α_i are in the Néron-Severi space and $k = 1, n-1, n$. Next, by testing on $\partial\bar{\partial}$ -closed smooth positive $(n-k, n-k)$ -forms, [BDPP13, Theorem 3.5], [BEGZ10, Definition 1.17, Proposition 1.18 and Proposition 1.20] and [Bou02a, Definition 3.2.1 and Lemma 3.2.5] imply that these three definitions give the same positive cohomology class in $H^{k,k}(X, \mathbb{R})$; see [Pri13, Proposition 1.10] for the detailed proof.

Inspired by [BDPP13, Definition 1.3, Theorem 1.5 and Conjecture 2.3], the definition of movable $(n-1, n-1)$ -classes in the Kähler setting can be formulated as following.

Definition 2.2. Let X be a compact Kähler manifold of dimension n , and let $\gamma \in H^{n-1, n-1}(X, \mathbb{R})$. Then γ is called a movable $(n-1, n-1)$ -class if it is in the closure of the convex cone generated by cohomology classes of the form $\langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \rangle$ with every α_i pseudo-effective.

Remark 2.2. When X is a smooth projective variety of dimension n , [BDPP13, Theorem 1.5] implies that the rational movable $(n-1, n-1)$ -classes are in the cone of movable curves.

3. PROOF OF THE MAIN RESULTS

Now let us begin to prove our main results. We first give a Morse-type bigness criterion for the difference of two pseudo-effective $(1, 1)$ -classes by using movable intersections. To this end, we need some properties of movable intersections.

Lemma 3.1. Let X be a compact Kähler manifold of dimension n , and let $\alpha_1, \dots, \alpha_{n-1}, \beta \in H^{1,1}(X, \mathbb{R})$ be pseudo-effective classes with β nef. Then we have

$$\langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \cdot \beta \rangle = \langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \rangle \cdot \beta.$$

Proof. By the continuity of positive products, by taking limits, we can assume that $\alpha_1, \dots, \alpha_{n-1}$ are big and β is Kähler.

First, note that we always have $\langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \cdot \beta \rangle \leq \langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \rangle \cdot \beta$ when β is only assumed to be big (or pseudo-effective). By [BDPP13, Theorem 3.5], there exists a sequence of simultaneous log-resolutions $\mu_m : X_m \rightarrow X$ with $\mu_m^*\alpha_i = \omega_{i,m} + [D_{i,m}]$ and $\mu_m^*\beta = \gamma_m + [E_m]$ such that

$$\langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \cdot \beta \rangle = \limsup_{m \rightarrow \infty} (\omega_{1,m} \cdot \dots \cdot \omega_{n-1,m} \cdot \gamma_m),$$

and

$$\langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \rangle \cdot \beta = \limsup_{m \rightarrow \infty} (\omega_{1,m} \cdot \dots \cdot \omega_{n-1,m}) \cdot \mu_m^* \beta.$$

When β is Kähler, the class γ_m is just $\mu_m^* \beta$. In conclusion, if β is nef then we have the desired equality

$$\langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \cdot \beta \rangle = \langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \rangle \cdot \beta.$$

□

Lemma 3.2. Let X be a compact Kähler manifold of dimension n . Let $\alpha_1, \dots, \alpha_{n-1}$ be pseudo-effective $(1,1)$ -classes, and let $\pi : Y \rightarrow X$ be a modification. Then for any Kähler class $\widehat{\omega}$ on Y we have

$$\langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \rangle \cdot \pi_* \widehat{\omega} \geq \langle \pi^* \alpha_1 \cdot \dots \cdot \pi^* \alpha_{n-1} \rangle \cdot \widehat{\omega}.$$

Proof. First, we have $\langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \rangle \cdot \pi_* \widehat{\omega} \geq \langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \cdot \pi_* \widehat{\omega} \rangle$ as noted in the proof of Lemma 3.1. On the other hand, we have $\langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \cdot \pi_* \widehat{\omega} \rangle = \langle \pi^* \alpha_1 \cdot \dots \cdot \pi^* \alpha_{n-1} \cdot \pi^*(\pi_* \widehat{\omega}) \rangle$. Since $\pi^*(\pi_* \widehat{\omega}) - \widehat{\omega}$ is pseudo-effective, we get that

$$\begin{aligned} \langle \alpha_1 \cdot \dots \cdot \alpha_{n-1} \rangle \cdot \pi_* \widehat{\omega} &\geq \langle \pi^* \alpha_1 \cdot \dots \cdot \pi^* \alpha_{n-1} \cdot \widehat{\omega} \rangle \\ &= \langle \pi^* \alpha_1 \cdot \dots \cdot \pi^* \alpha_{n-1} \rangle \cdot \widehat{\omega}. \end{aligned}$$

□

3.1. Theorem 1.1.

Proof of Theorem 1.1. Fix a Kähler metric ω on X , and denote the Kähler class by the same symbol. By continuity and the definition of movable intersections, we have

$$\lim_{\delta \rightarrow 0} (\langle (\alpha + \delta\omega)^n \rangle - n \langle (\alpha + \delta\omega)^{n-1} \rangle \cdot (\beta + \delta\omega)) = \langle \alpha^n \rangle - n \langle \alpha^{n-1} \rangle \cdot \beta.$$

So $\langle (\alpha + \delta\omega)^n \rangle - n \langle (\alpha + \delta\omega)^{n-1} \rangle \cdot (\beta + \delta\omega) > 0$ for small $\delta > 0$. Note also that $\alpha - \beta = (\alpha + \delta\omega) - (\beta + \delta\omega)$. Thus to prove the bigness of the class $\alpha - \beta$, we can assume α is big, and assume $\beta = \mu_* \widetilde{\omega}$ for some modification $\mu : Y \rightarrow X$ and some Kähler class $\widetilde{\omega}$ on Y at the beginning. By Lemma 3.1 and Lemma 3.2, our assumption then implies that

$$\langle (\mu^* \alpha)^n \rangle - n \langle (\mu^* \alpha)^{n-1} \rangle \cdot \widetilde{\omega} > 0.$$

We claim that this implies there exists a Kähler current in the class $\mu^* \alpha - \widetilde{\omega}$, which then implies the bigness of the class $\alpha - \beta = \mu_* (\mu^* \alpha - \widetilde{\omega})$.

Now it is reduced to prove the case when α is big and β is Kähler. Let ω be a Kähler metric in the class β . The definition of movable intersections implies there exists some Kähler current $T \in \alpha$ with analytic singularities along some subvariety V such that

$$\int_{X \setminus V} T^n - n \int_{X \setminus V} T^{n-1} \wedge \omega > 0.$$

Let $\pi : Z \rightarrow X$ be the log-resolution of the current T with $\pi^* T = \theta + [D]$ such that θ is a smooth positive $(1,1)$ -form on Z . By Proposition 2.1 we have

$$\int_Z \theta^n - n \int_Z \theta^{n-1} \wedge \pi^* \omega > 0.$$

The result of [Pop16] then implies that there exists a Kähler current in the class $\{\theta - \pi^* \omega\}$. As $\pi^* \alpha = \{\theta + [D]\}$, this proves the bigness of the class $\alpha - \beta$.

Thus we finish the proof that there exists a Kähler current in the general case when α is pseudo-effective and β is movable. □

Remark 3.1. Indeed, by the above argument we have the following bigness criterion: for any pseudo-effective $(1, 1)$ classes α, β ,

$$\langle \alpha^n \rangle - n \langle \alpha^{n-1} \cdot \beta \rangle > 0 \Rightarrow \text{vol}(Z(\alpha) - Z(\beta)) > 0.$$

Since the positive products $\langle \alpha^n \rangle$ and $\langle \alpha^{n-1} \cdot \beta \rangle$ depend only on the positive parts of α, β (see e.g. [Bou02a, Proposition 3.2.10]), we can assume that α, β are movable at the beginning. And by taking limits, we can also assume $\beta = \pi_* \hat{\omega}$ for some modification π . Then we have

$$\begin{aligned} \langle (\pi^* \alpha)^n \rangle - n \langle (\pi^* \alpha)^{n-1} \cdot \hat{\omega} \rangle &= \langle (\pi^* \alpha)^n \rangle - n \langle (\pi^* \alpha)^{n-1} \cdot \hat{\omega} \rangle \\ &\geq \langle (\pi^* \alpha)^n \rangle - n \langle (\pi^* \alpha)^{n-1} \cdot \pi^* \beta \rangle \\ &> 0, \end{aligned}$$

which implies that $\alpha - \beta = \pi_*(\pi^* \alpha - \hat{\omega})$ is big.

In the case of Example 1.1, since R is nef and E is exceptional, we have $\langle R^2 \rangle - 2 \langle R \cdot 2E \rangle = R^2 > 0$, we then get the bigness of $Z(R) - Z(2E) = R$.

3.2. Proposition 1.1. Towards the transcendental version of Theorem 3.1, we give the proof of Proposition 1.1 which is essentially already known in [BFJ09].

Proof of Proposition 1.1. It is obvious that (2) \Rightarrow (1). We will show that (1) \Rightarrow (2) and (1) \Leftrightarrow (3), which then proves the equivalence of the statements.

Firstly, we prove (1) \Rightarrow (2). To prove (2), we only need to consider the case when $\langle \alpha^n \rangle - n \langle \alpha^{n-1} \cdot \beta \rangle > 0$ and β is big and movable. By Theorem 1.1 we know that $\alpha - \beta$ is big. First assume that β is Kähler. By taking a sequence of suitable Fujita approximations μ_m satisfying $\mu_m^* \alpha = \omega_m + [E_m]$ and

$$\langle \alpha^n \rangle = \limsup_{m \rightarrow \infty} \omega_m^n, \quad \langle \alpha^{n-1} \rangle = \limsup_{m \rightarrow \infty} (\mu_m)_*(\omega_m^{n-1}).$$

Then (1) implies that $\text{vol}(\omega_m - \mu_m^* \beta) \geq \omega_m^n - n \omega_m^{n-1} \cdot \mu_m^* \beta$. Thus (1) implies that

$$\text{vol}(\alpha - \beta) \geq \text{vol}(\alpha) - n \langle \alpha^{n-1} \rangle \cdot \beta$$

holds in the case when β is Kähler. In the general case, we can assume that $\beta = \pi_* \hat{\omega}$. Then we have

$$\begin{aligned} \text{vol}(\alpha - \beta) &\geq \text{vol}(\pi^* \alpha - \hat{\omega}) \geq \text{vol}(\pi^* \alpha) - n \langle (\pi^* \alpha)^{n-1} \rangle \cdot \hat{\omega} \\ &\geq \text{vol}(\pi^* \alpha) - n \langle \alpha^{n-1} \rangle \cdot \pi_* \hat{\omega} \quad \text{by Lemma 3.2.} \end{aligned}$$

Next, note that the implication (3) \Rightarrow (1) is obvious, and the implication (1) \Rightarrow (3) is just [BFJ09, Section 3.2]. For reader's convenience, we briefly recall and repeat the arguments of [BFJ09, Section 3.2]. By [BFJ09, Corollary 3.4] (or the proof of [BDPP13, Theorem 4.1]), (1) implies

$$\text{vol}(\beta + t\gamma) \geq \beta^n + tn\beta^{n-1} \cdot \gamma - Ct^2$$

for an arbitrary nef class β , an arbitrary $(1, 1)$ -class γ and $t \in [0, 1]$. Here the constant C depends only on the class β, γ ; more precisely, the constant C depends on the volume of a big and nef class ω such that $\omega - \beta$ is pseudo-effective and $\omega \pm \gamma$ is nef.

Now take a log-resolution $\mu^* \alpha = \beta + [E]$, then we have

$$\begin{aligned} \text{vol}(\alpha + t\gamma) &\geq \text{vol}(\beta + t\mu^* \gamma) \\ &\geq \beta^n + tn\beta^{n-1} \cdot \mu^* \gamma - Ct^2 \\ &= \beta^n + tn\mu_*(\beta^{n-1}) \cdot \gamma - Ct^2. \end{aligned}$$

Note that the constant C does not depend on the resolution μ , since $\mu^* \omega - \beta$ is pseudo-effective and $\mu^* \omega \pm \mu^* \gamma$ is nef if ω has similar property with respect to α, γ . And we have $\text{vol}(\mu^* \omega) = \text{vol}(\omega)$.

By taking limits of some sequence of log-resolutions, we get

$$\text{vol}(\alpha + t\gamma) \geq \text{vol}(\alpha) + tn\langle\alpha^{n-1}\rangle \cdot \gamma - Ct^2.$$

Replace γ by $-\gamma$, we then get

$$\text{vol}(\alpha) \geq \text{vol}(\alpha + t\gamma) - tn\langle(\alpha + t\gamma)^{n-1}\rangle \cdot \gamma - Ct^2.$$

Since α is big, by the continuity of movable intersections (see e.g. [BDPP13, Theorem 3.5]) we have

$$\lim_{t \rightarrow 0} \langle(\alpha + t\gamma)^{n-1}\rangle = \langle\alpha^{n-1}\rangle.$$

Then (3) follows easily from the above inequalities. \square

Remark 3.2. It is shown in [Den15] that the C^1 differentiability of the volume function for transcendental $(1, 1)$ -classes holds on compact Kähler surfaces. And it is used to construct the Okounkov bodies of transcendental $(1, 1)$ -classes over compact Kähler surfaces.

3.3. Theorem 1.2. The algebraic Morse inequality tells us that, if L and F are two nef line bundles, then

$$\text{vol}(L - F) \geq L^n - nL^{n-1} \cdot F.$$

Recently, [Tra11] generalizes this result to the case when F is only movable. Assume that L is nef and F is pseudo-effective, and let $F = Z(F) + N(F)$ be the divisorial Zariski decomposition of F . Then [Tra11, Corollary 3.2] shows that

$$\text{vol}(L - Z(F)) \geq L^n - nL^{n-1} \cdot Z(F).$$

Moreover, if we write the negative part $N(F) = \sum_j \nu_j D_j$ where $\nu_j > 0$ and let u be a nef class on X such that $c_1(\mathcal{O}_{TX}(1)) + \pi^*u$ is a nef class on $\mathbb{P}(T^*X)$. Then [Tra11, Theorem 3.3] also gives a lower bound for $\text{vol}(L - F)$:

$$\text{vol}(L - F) \geq L^n - nL^{n-1} \cdot Z(F) - n \sum_j (L + \nu_j u)^{n-1} \cdot \nu_j D_j.$$

Our next result shows that L can be any pseudo-effective line bundle, where we restate Theorem 1.2.

Theorem 3.1. Let X be a smooth projective variety of dimension n , and let L, M be two pseudo-effective line bundles with M movable. Then we have

$$\text{vol}(L - M) \geq \text{vol}(L) - n\langle L^{n-1} \rangle \cdot M.$$

Proof. This follows directly from Proposition 1.1. \square

Remark 3.3. When M is nef and L is pseudo-effective, Theorem 3.1 can be also proved by using the singular Morse inequalities for line bundles (see [Bon98]). Without loss of generality, we can assume L is big and M is ample. Let $\omega \in c_1(M)$ be a Kähler metric. For any Kähler current $T \in c_1(L)$ with analytic singularities, $T - \omega$ is an almost positive curvature current of $L - M$ with analytic singularities. With the elementary pointwise inequality

$$\mathbf{1}_{X(\alpha - \beta, \leq 1)}(\alpha - \beta)^n \geq \alpha^n - n\alpha^{n-1} \wedge \beta$$

for positive $(1, 1)$ -forms, Theorem 3.1 then follows easily from [Bon98].

3.4. Theorem 1.3. Finally, inspired by the method in our previous work [Xia15] (see also [Chi16]), we show that Theorem 1.1 gives a Morse-type bigness criterion of the difference of two movable $(n-1, n-1)$ -classes, thus giving the proof of Theorem 1.3.

Proof of Theorem 1.3. Denote the Kähler cone of X by \mathcal{K} , and denote the cone generated by cohomology classes represented by positive $(n-1, n-1)$ -currents by \mathcal{N} . Then by the numerical characterization of Kähler cone of [DP04] (see also [BDPP13, Theorem 2.1]) we have the cone duality relation

$$\overline{\mathcal{K}}^\vee = \mathcal{N}.$$

Without loss of generality, we can assume that α, β are big. Then the existence of a strictly positive $(n-1, n-1)$ -current in the class $\langle \alpha^{n-1} \rangle - \langle \beta^{n-1} \rangle$ is equivalent to the existence of some positive constant $\delta > 0$ such that

$$\langle \alpha^{n-1} \rangle - \langle \beta^{n-1} \rangle \succeq \delta \langle \beta^{n-1} \rangle,$$

or equivalently,

$$\langle \alpha^{n-1} \rangle \succeq (1 + \delta) \langle \beta^{n-1} \rangle.$$

Here we denote $\gamma \succeq \eta$ if $\gamma - \eta$ contains a positive current.

In the following, we will argue by contradiction. By the cone duality relation $\overline{\mathcal{K}}^\vee = \mathcal{N}$, the class $\langle \alpha^{n-1} \rangle - \langle \beta^{n-1} \rangle$ does not contain any strictly positive $(n-1, n-1)$ -current is then equivalent to the statement: for any $\epsilon > 0$ there exists some non-zero class $N_\epsilon \in \overline{\mathcal{K}}$ such that

$$\langle \alpha^{n-1} \rangle \cdot N_\epsilon \leq (1 + \epsilon) \langle \beta^{n-1} \rangle \cdot N_\epsilon.$$

On the other hand, we claim that Theorem 1.1 implies

$$n(N \cdot \langle \alpha^{n-1} \rangle)(\alpha \cdot \langle \beta^{n-1} \rangle) \geq \langle \alpha^n \rangle (N \cdot \langle \beta^{n-1} \rangle)$$

for any nef $(1, 1)$ -class N . First note that both sides of the above inequality are of the same degree of each cohomology class. After scaling, we can assume

$$\alpha \cdot \langle \beta^{n-1} \rangle = N \cdot \langle \beta^{n-1} \rangle.$$

Then we need to prove $nN \cdot \langle \alpha^{n-1} \rangle \geq \langle \alpha^n \rangle$. Otherwise, we have $nN \cdot \langle \alpha^{n-1} \rangle < \langle \alpha^n \rangle$. And Theorem 1.1 implies that there must exist a Kähler current in the class $\alpha - N$. Then we must have

$$\langle \beta^{n-1} \rangle \cdot (\alpha - N) > 0,$$

which contradicts with our scaling equality $\langle \beta^{n-1} \rangle \cdot (\alpha - N) = 0$.

Let $N = N_\epsilon$, we get

$$\begin{aligned} (1 + \epsilon)n(N_\epsilon \cdot \langle \beta^{n-1} \rangle)(\alpha \cdot \langle \beta^{n-1} \rangle) &\geq n(N_\epsilon \cdot \langle \alpha^{n-1} \rangle)(\alpha \cdot \langle \beta^{n-1} \rangle) \\ &\geq \langle \alpha^n \rangle (N_\epsilon \cdot \langle \beta^{n-1} \rangle). \end{aligned}$$

This implies

$$(1 + \epsilon)n\alpha \cdot \langle \beta^{n-1} \rangle \geq \langle \alpha^n \rangle.$$

Since $\epsilon > 0$ is arbitrary, this contradicts with our assumption $\langle \alpha^n \rangle - n\alpha \cdot \langle \beta^{n-1} \rangle > 0$. Thus there must exist a strictly positive $(n-1, n-1)$ -current in the class $\langle \alpha^{n-1} \rangle - \langle \beta^{n-1} \rangle$. \square

Remark 3.4. Let X be a smooth projective variety of dimension n and let $\text{Mov}_1(X)$ be the closure of the cone generated by movable curve classes. In [LX15], we show that any interior point of $\text{Mov}_1(X)$ is the form $\langle L^{n-1} \rangle$ for a unique big and movable divisor class. And under Demailly's conjecture on transcendental Morse inequality, this also extends to transcendental movable $(n-1, n-1)$ -classes over compact Kähler manifold. In particular, this extends to compact hyperkähler manifolds.

REFERENCES

- [BDPP13] Sébastien Boucksom, Jean-Pierre Demailly, Mihai Păun, and Thomas Peternell. The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. *J. Algebraic Geom.*, 22(2):201–248, 2013.
- [BEGZ10] Sébastien Boucksom, Philippe Eyssidieux, Vincent Guedj, and Ahmed Zeriahi. Monge-Ampère equations in big cohomology classes. *Acta Math.*, 205(2):199–262, 2010.
- [BFJ09] Sébastien Boucksom, Charles Favre, and Mattias Jonsson. Differentiability of volumes of divisors and a problem of Teissier. *J. Algebraic Geom.*, 18(2):279–308, 2009.
- [Bon98] Laurent Bonavero. Inégalités de Morse holomorphes singulières. *J. Geom. Anal.*, 8(3):409–425, 1998.
- [Bou02a] Sébastien Boucksom. *Cônes positifs des variétés complexes compactes*. PhD thesis, Université Joseph-Fourier-Grenoble I, 2002.
- [Bou02b] Sébastien Boucksom. On the volume of a line bundle. *Internat. J. Math.*, 13(10):1043–1063, 2002.
- [Bou04] Sébastien Boucksom. Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. École Norm. Sup. (4)*, 37(1):45–76, 2004.
- [Chi16] Ionuț Chiose. The kähler rank of compact complex manifolds. *The Journal of Geometric Analysis*, 26(1):603–615, 2016.
- [Dem85] Jean-Pierre Demailly. Champs magnétiques et inégalités de Morse pour la d'' -cohomologie. *Ann. Inst. Fourier (Grenoble)*, 35(4):189–229, 1985.
- [Dem92] Jean-Pierre Demailly. Regularization of closed positive currents and intersection theory. *J. Algebraic Geom.*, 1(3):361–409, 1992.
- [Dem11] Jean-Pierre Demailly. Holomorphic Morse inequalities and the Green-Griffiths-Lang conjecture. *Pure Appl. Math. Q.*, 7(4, Special Issue: In memory of Eckart Viehweg):1165–1207, 2011.
- [Dem12] Jean-Pierre Demailly. Complex analytic and differential geometry. online book. *available at www-fourier.ujf-grenoble.fr/demailly/manuscripts/agbook.pdf*, Institut Fourier, Grenoble, 2012.
- [Den15] Ya Deng. Transcendental Morse inequality and generalized Okounkov bodies. *arXiv preprint arXiv:1503.00112*, 2015.
- [DMR10] Simone Diverio, Joël Merker, and Erwan Rousseau. Effective algebraic degeneracy. *Invent. Math.*, 180(1):161–223, 2010.
- [DP04] Jean-Pierre Demailly and Mihai Păun. Numerical characterization of the Kähler cone of a compact Kähler manifold. *Ann. of Math. (2)*, 159(3):1247–1274, 2004.
- [LX15] Brian Lehmann and Jian Xiao. Zariski decomposition of curves on algebraic varieties. *arXiv:1507.04316v1*, 2015.
- [Nak04] Noboru Nakayama. *Zariski-decomposition and abundance*, volume 14 of *MSJ Memoirs*. Mathematical Society of Japan, Tokyo, 2004.
- [Pop16] Dan Popovici. Sufficient bigness criterion for differences of two nef classes. *Math. Ann.*, 364(1-2):649–655, 2016.
- [Pri13] Mario Principato. Mobile product and zariski decomposition. *arXiv preprint arXiv:1301.1477*, 2013.
- [Siu93] Yum Tong Siu. An effective Matsusaka big theorem. *Ann. Inst. Fourier (Grenoble)*, 43(5):1387–1405, 1993.
- [Tra95] Stefano Trapani. Numerical criteria for the positivity of the difference of ample divisors. *Math. Z.*, 219(3):387–401, 1995.
- [Tra11] Stefano Trapani. Divisorial Zariski decomposition and algebraic Morse inequalities. *Proc. Amer. Math. Soc.*, 139(1):177–183, 2011.
- [WN16] David Witt Nyström. Duality between the pseudoeffective and the movable cone on a projective manifold. *with an appendix by S. Boucksom*, *arXiv:1602.03778*, 2016.
- [Xia15] Jian Xiao. Weak transcendental holomorphic Morse inequalities on compact Kähler manifolds. *Ann. Inst. Fourier (Grenoble)*, 65(3):1367–1379, 2015.

FUDAN UNIVERSITY & UNIVERSITÉ GRENOBLE ALPES

Email: jian.xiao@ujf-grenoble.fr