

Proving $\zeta(2)$ through an evolution of the Mengoli's series to the set of rational numbers

URIEL VALENTINIS RAMOS
email: valentinisramos@gmail.com

ABSTRACT.

The present paper is an evolution of the Mengoli's series to the set of rational numbers, which eventually will allow developing the summation, by limits, obtaining the value of $\zeta(2)$; problem which Mengoli himself was the first to postulate.

More specifically in this paper is postulated and demonstrated the function which resolves every infinite sum of the type $\sum_{n=1}^{\infty} \left(\frac{1}{(n+\frac{a}{w})(n+\frac{b}{w})\dots(n+\frac{z}{w})} \right)$ for all a, b, \dots, z, w belonging to the set of integers, where $a < b < \dots < z$; $\frac{a}{w}, \frac{b}{w}, \dots, \frac{z}{w} \neq -1, -2, -3, \dots$

Finally the following limit $\lim_{w \rightarrow \infty} \left\{ \sum_{n=1}^{\infty} \left(\frac{1}{(n-\frac{1}{w})(n+\frac{1}{w})} \right) \right\}$ is developed to demonstrate $\zeta(2)$.

Proposition 1.1.

$$\text{All } \sum_{n=1}^{\infty} \left(\frac{1}{(n+a)(n+b)} \right) = \frac{1}{(b-a)} \left[\sum_{n=1}^b \left(\frac{1}{n} \right) - \sum_{n=1}^a \left(\frac{1}{n} \right) \right] \text{ where } a, b \in \mathbb{Z}; \ a, b \geq 0; \ a \neq b.$$

Proof:

$$\begin{aligned} \sum_{n=1}^{\infty} \left(\frac{1}{(n+a)(n+b)} \right) &= \frac{1}{(b-a)} \left[\sum_{n=1}^{\infty} \left(\frac{n+b-(n+a)}{(n+a)(n+b)} \right) \right] = \frac{1}{(b-a)} \left[\sum_{n=1}^{\infty} \left(\frac{1}{n+a} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{n+b} \right) \right] = \\ &= \frac{1}{(b-a)} \left[\left(\sum_{n=1}^{\infty} \left(\frac{1}{n} \right) - \sum_{n=1}^a \left(\frac{1}{n} \right) \right) - \left(\sum_{n=1}^{\infty} \left(\frac{1}{n} \right) - \sum_{n=1}^b \left(\frac{1}{n} \right) \right) \right] = \frac{1}{(b-a)} \left[\sum_{n=1}^b \left(\frac{1}{n} \right) - \sum_{n=1}^a \left(\frac{1}{n} \right) \right] \end{aligned}$$

Proposition 1.2.

$$\begin{aligned} \text{All } \sum_{n=1}^{\infty} \left(\frac{1}{(n+a+\frac{x}{w})(n+b+\frac{y}{w})} \right) &= \\ &= \frac{w}{(x+wa-y-wb)} \left[\frac{\pi}{2} \left(\cot \left(\frac{\pi y}{w} \right) - \cot \left(\frac{\pi x}{w} \right) \right) + \sum_{n=1}^{w-1} \left(\ln \left(2 \sin \left(\frac{\pi n}{w} \right) \right) \left[\cos \left(\frac{2\pi n x}{w} \right) - \cos \left(\frac{2\pi n y}{w} \right) \right] \right) + \frac{w}{x} - \right. \\ &\left. \frac{w}{y} + \frac{|a|}{a(-1)^{2|a|}} \sum_{n=-\frac{1+|a|}{2}}^{|a|-\frac{1-|a|}{2}} \left(\frac{w}{x+\frac{|a|}{a}wn} \right) - \frac{|b|}{b(-1)^{2|b|}} \sum_{n=-\frac{1+|b|}{2}}^{|b|-\frac{1-|b|}{2}} \left(\frac{w}{y+\frac{|b|}{b}wn} \right) \right] \end{aligned}$$

where $x, y, w \in \mathbb{N}; \ 0 < x, y < w; \ a, b \in \mathbb{Z}; \ a + \frac{x}{w} \neq b + \frac{y}{w}.$

Proof:

From Gauss's Theorem for Digamma Function (studied by Jensen [2]) we know that

$$\sum_{n=0}^{\infty} \left(\frac{1}{n+1} - \frac{q}{p+nq} \right) = \frac{-\pi}{2} \cot \left(\frac{\pi p}{q} \right) - \ln(q) + \sum_{n=1}^{q-1} \left(\ln \left(2 \sin \left(\frac{\pi n}{q} \right) \right) \cos \left(\frac{2\pi n p}{q} \right) \right)$$

where $p, q \in \mathbb{N}; \ p < q.$

Increasing n from 0 to 1, the following summation is obtained:

$$\sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{q}{p+nq} \right) = \frac{-\pi}{2} \cot \left(\frac{\pi p}{q} \right) - \ln(q) + \sum_{n=1}^{q-1} \left(\ln \left(2 \sin \left(\frac{\pi n}{q} \right) \right) \cos \left(\frac{2\pi n p}{q} \right) \right) - 1 + \frac{q}{p}$$

From here two summations are generated:

Summation 1.2.1

If $a > 0$

$$\sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+w(n+a)} \right) = \sum_{n=1}^{\infty} \left(\frac{1}{n+1} \right) - \sum_{n=1}^{\infty} \left(\frac{w}{x+wn} \right) + \sum_{n=1}^a \left(\frac{w}{x+wn} \right)$$

If $a = 0$

$$\sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+w(n+a)} \right) = \sum_{n=1}^{\infty} \left(\frac{1}{n+1} \right) - \sum_{n=1}^{\infty} \left(\frac{w}{x+wn} \right)$$

If $a < 0$

$$\sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+w(n+a)} \right) = \sum_{n=1}^{\infty} \left(\frac{1}{n+1} \right) - \sum_{n=1}^{\infty} \left(\frac{w}{x+wn} \right) - \sum_{n=0}^{|a|-1} \left(\frac{w}{x-wn} \right)$$

Unifying the three possibilities:

$$\begin{aligned} \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+w(n+a)} \right) &= \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+wn} \right) + \frac{|a|}{a^{(-1)^{2|a|}}} \sum_{n=\frac{1+|a|}{2}}^{|a|-\frac{1-|a|}{2}} \left(\frac{w}{x+\frac{|a|}{a}wn} \right) = \\ &= \frac{-\pi}{2} \cot\left(\frac{\pi x}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \cos\left(\frac{2\pi n x}{w}\right) \right) - 1 + \frac{w}{x} + \frac{|a|}{a^{(-1)^{2|a|}}} \sum_{n=\frac{1+|a|}{2}}^{|a|-\frac{1-|a|}{2}} \left(\frac{w}{x+\frac{|a|}{a}wn} \right) \end{aligned} \quad (1.2.1)$$

Summation 1.2.2

$$\begin{aligned} \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{y+w(n+b)} \right) &= \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{y+wn} \right) + \frac{|b|}{b^{(-1)^{2|b|}}} \sum_{n=\frac{1+|b|}{2}}^{|b|-\frac{1-|b|}{2}} \left(\frac{w}{y+\frac{|b|}{b}wn} \right) = \\ &= \frac{-\pi}{2} \cot\left(\frac{\pi y}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \cos\left(\frac{2\pi n y}{w}\right) \right) - 1 + \frac{w}{y} + \frac{|b|}{b^{(-1)^{2|b|}}} \sum_{n=\frac{1+|b|}{2}}^{|b|-\frac{1-|b|}{2}} \left(\frac{w}{y+\frac{|b|}{b}wn} \right) \end{aligned} \quad (1.2.2)$$

From these expressions, the following operation can be defined **(1.2.1 – 1.2.2)**:

$$\begin{aligned}
 & \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+w(n+a)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{y+w(n+b)} \right) = \sum_{n=1}^{\infty} \left(\frac{w}{y+w(n+b)} - \frac{w}{x+w(n+a)} \right) = \\
 & = \sum_{n=1}^{\infty} \left(\frac{w(x+wn+wa) - w(y+wn+wb)}{(x+wn+wa)(y+wn+wb)} \right) = \sum_{n=1}^{\infty} \left(\frac{w(x+wa-y-wb)}{(x+wn+wa)(y+wn+wb)} \right) \\
 & = \sum_{n=1}^{\infty} \left(\frac{w(x+wa-y-wb)/w^2}{(x+wn+wa)(y+wn+wb)/w^2} \right) = \sum_{n=1}^{\infty} \left(\frac{(x+wa-y-wb)/w}{\frac{(x+wn+wa)}{w} \frac{(y+wn+wb)}{w}} \right) = \\
 & = \frac{(x+wa-y-wb)}{w} \sum_{n=1}^{\infty} \left(\frac{1}{\frac{(x+wn+wa)}{w} \frac{(y+wn+wb)}{w}} \right) = \frac{(x+wa-y-wb)}{w} \sum_{n=1}^{\infty} \left(\frac{1}{\left(n+a+\frac{x}{w}\right) \left(n+b+\frac{y}{w}\right)} \right)
 \end{aligned}$$

Using this last expression and substituting **1.2.1** and **1.2.2**, the following equation is obtained:

$$\begin{aligned}
 & \frac{(x+wa-y-wb)}{w} \sum_{n=1}^{\infty} \left(\frac{1}{\left(n+a+\frac{x}{w}\right) \left(n+b+\frac{y}{w}\right)} \right) = \\
 & = \frac{-\pi}{2} \cot\left(\frac{\pi x}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \cos\left(\frac{2\pi n x}{w}\right) \right) - 1 + \frac{w}{x} + \frac{|a|}{a^{(-1)^{2|a|}}} \sum_{n=\frac{1+|a|}{2}}^{\frac{1-|a|}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) \\
 & - \left[\frac{-\pi}{2} \cot\left(\frac{\pi y}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \cos\left(\frac{2\pi n y}{w}\right) \right) - 1 + \frac{w}{y} + \frac{|b|}{b^{(-1)^{2|b|}}} \sum_{n=\frac{1+|b|}{2}}^{\frac{1-|b|}{2}} \left(\frac{w}{y + \frac{|b|}{b} w n} \right) \right] = \\
 & = \frac{w}{(x+wa-y-wb)} \left[\frac{\pi}{2} \left(\cot\left(\frac{\pi y}{w}\right) - \cot\left(\frac{\pi x}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n x}{w}\right) - \cos\left(\frac{2\pi n y}{w}\right) \right] \right) + \frac{w}{x} - \frac{w}{y} + \frac{|a|}{a^{(-1)^{2|a|}}} \sum_{n=\frac{1+|a|}{2}}^{\frac{1-|a|}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) \right. \\
 & \left. - \frac{|b|}{b^{(-1)^{2|b|}}} \sum_{n=\frac{1+|b|}{2}}^{\frac{1-|b|}{2}} \left(\frac{w}{y + \frac{|b|}{b} w n} \right) \right]
 \end{aligned}$$

Proposition 1.3.

$$\begin{aligned}
 \text{All } \sum_{n=1}^{\infty} \left(\frac{1}{\left(n+a+\frac{x}{w}\right)(n+b)} \right) &= \\
 &= \frac{w}{(x+wa-wb)} \left[\frac{-\pi}{2} \cot\left(\frac{\pi x}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \cos\left(\frac{2\pi n x}{w}\right) \right) - 1 + \frac{w}{x} + \frac{|a|}{a} \sum_{n=\frac{1+|a|}{2}}^{\frac{1-|a|}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) \right. \\
 &\quad \left. + \frac{1}{b+1} - \sum_{n=1}^b \left(\frac{1}{n}\right) \right]
 \end{aligned}$$

where $x, w \in \mathbb{N}$; $x < w$; $a, b \in \mathbb{Z}$; $b \geq 0$.

Proof:

Using **Proposition 1**, the next summation is created:

$$\sum_{n=1}^{\infty} \left(\frac{1}{(n+a)(n+b)} \right) = \frac{1}{(b+1-b)} \left[\sum_{n=1}^{b+1} \left(\frac{1}{n}\right) - \sum_{n=1}^b \left(\frac{1}{n}\right) \right] = \frac{1}{b+1} \quad (1.1)$$

Using summation **1.2.1** (developed in **Proposition 1.2**) and adding the summation **1.1**, the following operation is presented:

$$\begin{aligned}
 &\sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+w(n+a)} \right) + \sum_{n=1}^{\infty} \left(\frac{1}{(n+b)(n+b+1)} \right) = \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+w(n+a)} \right) + \frac{1}{b+1} = \\
 &= \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+w(n+a)} \right) + \left[\sum_{n=1}^{\infty} \left(\frac{1}{n+b}\right) - \sum_{n=1}^{\infty} \left(\frac{1}{n+b+1}\right) \right] = \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x+w(n+a)} + \frac{1}{n+b} - \frac{1}{n+b+1} \right) = \\
 &= \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{1}{n+b+1} \right) + \sum_{n=1}^{\infty} \left(\frac{1}{n+b} - \frac{w}{x+w(n+a)} \right) = \sum_{n=1}^b \left(\frac{1}{n+1}\right) + \sum_{n=1}^{\infty} \left(\frac{(x+wn+wa) - w(n+b)}{(x+wn+wa)(n+b)} \right) = \\
 &= \sum_{n=1}^b \left(\frac{1}{n+1}\right) + \sum_{n=1}^{\infty} \left(\frac{(x+wa-wb)}{(x+wn+wa)(n+b)} \right) = \sum_{n=1}^b \left(\frac{1}{n+1}\right) + \sum_{n=1}^{\infty} \left(\frac{(x+wa-wb)/w}{(x+wn+wa)(n+b)/w} \right) = \\
 &= \sum_{n=1}^b \left(\frac{1}{n+1}\right) + \left(\frac{x}{w} + a - b\right) \sum_{n=1}^{\infty} \left(\frac{1}{\left(n+a+\frac{x}{w}\right)(n+b)} \right)
 \end{aligned}$$

hence,

$$\left(\frac{x}{w} + a - b\right) \sum_{n=1}^{\infty} \left(\frac{1}{(n + a + \frac{x}{w})(n + b)} \right) = \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{w}{x + w(n+a)} \right) + \sum_{n=1}^{\infty} \left(\frac{1}{(n+b)(n+b+1)} \right) - \sum_{n=1}^b \left(\frac{1}{n+1} \right)$$

and substituting both summation **1.1** and **1.2.1**, the next expression is achieved:

$$\begin{aligned} & \left(\frac{x}{w} + a - b\right) \sum_{n=1}^{\infty} \left(\frac{1}{(n + a + \frac{x}{w})(n + b)} \right) = \\ & = \frac{-\pi}{2} \cot\left(\frac{\pi x}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n x}{w}\right) \right] \right) - 1 + \frac{w}{x} + \frac{|a|}{a(-1)^{2|a|}} \sum_{n=\frac{1+|a|}{2}}^{\frac{|a|-\frac{1}{2}}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) + \frac{1}{b+1} - \sum_{n=1}^b \left(\frac{1}{n+1} \right) \end{aligned}$$

Therefore,

$$\begin{aligned} & \sum_{n=1}^{\infty} \left(\frac{1}{(n + a + x/w)(n + b)} \right) = \\ & = \frac{w}{(x + wa - wb)} \left[\frac{-\pi}{2} \cot\left(\frac{\pi x}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n x}{w}\right) \right] \right) + \frac{w}{x} + \frac{|a|}{a(-1)^{2|a|}} \sum_{n=\frac{1+|a|}{2}}^{\frac{|a|-\frac{1}{2}}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) - 1 + \frac{1}{b+1} - \sum_{n=1}^b \left(\frac{1}{n+1} \right) \right] = \\ & = \frac{w}{(x + wa - wb)} \left[\frac{-\pi}{2} \cot\left(\frac{\pi x}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n x}{w}\right) \right] \right) + \frac{w}{x} + \frac{|a|}{a(-1)^{2|a|}} \sum_{n=\frac{1+|a|}{2}}^{\frac{|a|-\frac{1}{2}}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) - \sum_{n=1}^b \left(\frac{1}{n} \right) \right] \end{aligned}$$

Proposition 1.4.

$$\begin{aligned}
 \text{All } \sum_{n=1}^{\infty} \left(\frac{1}{\left(n+a+\frac{x}{w}\right)\left(n+b+\frac{y}{w}\right)} \right) &= \\
 &= \frac{w}{(x+wa-y-wb)} \left[\frac{\pi}{2} \left(\frac{\cos\left(\frac{\pi y}{w}\right)}{\left[\sin\left(\frac{\pi y}{w}\right)\right]^{(-1)^{2y}}} - \frac{\cos\left(\frac{\pi x}{w}\right)}{\left[\sin\left(\frac{\pi x}{w}\right)\right]^{(-1)^{2x}}} \right) - \frac{x}{x^{(-1)^{2x}} \ln(w)} + \frac{y}{y^{(-1)^{2y}} \ln(w)} \right. \\
 &\quad + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\frac{x}{x^{(-1)^{2x}}} \cos\left(\frac{2\pi n x}{w}\right) - \frac{y}{y^{(-1)^{2y}}} \cos\left(\frac{2\pi n y}{w}\right) \right] \right) + \frac{w}{x^{(-1)^{2x}}} - \frac{w}{y^{(-1)^{2y}}} \\
 &\quad \left. + \frac{|a|}{a^{(-1)^{2|a|}}} \sum_{n=\frac{1+|a|}{2}}^{|a|-\frac{|a|}{2}} \left(\frac{w}{x+\frac{|a|}{a} w n} \right) - \frac{|b|}{b^{(-1)^{2|b|}}} \sum_{n=\frac{1+|b|}{2}}^{|b|-\frac{|b|}{2}} \left(\frac{w}{y+\frac{|b|}{b} w n} \right) \right] =
 \end{aligned}$$

where $a, b, x, y, w \in \mathbb{Z}$; $0 \leq x, y < w$; $a + \frac{x}{w} \neq b + \frac{y}{w}$; $\left(a + \frac{x}{w}\right), \left(b + \frac{y}{w}\right) \neq -1, -2, -3, \dots$

Proof:

Proposition 1.4 is obtained by consolidation of Propositions 1.1, 1.2 and 1.3.

- From Proposition 1.1: $\text{All } \sum_{n=1}^{\infty} \left(\frac{1}{(n+a)(n+b)} \right) = \frac{1}{(b-a)} \left[\sum_{n=1}^b \left(\frac{1}{n} \right) - \sum_{n=1}^a \left(\frac{1}{n} \right) \right]$

where $a, b \in \mathbb{Z}$; $a, b \geq 0$; $a \neq b$.

And $\sum_{n=1}^{\infty} \left(\frac{1}{(n+a)(n+b)} \right) = \sum_{n=1}^{\infty} \left(\frac{1}{\left(n+a+\frac{x}{w}\right)\left(n+b+\frac{y}{w}\right)} \right) \Rightarrow x, y = 0, w \neq 0$

Substituting $x, y = 0; w \neq 0; a, b \geq 0$ at **Proposition 1.4**:

$$\begin{aligned}
 &= \frac{w}{(0 + wa - 0 - wb)} \left[\frac{\pi}{2} \left(\frac{\cos\left(\frac{\pi 0}{w}\right)}{\left[\sin\left(\frac{\pi 0}{w}\right)\right]^{(-1)^{2^0}}} - \frac{\cos\left(\frac{\pi 0}{w}\right)}{\left[\sin\left(\frac{\pi 0}{w}\right)\right]^{(-1)^{2^0}}} \right) - \frac{0}{0^{(-1)^{2^{|0|}}}} \ln(w) + \frac{0}{0^{(-1)^{2^{|0|}}}} \ln(w) \right. \\
 &\quad + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\frac{0}{0^{(-1)^{2^{|0|}}}} \cos\left(\frac{2\pi n 0}{w}\right) - \frac{0}{0^{(-1)^{2^{|0|}}}} \cos\left(\frac{2\pi n 0}{w}\right) \right] \right) + \frac{w}{0^{(-1)^{2^{|0|}}}} - \frac{w}{0^{(-1)^{2^{|0|}}}} \\
 &\quad \left. + \frac{|a|}{+a^{(-1)^{2^{|a|}}}} \sum_{n=\frac{1+|a|}{2}}^{|a|-\frac{1-|a|}{2}} \left(\frac{w}{0 + \frac{|a|}{a} wn} \right) - \frac{|b|}{b^{(-1)^{2^{|b|}}}} \sum_{n=\frac{1+|b|}{2}}^{|b|-\frac{1-|b|}{2}} \left(\frac{w}{0 + \frac{|b|}{b} wn} \right) \right] = \\
 &= \frac{w}{w(a-b)} \left[\sum_{n=1}^a \left(\frac{1}{n}\right) - \sum_{n=1}^b \left(\frac{1}{n}\right) \right] = \frac{1}{(a-b)} \left[\sum_{n=1}^a \left(\frac{1}{n}\right) - \sum_{n=1}^b \left(\frac{1}{n}\right) \right]
 \end{aligned}$$

▪ From Proposition 1.2: $\text{All } \sum_{n=1}^{\infty} \left(\frac{1}{\left(n+a+\frac{x}{w}\right)\left(n+b+\frac{y}{w}\right)} \right)$

$$\begin{aligned}
 &= \frac{w}{(x + wa - y - wb)} \left[\frac{\pi}{2} \left(\cot\left(\frac{\pi y}{w}\right) - \cot\left(\frac{\pi x}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi nx}{w}\right) - \cos\left(\frac{2\pi ny}{w}\right) \right] \right) + \frac{w}{x} - \frac{w}{y} \right. \\
 &\quad \left. + \frac{|a|}{a^{(-1)^{2^{|a|}}}} \sum_{n=\frac{1+|a|}{2}}^{|a|-\frac{1-|a|}{2}} \left(\frac{w}{x + \frac{|a|}{a} wn} \right) - \frac{|b|}{b^{(-1)^{2^{|b|}}}} \sum_{n=\frac{1+|b|}{2}}^{|b|-\frac{1-|b|}{2}} \left(\frac{w}{y + \frac{|b|}{b} wn} \right) \right]
 \end{aligned}$$

where $x, y, w \in \mathbb{N}; 0 < x, y < w; a, b \in \mathbb{Z}; a + \frac{x}{w} \neq b + \frac{y}{w}$.

Substituting $0 < x, y < w$ at **Proposition 1.4**:

$$\begin{aligned}
 &= \frac{w}{(x + wa - y - wb)} \left[\frac{\pi}{2} \left(\frac{\cos\left(\frac{\pi y}{w}\right)}{\sin\left(\frac{\pi y}{w}\right)} - \frac{\cos\left(\frac{\pi x}{w}\right)}{\sin\left(\frac{\pi x}{w}\right)} \right) - \frac{x}{w} \ln(w) + \frac{y}{w} \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\frac{x}{w} \cos\left(\frac{2\pi n x}{w}\right) - \frac{y}{w} \cos\left(\frac{2\pi n y}{w}\right) \right] \right) \right. \\
 &\quad \left. + \frac{w}{x} - \frac{w}{y} + \frac{|a|}{a^{(-1)^{2|a|}}} \sum_{n=\frac{1+|a|}{2}}^{|a|-\frac{|a|}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) - \frac{|b|}{b^{(-1)^{2|b|}}} \sum_{n=\frac{1+|b|}{2}}^{|b|-\frac{|b|}{2}} \left(\frac{w}{y + \frac{|b|}{b} w n} \right) \right] = \\
 &= \frac{w}{(x + wa - y - wb)} \left[\frac{\pi}{2} \left(\cot\left(\frac{\pi y}{w}\right) - \cot\left(\frac{\pi x}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n x}{w}\right) - \cos\left(\frac{2\pi n y}{w}\right) \right] \right) + \frac{w}{x} - \frac{w}{y} \right. \\
 &\quad \left. + \frac{|a|}{a^{(-1)^{2|a|}}} \sum_{n=\frac{1+|a|}{2}}^{|a|-\frac{|a|}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) - \frac{|b|}{b^{(-1)^{2|b|}}} \sum_{n=\frac{1+|b|}{2}}^{|b|-\frac{|b|}{2}} \left(\frac{w}{y + \frac{|b|}{b} w n} \right) \right]
 \end{aligned}$$

- From Proposition 1.3: All $\sum_{n=1}^{\infty} \left(\frac{1}{(n+a+\frac{x}{w})(n+b)} \right) =$

$$\begin{aligned}
 &= \frac{w}{(x + wa - y - wb)} \left[\frac{-\pi}{2} \cot\left(\frac{\pi x}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \cos\left(\frac{2\pi n x}{w}\right) \right) - 1 + \frac{w}{x} \right. \\
 &\quad \left. + \frac{|a|}{a} \sum_{n=\frac{1+|a|}{2}}^{|a|-\frac{|a|}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) + \frac{1}{b+1} - \sum_{n=1}^b \left(\frac{1}{n} \right) \right] \quad \text{where } x, w \in \mathbb{N}; x < w; a, b \in \mathbb{Z}; b \geq 0.
 \end{aligned}$$

Substituting $y = 0$; $0 < x < w$; $b \geq 0$ at **Proposition 1.4**:

$$\begin{aligned}
 &= \frac{w}{(x + wa - 0 - wb)} \left[\frac{\pi}{2} \left(0 - \frac{\cos\left(\frac{\pi x}{w}\right)}{\sin\left(\frac{\pi x}{w}\right)} \right) - \frac{x}{x} \ln(w) + 0 \ln(w) \right. \\
 &\quad + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\frac{x}{x} \cos\left(\frac{2\pi n x}{w}\right) - 0 \cos\left(\frac{2\pi n y}{w}\right) \right] \right) + \frac{w}{x} - 0 w \\
 &\quad \left. + \frac{|a|}{a^{(-1)^{2|a|}}} \sum_{n=\frac{1+|a|}{2}}^{\frac{1-|a|}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) - \sum_{n=1}^{|b|} \left(\frac{w}{0 + w n} \right) \right] = \\
 &= \frac{w}{(x + wa - wb)} \left[\frac{-\pi}{2} \cot\left(\frac{\pi x}{w}\right) - \ln(w) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n x}{w}\right) \right] \right) + \frac{w}{x} + \frac{|a|}{a^{(-1)^{2|a|}}} \sum_{n=\frac{1+|a|}{2}}^{\frac{1-|a|}{2}} \left(\frac{w}{x + \frac{|a|}{a} w n} \right) \right. \\
 &\quad \left. - \sum_{n=1}^b \left(\frac{1}{n} \right) \right]
 \end{aligned}$$

An example of the proposition 1.4 can be found in the development of $\zeta(2)$, which is proved by limits, as follows:

$$\begin{aligned}
 &\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w} \right) \left(n + 0 + \frac{1}{w} \right)} \right) = \\
 &= \frac{w}{(w-1-w-1)} \left[\frac{\pi}{2} \left(\cot\left(\frac{\pi}{w}\right) - \cot\left(\frac{\pi(w-1)}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n(w-1)}{w}\right) - \cos\left(\frac{2\pi n}{w}\right) \right] \right) + \frac{w}{(w-1)} - \frac{w}{1} - \frac{w}{(w-1)} \right] = \\
 &= \frac{w}{(-2)} \left[\frac{\pi}{2} \left(\cot\left(\frac{\pi}{w}\right) - \cot\left(\frac{\pi(w-1)}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n(w-1)}{w}\right) - \cos\left(\frac{2\pi n}{w}\right) \right] \right) - \frac{w}{1} \right] = \\
 &= \frac{w}{2} \left[-\frac{\pi}{2} \left(2 \cot\left(\frac{\pi}{w}\right) \right) - \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) [0] \right) + w \right] = w \left[\frac{w}{2} - \frac{\pi}{2} \cot\left(\frac{\pi}{w}\right) \right]
 \end{aligned}$$

Applying limits:

$$\lim_{w \rightarrow \infty} \left\{ \sum_{n=1}^{\infty} \left(\frac{1}{\left((n+(-1) + \frac{w-1}{w}) \left(n + \frac{1}{w} \right) \right)} \right) \right\} = \sum_{n=1}^{\infty} \left(\frac{1}{n^2} \right) = \lim_{w \rightarrow \infty} \left\{ w \left[\frac{w}{2} - \frac{\pi}{2} \cot \left(\frac{\pi}{w} \right) \right] \right\}$$

Due to $\lim_{w \rightarrow \infty} \left\{ w \left[\frac{w}{2} - \frac{\pi}{2} \cot \left(\frac{\pi}{w} \right) \right] \right\} = c$, $f(w) = \left[\frac{w^2}{2} - \frac{\pi w}{2} \cot \left(\frac{\pi}{w} \right) \right]$ being continuous and $\lim_{w \rightarrow 0} \{1/w\} = \infty$ these imply $\lim_{w \rightarrow \infty} \left\{ w \left[\frac{w}{2} - \frac{\pi}{2} \cot \left(\frac{\pi}{w} \right) \right] \right\} = \lim_{w \rightarrow 0} \left\{ 1/w \left[\frac{1/w}{2} - \frac{\pi}{2} \cot \left(\frac{\pi}{1/w} \right) \right] \right\}$

$$\lim_{w \rightarrow 0} \left\{ 1/w \left[\frac{1/w}{2} - \frac{\pi}{2} \cot \left(\frac{\pi}{1/w} \right) \right] \right\} = \lim_{w \rightarrow 0} \left\{ 1/w \left[\frac{1/w}{2} - \frac{\pi}{2} \cot \left(\frac{\pi}{1/w} \right) \right] \right\} = \lim_{w \rightarrow 0} \left\{ \frac{1}{2w^2} \right\} = \lim_{w \rightarrow 0} \left\{ \frac{\tan(\pi w) - \pi w}{2w^2 \tan(\pi w)} - \frac{\pi}{2w} \cot(\pi w) \right\} = \frac{0}{0}$$

L'Hopital is applied

$$\begin{aligned} \lim_{w \rightarrow 0} \left\{ \frac{\frac{\pi[\cos^2(\pi w) + \sin^2(\pi w)] - \pi}{\cos^2(\pi w)}}{2 \left[2w \tan(\pi w) + \frac{w^2 \pi [\cos^2(\pi w) + \sin^2(\pi w)]}{\cos^2(\pi w)} \right]} \right\} &= \pi/2 * \lim_{w \rightarrow 0} \left\{ \frac{\frac{1}{\cos^2(\pi w)} - 1}{2w \tan(\pi w) + \frac{w^2 \pi}{\cos^2(\pi w)}} \right\} \\ &= \pi/2 * \lim_{w \rightarrow 0} \left\{ \frac{\cos^2(\pi w) \frac{1}{\cos^2(\pi w)} - 1}{\cos^2(\pi w) \left(2w \tan(\pi w) + \frac{w^2 \pi}{\cos^2(\pi w)} \right)} \right\} \\ &= \pi/2 * \lim_{w \rightarrow 0} \left\{ \frac{1 - \cos^2(\pi w)}{2w \sin(\pi w) \cos(\pi w) + w^2 \pi} \right\} = \pi/2 * \lim_{w \rightarrow 0} \left\{ \frac{1 - \cos^2(\pi w)}{w \sin(2\pi w) + w^2 \pi} \right\} = \frac{0}{0} \end{aligned}$$

L'Hopital is applied

$$\begin{aligned} \pi/2 * \lim_{w \rightarrow 0} \left\{ \frac{-[-2\pi \sin(\pi w) \cos(\pi w)]}{\sin(2\pi w) + 2\pi w \cos(2\pi w) + 2w\pi} \right\} &= \pi/2 * \lim_{w \rightarrow 0} \left\{ \frac{\pi \sin(2\pi w)}{\sin(2\pi w) + 2\pi w \cos(2\pi w) + 2w\pi} \right\} \\ &= \pi^2/2 * \lim_{w \rightarrow 0} \left\{ \frac{\sin(2\pi w)}{\sin(2\pi w) + 2\pi w \cos(2\pi w) + 2w\pi} \right\} = \frac{0}{0} \end{aligned}$$

L'Hopital is applied

$$\begin{aligned} \pi^2/2 * \lim_{w \rightarrow 0} \left\{ \frac{2\pi \cos(2\pi w)}{2\pi \cos(2\pi w) + 2\pi \cos(2\pi w) - 4\pi^2 w \sin(2\pi w) + 2\pi} \right\} &= \\ &= \pi^2/2 * \lim_{w \rightarrow 0} \left\{ \frac{\cos(2\pi w)}{\cos(2\pi w) + \cos(2\pi w) - 2\pi w \sin(2\pi w) + 1} \right\} = \pi^2/2 * \left[\frac{1}{1+1-0+1} \right] = \frac{\pi^2}{6} \end{aligned}$$

Proposition 2.

Every infinite sum with z divisors: $\left(n + \frac{a}{w}\right)\left(n + \frac{b}{w}\right) \dots \left(n + \frac{z}{w}\right)$ can be solved based on a finite set of infinite sums with 2 divisors. This means, that every

$$\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{a}{w}\right)\left(n + \frac{b}{w}\right) \dots \left(n + \frac{z}{w}\right)} \right)$$

Can be solved as follows

$$\frac{\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{a}{w}\right) \dots \left(n + \frac{z}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{b}{w}\right) \dots \left(n + \frac{z}{w}\right)} \right)}{\frac{b}{w} - \frac{a}{w}}$$

Proof:

$$\begin{aligned} \frac{1}{\left(n + \frac{a}{w}\right)\left(n + \frac{b}{w}\right) \dots \left(n + \frac{z}{w}\right)} &= \frac{1}{\frac{b}{w} - \frac{a}{w}} \left[\frac{1}{\left(n + \frac{a}{w}\right) \dots \left(n + \frac{z}{w}\right)} - \frac{1}{\left(n + \frac{b}{w}\right) \dots \left(n + \frac{z}{w}\right)} \right] = \\ &= \frac{1}{\frac{b}{w} - \frac{a}{w}} \left[\frac{\left(n + \frac{b}{w}\right) - \left(n + \frac{a}{w}\right)}{\left(n + \frac{a}{w}\right)\left(n + \frac{b}{w}\right) \dots \left(n + \frac{z}{w}\right)} \right] = \frac{1}{\frac{b}{w} - \frac{a}{w}} \left[\frac{\frac{b}{w} - \frac{a}{w}}{\left(n + \frac{a}{w}\right)\left(n + \frac{b}{w}\right) \dots \left(n + \frac{z}{w}\right)} \right] = \\ &= \frac{1}{\left(n + \frac{a}{w}\right)\left(n + \frac{b}{w}\right) \dots \left(n + \frac{z}{w}\right)} \end{aligned}$$

An example of the proposition 2 can be found in the development of $\zeta(4)$, as follows:

$$\begin{aligned} \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{-2}{w}\right)\left(n + \frac{-1}{w}\right)\left(n + \frac{1}{w}\right)\left(n + \frac{2}{w}\right)} \right) &= \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right)\left(n + (-1) + \frac{w-1}{w}\right)\left(n + \frac{1}{w}\right)\left(n + \frac{2}{w}\right)} \right) = \\ &= \frac{\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right)\left(n + \frac{1}{w}\right)\left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right)\left(n + \frac{1}{w}\right)\left(n + \frac{2}{w}\right)} \right)}{\left(-1 + \frac{w-1}{w}\right) - \left(-1 + \frac{w-2}{w}\right)} = \\ &= \frac{\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right)\left(n + \frac{1}{w}\right)\left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right)\left(n + \frac{1}{w}\right)\left(n + \frac{2}{w}\right)} \right)}{\frac{w-1-w+2}{w}} = \end{aligned}$$

$$\begin{aligned}
 &= w \left[\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right) \left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right) \left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right) \right] = \\
 &= w \left[\frac{\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right) \left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right)}{\left(\frac{1}{w}\right) - \left(-1 + \frac{w-2}{w}\right)} - \frac{\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right) \left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right)}{\left(\frac{1}{w}\right) - \left(-1 + \frac{w-1}{w}\right)} \right] = \\
 &= w \left[\frac{\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right) \left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right)}{1 + \left(\frac{1-w+2}{w}\right)} - \frac{\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right) \left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right)}{1 + \frac{1-w+1}{w}} \right] = \\
 &= w \left[\frac{\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right) \left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right)}{\frac{3}{w}} - \frac{\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right) \left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right)}{\frac{2}{w}} \right] = \\
 &= \frac{w^2}{6} \left[2 \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right) \left(n + \frac{2}{w}\right)} \right) - 2 \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right) - 3 \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right) \left(n + \frac{2}{w}\right)} \right) + 3 \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right) \right] = \\
 &= \frac{w^2}{6} \left[2 \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right) \left(n + \frac{2}{w}\right)} \right) + \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + \frac{1}{w}\right) \left(n + \frac{2}{w}\right)} \right) - 3 \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right) \left(n + \frac{2}{w}\right)} \right) \right] = \\
 &= \frac{w^2}{3} \left[\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right) \left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right) \left(n + \frac{1}{w}\right)} \right) \right]
 \end{aligned}$$

Proposition 1 is applied

$$\begin{aligned}
 &\frac{w^2}{3} \left[\sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w}\right) \left(n + \frac{2}{w}\right)} \right) - \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-1}{w}\right) \left(n + \frac{1}{w}\right)} \right) \right] = \\
 &= \frac{w^2}{6} \left[2 \left(\frac{w}{((w-2) - w - 2)} \left[\frac{\pi}{2} \left(\cot\left(\frac{2\pi}{w}\right) - \cot\left(\frac{\pi(w-2)}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n(w-2)}{w}\right) - \cos\left(\frac{2\pi n 2}{w}\right) \right] \right) + \frac{w}{(w-2)} - \frac{w}{2} - \frac{w}{(w-2)} \right) \right. \right. \\
 &\quad \left. \left. + \left(\frac{w}{(1-2)} \left[\frac{\pi}{2} \left(\cot\left(\frac{2\pi}{w}\right) - \cot\left(\frac{\pi}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n}{w}\right) - \cos\left(\frac{2\pi n 2}{w}\right) \right] \right) + \frac{w}{1} - \frac{w}{2} \right] \right) \right. \right. \\
 &\quad \left. \left. - 3 \left(\frac{w}{((w-1) - w - 2)} \left[\frac{\pi}{2} \left(\cot\left(\frac{2\pi}{w}\right) - \cot\left(\frac{\pi(w-1)}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n(w-1)}{w}\right) - \cos\left(\frac{2\pi n 2}{w}\right) \right] \right) + \frac{w}{(w-1)} - \frac{w}{2} \right. \right. \right. \right. \\
 &\quad \left. \left. \left. - \frac{w}{(w-1)} \right] \right) \right] \right] =
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{w^2}{6} \left[2 \left(\frac{w}{(-4)} \left[\frac{\pi}{2} \left(\cot\left(\frac{2\pi}{w}\right) - \cot\left(\frac{\pi(w-2)}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) [0] \right) - \frac{w}{2} \right) \right. \right. \\
 &\quad \left. \left. + \left(\frac{w}{(-1)} \left[\frac{\pi}{2} \left(\cot\left(\frac{2\pi}{w}\right) - \cot\left(\frac{\pi}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n}{w}\right) - \cos\left(\frac{2\pi n 2}{w}\right) \right] \right) + \frac{w}{2} \right] \right) \right. \right. \\
 &\quad \left. \left. - 3 \left(\frac{w}{(-3)} \left[\frac{\pi}{2} \left(\cot\left(\frac{2\pi}{w}\right) - \cot\left(\frac{\pi(w-1)}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n(w-1)}{w}\right) - \cos\left(\frac{2\pi n 2}{w}\right) \right] \right) - \frac{w}{2} \right] \right) \right] =
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{w^3}{6} \left[\frac{\pi}{4} \left(\cot\left(\frac{\pi(w-2)}{w}\right) - \cot\left(\frac{2\pi}{w}\right) \right) + \frac{w}{4} - \frac{\pi}{2} \left(\cot\left(\frac{2\pi}{w}\right) - \cot\left(\frac{\pi}{w}\right) \right) - \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n}{w}\right) - \cos\left(\frac{2\pi n 2}{w}\right) \right] \right) - \frac{w}{2} \right. \\
 &\quad \left. + \frac{\pi}{2} \left(\cot\left(\frac{2\pi}{w}\right) - \cot\left(\frac{\pi(w-1)}{w}\right) \right) + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n(w-1)}{w}\right) - \cos\left(\frac{2\pi n 2}{w}\right) \right] \right) - \frac{w}{2} \right] =
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{w^3}{6} \left[\frac{\pi}{4} \left(\cot\left(\frac{\pi(w-2)}{w}\right) - \cot\left(\frac{2\pi}{w}\right) + 2 \cot\left(\frac{\pi}{w}\right) - 2 \cot\left(\frac{\pi(w-1)}{w}\right) \right) + \frac{w}{4} - w \right. \\
 &\quad \left. + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) \left[\cos\left(\frac{2\pi n(w-1)}{w}\right) - \cos\left(\frac{2\pi n 2}{w}\right) - \cos\left(\frac{2\pi n}{w}\right) + \cos\left(\frac{2\pi n 2}{w}\right) \right] \right) \right] =
 \end{aligned}$$

$$= \frac{w^3}{6} \left[\frac{\pi}{4} \left(4 \cot\left(\frac{\pi}{w}\right) - 2 \cot\left(\frac{2\pi}{w}\right) \right) - \frac{3w}{4} + \sum_{n=1}^{w-1} \left(\ln\left(2 \sin\left(\frac{\pi n}{w}\right)\right) [0] \right) \right] = \frac{w^3}{24} \left[\pi \left(4 \cot\left(\frac{\pi}{w}\right) - 2 \cot\left(\frac{2\pi}{w}\right) \right) - 3w \right]$$

Finally, applying limits:

$$\begin{aligned}
 \lim_{w \rightarrow \infty} \left\{ \sum_{n=1}^{\infty} \left(\frac{1}{\left(n + (-1) + \frac{w-2}{w} \right) \left(n + (-1) + \frac{w-1}{w} \right) \left(n + \frac{1}{w} \right) \left(n + \frac{2}{w} \right)} \right) \right\} &= \sum_{n=1}^{\infty} \left(\frac{1}{n^4} \right) \\
 &= \lim_{w \rightarrow \infty} \left\{ \frac{w^3}{24} \left[\pi \left(4 \cot\left(\frac{\pi}{w}\right) - 2 \cot\left(\frac{2\pi}{w}\right) \right) - 3w \right] \right\} = \frac{\pi^4}{90}
 \end{aligned}$$

References

- [1] Pietro Mengoli (1650). *Novae quadraturae arithmeticae, seu de additione fractionum*, Bologna.
- [2] J.L. Jensen (1915-1916). *An elementary exposition of the theory of the gamma function*, *Ann. Math.* **17**, 124-166.