

ONE POSITIVE AND TWO NEGATIVE RESULTS FOR DERIVED CATEGORIES OF ALGEBRAIC STACKS

JACK HALL, AMNON NEEMAN, AND DAVID RYDH

ABSTRACT. Let X be a quasi-compact and quasi-separated scheme. There are two fundamental and pervasive facts about the unbounded derived category of X : (1) $D_{\text{qc}}(X)$ is compactly generated by perfect complexes and (2) if X is noetherian or has affine diagonal, then the functor $\Psi_X: D(\text{QCoh}(X)) \rightarrow D_{\text{qc}}(X)$ is an equivalence. Our main results are that for algebraic stacks in positive characteristic, the assertions (1) and (2) are typically false.

1. INTRODUCTION

Fix a field k and an algebraic group G over k . Ben-Zvi posed the following question [BZ09]: if k has positive characteristic, then is the unbounded derived category of representations of G compactly generated?

The second author recently answered Ben-Zvi's question negatively in the case of \mathbb{G}_a [Nee14, Rem. 4.2]. We establish a much stronger version of this result: in the unbounded derived category of representations of \mathbb{G}_a in positive characteristic, there are no compact objects besides 0 (Proposition 3.1).

We say that G is *poor* if k has positive characteristic and $\overline{G} = G \otimes_k \overline{k}$ has a subgroup isomorphic to \mathbb{G}_a , or, equivalently, if $\overline{G}_{\text{red}}^0$ is not semi-abelian (Lemma 4.2). Examples of poor groups are \mathbb{G}_a and GL_n . The results of this article imply that in positive characteristic, the derived category of representations of G is not compactly generated if G is poor. Conversely, when G is not poor, the first and third author show that its derived category of representations is compactly generated [HR14a, Thm. A]. Ben-Zvi's question is thus completely resolved.

A somewhat subtle point that we have suppressed so far is that there are two potential ways to look at the unbounded derived category of representations of G . First, there is $D(\text{Rep}(G))$; second, there is $D_{\text{qc}}(BG)$, the unbounded derived category of lisse-étale \mathcal{O}_{BG} -modules with quasi-coherent cohomology. There is a natural functor $D(\text{Rep}(G)) \rightarrow D_{\text{qc}}(BG)$ and if G is affine, then this functor induces an equivalence on bounded below derived categories.

In the present article, we will show that in positive characteristic if G is affine and poor, then this functor is not full. We also prove that if G is poor, then neither $D(\text{Rep}(G))$ nor $D_{\text{qc}}(BG)$ is compactly generated.

The results above are actually special cases of some general results for unbounded derived categories of quasi-coherent sheaves on algebraic stacks. We say that an algebraic stack is *poorly stabilized* (see §4) if it has a point with poor stabilizer group. Our first main result is the following.

Theorem 1.1. *Let X be an algebraic stack that is quasi-compact, quasi-separated and poorly stabilized.*

- (1) *The triangulated category $D_{\text{qc}}(X)$ is not compactly generated.*

Date: December 3, 2015.

2010 Mathematics Subject Classification. Primary 14F05; secondary 13D09, 14A20, 18G10.

Key words and phrases. Derived categories, algebraic stacks.

The third author is supported by the Swedish Research Council (VR), grant number 2011-5599.

- (2) *If X has affine diagonal or is noetherian and affine-pointed, then $D(\mathrm{QCoh}(X))$ is not compactly generated.*

An algebraic stack X is *affine-pointed* if every morphism $\mathrm{Spec} k \rightarrow X$, where k is a field, is affine. If X has quasi-affine or quasi-finite diagonal, then X is affine-pointed [HR14b, Lem. 4.5].

We wish to point out that Theorem 1.1 is counter to the prevailing wisdom. Indeed, let X be a quasi-compact and quasi-separated algebraic stack. If X is a scheme, then it is well-known that $D_{\mathrm{qc}}(X)$ is compactly generated by perfect complexes [BB03, Thm. 3.1.1(b)]. More generally, recent work of Krishna [Kri09, Lem. 4.8], Ben-Zvi–Francis–Nadler [BZFN10, §3.3], Toën [Toë12, Cor. 5.2], and the first and third authors [HR14c], has shown that the unbounded derived category $D_{\mathrm{qc}}(X)$ is compactly generated by perfect complexes if X is a Deligne–Mumford stack or a stack in characteristic zero that étale-locally has the resolution property [Tot04, Gro13].

Also recall that if X is a scheme that is either quasi-compact with affine diagonal or noetherian, then the functor $\Psi_X: D(\mathrm{QCoh}(X)) \rightarrow D_{\mathrm{qc}}(X)$ is an equivalence of triangulated categories—see [BN93, Cor. 5.5] for the separated case (the argument adapts trivially to the case of affine diagonal) and [Stacks, Tags 08H1 & 09TN] in the setting of algebraic spaces. Our second main result is a partial extension of this to algebraic stacks.

Theorem 1.2. *Let X be an algebraic stack that is quasi-compact with affine diagonal or noetherian and affine-pointed. If $D_{\mathrm{qc}}(X)$ is compactly generated, then the functor $\Psi_X: D(\mathrm{QCoh}(X)) \rightarrow D_{\mathrm{qc}}(X)$ is an equivalence of categories.*

In particular, Ψ_X is an equivalence for every Deligne–Mumford stack with affine diagonal, every noetherian Deligne–Mumford stack, and every stack in characteristic zero with affine diagonal that étale-locally has the resolution property [HR14c]. This is a vast extension of work of Krishna [Kri09, Cor. 3.7], where Ψ_X was proved to be an equivalence for every Deligne–Mumford stack that is separated, of finite type over a field of characteristic 0, has the resolution property and whose coarse moduli space is a scheme.

It is natural to ask whether Ψ_X is always an equivalence of categories. On the positive side, we prove that the restricted functor $\Psi_X^+: D^+(\mathrm{QCoh}(X)) \rightarrow D_{\mathrm{qc}}^+(X)$ is an equivalence of triangulated categories when either X is quasi-compact with affine diagonal or noetherian and affine-pointed (Theorem C.1, also see [Lur04, Thm. 3.8] and [SGA6, Prop. II.3.5]). On the negative side, we have the following result.

Theorem 1.3. *Let X be an algebraic stack that is quasi-compact with affine diagonal or noetherian and affine-pointed. If X is poorly stabilized, then the functor $\Psi_X: D(\mathrm{QCoh}(X)) \rightarrow D_{\mathrm{qc}}(X)$ is not full.*

We were unable to determine whether the functor Ψ_X in Theorem 1.3 is faithful or not. For stacks with non-affine stabilizer groups the situation is even worse: if $X = BE$, where E is an elliptic curve over \mathbb{C} , then the functor $\Psi_X^b: D^b(\mathrm{Coh}(X)) \rightarrow D_{\mathrm{Coh}}^b(X)$ is neither essentially surjective nor full.

Note that when X has affine diagonal or is noetherian and affine-pointed, the first claim in Theorem 1.1 is a trivial consequence of Theorems 1.2 and 1.3.

Left-completeness. In the course of proving Theorem 1.3, we will prove that the triangulated category $D(\mathrm{QCoh}(X))$ is not left-complete whenever X is poorly stabilized with affine diagonal. This generalizes an example of Neeman [Nee11] and amplifies some observations of Drinfeld–Gaitsgory [DG13, Rem. 1.2.10].

In Appendix B, we will prove that $D_{\mathrm{qc}}(X)$ is left-complete for all algebraic stacks X . An analogous assertion in the context of derived algebraic geometry has been

addressed by Drinfeld–Gaitsgory [DG13, Lem. 1.2.8]. In the Stacks Project [Stacks, 081Y] a similar result has been proved, albeit in a different context.

A requirement for a triangulated category to be left-complete is that it admits countable products. We were unable to locate a proof in the literature that $D_{\text{qc}}(X)$ admits countable products, however. Thus we also address this in Appendix B. By [Nee01b, Cor. 1.18], it suffices to prove that $D_{\text{qc}}(X)$ is well generated.

Well generation. If X is an algebraic stack, then it is known that $D_{\text{qc}}(X)$ is well generated in the following cases:

- X is Deligne–Mumford [Lur11, Prop. 2.3.13(3)]; or
- X is quasi-compact with affine diagonal [Lur11, Prop. 3.4.17(1)]; or
- X is a \mathbb{Q} -stack that is quasi-compact and quasi-separated with affine stabilizers and finitely presented inertia [DG13, Thm. 1.4.10].

This list is quite extensive, covering the majority of algebraic stacks encountered in this article and in practice. We are not aware of a reference that $D_{\text{qc}}(X)$ is well generated in complete generality, however.

In Appendix A we show that if \mathcal{A} is a Grothendieck abelian category and $\mathcal{M} \subseteq \mathcal{A}$ is a weak Serre subcategory that is closed under coproducts and is Grothendieck abelian, then $D_{\mathcal{M}}(\mathcal{A})$ is a well generated triangulated category—a result we expect to be of independent interest. Since the inclusion $\text{QCoh}(X) \subseteq \text{Mod}(X)$ has these properties, this establishes that $D_{\text{qc}}(X)$ is well generated.

While we hoped that the well generation of $D_{\mathcal{M}}(\mathcal{A})$, which is essentially a set-theoretic issue, would follow from some general results in the theory of presentable categories (even of the ∞ -category type), we did not succeed along those lines. Instead, we give a direct proof using the Gabriel–Popescu Theorem. We also wish to point out that while [KS06, Prop. 14.2.4] is quite general, it does not apply in our situation. Indeed, they require that the embedding $\mathcal{M} \subseteq \mathcal{A}$ is closed under \mathcal{A} -subquotients (i.e., \mathcal{M} is a Serre subcategory of \mathcal{A}), which is not the case for $\text{QCoh}(X) \subseteq \text{Mod}(X)$.

Note that we do not understand, in general, under what circumstances localizing subcategories of well generated categories are generated by sets of objects. The question goes back to the early 1970s, when Adams claimed to prove that it is possible to localize spectra with respect to arbitrary homology and cohomology theories. Bousfield pointed out the gap in Adams’ argument and half-fixed it: he showed that it is possible to localize spectra with respect to homology, see [Bou75, Bou79].

We restate the problem in the language here: let \mathcal{T} be the homotopy category of spectra, which is well-known to be well generated (even compactly generated). Given a homology theory E and a cohomology theory F , we can define localizing subcategories

$$\mathcal{L}(E) = \{X \in \mathcal{T} : E(X) = 0\}, \quad \mathcal{L}(F) = \{X \in \mathcal{T} : F(X) = 0\}.$$

Bousfield proved that $\mathcal{L}(E)$ has a set of generators, and conjectured that so does $\mathcal{L}(F)$. This conjecture remains open, despite considerable effort over the intervening four decades. Some recent highlights of the literature are [CSS05, BF03, Cho05].

2. PRELIMINARIES

Let $\phi: X \rightarrow Y$ be a quasi-compact and quasi-separated morphism of algebraic stacks. Then the restriction of the functor $(\phi_{\text{lis-ét}})_*: \text{Mod}(X) \rightarrow \text{Mod}(Y)$ to $\text{QCoh}(X)$ factors through $\text{QCoh}(Y)$ [Ols07, Lem. 6.5(i)], giving rise to a functor $(\phi_{\text{QCoh}})_*: \text{QCoh}(X) \rightarrow \text{QCoh}(Y)$. Since the categories $\text{Mod}(X)$ and $\text{QCoh}(X)$ are Grothendieck abelian [Stacks, 0781], the unbounded derived functors of $(\phi_{\text{lis-ét}})_*$ and $(\phi_{\text{QCoh}})_*$ exist [Stacks, 079P & 070K], and we denote these as $R(\phi_{\text{lis-ét}})_*$ and

$R(\phi_{\text{QCoh}})_*$, respectively. By [Ols07, Lem. 6.20], the restriction of $R(\phi_{\text{lis-ét}})_*$ to $D_{\text{qc}}^+(X)$ factors uniquely through $D_{\text{qc}}^+(Y)$. If, in addition, ϕ is concentrated (e.g., representable), then the restriction of $R(\phi_{\text{lis-ét}})_*$ to $D_{\text{qc}}(X)$ factors through $D_{\text{qc}}(Y)$ (see [Hal14, Lem. 2.1] for the representable case and [HR14c, Thm. 2.6(2)] in general).

For an algebraic stack W let $\Psi_W: D(\text{QCoh}(W)) \rightarrow D_{\text{qc}}(W)$ denote the natural functor. The universal properties of right-derived functors provide a diagram:

$$\begin{array}{ccc} D(\text{QCoh}(X)) & \xrightarrow{R(\phi_{\text{QCoh}})_*} & D(\text{QCoh}(Y)) \\ \Psi_X \downarrow & & \downarrow \Psi_Y \\ D_{\text{qc}}(X) & \xrightarrow{R(\phi_{\text{lis-ét}})_*} & D_{\text{qc}}(Y), \end{array}$$

together with a natural transformation of functors:

$$(2.1) \quad \epsilon_\phi: \Psi_Y \circ R(\phi_{\text{QCoh}})_* \Rightarrow R(\phi_{\text{lis-ét}})_* \circ \Psi_X.$$

The following result, for schemes, is well-known [TT90, B.8]; for algebraic spaces, see [Stacks, Tags 09TH & 08GX].

Proposition 2.1. *Let $\phi: X \rightarrow Y$ be a morphism of algebraic stacks. Suppose that both X and Y are quasi-compact with affine diagonal or noetherian and affine-pointed. If $M \in D^+(\text{QCoh}(X))$, then the morphism induced by (2.1):*

$$\epsilon_\phi(M): \Psi_Y \circ R(\phi_{\text{QCoh}})_*(M) \rightarrow R(\phi_{\text{lis-ét}})_* \circ \Psi_X(M)$$

is an isomorphism. In particular, since $\Psi_Y^+: D^+(\text{QCoh}(Y)) \rightarrow D_{\text{qc}}^+(Y)$ is an equivalence (Theorem C.1), it follows that there is a natural isomorphism for each $M \in D^+(\text{QCoh}(X))$:

$$R(\phi_{\text{QCoh}})_*(M) \rightarrow (\Psi_Y^+)^{-1} \circ R(\phi_{\text{lis-ét}})_* \circ \Psi_X^+(M).$$

Proof. The functors $(\phi_{\text{QCoh}})_*$ and $(\phi_{\text{lis-ét}})_*$ are left-exact, thus the functors $R(\phi_{\text{QCoh}})_*$ and $R(\phi_{\text{lis-ét}})_*$ are bounded below. Via standard “way-out” arguments, one readily reduces to proving the isomorphism above in the case $M \simeq N[0]$, where $N \in \text{QCoh}(X)$. The isomorphism, in this case, reduces to proving that if $N \in \text{QCoh}(X)$, then the natural morphism $R^i(\phi_{\text{QCoh}})_*N \rightarrow R^i(\phi_{\text{lis-ét}})_*N$ is an isomorphism for all integers $i \geq 0$, where $R^i(\phi_{\text{QCoh}})_*$ (resp. $R^i(\phi_{\text{lis-ét}})_*$) denotes the i th right-derived functor of $(\phi_{\text{QCoh}})_*$ (resp. $(\phi_{\text{lis-ét}})_*$). A standard δ -functor argument shows that it is sufficient to prove that $R^i(\phi_{\text{lis-ét}})_*I = 0$ for every $i > 0$ and injective I of $\text{QCoh}(X)$. This now follows from Lemma C.3(2). \square

Corollary 2.2. *Let $\phi: X \rightarrow Y$ be a concentrated morphism of algebraic stacks. If X and Y are quasi-compact with affine diagonal or noetherian and affine-pointed, then there exists an integer $r \geq 0$ such that for all $M \in D(\text{QCoh}(X))$ and integers n the natural map:*

$$\tau^{\geq n} R(\phi_{\text{QCoh}})_*M \rightarrow \tau^{\geq n} R(\phi_{\text{QCoh}})_* \tau^{\geq n-r} M$$

is a quasi-isomorphism. It follows that

- (1) $R(\phi_{\text{QCoh}})_*$ preserves small coproducts,
- (2) for all $M \in D(\text{QCoh}(X))$ the natural morphism induced by (2.1):

$$\epsilon_\phi(M): \Psi_Y \circ R(\phi_{\text{QCoh}})_*M \rightarrow R(\phi_{\text{lis-ét}})_* \circ \Psi_X(M)$$

is an isomorphism, and

- (3) if ϕ is quasi-affine, then $R(\phi_{\text{QCoh}})_*$ is conservative.

Proof. The claims (1)–(3) are all simple consequences of the main claim and Proposition 2.1. Since ϕ is a concentrated morphism and Y is quasi-compact and quasi-separated, there exists an integer $r \geq 0$ such that if $N \in \mathbf{QCoh}(X)$, then $R^i(\phi_{\text{lis-ét}})_* N = 0$ for all $i > r$. By Proposition 2.1 it follows that $R^i(\phi_{\mathbf{QCoh}})_* N = 0$ for all $i > r$ too. The result now follows from [Stacks, 07K7]. \square

Corollary 2.3. *Let X be an algebraic stack that is quasi-compact with affine diagonal or noetherian and affine-pointed. If C is a compact object of either $\mathbf{D}(\mathbf{QCoh}(X))$ or $\mathbf{D}_{\text{qc}}(X)$, then C is perfect. Moreover if X is noetherian, then C is quasi-isomorphic to a bounded complex of coherent sheaves on X .*

Proof. Let C be a compact object of $\mathbf{D}_{\text{qc}}(X)$. By [HR14c, Lem. 4.4(1)], C is a perfect complex and in particular belongs to $\mathbf{D}_{\text{qc}}^b(X) \subseteq \mathbf{D}_{\text{qc}}^+(X)$. By Theorem C.1, it follows that $C \simeq \Psi_X(\tilde{C})$ for some $\tilde{C} \in \mathbf{D}(\mathbf{QCoh}(X))$. If X is noetherian, \tilde{C} even belongs to $\mathbf{D}_{\text{Coh}(X)}^b(\mathbf{QCoh}(X))$. Combining [LMB, Prop. 15.4] with [SGA6, II.2.2], we deduce that C belongs to the image of $\mathbf{D}(\mathbf{Coh}(X)) \rightarrow \mathbf{D}_{\text{qc}}(X)$.

Now let C be a compact object of $\mathbf{D}(\mathbf{QCoh}(X))$. Let $p: U \rightarrow X$ be a smooth surjection from an affine scheme U . The functor $p_{\mathbf{QCoh}}^*: \mathbf{QCoh}(X) \rightarrow \mathbf{QCoh}(U)$ is exact and gives rise to a derived functor $\mathbf{L}p_{\mathbf{QCoh}}^*: \mathbf{D}(\mathbf{QCoh}(X)) \rightarrow \mathbf{D}(\mathbf{QCoh}(U))$. Its right adjoint $R(p_{\mathbf{QCoh}})_*$ preserves small coproducts by Corollary 2.2(1), so $\mathbf{L}p_{\mathbf{QCoh}}^* C \in \mathbf{D}(\mathbf{QCoh}(U))$ is compact. Since $U = \text{Spec } A$ is affine, it follows that $\mathbf{QCoh}(U) \cong \text{Mod}(A)$ and so $\mathbf{L}p_{\mathbf{QCoh}}^* C$ is a perfect complex [Stacks, 07LT]. If X is noetherian, then $C \in \mathbf{D}_{\text{Coh}(X)}^b(\mathbf{QCoh}(X))$. Arguing as before, we deduce that C belongs to the image of $\mathbf{D}(\mathbf{Coh}(X)) \rightarrow \mathbf{D}(\mathbf{QCoh}(X))$. \square

In the following Lemma we will give a sufficient condition for compactness of a perfect object in $\mathbf{D}(\mathbf{QCoh}(X))$. We do not know if this condition is necessary. The analogous condition in $\mathbf{D}_{\text{qc}}(X)$ is necessary [HR14c, Lem. 4.5].

Lemma 2.4. *Let X be an algebraic stack that is quasi-compact with affine diagonal or noetherian and affine-pointed. Let $P \in \mathbf{D}(\mathbf{QCoh}(X))$ be a perfect complex. Consider the following conditions*

- (1) P is a compact object of $\mathbf{D}(\mathbf{QCoh}(X))$.
- (2) There exists an integer $r \geq 0$ such that $\text{Hom}_{\mathcal{O}_X}(P, N[i]) = 0$ for all $N \in \mathbf{QCoh}(X)$ and $i > r$.
- (3) There exists an integer $r \geq 0$ such that the natural map

$$\tau^{\geq j} \mathbf{RHom}_{\mathbf{D}(\mathbf{QCoh}(X))}(P, M) \rightarrow \tau^{\geq j} \mathbf{RHom}_{\mathbf{D}(\mathbf{QCoh}(X))}(P, \tau^{\geq j-r} M)$$

is a quasi-isomorphism for all $M \in \mathbf{D}(\mathbf{QCoh}(X))$ and integers j .

Then (2) and (3) are equivalent and imply (1).

Proof. Condition (2) is a special case of (3): let $M = N[i]$ and $j = 0$.

Conversely, assume that condition (2) holds and let $M \in \mathbf{D}(\mathbf{QCoh}(X))$. Since $\mathbf{QCoh}(X)$ is Grothendieck abelian, there is a quasi-isomorphism $M \rightarrow I^\bullet$ in $\mathbf{D}(\mathbf{QCoh}(X))$, where I^\bullet is K-injective and I^j is injective for every integer j [Ser03].

Let $p \geq r + 1$ be an integer with the property that $P \in \mathbf{D}^{\geq -p+1}(\mathbf{QCoh}(X))$. Then the natural morphism of chain complexes:

$$(2.2) \quad \tau^{\geq j} \text{Hom}_{\mathbf{K}(\mathbf{QCoh}(X))}^\bullet(P, I^\bullet) \rightarrow \tau^{\geq j} \text{Hom}_{\mathbf{K}(\mathbf{QCoh}(X))}^\bullet(P, \sigma^{\geq j-p} I^\bullet),$$

where σ is the brutal truncation, is a quasi-isomorphism. For every integer j there is also a morphism $s_j: \sigma^{\geq j} I^\bullet \rightarrow \tau^{\geq j} M$. If C_j^\bullet is the mapping cone of s_j , then $C_j^\bullet \simeq d(I^{j-1})[-(j-1)]$. Thus, by condition (2), we have for every integer j

$$\tau^{\geq j+r} \mathbf{RHom}_{\mathbf{D}(\mathbf{QCoh}(X))}(P, C_j^\bullet) \simeq 0.$$

Since there is also distinguished triangle for every integer j :

$$\mathrm{RHom}_{\mathcal{O}_X}(P, \sigma^{\geq j-p} I^\bullet) \rightarrow \mathrm{RHom}_{\mathcal{O}_X}(P, \tau^{\geq j-p} M) \rightarrow \mathrm{RHom}_{\mathcal{O}_X}(P, C_{j-p}^\bullet),$$

it follows that for every integer j there is a quasi-isomorphism:

$$\tau^{\geq j} \mathrm{RHom}_{\mathcal{O}_X}(P, \sigma^{\geq j-p} I^\bullet) \simeq \tau^{\geq j} \mathrm{RHom}_{\mathcal{O}_X}(P, \tau^{\geq j-p} M).$$

For every integer j , we also have a distinguished triangle

$$H^{j-r-1}(M)[- (j-r-1)] \rightarrow \tau^{\geq j-r-1} M \rightarrow \tau^{\geq j-r} M.$$

As before, it follows that $\tau^{\geq j} \mathrm{RHom}_{\mathcal{O}_X}(P, \tau^{\geq j-r-1} M) \simeq \tau^{\geq j} \mathrm{RHom}_{\mathcal{O}_X}(P, \tau^{\geq j-r} M)$ and thus by induction:

$$\tau^{\geq j} \mathrm{RHom}_{\mathcal{O}_X}(P, \tau^{\geq j-p} M) \rightarrow \tau^{\geq j} \mathrm{RHom}_{\mathcal{O}_X}(P, \tau^{\geq j-r} M).$$

Combining these quasi-isomorphisms with (2.2) gives (3).

For (3) implies (1): this follows from Theorem C.1 and [HR14c, Lem. 1.2(3)]. \square

We now relate compact generation in $\mathrm{D}(\mathrm{QCoh}(X))$ with compact generation in $\mathrm{D}_{\mathrm{qc}}(X)$.

Lemma 2.5. *Let X be an algebraic stack that is quasi-compact with affine diagonal or noetherian and affine-pointed.*

- (1) *If $P \in \mathrm{D}(\mathrm{QCoh}(X))$ is a perfect complex such that $\Psi(P)$ is compact in $\mathrm{D}_{\mathrm{qc}}(X)$, then P is compact in $\mathrm{D}(\mathrm{QCoh}(X))$.*
- (2) *If X has finite cohomological dimension, then every perfect complex is compact in both $\mathrm{D}(\mathrm{QCoh}(X))$ and $\mathrm{D}_{\mathrm{qc}}(X)$.*
- (3) *If a set of objects $\{P_i\}$ of $\mathrm{D}(\mathrm{QCoh}(X))$ has the property that $\{\Psi(P_i)\}$ compactly generates $\mathrm{D}_{\mathrm{qc}}(X)$, then $\{P_i\}$ compactly generates $\mathrm{D}(\mathrm{QCoh}(X))$.*

Proof. For (1), by [HR14c, Lem. 4.5], since $\Psi(P)$ is compact, there exists an integer r such that if $i > r$ and $N \in \mathrm{QCoh}(X)$, then $\mathrm{Hom}_{\mathcal{O}_X}(\Psi(P), N[i]) = 0$. The functor Ψ^+ is an equivalence (Theorem C.1), so $\mathrm{Hom}_{\mathcal{O}_X}(P, N[i]) = 0$ for all $i > r$ and $N \in \mathrm{QCoh}(X)$. It follows that P is compact by Lemma 2.4.

Statement (2) is a direct consequence of (1) and [HR14c, Lem. 4.4(3)].

For (3), let $M \in \mathrm{D}(\mathrm{QCoh}(X))$. If P is perfect and $\Psi(P)$ is compact, then $\mathrm{RHom}(P, M) = \mathrm{RHom}(\Psi(P), \Psi(M))$. Indeed, there exists an integer r such that for all integers j

$$\begin{aligned} \tau^{\geq j} \mathrm{RHom}(P, M) &\simeq \tau^{\geq j} \mathrm{RHom}(P, \tau^{\geq j-r} M) \\ &\simeq \tau^{\geq j} \mathrm{RHom}(\Psi(P), \tau^{\geq j-r} \Psi(M)) \simeq \tau^{\geq j} \mathrm{RHom}(\Psi(P), \Psi(M)), \end{aligned}$$

by Lemma 2.4 and [HR14c, Lem. 4.5] since Ψ^+ is an equivalence of triangulated categories (Theorem C.1) and Ψ is t-exact. Thus, if $\mathrm{Hom}_{\mathrm{D}(\mathrm{QCoh}(X))}(P_i[l], M) = 0$ for all i and integers l , then $\mathrm{Hom}_{\mathcal{O}_X}(\Psi(P_i)[l], \Psi(M)) = 0$ for all i and l . It follows that $\Psi(M) = 0$ and, since Ψ is conservative, that $M = 0$. \square

The following lemma, while technical, gives an explicit description of an adjunction that is useful in the article.

Lemma 2.6. *Let X be an algebraic stack and let $M \in \mathrm{D}(\mathrm{QCoh}(X))$.*

- (1) *The functor $\Psi_X: \mathrm{D}(\mathrm{QCoh}(X)) \rightarrow \mathrm{D}_{\mathrm{qc}}(X)$ admits a right adjoint $\Phi_X: \mathrm{D}_{\mathrm{qc}}(X) \rightarrow \mathrm{D}(\mathrm{QCoh}(X))$.*
- (2) *If X is a quasi-compact with affine diagonal or noetherian and affine-pointed, then there exists a compatible quasi-isomorphism:*

$$\Phi_X \Psi_X(M) \simeq \mathrm{holim}_n \tau^{\geq -n} M.$$

Proof. We suppress the subscript X from Ψ and Φ throughout. Since Ψ preserves small coproducts and $\mathrm{D}(\mathrm{QCoh}(X))$ is well generated [Nee01a, Thm. 0.2], Ψ admits a right adjoint $\Phi: \mathrm{D}_{\mathrm{qc}}(X) \rightarrow \mathrm{D}(\mathrm{QCoh}(X))$ [Nee01b, Prop. 1.20]. This proves (1).

To prove (2), by left-completeness of $\mathrm{D}_{\mathrm{qc}}(X)$ (Theorem B.1),

$$\Phi\Psi(M) \rightarrow \Phi(\mathrm{holim}_n \tau^{\geq -n}\Psi(M))$$

is a quasi-isomorphism. Since Φ is a right adjoint, it preserves homotopy limits. Also, Ψ is t-exact. Hence, there is a quasi-isomorphism

$$\Phi(\mathrm{holim}_n \tau^{\geq -n}\Psi(M)) \simeq \mathrm{holim}_n \Phi\Psi(\tau^{\geq -n}M).$$

By Theorem C.1, however, $\tau^{\geq -n}M \simeq \Phi\Psi(\tau^{\geq -n}M)$. This proves the claim. \square

Remark 2.7. From Lemma 2.6(2) it is immediate that when X is quasi-compact with affine diagonal or noetherian and affine-pointed, the left-completeness of $\mathrm{D}(\mathrm{QCoh}(X))$ is equivalent to Ψ_X being fully faithful.

We now prove Theorem 1.2 using an argument similar to [BIK11, Lem. 4.5].

Proof of Theorem 1.2. By Lemma 2.5(3), both $\mathrm{D}(\mathrm{QCoh}(X))$ and $\mathrm{D}_{\mathrm{qc}}(X)$ are compactly generated and Ψ takes a set of compact generators to a set of compact generators. In particular, the right adjoint $\Phi: \mathrm{D}_{\mathrm{qc}}(X) \rightarrow \mathrm{D}(\mathrm{QCoh}(X))$ of Ψ preserves small coproducts [Nee96, Thm. 5.1].

Consider the unit $\eta_M: M \rightarrow \Phi\Psi(M)$ and the counit $\epsilon_M: \Psi\Phi(M) \rightarrow M$ of the adjunction. Since Ψ^+ is an equivalence, we have that η_P and ϵ_P are isomorphisms for every compact object P . Since η and ϵ are triangulated functors that preserve small coproducts and $\mathrm{D}_{\mathrm{qc}}(X)$ and $\mathrm{D}(\mathrm{QCoh}(X))$ are compactly generated, it follows that η and ϵ are equivalences. We conclude that Ψ is an equivalence. \square

3. THE CASE OF $B_k\mathbb{G}_a$ IN POSITIVE CHARACTERISTIC

Throughout this section we let k denote a field of characteristic $p > 0$. Let $B_k\mathbb{G}_a$ be the algebraic stack classifying \mathbb{G}_a -torsors over k . We remind ourselves that the category of quasi-coherent sheaves on $B_k\mathbb{G}_a$ is the category of \mathbb{G}_a -modules, which is equivalent to the category of locally small modules over a certain ring R . In fact R is the ring

$$R = \frac{k[x_1, x_2, x_3, \dots]}{(x_1^p, x_2^p, x_3^p, \dots)}$$

and a module is locally small if every element is annihilated by all but finitely many x_i . Let us write $\mathrm{D}(R^{\mathrm{ls}})$ for the derived category of the category of locally small R -modules, and observe that $\mathrm{D}(R^{\mathrm{ls}}) \cong \mathrm{D}(\mathrm{QCoh}(B_k\mathbb{G}_a))$.

Proposition 3.1. *The only compact objects, in either $\mathrm{D}(\mathrm{QCoh}(B_k\mathbb{G}_a))$ or $\mathrm{D}_{\mathrm{qc}}(B_k\mathbb{G}_a)$, are the zero objects.*

Proof. The algebraic stack $B_k\mathbb{G}_a$ is noetherian with affine diagonal and so, by Corollary 2.3, every compact object is the image of a bounded complex of coherent sheaves. Let C be a compact object; we need to show that C vanishes.

Our compact object C is the image of a finite complex of finitely generated modules in $\mathrm{D}(R^{\mathrm{ls}})$. In particular, there exists an integer $n > 1$ such that x_i annihilates C for all $i \geq n$. Let us put this slightly differently: consider the ring homomorphisms $S \xrightarrow{\alpha} T \xrightarrow{\beta} R \xrightarrow{\gamma} T$ where

$$S = k[x_n]/(x_n^p), \quad T = \frac{k[x_1, x_2, \dots, x_{n-1}, x_n]}{(x_1^p, x_2^p, \dots, x_{n-1}^p, x_n^p)}$$

where the maps $S \xrightarrow{\alpha} T \xrightarrow{\beta} R$ are the natural inclusions, and where $\gamma: R \rightarrow T$ is defined by

$$\gamma(x_i) = \begin{cases} x_i & \text{if } i \leq n \\ 0 & \text{if } i > n. \end{cases}$$

Note that $\gamma\beta = \text{id}$. Restriction of scalars gives induced maps of derived categories, which we write as $\mathbf{D}(T) \xrightarrow{\gamma_*} \mathbf{D}(R^{\text{ls}}) \xrightarrow{\beta_*} \mathbf{D}(T) \xrightarrow{\alpha_*} \mathbf{D}(S)$, and $\beta_*\gamma_* = \text{id}$. Our complex C , which is a bounded complex annihilated by x_i for all $i \geq n$, is of the form γ_*B where $B \in \mathbf{D}^b(T)$ is a bounded complex of finite T -modules. And the fact that x_n annihilates C translates to saying that α_*B is a complex of modules annihilated by x_n , that is a complex of k -vector spaces. We wish to show that $C = 0$ or, equivalently, that α_*B is acyclic. We will show that if C is non-zero, then this gives rise to a contradiction.

Thus, assume that the cohomology of α_*B is non-trivial: in $\mathbf{D}(S)$ the complex α_*B is isomorphic to a non-zero sum of suspensions $k[\ell]$ of k . Then there are infinitely many integers m and non-zero maps in $\mathbf{D}(S)$ of the form $\alpha_*B \rightarrow k[m]$. Indeed, $\text{Ext}_S^m(k, k) \neq 0$ for all $m \geq 0$. But α_* has a right adjoint $\alpha^\times = \text{RHom}_S(T, -)$, and we deduce infinitely many non-zero maps in $\mathbf{D}(T)$ of the form $B \rightarrow \alpha^\times k[m] = \text{Hom}_S(T, k)[m]$. Since $\mathbf{D}(T)$ is left-complete, these combine to a map in $\mathbf{D}(T)$

$$B \xrightarrow{\Psi} \prod_m \text{Hom}_S(T, k)[m] \cong \prod_m \text{Hom}_S(T, k)[m]$$

for which the composites

$$B \xrightarrow{\Psi} \prod_m \text{Hom}_S(T, k)[m] \xrightarrow{\pi_m} \text{Hom}_S(T, k)[m]$$

are non-zero. Applying γ_* , which preserves coproducts, we deduce maps

$$\gamma_*B \xrightarrow{\gamma_*\Psi} \prod_m \gamma_* \text{Hom}_S(T, k)[m] \xrightarrow{\gamma_*\pi_m} \gamma_* \text{Hom}_S(T, k)[m]$$

whose composites cannot vanish in $\mathbf{D}(R^{\text{ls}})$, since β_* takes them to non-zero maps. The equivalence $\mathbf{D}(R^{\text{ls}}) \cong \mathbf{D}(\text{QCoh}(B_k\mathbb{G}_a))$ gives us that the composites in $\mathbf{D}(\text{QCoh}(B_k\mathbb{G}_a))$ do not vanish. Furthermore, the composites lie in $\mathbf{D}^+(\text{QCoh}(B_k\mathbb{G}_a)) \subseteq \mathbf{D}(\text{QCoh}(B_k\mathbb{G}_a))$, and on $\mathbf{D}^+(\text{QCoh}(B_k\mathbb{G}_a))$ the map to $\mathbf{D}_{\text{qc}}(B_k\mathbb{G}_a)$ is fully faithful [Lur04, Thm. 3.8]. Hence the images of the composites are non-zero in $\mathbf{D}_{\text{qc}}(B_k\mathbb{G}_a)$ as well. But this contradicts the compactness of $C = \gamma_*B$. \square

4. THE GENERAL CASE

In this section we extend the results of the previous section and show that the presence of \mathbb{G}_a in the stabilizer groups of an algebraic stack X is an obstruction to compact generation in positive characteristic. The existence of finite unipotent subgroups such as $\mathbb{Z}/p\mathbb{Z}$ and α_p is an obstruction to the compactness of the structure sheaf \mathcal{O}_X but does not rule out compact generation [HR14a]. The only connected groups in characteristic p without unipotent subgroups are the groups of multiplicative type. The following well-known lemma characterizes the groups without \mathbb{G}_a 's.

Lemma 4.1. *Let G be a group scheme of finite type over an algebraically closed field k . Then the following are equivalent:*

- (1) G_{red}^0 is semiabelian, that is, a torus or the extension of an abelian variety by a torus;
- (2) there is no subgroup $\mathbb{G}_a \hookrightarrow G$.

Proof. By Chevalley's Theorem [Con02, Thm. 1.1] there is an extension $1 \rightarrow H \rightarrow G_{\text{red}}^0 \rightarrow A \rightarrow 1$ where H is smooth, affine and connected and A is an abelian variety. A subgroup $\mathbb{G}_a \hookrightarrow G$ would have to be contained in H which implies that H is not a torus. Conversely, recall that $H(k)$ is generated by its semi-simple and unipotent elements by the Jordan Decomposition Theorem [Bor91, Thm. 4.4]. If H is not a torus, then there exist non-trivial unipotent elements in $H(k)$. But any non-trivial unipotent element of $H(k)$ lies in a subgroup $\mathbb{G}_a \hookrightarrow G$. The result follows. \square

If k is of positive characteristic, then we say that G is poor if G_{red}^0 is not semi-abelian. We say that an algebraic stack X is *poorly stabilized* if there exists a geometric point x of X whose residue field $\kappa(x)$ is of characteristic $p > 0$ and stabilizer group scheme G_x is poor. In particular, the algebraic stacks $B_k\mathbb{G}_a$ and $B_k\text{GL}_n$ for $n > 1$ are poorly stabilized in positive characteristic. The following characterization of poorly stabilized algebraic stacks will be useful.

Lemma 4.2. *Let X be a quasi-separated algebraic stack.*

- (1) *The stack X is poorly stabilized if and only if there exists a field k of characteristic $p > 0$ and a representable morphism $\phi: B_k\mathbb{G}_a \rightarrow X$.*
- (2) *If X has affine stabilizers, then every representable morphism $\phi: B_k\mathbb{G}_a \rightarrow X$ is quasi-affine.*

Proof. Let k be an algebraically closed field and let $x: \text{Spec } k \rightarrow X$ be a geometric point with stabilizer group scheme G . This induces a representable morphism $BG \rightarrow X$. If X is poorly stabilized, then there exists a point x such that G_{red}^0 is not semiabelian. By the previous lemma, there is a subgroup $\mathbb{G}_a \hookrightarrow G$ and hence a representable morphism $B\mathbb{G}_a \rightarrow BG$.

Conversely, given a representable morphism $\phi: B_k\mathbb{G}_a \rightarrow X$, there is an induced representable morphism $\psi: B_k\mathbb{G}_a \rightarrow B_kG$. The morphism ψ is induced by some subgroup $\mathbb{G}_a \hookrightarrow G$ (unique up to conjugation) so X is poorly stabilized.

The structure morphism $\iota_x: \mathcal{G}_x \hookrightarrow X$ of the residual gerbe \mathcal{G}_x at x is quasi-affine [Ryd11, Thm. B.2] and $\phi = \iota_x \circ \rho \circ \psi$ where $\rho: B_kG \rightarrow \mathcal{G}_x$ is affine. If X has affine stabilizers, then G is affine and it follows that the quotient G/\mathbb{G}_a is quasi-affine since \mathbb{G}_a is unipotent [Ros61, Thm. 3]. We conclude that the morphism $\psi: B_k\mathbb{G}_a \rightarrow B_kG$, as well as ϕ , is quasi-affine. \square

We now prove Theorems 1.1 and 1.3.

Proof of Theorem 1.1. By Lemma 4.2, there exists a field of characteristic $p > 0$ and a quasi-compact, quasi-separated and representable morphism $\phi: B_k\mathbb{G}_a \rightarrow X$.

If $D_{\text{qc}}(X)$ is compactly generated, then there is a compact object $M \in D_{\text{qc}}(X)$ and a non-zero map $M \rightarrow R(\phi_{\text{lis-ét}})_* \mathcal{O}_{B_k\mathbb{G}_a}$. Indeed, $R(\phi_{\text{lis-ét}})_* \mathcal{O}_{B_k\mathbb{G}_a} \in D_{\text{qc}}(X)$ and is non-zero. By adjunction, there is a non-zero map $L\phi^*M \rightarrow \mathcal{O}_{B_k\mathbb{G}_a}$. But the functor $L\phi^*$ sends compact objects of $D_{\text{qc}}(X)$ to compact objects of $D_{\text{qc}}(B_k\mathbb{G}_a)$ [HR14c, Ex. 3.8 & Thm. 2.6(3)]. By Proposition 3.1, it follows that $L\phi^*M \simeq 0$ and we have a contradiction. Hence $D_{\text{qc}}(X)$ is not compactly generated.

If X has affine diagonal or is noetherian and affine-pointed, and $D(\text{QCoh}(X))$ is compactly generated, then there is a compact object $Q \in D(\text{QCoh}(X))$ and a non-zero map $Q \rightarrow R(\phi_{\text{QCoh}})_* \mathcal{O}_{B_k\mathbb{G}_a}$. Indeed, $R(\phi_{\text{QCoh}})_* \mathcal{O}_{B_k\mathbb{G}_a} \in D(\text{QCoh}(X))$ and is non-zero. Since $(\phi_{\text{QCoh}})_*$ preserves small products, $R(\phi_{\text{QCoh}})_*$ preserves small products. Furthermore, $D(\text{QCoh}(B_k\mathbb{G}_a))$ is well-generated and so $R(\phi_{\text{QCoh}})_*$ admits a left adjoint $F: D(\text{QCoh}(X)) \rightarrow D(\text{QCoh}(B_k\mathbb{G}_a))$. By Corollary 2.2(1) and [HR14c, Ex. 3.8], the functor F preserves compact objects. Hence, Proposition 3.1 implies that $F(Q) \simeq 0$. We have a contradiction and $D(\text{QCoh}(X))$ is not compactly generated. \square

Proof of Theorem 1.3. By Lemma 4.2, there exists a field of characteristic $p > 0$ and a quasi-affine morphism $\phi: B_k\mathbb{G}_a \rightarrow X$. By Corollary 2.2, there exists an integer $n \geq 1$ such that if $N \in \mathrm{QCoh}(B_k\mathbb{G}_a)$, then $R(\phi_{\mathrm{QCoh}})_*N \in \mathrm{D}^{[0, n-1]}(\mathrm{QCoh}(X))$. By [Nee11, Thm. 1.1], there exists $M \in \mathrm{QCoh}(B_k\mathbb{G}_a)$ such that the natural map in $\mathrm{D}(\mathrm{QCoh}(B_k\mathbb{G}_a))$:

$$\bigoplus_{i \geq 0} M[in] \rightarrow \prod_{i \geq 0} M[in]$$

is not a quasi-isomorphism—note that while [Nee11, Thm. 1.1] only proves the above assertion in the case where $n = 1$, a simple argument by induction on n gives the claim above. Corollary 2.2(3) now implies that the natural map:

$$\bigoplus_{i \geq 0} R(\phi_{\mathrm{QCoh}})_*M[in] \rightarrow \prod_{i \geq 0} R(\phi_{\mathrm{QCoh}})_*M[in]$$

is not a quasi-isomorphism. Since $R(\phi_{\mathrm{QCoh}})_*M \in \mathrm{D}^{[0, n-1]}(\mathrm{QCoh}(X))$, it follows that $\mathrm{D}(\mathrm{QCoh}(X))$ is not left-complete. By Remark 2.7, we have established that Ψ_X is not fully faithful. To prove that Ψ_X is not full, we will have to argue further.

Let $L = R(\phi_{\mathrm{QCoh}})_*M$, $S = \bigoplus_{i \geq 0} L[in]$, and $P = \prod_{i \geq 0} L[in]$. Also, $\Phi_X\Psi_X(S) \simeq P$ (Lemma 2.6(2)). If Ψ_X is full, then there exists a map $P \rightarrow S$ such that the induced map $P \rightarrow S \rightarrow \Phi_X\Psi_X(S) \simeq P$ is the identity morphism. That is, P is a direct summand of S . Since $\prod_{i \geq 0} M[in]$ is not bounded above [Nee11, Rem. 1.2] and ϕ is quasi-affine, it follows that P is not bounded above. But S is bounded above, so P cannot be a direct summand of S ; hence, we have a contradiction and Ψ_X is not full. \square

APPENDIX A. $\mathrm{D}_{\mathcal{M}}(\mathcal{A})$ IS WELL GENERATED

We begin with a general lemma.

Lemma A.1. *Let \mathcal{T} be a well generated triangulated category and let $\mathcal{S} \subseteq \mathcal{T}$ be a localizing subcategory. The category \mathcal{S} is well generated if and only if there is a set of generators in \mathcal{S} . That is: \mathcal{S} is well generated if and only if there is a set of objects $S \subseteq \mathcal{S}$ such that any nonzero object $y \in \mathcal{S}$ admits a nonzero map $s \rightarrow y$, with $s \in S$.*

Proof. If \mathcal{S} is well generated then it has a set of generators S satisfying a bunch of properties, one of which is that S detects nonzero objects—see the definitions in [Nee01b, pp. 273-274]. What needs proof is the reverse implication.

Suppose therefore that \mathcal{S} contains a set of objects S as in the Lemma, that is every nonzero object $y \in \mathcal{S}$ admits a nonzero map $s \rightarrow y$ with $s \in S$. By [Nee01b, Prop. 8.4.2] the set S is contained in \mathcal{T}^α for some regular cardinal α . If $\mathcal{L} = \mathrm{Loc}(S)$ is the localizing subcategory generated by S then [Nee01b, Thm. 4.4.9] informs us that \mathcal{L} is well generated. Since $S \subseteq \mathcal{S}$ and \mathcal{S} is localizing it follows that $\mathcal{L} \subseteq \mathcal{S}$.

We know that \mathcal{L} is well generated; to finish the proof it suffices to show that the inclusion $\mathcal{L} \subseteq \mathcal{S}$ is an equality. In any case the inclusion is a coproduct-preserving functor from the well generated category \mathcal{L} and must have a right adjoint. For every object $y \in \mathcal{S}$, [Nee01b, Prop. 9.1.8] tells us that there is a triangle in \mathcal{S}

$$x \longrightarrow y \longrightarrow z \longrightarrow \Sigma x$$

with $x \in \mathcal{L}$ and $z \in \mathcal{L}^\perp$. Since $z \in \mathcal{L}^\perp \subseteq S^\perp$ we have that the morphisms $s \rightarrow z$, with $s \in S$, all vanish. By the hypothesis of the Lemma it follows that $z = 0$, and hence $y \cong x$ belongs to \mathcal{L} . \square

Remark A.2. We specialize Lemma A.1 to the situation where $\mathcal{T} = \mathrm{D}(\mathcal{A})$ is the derived category of a Grothendieck abelian category \mathcal{A} ; by [Nee01a, Thm. 0.2] we know that \mathcal{T} is well generated, and Lemma A.1 informs us that a localizing subcategory of $\mathrm{D}(\mathcal{A})$ is well generated if and only if it has a set of generators.

Let \mathcal{A} be an abelian category and fix a fully faithful subcategory $\mathcal{C} \subseteq \mathcal{A}$. Following [Stacks, Tag 02MO] we say that

- (1) \mathcal{C} is a *Serre* subcategory if it is non-empty and if $C_1 \rightarrow A \rightarrow C_2$ is an exact sequence in \mathcal{A} with $C_1, C_2 \in \mathcal{C}$, then $A \in \mathcal{C}$;
- (2) \mathcal{C} is a *weak Serre* subcategory if it is non-empty and if

$$C_1 \longrightarrow C_2 \longrightarrow A \longrightarrow C_3 \longrightarrow C_4,$$

is an exact sequence in \mathcal{A} , where the $C_i \in \mathcal{C}$ and $A \in \mathcal{A}$, then $A \in \mathcal{C}$.

Clearly, Serre subcategories are weak Serre subcategories. Also, weak Serre subcategories are automatically abelian and the inclusion $\mathcal{C} \subseteq \mathcal{A}$ is exact [Stacks, Tags 02MP & 0754]. Moreover, the subcategory $D_{\mathcal{C}}(\mathcal{A})$ of $D(\mathcal{A})$, consisting of complexes in \mathcal{A} with cohomology in \mathcal{C} , is triangulated [Stacks, Tag 06UQ].

The main result of this appendix is

Theorem A.3. *Let \mathcal{A} be a Grothendieck abelian category and let $\mathcal{M} \subseteq \mathcal{A}$ be a weak Serre subcategory closed under coproducts. If \mathcal{M} is Grothendieck abelian, then $D_{\mathcal{M}}(\mathcal{A})$ is well generated.*

The example we have in mind is where X is an algebraic stack, \mathcal{A} is the category of lisse-étale sheaves of \mathcal{O}_X -modules, and \mathcal{M} is the subcategory of quasi-coherent sheaves.

If λ is a cardinal and \mathcal{B} is a cocomplete category, then we let \mathcal{B}^λ denote the subcategory of λ -presentable objects. If \mathcal{B} is locally presentable, then \mathcal{B}^λ is always a set.

It is clear that $D_{\mathcal{M}}(\mathcal{A})$ is a localizing subcategory of the well generated triangulated category $D(\mathcal{A})$; Remark A.2 tells us that to prove Theorem A.3 it suffices to exhibit a set S of generators for $D_{\mathcal{M}}(\mathcal{A})$. The idea is simple enough: we will find a cardinal λ such that $S = D_{\mathcal{M}^\lambda}(\mathcal{A}^\lambda) \subseteq D_{\mathcal{M}}(\mathcal{A})$, which is obviously essentially small, suffices. Thus, the problem becomes to better understand the category of λ -presentable objects in \mathcal{A} . The results below are easy to extract from [Nee14], but for the reader's convenience we give a self-contained treatment.

Remark A.4. Let \mathcal{A} be a Grothendieck abelian category. By the Gabriel–Popescu theorem, there exists a ring R and a pair of adjoint additive functors

$$F: \text{Mod}(R) \rightleftarrows \mathcal{A} : G$$

such that F is exact and $FG \simeq \text{id}$. Let μ be an infinite cardinal \geq to the cardinality of R .

Lemma A.5. *With notation as in Remark A.4, let $\lambda > \mu$ be a regular cardinal. Then the λ -presentable objects of \mathcal{A} are precisely the objects of \mathcal{A} isomorphic to some FN , where N is an R -module of cardinality $< \lambda$.*

Proof. Let us first prove that, if N is an R -module of cardinality $< \lambda$, then FN is λ -presentable. Suppose $\{x_i, i \in I\}$ is a λ -filtered system in \mathcal{A} , and suppose that in the category \mathcal{A} we are given a map $\phi: FN \rightarrow \text{colim } x_i$. We need to show that ϕ factors through some x_i . In the category of R -modules, there is a natural map

$$\text{colim } Gx_i \xrightarrow{\rho} G[\text{colim } x_i].$$

Since F respects colimits and $FG \simeq \text{id}$, the map $F\rho$ is an isomorphism. As F is exact it must annihilate the kernel and cokernel of ρ . Form the pullback square

$$\begin{array}{ccc} P & \xrightarrow{\phi} & N \\ \downarrow & & \downarrow \theta \\ \text{colim } Gx_i & \xrightarrow{\rho} & G[\text{colim } x_i]. \end{array}$$

The image of ϕ is a submodule of N , hence has cardinality $< \lambda$. If we lift every element of $\text{Image}(\phi)$ arbitrarily to an element of P , the lifts generate a submodule $M \subseteq P$ of cardinality $< \lambda$. The kernel (respectively cokernel) of the map $M \rightarrow N$ is a submodule of $\text{Kernel}(\rho)$ (respectively $\text{Coker}(\rho)$), and hence both are annihilated by F . Summarizing: we have produced a morphism $M \rightarrow N$ of R -modules, with M of cardinality $< \lambda$ and $FM \rightarrow FN$ an isomorphism in \mathcal{A} , and such that the composite $M \rightarrow N \rightarrow G[\text{colim } x_i]$ factors through $\text{colim } Gx_i$. But $\{Gx_i, i \in I\}$ is a λ -filtered system in $\text{Mod}(R)$ and M is of cardinality $< \lambda$, and hence the map factors as $M \rightarrow Gx_i$ for some $i \in I$.

It remains to prove the converse: suppose $a \in \mathcal{A}$ is a λ -presentable object. Then Ga is an R -module, hence it is the λ -filtered colimit of all its submodules N_i of cardinality $< \lambda$. But then the identity map $a \rightarrow a$ is a map from the λ -presentable object a to the λ -filtered colimit of the FN_i , and therefore factors through some FN_i . Thus a is a direct summand of an object FN_i where the cardinality of N_i is $< \lambda$. On the other hand the map $N_i \rightarrow Ga$ is injective, hence so is $FN_i \rightarrow FGa = a$. Thus $a \cong FN_i$. \square

Lemma A.6. *Let \mathcal{A} be a Grothendieck abelian category. There is an infinite cardinal ν with the following properties: if $\lambda \geq \nu$ is a regular cardinal, then*

- (1) \mathcal{A}^λ is a Serre subcategory of \mathcal{A} ;
- (2) every object of \mathcal{A} is a λ -filtered colimit of subobjects belonging to \mathcal{A}^λ ;
- (3) an object belongs to \mathcal{A}^λ if and only if it is the quotient of a coproduct of $< \lambda$ objects of \mathcal{A}^ν ; and
- (4) any pair of morphisms $x \rightarrow y \leftarrow n$ in \mathcal{A} , where $x \rightarrow y$ is epi and $n \in \mathcal{A}^\lambda$, may be completed to a commutative square

$$\begin{array}{ccc} m & \xrightarrow{\quad} & n \\ \downarrow & & \downarrow \\ x & \xrightarrow{\quad} & y \end{array}$$

with $m \in \mathcal{A}^\lambda$ and $m \rightarrow n$ epi. Moreover, if $n \rightarrow y$ is mono, then $m \rightarrow x$ can be chosen to be mono.

Proof. Let $\nu = \mu + 1$ be the successor of the infinite cardinal μ of Remark A.4. By Lemma A.5 the objects of \mathcal{A}^λ are precisely the ones isomorphic to FM where M is of cardinality $< \lambda$.

For (1), it is readily verified that a subobject (resp. a quotient) of FM can be expressed as FN where N is a submodule (resp. a quotient) of M . This shows that \mathcal{A}^λ is closed under taking subobjects and quotients; we will later see that it is also closed under extensions.

For (2), if a is an object of \mathcal{A} then Ga is the λ -filtered colimit of all the submodules $M_i \subseteq Ga$ of cardinality $< \lambda$, and $a \cong FGa$ is the λ -filtered colimit of $FM_i \in \mathcal{A}^\lambda$.

For (3), observe that any coproduct of $< \lambda$ objects in \mathcal{A}^λ belongs to \mathcal{A}^λ , and if M is a module of cardinality $< \lambda$ then M is a quotient of the free module on all its elements, which is a coproduct of $< \lambda$ copies of R . Thus FM is the quotient of a coproduct of $< \lambda$ copies of $FR \in \mathcal{A}^\nu$.

For (4), let $x \rightarrow y \leftarrow n$ be a pair of morphisms in \mathcal{A} , with $x \rightarrow y$ epi and $n \in \mathcal{A}^\lambda$. Let \tilde{m} be the pullback of $n \rightarrow y$ along $x \rightarrow y$. It is sufficient to find a subobject m of \tilde{m} belonging to \mathcal{A}^λ such that $m \rightarrow \tilde{m} \rightarrow n$ is epi. By (2), we may express $\tilde{m} = \operatorname{colim} m_i$ as a λ -filtered colimit of subobjects belonging to \mathcal{A}^λ . If $n_i \subseteq n$ is the image of m_i in n , then (1) implies that $n_i \in \mathcal{A}^\lambda$. Since $n \in \mathcal{A}^\lambda$, there is an i such that $n_i = n$. Taking $m = m_i$ does the job. By construction, if $n \rightarrow y$ is mono, then $m \rightarrow x$ is mono.

Finally, to show that \mathcal{A}^λ is closed under extensions, we note that if $0 \rightarrow k \rightarrow x \rightarrow n \rightarrow 0$ is an exact sequence with $k, n \in \mathcal{A}^\lambda$, then (4) implies that there is a subobject m of x such that $m \in \mathcal{A}^\lambda$ and $m \rightarrow n$ is epi. It follows that $k \oplus m \rightarrow x$ is epi and consequently, $x \in \mathcal{A}^\lambda$, as required. \square

Proof of Theorem A.3. Because \mathcal{M} and \mathcal{A} are both Grothendieck abelian categories we may choose regular cardinals ν for \mathcal{M} and ν' for \mathcal{A} as in Lemma A.6. The category \mathcal{M}^ν is an essentially small subcategory of \mathcal{A} , hence must be contained in \mathcal{A}^β for some regular cardinal β . Let λ be a regular cardinal $> \max(\beta, \nu, \nu')$. By construction $\mathcal{M}^\nu \subseteq \mathcal{A}^\lambda$, and by Lemma A.6 every object in \mathcal{M}^λ is the quotient of a coproduct of $< \lambda$ objects in \mathcal{M}^ν . Hence $\mathcal{M}^\lambda \subseteq \mathcal{A}^\lambda$. But since every object in $\mathcal{M} \cap \mathcal{A}^\lambda$ is λ -presentable in \mathcal{A} it must be λ -presentable in the smaller \mathcal{M} , and we conclude that $\mathcal{M} \cap \mathcal{A}^\lambda = \mathcal{M}^\lambda$.

We have now made our choice of λ and we let $\mathcal{B} = \mathcal{A}^\lambda$. By Remark A.2 it suffices to show that, given any non-zero object $Z \in \mathbf{D}_{\mathcal{M}}(\mathcal{A})$, there is an object $N \in \mathbf{D}_{\mathcal{B} \cap \mathcal{M}}(\mathcal{B})$ and a non-zero map $N \rightarrow Z$. If Z is the chain complex

$$\dots \longrightarrow Z^{i-1} \xrightarrow{\partial} Z^i \xrightarrow{\partial} Z^{i+1} \longrightarrow \dots,$$

we let $Y^i \subseteq Z^i$ be the cycles, in other words the kernel of $\partial: Z^i \rightarrow Z^{i+1}$, and $X^i \subseteq Y^i$ be the boundaries, that is the image of $\partial: Z^{i-1} \rightarrow Z^i$. We are assuming that $Z \in \mathbf{D}_{\mathcal{M}}(\mathcal{A})$ is non-zero, meaning its cohomology is not all zero; without loss of generality we may assume $H^0(Z) \neq 0$. Thus Y^0/X^0 is a non-zero object of \mathcal{M} .

By Lemma A.6, applied to $\mathcal{B} \cap \mathcal{M} = \mathcal{M}^\lambda \subseteq \mathcal{M}$, the object $Y^0/X^0 \in \mathcal{M}$ is a λ -filtered colimit of its subobjects belonging to $\mathcal{B} \cap \mathcal{M}$; since $Y^0/X^0 \neq 0$ we may choose a subobject $M \subseteq Y^0/X^0$, with $M \in \mathcal{B} \cap \mathcal{M}$ and $M \neq 0$. By Lemma A.6, applied to the pair of maps $Y^0 \rightarrow Y^0/X^0 \leftarrow M$ in \mathcal{A} , we may complete to a commutative square

$$\begin{array}{ccc} N^0 & \xrightarrow{\phi} & M \\ \downarrow & & \downarrow \\ Y^0 & \longrightarrow & Y^0/X^0 \end{array}$$

with $N^0 \in \mathcal{B}$. Since Y^0 is the kernel of $Z^0 \rightarrow Z^1$ this gives us a commutative square

$$\begin{array}{ccc} N^0 & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ Z^0 & \longrightarrow & Z^1 \end{array}$$

such that the image of the map $N^0 \rightarrow Y^0/X^0 = H^0(Z)$ is non-zero and belongs to $\mathcal{B} \cap \mathcal{M}$.

We propose to inductively extend this to the left. We will define a commutative diagram

$$\begin{array}{ccccccccccccccc} & & & & N^i & \longrightarrow & N^{i+1} & \longrightarrow & \dots & \longrightarrow & N^{-1} & \longrightarrow & N^0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow & & \\ \dots & \longrightarrow & Z^{i-1} & \longrightarrow & Z^i & \longrightarrow & Z^{i+1} & \longrightarrow & \dots & \longrightarrow & Z^{-1} & \longrightarrow & Z^0 & \longrightarrow & Z^1 & \longrightarrow & \dots \end{array}$$

where

- (1) The subobjects $N^j \subseteq Z^j$ belong to \mathcal{B} .
- (2) For $j > i$ the cohomology of $N^{j-1} \rightarrow N^j \rightarrow N^{j+1}$ belongs to $\mathcal{B} \cap \mathcal{M}$.
- (3) Let K^i be the kernel of the map $N^i \rightarrow N^{i+1}$. Then the image of the natural map $K^i \rightarrow H^i(Z)$ belongs to $\mathcal{B} \cap \mathcal{M}$.

Since we have constructed N^0 we only need to prove the inductive step. Let us therefore suppose we have constructed the diagram as far as i ; we need to extend it to $i-1$. We first form the pullback square

$$\begin{array}{ccc} L^i & \longrightarrow & K^i \\ \downarrow & & \downarrow \\ X^i & \longrightarrow & Y^i \end{array}$$

Since $X^i \rightarrow Y^i$ and $K^i \rightarrow Y^i$ are monomorphisms so are $L^i \rightarrow K^i$ and $L^i \rightarrow X^i$. Since N^i belongs to \mathcal{B} so do its subobjects $L^i \subseteq K^i$. The cokernel of $L^i \rightarrow K^i$ is the image of $K^i \rightarrow Y^i/X^i = H^i(Z)$, and belongs to $\mathcal{B} \cap \mathcal{M}$ by (3). Next we apply Lemma A.6 to the pair of maps $Z^{i-1}/X^{i-1} \rightarrow X^i \leftarrow L^i$ in \mathcal{A} , completing to a commutative square

$$\begin{array}{ccc} M^i & \longrightarrow & L^i \\ \downarrow & & \downarrow \\ Z^{i-1}/X^{i-1} & \longrightarrow & X^i \end{array}$$

with $M^i \in \mathcal{B}$. Form the pullback

$$\begin{array}{ccc} \widetilde{M}^i & \longrightarrow & M^i \\ \downarrow & & \downarrow \\ Y^{i-1}/X^{i-1} & \longrightarrow & Z^{i-1}/X^{i-1} \end{array}$$

Since $Y^{i-1}/X^{i-1} \rightarrow Z^{i-1}/X^{i-1}$ is injective so is $\widetilde{M}^i \rightarrow M^i$, making \widetilde{M}^i a subobject of $M^i \in \mathcal{B}$. Hence \widetilde{M}^i belongs to \mathcal{B} . But now the map $\widetilde{M}^i \rightarrow Y^{i-1}/X^{i-1} = H^{i-1}(Z)$ is a morphism from the λ -presentable object $\widetilde{M}^i \in \mathcal{B} = \mathcal{A}^\lambda$ to the object $H^{i-1}(Z) \in \mathcal{M}$, which by Lemma A.6 is a λ -filtered colimit of its subobjects in $\mathcal{M}^\lambda = \mathcal{B} \cap \mathcal{M}$. Hence the map $\widetilde{M}^i \rightarrow Y^{i-1}/X^{i-1}$ factors as $\widetilde{M}^i \rightarrow P^i \rightarrow Y^{i-1}/X^{i-1}$ with $P^i \in \mathcal{B} \cap \mathcal{M}$ a subobject of Y^{i-1}/X^{i-1} . Form the pushout square in \mathcal{B}

$$\begin{array}{ccc} \widetilde{M}^i & \longrightarrow & M^i \\ \downarrow & & \downarrow \\ P^i & \longrightarrow & Q^i \end{array}$$

and let $Q^i \rightarrow Z^{i-1}/X^{i-1}$ be the natural map. We have a commutative square

$$\begin{array}{ccc} Q^i & \xrightarrow{\phi} & L^i \\ \downarrow & & \downarrow \\ Z^{i-1}/X^{i-1} & \longrightarrow & X^i \end{array}$$

and the kernel of ϕ maps isomorphically to the subobject $P^i \subseteq H^{i-1}(Z)$, with $P^i \in \mathcal{B} \cap \mathcal{M}$. Finally apply Lemma A.6 to the pair of maps $Z^{i-1} \rightarrow Z^{i-1}/X^{i-1} \leftarrow Q^i$ to complete to a square

$$\begin{array}{ccc} N^{i-1} & \longrightarrow & Q^i \\ \downarrow & & \downarrow \\ Z^{i-1} & \longrightarrow & Z^{i-1}/X^{i-1} \end{array}$$

with $N^i \in \mathcal{B}$. We leave it to the reader to check that the diagram

$$\begin{array}{cccccccccccc} N^{i-1} & \longrightarrow & N^i & \longrightarrow & \dots & \longrightarrow & N^{-1} & \longrightarrow & N^0 & \longrightarrow & 0 & \longrightarrow & \dots \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \dots & \longrightarrow & Z^{i-2} & \longrightarrow & Z^{i-1} & \longrightarrow & Z^i & \longrightarrow & \dots & \longrightarrow & Z^{-1} & \longrightarrow & Z^0 & \longrightarrow & Z^1 & \longrightarrow & \dots \end{array}$$

satisfies hypotheses (1), (2) and (3) of our induction. \square

APPENDIX B. $D_{\text{qc}}(X)$ IS LEFT-COMPLETE

In this section we prove the following Theorem.

Theorem B.1. *If X is an algebraic stack, then $D_{\text{qc}}(X)$ is well generated. In particular, it admits small products. Moreover, $D_{\text{qc}}(X)$ is left-complete.*

Proof. The subcategory $\text{QCoh}(X) \subseteq \text{Mod}(X_{\text{lis-ét}})$ is weak Serre and the inclusion is coproduct preserving. Since $\text{QCoh}(X)$ and $\text{Mod}(X_{\text{lis-ét}})$ are Grothendieck abelian categories [Stacks, 07A5 & 0781], it follows that $D_{\text{qc}}(X)$ is well generated (Theorem A.3). By [Nee01b, Cor. 1.18], $D_{\text{qc}}(X)$ admits small products.

It remains to prove that $D_{\text{qc}}(X)$ is left-complete. Let $p: U \rightarrow X$ be a smooth surjection from an algebraic space U . Let $U_{\bullet, \text{ét}}^+$ denote the resulting strictly simplicial algebraic space [Ols07, §4.1]. By [LO08, Ex. 2.2.5], there is an equivalence of triangulated categories $D_{\text{qc}}(X) \simeq D_{\text{qc}}(U_{\bullet, \text{ét}}^+)$. The inclusion $\text{QCoh}(U_{\bullet, \text{ét}}^+) \subseteq \text{Mod}(U_{\bullet, \text{ét}}^+)$ is exact, stable under extensions, and coproduct preserving. It follows that the functor $\omega: D_{\text{qc}}(U_{\bullet, \text{ét}}^+) \rightarrow D(U_{\bullet, \text{ét}}^+)$ is exact and coproduct preserving. As we already have seen, the category $D_{\text{qc}}(X) \simeq D_{\text{qc}}(U_{\bullet, \text{ét}}^+)$ is well generated. Thus the functor ω admits a right adjoint λ [Nee01b, Prop. 1.20]. Because the functor ω is fully faithful, the adjunction $\text{id} \Rightarrow \lambda \circ \omega$ is an isomorphism of functors.

Note that because λ is a right adjoint, it preserves products. In particular, it remains to prove that if $K \in D_{\text{qc}}(U_{\bullet, \text{ét}}^+)$, then there exists a distinguished triangle in $D(U_{\bullet, \text{ét}}^+)$ (where we also take the products in $D(U_{\bullet, \text{ét}}^+)$):

$$\omega(K) \longrightarrow \prod_{n \geq 0} \tau^{\geq -n} \omega(K) \xrightarrow{1\text{-shift}} \prod_{n \geq 0} \tau^{\geq -n} \omega(K) \longrightarrow \omega(K)[1].$$

Indeed, this follows from the observation that $\tau^{\geq -n} \omega(K) \simeq \omega(\tau^{\geq -n} K)$ for all integers n and $K \rightarrow \lambda \circ \omega(K)$ is an isomorphism.

Let $(W \rightarrow U_n)$ be an object of $U_{\bullet, \text{ét}}^+$. The resulting slice $U_{\bullet, \text{ét}}^+/(W \rightarrow U_n)$ is equivalent to the small étale site on W . In particular, it follows that if W is an affine scheme and $M \in \text{QCoh}(U_{\bullet, \text{ét}}^+) \cong \text{QCoh}(X)$, then $H^p(U_{\bullet, \text{ét}}^+/(W \rightarrow U_n), M) = 0$ for

all $p > 0$ [Stacks, 01XB & 0756]. Now let $\mathcal{B} \subseteq U_{\bullet, \text{ét}}^+$ denote the full subcategory consisting of those objects $(W \rightarrow U_n)$, where W is an affine scheme. It follows that \mathcal{B} satisfies the requirements of [Stacks, 08U3] and we deduce the result. \square

APPENDIX C. THE BOUNDED BELOW DERIVED CATEGORY

In this section, we prove an analog of [Har66, Cor. II.7.19] for noetherian algebraic stacks that are affine-pointed. Essentially for free, we will also establish Lurie's result [Lur04, Thm. 3.8].

Theorem C.1. *Let X be an algebraic stack. If X is either quasi-compact with affine diagonal or noetherian and affine-pointed, then the natural functor*

$$\Psi_X^{\dagger}: D^+(\text{QCoh}(X)) \rightarrow D_{\text{qc}}^+(X)$$

is an equivalence.

For noetherian algebraic spaces, a version of Theorem C.1 for the unbounded derived category was proved in [Stacks, Tag 09TN] and we will closely follow this approach. The following two lemmas do most of the work.

Lemma C.2 (cf. [Stacks, Tag 09TJ]). *Let X be a quasi-compact and quasi-separated algebraic stack and let I be an injective object of $\text{QCoh}(X)$.*

- (1) *Then I is a direct summand of p_*J , where $p: \text{Spec } A \rightarrow X$ is smooth and surjective and J is an injective A -module.*
- (2) *If X is noetherian, then I is a direct summand of a filtered colimit $\text{colim}_i F_i$ of quasi-coherent sheaves of the form $F_i = \gamma_*G_i$, where $\gamma: Z_i \rightarrow X$ is a morphism from an artinian scheme Z_i and $G_i \in \text{Coh}(Z_i)$.*

Proof. Let $p: U \rightarrow X$ be a smooth and surjective morphism, where $U = \text{Spec } A$ is an affine scheme. Let I be an injective object of $\text{QCoh}(X)$. Choose an injective object J of $\text{QCoh}(U)$ and an injection $p^*I \subseteq J$. By adjunction, we have an inclusion $I \subseteq p_*J$. Since p^* is exact, p_*J is injective in $\text{QCoh}(X)$ and I is a direct summand of p_*J . This proves (1). For (2): we may now reduce to the case where $X = U$. The result is now well-known (e.g., [Stacks, Tag 09TI]). \square

Lemma C.3 (cf. [Stacks, Tag 09TL]). *Let X be an algebraic stack and let I be an injective object of $\text{QCoh}(X)$. If X is quasi-compact with affine diagonal (resp. noetherian and affine-pointed), then*

- (1) *$H^q(U_{\text{lis-ét}}, I) = 0$ for every $q > 0$ and smooth morphism $u: U \rightarrow X$ that is affine (resp. has affine fibers);*
- (2) *for any morphism $f: X \rightarrow Y$ of algebraic stacks, where Y has affine diagonal (resp. Y is affine-pointed) we have $R^q(f_{\text{lis-ét}})_*I = 0$ for $q > 0$.*

Proof. Let W be an affine (resp. artinian) scheme and let $M \in \text{QCoh}(W)$ be injective (resp. $M \in \text{Coh}(W)$). Let $w: W \rightarrow X$ be a smooth and surjective morphism (resp. a morphism). By Lemma C.2, it is sufficient to prove the result for $I = w_*M$. Since X has affine diagonal (resp. X is affine-pointed), w is affine. In particular, the natural map $(w_*M)[0] \rightarrow R(w_{\text{lis-ét}})_*M$ is a quasi-isomorphism.

We now prove (1). Let $u_W: W_U \rightarrow W$ be the pull back of u along w and let $w_U: W_U \rightarrow U$ be the pull back of w along u . In both cases, u_W is smooth and affine and w_U is affine; in particular, W_U is an affine scheme. Since u is smooth,

$$\begin{aligned} \text{R}\Gamma(U_{\text{lis-ét}}, I) &\simeq \text{R}\Gamma(U_{\text{lis-ét}}, u^*R(w_{\text{lis-ét}})_*M) \simeq \text{R}\Gamma(U_{\text{lis-ét}}, R((w_U)_{\text{lis-ét}})_*(u_W^*M)) \\ &\simeq \text{R}\Gamma((W_U)_{\text{lis-ét}}, M). \end{aligned}$$

The result now follows from the affine case (e.g., [EGA, III.1.3.1]).

For (2): let $v: V \rightarrow Y$ be a smooth morphism, where V is an affine scheme. Since Y has affine diagonal (resp. is affine-pointed), v is affine (resp. has affine fibers). By (1), $H^q((V \times_Y X)_{\text{lis-ét}}, I) = 0$. But $R^q(f_{\text{lis-ét}})_* I$ is the sheafification of the presheaf $V \mapsto H^q((V \times_Y X)_{\text{lis-ét}}, I)$; the result follows. \square

Proof of Theorem C.1. We first establish the full faithfulness. Let $F, G \in D^+(\text{QCoh}(X))$; then we wish to prove that the natural map

$$\text{Hom}_{D(\text{QCoh}(X))}(F, G) \rightarrow \text{Hom}_{D(X)}(F, G)$$

is an isomorphism. A standard way-out argument shows that it is sufficient to prove that the natural map

$$\text{Ext}_{\text{QCoh}(X)}^q(N, M) \rightarrow \text{Ext}_{\mathcal{O}_X}^q(N, M)$$

is an isomorphism for every $q \in \mathbb{Z}$ and $M, N \in \text{QCoh}(X)$. For $q < 0$ both sides vanish and for $q = 0$ we clearly have an isomorphism. For $q > 0$, since every M embeds in a quasi-coherent injective I , a standard δ -functor argument shows that it is sufficient to prove that if I is an injective object of $\text{QCoh}(X)$, then

$$\text{Ext}_{\mathcal{O}_X}^q(N, I) = 0$$

for all $q > 0$ and $N \in \text{QCoh}(X)$. To see this we note that by Lemma C.2(1), I is a direct summand of $(p_{\text{QCoh}})_* J$, where $p: \text{Spec } A \rightarrow X$ is smooth and surjective and J is an injective A -module. Thus, it suffices to prove the result when $I = (p_{\text{QCoh}})_* J$. By Lemma C.3(2), the natural map $((p_{\text{QCoh}})_* J)[0] \rightarrow R(p_{\text{lis-ét}})_* J$ is a quasi-isomorphism. Hence, there are natural isomorphisms:

$$\text{Ext}_{\mathcal{O}_X}^q(N, (p_{\text{QCoh}})_* J) \cong \text{Ext}_{\mathcal{O}_X}^q(N, R(p_{\text{lis-ét}})_* J) \cong \text{Ext}_{\mathcal{O}_{\text{Spec } A}}^q(p^* N, J).$$

We are now reduced to the affine case, which is well-known (e.g., [BN93, Lem. 5.4]).

For the essential surjectivity, we argue as follows: by induction and using the full faithfulness, one easily sees that $D^b(\text{QCoh}(X)) \simeq D_{\text{qc}}^b(X)$. Passing to homotopy colimits, we obtain the claim. \square

REFERENCES

- [BB03] A. Bondal and M. van den Bergh, *Generators and representability of functors in commutative and noncommutative geometry*, Mosc. Math. J. **3** (2003), no. 1, 1–36, 258.
- [BF03] A. J. Berrick and E. D. Farjoun, *Fibrations and nullifications*, Israel J. Math. **135** (2003), 205–220.
- [BIK11] D. J. Benson, S. B. Iyengar, and H. Krause, *Stratifying modular representations of finite groups*, Ann. of Math. (2) **174** (2011), no. 3, 1643–1684.
- [BN93] M. Bökstedt and A. Neeman, *Homotopy limits in triangulated categories*, Compositio Math. **86** (1993), no. 2, 209–234.
- [Bor91] A. Borel, *Linear algebraic groups*, second ed., Graduate Texts in Mathematics, vol. 126, Springer-Verlag, New York, 1991.
- [Bou75] A.K. Bousfield, *The localization of spaces with respect to homology*, Topology **14** (1975), 133–150.
- [Bou79] A.K. Bousfield, *The localization of spectra with respect to homology*, Topology **18** (1979), 257–281.
- [BZ09] D. Ben-Zvi, *Compact generation for modular representations*, MathOverflow, Nov 2009, <http://mathoverflow.net/q/4965>.
- [BZFN10] D. Ben-Zvi, J. Francis, and D. Nadler, *Integral transforms and Drinfeld centers in derived algebraic geometry*, J. Amer. Math. Soc. **23** (2010), no. 4, 909–966.
- [Cho05] B. Chorny, *Localization with respect to a class of maps. I. Equivariant localization of diagrams of spaces*, Israel J. Math. **147** (2005), 93–139.
- [Con02] B. Conrad, *A modern proof of Chevalley’s theorem on algebraic groups*, J. Ramanujan Math. Soc. **17** (2002), no. 1, 1–18.
- [CSS05] C. Casacuberta, D. Scevenels, and J. H. Smith, *Implications of large-cardinal principles in homotopical localization*, Adv. Math. **197** (2005), no. 1, 120–139.
- [DG13] V. Drinfeld and D. Gaitsgory, *On some finiteness questions for algebraic stacks*, Geom. Funct. Anal. **23** (2013), no. 1, 149–294.

- [EGA] A. Grothendieck, *Éléments de géométrie algébrique*, I.H.E.S. Publ. Math. **4**, **8**, **11**, **17**, **20**, **24**, **28**, **32** (1960, 1961, 1961, 1963, 1964, 1965, 1966, 1967).
- [Gro13] P. Gross, *Tensor generators on schemes and stacks*, preprint, June 2013, arXiv:1306.5418.
- [Hal14] J. Hall, *Cohomology and base change for algebraic stacks*, Math. Z. **278** (2014), no. 1-2, 401–429.
- [Har66] R. Hartshorne, *Residues and duality*, Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20, Springer-Verlag, Berlin, 1966.
- [HR14a] J. Hall and D. Rydh, *Algebraic groups and compact generation of their derived categories of representations*, Indiana Univ. Math. J. (2014), accepted for publication.
- [HR14b] J. Hall and D. Rydh, *Coherent Tannaka duality and algebraicity of Hom-stacks*, preprint, May 2014, arXiv:1405.7680.
- [HR14c] J. Hall and D. Rydh, *Perfect complexes on algebraic stacks*, preprint, May 2014, arXiv:1405.1887v2.
- [Kri09] A. Krishna, *Perfect complexes on Deligne-Mumford stacks and applications*, J. K-Theory **4** (2009), no. 3, 559–603.
- [KS06] M. Kashiwara and P. Schapira, *Categories and sheaves*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 332, Springer-Verlag, Berlin, 2006.
- [LMB] G. Laumon and L. Moret-Bailly, *Champs algébriques*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge., vol. 39, Springer-Verlag, Berlin, 2000.
- [LO08] Y. Laszlo and M. Olsson, *The six operations for sheaves on Artin stacks. I. Finite coefficients*, Publ. Math. Inst. Hautes Études Sci. (2008), no. 107, 109–168.
- [Lur04] J. Lurie, *Tannaka duality for geometric stacks*, preprint, December 2004, arXiv:math/0412266, p. 14.
- [Lur11] J. Lurie, *Derived Algebraic Geometry VIII: Quasi-coherent sheaves and Tannaka Duality Theorems*, available on homepage, May 2011.
- [Nee96] A. Neeman, *The Grothendieck duality theorem via Bousfield's techniques and Brown representability*, J. Amer. Math. Soc. **9** (1996), no. 1, 205–236.
- [Nee01a] A. Neeman, *On the derived category of sheaves on a manifold*, Doc. Math. **6** (2001), 483–488 (electronic).
- [Nee01b] A. Neeman, *Triangulated categories*, Annals of Mathematics Studies, vol. 148, Princeton University Press, Princeton, NJ, 2001.
- [Nee11] A. Neeman, *Non-left-complete derived categories*, Math. Res. Lett. **18** (2011), no. 5, 827–832.
- [Nee14] A. Neeman, *The homotopy category of injectives*, Algebra Number Theory **8** (2014), no. 2, 429–456.
- [Ols07] M. Olsson, *Sheaves on Artin stacks*, J. Reine Angew. Math. **603** (2007), 55–112.
- [Ros61] M. Rosenlicht, *On quotient varieties and the affine embedding of certain homogeneous spaces*, Trans. Amer. Math. Soc. **101** (1961), 211–223.
- [Ryd11] D. Rydh, *Étale dévissage, descent and pushouts of stacks*, J. Algebra **331** (2011), 194–223.
- [Ser03] C. Serpé, *Resolution of unbounded complexes in Grothendieck categories*, J. Pure Appl. Algebra **177** (2003), no. 1, 103–112.
- [SGA6] *Théorie des intersections et théorème de Riemann-Roch*, Lecture Notes in Mathematics, Vol. 225, Springer-Verlag, Berlin, 1971, Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6), Dirigé par P. Berthelot, A. Grothendieck et L. Illusie. Avec la collaboration de D. Ferrand, J. P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud et J. P. Serre.
- [Stacks] The Stacks Project Authors, *Stacks Project*, http://math.columbia.edu/algebraic_geometry/stacks-git.
- [Toë12] B. Toën, *Derived Azumaya algebras and generators for twisted derived categories*, Invent. Math. **189** (2012), no. 3, 581–652.
- [Tot04] B. Totaro, *The resolution property for schemes and stacks*, J. Reine Angew. Math. **577** (2004), 1–22.
- [TT90] R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, The Grothendieck Festschrift, Vol. III, Progr. Math., vol. 88, Birkhäuser Boston, Boston, MA, 1990, pp. 247–435.

MATHEMATICAL SCIENCES INSTITUTE, THE AUSTRALIAN NATIONAL UNIVERSITY, ACTON, ACT,
2601, AUSTRALIA

E-mail address: `jack.hall@anu.edu.au`

MATHEMATICAL SCIENCES INSTITUTE, THE AUSTRALIAN NATIONAL UNIVERSITY, ACTON, ACT,
2601, AUSTRALIA

E-mail address: `Amnon.Neeman@anu.edu.au`

KTH ROYAL INSTITUTE OF TECHNOLOGY, DEPARTMENT OF MATHEMATICS, SE-100 44 STOCK-
HOLM, SWEDEN

E-mail address: `dary@math.kth.se`