Convergence properties of the Gronwall area formula for quadratic Julia sets

Alexandre Dezotti

Abstract

Gronwall area formula can be used to express the area of the filled Julia set of connected quadratic Julia sets. Using parabolic enrichment, it is shown that there cannot be a nice approximation of this formula by partial sums which is uniform along the boundary of the main cardioid of the Mandelbrot set.

1 Introduction

One important aspect of the study of the iteration of holomorphic functions is the study of the Julia set. The Julia set of a holomorphic mapping corresponds to the chaotic part of the dynamical system consisting in iterating the corresponding mapping.

The question of the measure of Julia sets is a very natural question, already raised by Fatou in his memoir [7], p.243. For some time it has been possible to conjecture that all of the Julia sets of quadratic polynomials have zero area. Indeed this is the case in many situations (see, for example [5], [15], [18]) and, if this conjecture were true, it would have implied the conjecture stating that hyperbolic dynamical systems are dense in the family of complex quadratic polynomials ¹. No proof of the latter conjecture has been published yet, except for real polynomials ([9] or [10]), but in their paper [3], Buff and Chéritat showed the existence of quadratic polynomials for which the Julia set has positive area. Hence the proof of the hyperbolicity density conjecture cannot relies on this argument.

Interestingly, some of those examples are period one Cremer parameters. Recall that, in the quadratic family, a period one Cremer parameter is a value of $c \in \mathbb{C}$ such that the polynomial $P_c(z) = z^2 + c$ has a non hyperbolic fixed point ² on the neighbourhood of which P_c cannot be conjugated to its linear part and such that the linear part is not periodic. Those parameters lie in the boundary of the main cardioid of the Mandelbrot set. An equivalent definition of Cremer parameters is the following. If z_0 denotes the corresponding fixed

¹ This conjecture could, in some way, be dated back to Fatou's discussion [8], p.73. Nevertheless, a likely interpretation of his discussion would lead to a slightly different conjecture, which has been proven false. On this matter, compare McMuller [16], chapter 4.

 $^{^{2}}$ Which means, here, by a slight abuse of terminology, that the derivative has modulus one.

point, $|P'_c(z_0)| = 1$, $P'_c(z_0)$ is not a root of unity and P_c is not linearisable around z_0 . The point z_0 is called a (period one) Cremer point and one can also talk about periodic Cremer point in a similar way for any period ³.

For polynomials, one can define the filled Julia set which is the union of the Julia set and all the bounded component of its complement in the complex plane. The filled Julia sets of other parameters such as Siegel parameters and parabolic parameters have positive area. The Siegel parameters correspond to the linearisable case of a non hyperbolic fixed (or periodic) point and parabolic parameters are the values of c such that P_c has a periodic point z_0 for which $(P_c^{\circ p})'(z_0)$ is a root of unity where p denotes the period of z_0 . In both case, the filled Julia set has non empty interior.

As the boundary of the main cardioid correponds to the set of quadratic polynomials having a non hyperbolic fixed point, Hubbard asked whether there would be a lower bound on the area of the filled Julia set along the boundary of the main cardioid. For example, the question is stated for the Cremer parameters in [2]. The question is more critical for Cremer parameters as we do not know if some of them have a Julia set with zero area. Indeed, the complexity of the situation is reflected by the fact that, due to the "close proximity" of Cremer parameters to parabolic parameters, standard algorithms cannot distinguish between Cremer parameters and parabolic parameters, see for example figure 1.

It is well known that Cremer parameters are generic on the boundary of a hyperbolic component of the Mandelbrot set [17]. Moreover Lyubich proved that for generic parameters (in the sense of Baire) on the boundary of the Mandelbrot set the area of the Julia set is zero [14]. Unfortunately this is insufficient to prove that some of this Cremer Julia set have zero area, as the complement of the union of the boundaries of hyperbolic components is a generic subset of the boundary of the Mandelbrot set.

This article concerns the Gronwall area formula applied to the area of quadratic filled Julia sets. Let's recall its content.

For a closed subset of the Riemann sphere \mathbb{C} containing at least two points and whose complement is simply connected and contains ∞ , the Gronwall area formula provides a way to compute its area, that is its Lebesgue two dimensional measure. Namely[11], if $\psi(w) = \sum_{n \leq 1} b_n w^n$ is the Laurent series of a conformal

isomorphism between the complement of the disk of radius $r_0 \ge 1$ centred at 0 in $\widehat{\mathbb{C}}$ and the complement of the compact set $K \subset \mathbb{C}$, then, for any $r \ge r_0$,

Area
$$(\{z : |\psi^{-1}(z)| \le r\}) = \pi \sum_{n \le 1} n |b_n|^2 r^{2n}.$$
 (1)

Gronwall area formula is essentially a consequence of Green's theorem.

As a consequence, this result yields an expression for the area of the filled Julia set of any polynomial map with connected Julia set. Indeed, by a classical

³And also the fact that the function is a quadratic polynomial has nothing to do with the general definition, valid for any holomoprhic function around z_0 .



Figure 1: Computer representations of the filled Julia sets of $\lambda z + z^2$ for the parameter $\lambda = e^{2\pi i/3}$ of the fat Douady's rabbit on the left and, on the right, for the Cremer parameter $\lambda = e^{2\pi i\theta}$, with $\theta = [0, 3, 10^{16}, a_3, \ldots, a_n, \ldots]$ where $[a_0, a_1, a_2, \ldots]$ is the notation for the continued fraction $a_0 + 1/(a_1 + 1/(a_2 + \ldots)))$ with coefficients a_n , and a_n satisfies, for $n \geq 2$, $a_{n+1} = 2^{q_n}$, where q_n is the denominator of the *n*th convergent.

theorem of Böttcher (compare for example [17]), the dynamics of any polynomial of degree d is conjugated to the dynamics of $w \mapsto w^d$ near ∞ . This provides a natural isomorphism between the basin of attraction of ∞ and the complement of the unit disk.

It is possible to explicitly compute the coefficients of the Laurent series of the inverse of the Böttcher map of a polynomial and then use them in order to numerically evaluate an approximation of the formula (1) via a finite summation.

We are particularly interested in the case of quadratic polynomials. If we denote by G_{λ} the Green function, i.e. the logarithm of the modulus of the Böttcher map of the quadratic polynomial $Q_{\lambda}(z) = \lambda z + z^2$, we can define an approximation of the area of the sublevel set $\{z \in \mathbb{C} : G_{\lambda}(z) \leq \log r\}$ by a finite sum with

$$A(\lambda, r, N) = \pi \sum_{n=-N}^{1} n |b_n|^2 r^{2n}.$$
 (2)

It is natural to ask how close the value of $A(\lambda, r, N)$ is to the actual area of the sublevel set of the Green function, and, more interestingly, how close is the value of $A(\lambda, 1, N)$ to the area of the filled Julia set?

An example of a numerical computation using the Gronwall area formula is given in the figure 2 (see also [17], appendix A). This example shows, for quadratic polynomials in the form $P_c(z) = z^2 + c$, the value of the approximation when the parameter c varies on the upper half of the boundary of the main cardioid of the Mandelbrot set. Similar methods can be applied to compute an estimate of the area of the Mandelbrot set, see, for example, the work of Ewing and Schober [6] where they also compare the result with pixel counting methods and lower bounds using the area of the biggest hyperbolic components.



Figure 2: Plot of computed values of the truncated area formula for filled Julia sets (r = 1) along the upper boundary of the main cardioid. The different graphs represents different level of truncations: 1, 20, 200, 2000, 20000 and 200000 terms (due to the definition of B_c , half of the terms are 0). The result of the computations decreases as the level increases. The values on the horizontal axis represent the rotation number.

Testing the lower bound hypothesis with numerical experiments requires to know the answer to the previous questions. While it can give the impression of a lower bound, the figure 2 gives some hint of slow convergence of the terms appearing in the sum for some parameters, for example on the side of the 1/2 hill on the right. This could be an indication that one cannot relies on such numerical experiments for investigating the existence of a lower bound on the area.

Indeed, the theorem below states that the area of the filled Julia set is discontinuous near the parabolic parameter 1. This prevents a uniform approximation of the area by the formula.

In the following statement, mappings on the form $Q_{\lambda}(z) = \lambda z + z^2$ are considered with $\lambda \in \mathbb{C}$. Those are conjugated to the family of mappings $P_c(z) = z^2 + c$ by affine maps, the correspondence between the two parameters being $c = \frac{\lambda}{2} \left(1 - \frac{\lambda}{2}\right)$. The connectedness locus of the family $(P_c)_c$ is the Mandelbrot set, its counterpart for the family $(Q_{\lambda})_{\lambda}$, called the double Mandelbrot set ([17], figure 29), is represented on figure 3.

Theorem 1.1. Let K_{λ} denote the filled Julia set of the quadratic polynomial $Q_{\lambda}(z) = \lambda z + z^2$.

Then

$$\limsup_{\lambda \to 1, |\lambda|=1} \operatorname{Area}(K_{\lambda}) < \operatorname{Area}(K_{1}),$$

In a more precise way, the theorem below shows that approximation of the area using a truncated Gronwall area formula fails to provide insight into the complexity of the variation of the filled Julia set. Even if the number of terms used in finite sum approximations increases at a very fast rate as the parameter approaches 1, there will be a definite discrepancy between the numerical result and the actual value of the area.

In what follows, we use the notation $A(\lambda, r, N)$ for the approximation of the area of Green sublevels given by (2) and $\{x\}$ denotes the fractional part of the real number x, that is, $\{x\} = x - n$ where n is the largest integer smaller than or equal to x.

Theorem 1.2. If $(N_n)_n$ is a sequence of natural numbers converging to $+\infty$ such that $N_n \in o(2^{1/\alpha_n})$, then

$$A(\lambda_n, 1, N_n) \to \operatorname{Area}(K_1).$$

Theorem 1.3. For any $\gamma > 0$ and $\tau \in [0,1]$, there exists C > 0 satisfying the following.

Let $(\alpha_n)_n$ be a sequence of positive real numbers converging to 0 and such that $\left\{\frac{1}{\alpha_n}\right\} \to \tau$, and let $(N_n)_n$ be a sequence of natural numbers. Define $\lambda_n = e^{2i\pi\alpha_n}$ and suppose that $\log N_n \leq \frac{\gamma}{\alpha_n}$. Then, there exists $n_0 \in \mathbb{N}$ such that, for all $n \geq n_0$,

$$A(\lambda_n, 1, N_n) \ge C + \operatorname{Area} K_{\lambda_n}.$$

Suppose we are to compute an approximation of the area of the filled Julia set of a parameter close to 1. The above means that, in order to see any difference with the area of K_1 , it requires a number of terms which increases at least as fast as the exponential of the inverse of the distance of the parameter to 1.

Thanks to the continuity of the straightening map of Mandelbrot-like family and to the uniform estimate on area distortion by quasiconformal maps in [1], the theorems 1.2 and 1.3 are also true for the cusp and the boundary of the main component of any Mandelbrot-like families and in particular of primitve copy of the Mandelbrot set itself.

Finally, techniques similar to those used in the proofs allow, with some minor modifications, the extension of those results to the root of any hyperbolic components of the Mandelbrot set. See, e.g. [19], section 7.

The theory of perturbed Fatou coordinates ([5], [19]) plays an essential role in the proof of this theorem. The proof relies on the fact that, truncating the series amounts essentially to compute the area of points that spend a long time in a bounded domain close to the filled Julia set. For parameters close to the cusp of \mathcal{M} , this set can contain many points in the basin of infinity.

The article is organised as follows. Section 2 contains preliminary materials, including some results based on the bifurcation of parabolic fixed points and near parabolic fixed point theory. Section 3 contains the main argument of the proof of theorem 1.1 and theorem 1.3.



Figure 3: The double Mandelbrot set. The big disk on the left is the unit disk. The point of tangency of the two big disks is the point $\lambda = 1$.

Acknowledgements

I would like to thank Prof. Xavier Buff for his help. I would also like to thank Prof. Lasse Rempe-Gillen.

2 Preliminaries

2.1 Some notations

The following introduces some notations and basic facts. For r > 0, \mathbb{D}_r denotes the disk of centre 0 and radius r.

Let $Q_{\lambda}(z) = \lambda z + z^2$ and $\mathcal{DM} = \{\lambda \in \mathbb{C} : J_{\lambda} \text{ is connected}\}\$ be the double Mandelbrot set.

The Böttcher isomorphism of Q_{λ} is denoted by $B_{\lambda} : U_{\lambda} \to \mathbb{C} \setminus D_{\lambda}$. It is defined in a neighbourhood U_{λ} of ∞ in the Riemann sphere $\widehat{\mathbb{C}}$ onto the complementary of a closed disk D_{λ} with centre at 0. Its inverse will be denoted ψ_{λ} and G_{λ} will denote the Green function.

The notation $\mathcal{N}_g = \{\lambda \in \mathbb{C}, G_\lambda(c_\lambda) \leq g\}$ will be used. Thus $\mathcal{DM} = \mathcal{N}_0$.

For $\lambda \in \mathcal{DM}$, $V_{\lambda}(g)$ will be the set $\{z : G_{\lambda}(z) \leq g\}$. The mapping $g \in \mathbb{R}_+ \mapsto V_{\lambda}(g)$ is continuous with respect to the Hausdorff metric and $V_{\lambda}(0) = K_{\lambda}$.

It is well known that the mapping $\lambda \mapsto K_{\lambda}$ is upper semi-continuous. As a consequence, $\limsup_{\lambda \to 1} \operatorname{Area} K_{\lambda} \leq \operatorname{Area} K_1$ (see also ??).

Given $x \in \mathbb{R}$, let's denote by $\{x\}$ the fractional part of x, that is $\{x\} = x - \lfloor x \rfloor$ where $\lfloor x \rfloor = \max\{n \in \mathbb{Z} : n \leq x\}$ is the integer part of x. We will also need a notation for $\lceil x \rceil = \min\{n \in \mathbb{Z} : n \geq x\}$.

2.2Estimates on the Green function

This section contains some elementary yet useful estimates on the Green function.

Lemma 2.1. For all λ such that $|\lambda| \leq 5$ and all $\xi \in \mathbb{C}$,

$$\log |\xi| - \log 6 \le G_{\lambda}(\xi) \le \max\{\log 11, \log |\xi| + \log 11/6\}.$$

Proof. For $|\xi| \ge 6$, the value of $|Q_{\lambda}^{\circ p}(\xi)|$ increases and tends to ∞ as $p \to \infty$. For such ξ , we have $|\xi|^2 \left(1 + \left|\frac{\lambda}{\xi}\right|\right) \le Q_{\lambda}(\xi) \le |\xi|^2 \left(1 - \left|\frac{\lambda}{\xi}\right|\right)$. Hence,

$$\log |\xi| - \log 6 \left(1 - \frac{1}{2^p} \right) \le \frac{\log |Q_{\lambda}^{\circ p}(\xi)|}{2^p} \le \log |\xi| - \log 11/6 \left(1 - \frac{1}{2^p} \right).$$

This solve the case $|\xi| \leq 6$. In the case where $|\xi| \leq 6$, the result follows from maximum principle for harmonic functions.

Lemma 2.2. Let R > 6, g > 0 and $\lambda \in \mathbb{C}$ such that $|\lambda| \leq 5$. If $p = \left\lceil \frac{\log \log 11R/6 - \log g}{\log 2} \right\rceil \geq 0$ and if $z \in \mathbb{C}$ is such that $|Q_{\lambda}^{\circ p}(z)| \leq R$ then $|G_{\lambda}(z)| \leq g$.

Proof. Using lemma 2.1 with $\xi = Q_{\lambda}^{\circ p}(z)$ and the fact that $G_{\lambda} \circ Q_{\lambda} = 2G_{\lambda}$, we get $G_{\lambda}(z) \leq \frac{1}{2^{p}} \max \{ \log 11, \log 11R/6 \}$. The results then follows from the fact that R > 6 and the definition of p.

2.3The area formula

In the present situation, the area formula (1) can be restated as

Area
$$V_{\lambda}(\log r) = \pi \left(r^2 - \sum_{k=1}^{\infty} k \frac{|b_k^{\lambda}|^2}{r^{2k}} \right),$$
 (3)

where $\psi_{\lambda}(w) = w + \sum_{k=1}^{\infty} \frac{b_k^{\lambda}}{w^k}, r \ge 1 \text{ and } \lambda \in \mathcal{N}_{\log r} = \{\lambda \in \mathbb{C}, G_{\lambda}(c_{\lambda}) \le \log r\}.$ Define, for $\lambda \in \mathbb{C}, r > 1$ and $N \in \mathbb{N}^*$,

$$A(\lambda, r, N) = \pi \left(r^2 - \sum_{k=1}^{N} \frac{|b_k^{\lambda}|^2}{r^{2k}} \right)$$

$$\tag{4}$$

and

$$A(\lambda, r, \infty) = \operatorname{Area} V_{\lambda}(\log r).$$

Moreover, the following mappings are continuous:

- For $\lambda \in \mathcal{DM}, g \in \mathbb{R}_+ \mapsto A(\lambda, e^g, \infty),$
- For $g > 0, \lambda \in \mathring{\mathcal{N}}_{q} \mapsto A(\lambda, e^{g}, \infty),$

• For $N \in \mathbb{N}^*$, $(\lambda, g) \in \mathbb{C} \times \mathbb{R} \mapsto A(\lambda, e^g, N)$.

The following lemmas will allow us to relate truncated approximations (4) with sublevel sets of the Green function.

Lemma 2.3. Let $N \in \mathbb{N}^*$, r > 1 and $\lambda \in \mathbb{C}$ such that $\lambda \in \mathcal{N}_{\log r}$, then

$$A(\lambda, 1, N) \ge \pi(1 - r^{2N+2}) + r^{2N}A(\lambda, r, \infty).$$

Proof. Recall that $A(\lambda, 1, N) = \pi \left(1 - \sum_{k=1}^{N} k|b_k^{\lambda}|^2\right)$. From

$$\sum_{k=1}^{N} k |b_k^{\lambda}|^2 \le r^{2N} \sum_{k=1}^{N} k \frac{|b_k^{\lambda}|^2}{r^{2k}} \le r^{2N} \sum_{k=1}^{\infty} k \frac{|b_k^{\lambda}|^2}{r^{2k}},$$

it follows that $A(\lambda, 1, N) \ge \pi \left(1 - r^{2N} \sum_{k=1}^{\infty} k \frac{|b^{\lambda}k|^2}{r^{2k}}\right)$, which is equivalent to the statement of the lemma.

Lemma 2.4. Let R > 6, $N \in \mathbb{N}^*$, $r \in]1,11R/6[$ and $\lambda \in \mathbb{C}$. Suppose that $|\lambda| < 5$ and $\lambda \in \mathcal{N}_{\log r}$. Define $p = \left\lceil \frac{\log \log 11R/6 - \log \log r}{\log 2} \right\rceil$. Then $p \ge 0$ and

$$A(\lambda, 1, N) \ge \pi (1 - r^{2N+2}) + r^{2N} \operatorname{Area} \left(\left\{ z : |Q_{\lambda}^{\circ p}(z)| \le R \right\} \right).$$

Proof. This lemma is a direct consequence of lemmas 2.2 and 2.3.

Lemma 2.5. Let $N \in \mathbb{N}^*$, r > 1 and $\lambda \in \mathcal{N}_{\log r}$. Then,

$$A(\lambda, 1, N) \le \operatorname{Area}\left(V_{\log r}\right) + \pi r^{2N+2}.$$
(5)

Proof. Indeed, for any r > 1,

$$\pi\left(1 - \sum_{k=1}^{N} k |b_k^{\lambda}|^2\right) \le \pi\left(r - \sum_{k=1}^{N} \frac{k |b_k^{\lambda}|^2}{r^{2k}}\right).$$

But, since $\sum_{k=1}^{\infty} k|b_k|^2 \le 1$,

$$\sum_{k \ge N+1} \frac{k |b_k^{\lambda}|^2}{r^{2k}} \le \frac{1}{r^{2(N+1)}}.$$

2.4 Bifurcation of parabolic fixed points

We recall some results that appear in [5], [4] and [19]. A good introduction to the classical part of the theory can also be found in [21]. See also [3] for the Inou-Shishikura part.

2.4.1 Convergence with domain

We consider families of analytic maps $\varphi_{\lambda} : \mathcal{D}(\varphi_{\lambda}) \to \mathbb{C}$ defined on some open subsets $\mathcal{D}(\varphi_{\lambda})$ of a Riemann surface S, where λ belongs to a subset Λ of a metric topological space. Let λ_0 be in the closure of Λ .

Let $\psi : \mathcal{D}(\psi) \to \mathbb{C}$ be a holomorphic mapping and let U an open subset of the domain $\mathcal{D}(\psi)$ of the mapping ψ .

In what follow we will say that a family of mappings $\varphi_{\lambda} : \mathcal{D}(\varphi_{\lambda}) \to \mathbb{C}$ converge to ψ on U when $\lambda \to \lambda_0$ if, for all compact subset K of U, there is a neighbourhood N of λ_0 in $\Lambda \cup \{\lambda_0\}$ such that, for all $\lambda \in N \setminus \{\lambda_0\}$, $K \subset \mathcal{D}(\varphi_{\lambda})$, and φ_{λ} converges uniformly on K to ψ when $\lambda \to \lambda_0$ inside $N \setminus \{\lambda_0\}$, see also [19].

Note that, if $(\varphi_{\lambda})_{\lambda}$ converges to a non constant ψ on U as $\lambda \to \lambda_0$, then for any non empty open subset $V \subset U$, and for λ close enough to λ_0 (depending on V), the intersection $\psi(V) \cap \varphi_{\lambda}(V)$ has non empty interior.

We will use the following elementary lemma in conjunction with the above local uniform convergence.

Lemma 2.6. Let X be a compact metric space and Y a metric space. Let $f: X \to Y$ be a continuous mapping. Suppose that there is a compact K of Y with non empty interior, such that $f(X) \subset K$ and denote by d the distance on Y.

Then, there exists an $\varepsilon_0 > 0$ such that, for all $\varepsilon \in [0, \varepsilon_0[$, there is a compact set X_{ε} with non empty interior such that for any mapping $g: X \to Y$ such that $\sup\{d(g(x), f(x)), x \in X\} \le \varepsilon$, we have $g(X_{\varepsilon}) \subset K$.

Proof. For $\varepsilon > 0$, denote by K_{ε} the compact set $\{y \in K : d(y, \partial K) \ge \varepsilon\}$. Let ε_0 be the supremum of $\varepsilon > 0$ for which the set K_{ε} has non empty interior.

Let $\varepsilon \in [0, \varepsilon_0[$. Pick $\varepsilon' \in]\varepsilon, \varepsilon_0[$ and let $X_{\varepsilon} = f^{-1}(K_{\varepsilon'})$. Since f is continuous, X_{ε} has non empty interior.

Moreover, if $g: X_{\varepsilon} \to Y$ is such that $\sup\{d(g(x), f(x)), x \in X_{\varepsilon}\} \leq \varepsilon$, then, for all $x \in X_{\varepsilon}$, $d(g(x), K'_{\varepsilon}) \leq \varepsilon$. But if $x \in X_{\varepsilon}$ were such that $g(x) \notin K$, we would have $d(g(x), K'_{\varepsilon}) \geq \varepsilon' > \varepsilon$.

2.4.2 Fatou coordinates for parabolic and near parabolic mappings

In this section we recall some important well known results on Fatou coordinates.

The following theorem is classical. The bulk of these results can be found in [20].

In the present section, unless otherwise specified, phrases such as "open sets", "Jordan domains", etc. will refer to subsets of the complex plane.

Theorem 2.7 (Extended fatou coordinates). Let $Q_1(z) = z + z^2$, and let K_1 be its filled Julia set.

Then, there exist an open set B_1^r , Jordan domains Ω_1^a , Ω_1^r , points z^a , z^r , a simply connected domain Ω_1 , holomorphic mappings $\Phi_1^a : \mathring{K_1} \to \mathbb{C}, \ \Phi_1^r : B_1^r \to \mathbb{C}, \ \varphi_1 : \mathbb{C} \to \mathbb{C}$, simple arcs ℓ_1^a , ℓ_1^r , and closed Jordan domains S_1^a and S_1^r , satisfying the following properties.

- 1. (Petals)
 - $\begin{array}{ll} (a) \ z^{a} \in \Omega_{1}^{a}, \\ (b) \ z^{r} \in \Omega_{1}^{r} \ and \ Q_{1}(z^{r}) \in \Omega_{1}^{r}, \\ (c) \ \Omega_{1}^{a} \subset \mathring{K_{1}}, \\ (d) \ \Omega_{1} = \Omega_{1}^{a} \cup \Omega_{1}^{r} \cup \{0\} \ is \ a \ neighbourhood \ of \ 0, \\ (e) \ B_{1}^{r} = \bigcup_{n \geq 0} Q_{1}^{-n}(\Omega_{1}^{r}). \end{array}$
- 2. (Properties of the mapping on the petals)
 - (a) The mapping Q_1 is univalent on Ω_1 ,
 - (b) $Q_1(\Omega_1^a) \subset \Omega_1^a$,
 - (c) $\Omega_1^r \subset Q_1(\Omega_1^r),$
 - (d) For all $z \in \mathring{K}_1$, there is an $n \in \mathbb{N}$ such that $Q_1^{\circ n}(z) \in \Omega_1^a$.
- 3. (Fatou coordinates)
 - (a) The mapping Φ_1^a is univalent on Ω_1^a ,
 - (b) The mapping Φ_1^r is univalent on Ω_1^r ,
 - (c) The mapping $\varphi_1 : \mathbb{C} \to \varphi_1(\mathbb{C})$ coincides with the inverse of Φ_1^r on $\Phi_1^r(\Omega_1^r) \cap \mathbb{C}$,
 - (d) $\Phi_1^a(K_1) = \mathbb{C}$,
 - (e) $\varphi_1(\mathbb{C}) = \mathbb{C}$.
- 4. (Semi-conjugacy)
 - (a) $\Phi_1^a \circ Q_1 = \Phi_1^a + 1$,
 - (b) $\Phi_1^r \circ Q_1 = \Phi_1^r + 1 \text{ on } Q_1^{-1}(\Omega_1^r) \cap \Omega_1^r.$
- 5. (Normalisation)
 - $(a) \ \Phi_1^a(z) \Phi_1^r(z) \to 0 \ \text{as } z \to 0 \ \text{with } z \in \Omega_1^a \cap \Omega_1^r \ \text{and } \operatorname{Im} -1/z \to +\infty.$
- 6. (Crescents)
 - (a) $S_1^a = (\Phi_1^a)^{-1} (\{ w : \operatorname{Re} w \in [\operatorname{Re} \Phi_1^a(z^a), \operatorname{Re} \Phi_1^a(z^a) + 1] \}),$
 - (b) $S_1^r = \varphi_1(\{w : \operatorname{Re} w \in [\operatorname{Re} \Phi_1^r(z^r), \operatorname{Re} \Phi_1^r(z^r) + 1]\}),$
 - (c) $S_1^a \subset \Omega_1^a \cup \{0\},\$
 - (d) $S_1^r \subset \Omega_1^r \cup \{0\},\$
 - (e) The arcs ℓ_1^a and ℓ_1^r join 0 to itself,
 - (f) The closed domain S_1^a is bounded by ℓ_1^a and its image $Q_1(\ell_1^a)$,
 - (g) The closed domain S_1^r is bounded by ℓ_1^r and its image $Q_1(\ell_1^r)$,
 - (h) $S_1^a \cap S_1^r = \{0\}.$

Proof. From section 2.1 and propositions 2.21 and 3.2.3 of [20], there exists $\xi > 0$, such that we have domains Ω_1^a and Ω_1^r and mappings $\Phi_1^a : \Omega_1^a \to \mathbb{C}$ and $\Phi_1^r : \Omega_1^r \to \mathbb{C}$ satisfying 1c, 1d, 2a, 2b, 2c, 3a, 3b, 4 and 5a.

Moreover we may suppose that

$$\Omega_1^a = \Phi_1^a \left(\left\{ w : |\arg(w - \xi)| < \frac{2\pi}{3} \right\} \right),\tag{6}$$

$$\Omega_1^r = \Phi_1^r\left(\left\{w : |\arg(w+\xi)| > \frac{\pi}{3}\right\}\right),\tag{7}$$

where arg denotes the argument in $]-\pi,\pi]$.

Since vertical lines in $(\Phi_1^a)^{-1}(\Omega_1^a)$ are perpendicular to orbits, and since the basin of attraction of the parabolic fixed point 1 consists of points converging to 0 from the direction -1 (compare [17]), any orbit converging non trivially to 0 must intersect Ω_1^a . This shows 2d.

We can use a similar procedure as in section 4.2.3 of [20] to extend the domain of the mappings $\Phi_1^a: \Omega_1^a \to \mathbb{C}$ and $\Phi_1^r: \Omega_1^r \to \mathbb{C}$, respectively on \mathring{K}_1 and B_1^r . These extended mappings still satisfy 4.

Let $w \in \mathbb{C}$. From equation (6), it follows that there exists an $n \geq 0$ such that $w + n \in \Phi_1^a(\Omega_1^a)$. Let $z' \in \Omega_1^a$ such that $\Phi_1^a(z') = w + n$. Pick $z \in \mathbb{C}$ such that $Q_1^{\circ n}(z) = z'$. Then, $z \in K_1$ and $w = \Phi_1^a(z') - n = \Phi_1^a(z)$. Thus 3d.

Let's define $\varphi_1 = (\Phi_1^r)_{|\Omega_1^a}^{-1}$. The domain of the mapping φ_1 can be extended to \mathbb{C} (compare section 4.2.2 of Shishikura's article [20]). The mapping $\varphi_1 : \mathbb{C} \to \mathbb{C}$ is onto since the mapping Q_1 is well defined from \mathbb{C} to \mathbb{C} .

We can fix $z^a \in \Omega_1^a$ and $z^r \in \Omega_1^r$ so that points 1a and 1b are verified. Let's define S_1^a and S_1^r as in 6a and 6b. As in [20], it is possible to choose z^a and z^r so that all the properties of item 6 are satisfied.

The following can be found in [19] and has its origin in works of Ecalle, Lavaurs, Sentenac or Douady. The modern version of this is at the basis of Inou-Shishikura near parabolic renormalisation theory [12].

Theorem 2.8 (Perturbed Fatou coordinates). Let $Q_{\lambda}(z) = \lambda z + z^2$ and, for $\lambda \neq 1$, denote by $\{0, \sigma\}$ the set of fixed points of Q_{λ} . We will only consider λ such that $\lambda \neq 1$ and such that there is $\alpha \in \mathbb{C}$ satisfying $|\arg \alpha| \leq \pi/4$ and $\lambda = e^{2i\pi\alpha}$.

Then, there exist a positive real number ξ and points z^a and z^r such that the conclusions of the theorem 2.7 hold and such that we have the following.

There exists a positive real number α_0 , such that if λ and α are as above with $|\alpha| \leq \alpha_0$, there exist open sets B^a_{λ} , B^r_{λ} , simply connected domains Ω_{λ} and H_{λ} , Jordan domains Ω^a_{λ} , Ω^r_{λ} , holomorphic mappings $\Phi^a_{\lambda} : B^a_{\lambda} \to \mathbb{C}$, $\Phi^r_{\lambda} : B^r_{\lambda} \to \mathbb{C}$, $\varphi_{\lambda} : H_{\lambda} \to \mathbb{C}$, simple arcs ℓ^a_{λ} , ℓ^r_{λ} and closed Jordan domains S^a_{λ} and S^r_{λ} satisfying the following.

1. (Petals)

(a)
$$z^a \in \Omega^a_\lambda$$
,

- (b) $z^r \in \Omega^r_{\lambda}$ and $Q_{\lambda}(z^r) \in \Omega^r_{\lambda}$,
- (c) The set $\Omega_{\lambda} = \Omega_{\lambda}^{a} \cup \Omega_{\lambda}^{r} \cup \{0, \sigma\}$ is a simply connected neighbourhood of $\{0, \sigma\}$,
- $(d) \ B^a_\lambda = \bigcup_{n \ge 0} Q^{-n}_\lambda(\Omega^a_\lambda),$ $(e) \ B^r_{\lambda} = \bigcup_{n \ge 0} Q^{-n}_{\lambda}(\Omega^r_{\lambda}),$

(f)
$$H_{\lambda} = \left\{ w : \operatorname{Re} w > \xi - \operatorname{Re} \frac{1}{\alpha} \right\}$$

- 2. (a) The mapping Q_{λ} is univalent on Ω_{λ}^{a} and on Ω_{λ}^{r} .
- 3. (Perturbed Fatou coordinates)
 - (a) The mapping Φ^a_{λ} is univalent on Ω^a_{λ} ,
 - (b) The mapping Φ_{λ}^{r} is univalent on Ω_{λ}^{r} ,
 - (c) The restriction of the mapping φ_{λ} on $\left\{w: \operatorname{Re} w \in \left] -\frac{1}{2\alpha} \xi, -\xi\right[\right\}$ coincides with the inverse of Φ^r_{λ} on its image.
- 4. (Semi-conjugacy)
 - (a) If $z \in \Omega^a_{\lambda}$ is such that $Q_{\lambda}(z) \in \Omega^a_{\lambda}$, then $\Phi^a_{\lambda}(Q_{\lambda}(z)) = \Phi^a_{\lambda}(z) + 1$,
 - (b) If $z \in \Omega^r_{\lambda}$ is such that $Q_{\lambda}(z) \in \Omega^r_{\lambda}$, then $\Phi^r_{\lambda}(Q_{\lambda}(z)) = \Phi^r_{\lambda}(z) + 1$.
- 5. (Normalisation)
 - (a) There exists lifts $\widetilde{\Phi_{\lambda}^{a}}$ and $\widetilde{\Phi_{\lambda}^{r}}$ of Φ_{λ}^{a} and Φ_{λ}^{r} by $w \mapsto \frac{\sigma}{1-e^{-2i\pi\alpha w}}$ such that, $\widetilde{\Phi^r_\lambda} \circ T_\alpha = T_\alpha \circ \widetilde{\Phi^a_\lambda},$

where T_{α} denotes the translation by $-1/\alpha$.

- 6. (Crescents)
 - (a) $S^{a}_{\lambda} = (\Phi^{a}_{\lambda})^{-1} (\{ w : \operatorname{Re} w \in [\operatorname{Re} \Phi^{a}_{\lambda}(z^{a}), \operatorname{Re} \Phi^{a}_{\lambda}(z^{a}) + 1] \}),$
 - $(b) \ S^r_{\lambda} = \varphi_{\lambda} \left(\{ w : \operatorname{Re} w \in [\operatorname{Re} \Phi^r_{\lambda}(z^r), \operatorname{Re} \Phi^r_{\lambda}(z^r) + 1] \} \right),$
 - (c) $S^a_{\lambda} \subset \Omega^a_{\lambda} \cup \{0, \sigma\},\$
 - (d) $S_{\lambda}^r \subset \Omega_{\lambda}^r \cup \{0, \sigma\},\$
 - (e) Both arcs ℓ^a_{λ} and ℓ^r_{λ} join 0 to σ ,
 - (f) The closed domain S^a_{λ} is bounded by ℓ^a_{λ} and its image $Q_{\lambda}(\ell^a_{\lambda})$,
 - (g) The closed domain S_{λ}^{r} is bounded by ℓ_{λ}^{r} and its image $Q_{\lambda}(\ell_{\lambda}^{r})$,
 - (h) $S^a_{\lambda} \cap S^r_{\lambda} = \{0, \sigma\}.$

7. For all $z \in S^a_{\lambda}$, there is a $p \ge 1$ such that $Q^{\circ p}_{\lambda}(z) \in S^r_{\lambda}$, and for the smallest such p we have:

$$\Phi_{\lambda}^{r}\left(Q_{\lambda}^{\circ p}(\zeta)\right) = \Phi_{\lambda}^{a}(\zeta) - \frac{1}{\alpha} + p, \qquad (8)$$

for all ζ in $(\Phi^a_\lambda)^{-1}\left(\left\{w: \operatorname{Re} w \in \left]\xi - p, \xi + \operatorname{Re} \frac{1}{2\alpha}\right[\right\}\right) \supset \Omega^a_\lambda.$

- 8. When $\lambda \to 1$ with λ and α satisfying the above hypothesis, then we have the following convergences.
 - (a) With respect to Hausdorff metric:
 - $\begin{array}{l} i. \ \ell^a_\lambda \to \ell^a_1, \\ ii. \ \ell^r_\lambda \to \ell^r_1, \\ iii. \ S^a_\lambda \to S^a_1, \\ iv. \ S^r_\lambda \to S^r_1. \end{array}$
 - (b) With respect to Hausdorff pseudometric:
 - $i: \ \Omega^a_{\lambda} \to \Omega^a_1,$ $ii: \ \Omega^r_{\lambda} \to \Omega^r_1,$ $iii: \ \Omega_{\lambda} \to \Omega_1.$
 - (c) As mappings with domains (compare section 2.4.1):
 - $\begin{array}{ll} i. \ \Phi^a_\lambda \to \Phi^a_1 \ on \ \Omega^a_1, \\ ii. \ \Phi^r_\lambda \to \Phi^r_1 \ on \ \Omega^r_1, \\ iii. \ \varphi_\lambda \to \varphi_1 \ on \ \mathbb{C}. \end{array}$

The perturbed Fatou coordinates is also called Fatou-Douady coordinates.

Proof. Let ξ be as in theorem 2.7. One may increase the value of ξ if necessary without changing the final conclusions. Let $\tau_{\lambda}(w) = \frac{\sigma}{1-e^{-2i\pi\alpha w}}$ and let τ_{λ}^{a} and τ_{λ}^{r} be the restrictions of the mapping τ_{λ} on the respective domains $\{w: \Re e w \in]-\Re e \frac{1}{2\alpha} + \xi, \Re e \frac{1}{2\alpha} + \xi[\}$ and $\{w: \Re e w \in]-\Re e \frac{1}{2\alpha} - \xi, \Re e \frac{1}{2\alpha} - \xi[\}$. The mappings τ_{λ}^{a} and τ_{λ}^{r} are analytic diffeomorphisms onto their respective images.

Let Ω^a_{λ} be the image of $\{w : \operatorname{Re}(w - \xi) > -|\operatorname{Im} w|\}$ by τ^a_{λ} and Ω^r_{λ} the image of $\{w : \operatorname{Re}(w + \xi) < |\operatorname{Im} w|\}$ by τ^r_{λ} .

Then, as in [20] (see proposition 3.2.2 and sections 3.4.1, 3.4.3 and 4.2.3), the perturbed Fatou coordinates Φ_{λ}^{a} and Φ_{λ}^{r} are defined on Ω_{λ}^{a} and Ω_{λ}^{r} respectively. One can also define φ_{λ} the same way as before.

One can easily check that the domain of these mappings can be extended so that items 1, 2, 3 and 4 are true.

The item 5 follows from 3.4.3 of [20].

And the construction of crescents satisfying item 6 is similar to the proof of theorem 2.7. $\hfill \Box$

Remark 2.9. One can choose $\xi > 0$ so that $\Omega_1^a \subset \mathbb{D}_6$ and $\Omega_{\lambda}^a \subset \mathbb{D}_6$ for all λ close enough to 1.

2.4.3Lavaurs maps

The following result comes mainly from the work of Lavaurs [13] and can be found in Douady's paper [4].

Due to our purpose, the statement is only given with α real.

Lemma 2.10 ([4], section 1.8). Let $\alpha_n \to 0$ be a sequence of positive real numbers. Suppose that the fractional part $\left\{\frac{1}{\alpha_n}\right\}$ converges to τ . Let $\lambda_n = e^{2i\pi\alpha_n}$ and $p_n = \frac{1}{\alpha_n} - \left\{\frac{1}{\alpha_n}\right\}$. Then $Q_{\lambda_n}^{\circ p_n}$ converges on compact subsets of $\mathring{K_1}$ to the mapping

$$L_{\tau} = \varphi_1 \left(\Phi_1^a - \tau \right) : \mathring{K}_1 \to \mathbb{C}$$

Moreover, L_{τ} commutes with Q_1 and $L_{\tau}(\Omega_1^a) = \mathbb{C}$.

In [4], a mapping L_{τ} of the form $L_{\tau} = \varphi_1 (\Phi_1^a - \tau)$ is called a Lavaurs map. The above proposition can be proven by using theorem 2.8.

Proof. Let λ be as in theorem 2.8 and let $z_{\lambda} \in S_{\lambda}^{a} \setminus \{0, \sigma\}$. Then, from item 7 of that theorem, there exists $p' = p'(\lambda, z_{\lambda}) \ge 1$ minimal such that $Q_{\lambda}^{\circ p'}(z_{\lambda}) \in S_{\lambda}^{r}$. Moreover, for this p', the identity $Q_{\lambda}^{\circ p'}(z) = \varphi_{\lambda} \left(\Phi_{\lambda}^{a}(z) - \frac{1}{\alpha} + p' \right)$ is true for all z in a domain that converges to \mathring{K}_1 as $\lambda \to 1$.

For λ fixed and z close to 0, the mapping $Q_{\lambda}^{\circ k}(z)$ is close to $z \mapsto \lambda^k z$. Since the closed sets S^a_{λ} and S^r_{λ} converge respectively to S^a_1 and S^r_1 as $\lambda \to 1$, it follows that if $z_{\lambda} \in S_{\lambda}^{a}$ is close enough to 0, and if $p' \geq 1$ is minimal such that $Q_{\lambda}^{\circ p'}(z_{\lambda}) \in S_{\lambda}^{r}$, then, $p' < \frac{1}{\alpha}$.

Moreover, for this $p'(\lambda)$, equation (8) is verified. Since $p'(\lambda)$ does not depend

on z, it follows from the convergence part of theorem 2.8 that $\frac{1}{\alpha} - p'$ is bounded. Now, consider the sequence of λ_n and suppose that z_{λ_n} is chosen so that $p'(\lambda_n, z_{\lambda_n}) < \frac{1}{\alpha_n}$. Then $p_n = p'_n + k_n$ with $p'_n = p'(\lambda_n, z_{\lambda_n})$ and $k_n \ge 0$. Since the sequence $(k_n)_n$ is bounded, let k_{max} be an upper bound for k_n . Let $z \in \mathring{K}_1$. Then, for n large enough, $z, Q_{\lambda_n}(z), \ldots, Q_{\lambda_n}^{\circ k_{max}}(z)$ all belong

to $\left(\Phi_{\lambda_n}^a\right)^{-1}\left(\left\{w: \operatorname{Re} w \in \left[\xi - p'_n, \xi + \frac{1}{2\alpha_n}\right]\right\}\right)$. Then,

$$\begin{aligned} Q_{\lambda_n}^{\circ p_n}(z) &= \varphi_{\lambda_n} \left(\Phi_{\lambda_n}^a \left(Q_{\lambda_n}^{\circ k_n}(z) \right) - \frac{1}{\alpha_n} + p'_n \right) \\ &= \varphi_{\lambda_n} \left(\Phi_{\lambda_n}^a(z) - \frac{1}{\alpha_n} + p_n \right). \end{aligned}$$

Finally, the last part follows from the fact that Φ_1^a and φ_1 are onto.

Lemma 2.11. There exists C > 0 and P > 0 such that the following holds.

1. Let R > 6 and let $(\alpha_n)_n$ be a sequence of real numbers such that $\alpha_n \to 0$. Then

$$\operatorname{Area}(K_1) > \operatorname{Area}(K_{\alpha_n}) + C.$$

- 2. More precisely, if the fractional part $\left\{\frac{1}{\alpha_n}\right\}$ converges as $n \to \infty$, then there exists a non empty open set V, relatively compact in $\mathring{K_1}$, an integer n_0 and a sequence of whole numbers $(p_n)_{n\geq n_0}$ such that for all $n \geq n_0$,
 - (a) the set V is a subset of the basin of attraction of ∞ of $Q_{e^{2i\pi\alpha_n}}$,
 - (b) $Q_{e^{2i\pi\alpha_n}}^{\circ p_n}(V) \subset \mathbb{D}_R,$
 - (c) $\left| p_n \frac{1}{\alpha_n} \right| < P$,
 - (d) $\operatorname{Area}(V) > C$.

Proof. Choose a non empty open subset Ω , relatively compact in the intersection of the basin of attraction of ∞ of Q_1 with the disk of radius R centered at 0.

Let $\tau \in [0, 1]$ and let's first suppose that the sequence of fractional parts $\left\{\frac{1}{\alpha_n}\right\}$ converges to τ .

Let $V = L_{\tau}^{-1}(\Omega)$. By definition V is a non empty open subset relatively compact in \mathring{K}_1 .

The point 2a follows from the fact that $\alpha_n \to 0$. The two following points then follow from lemma 2.10 by letting $p_n = \frac{1}{\alpha_n} - \left\{\frac{1}{\alpha_n}\right\}$ and P = 2. Finally the uniform lower bound 2d on the area comes form the continuity

Finally the uniform lower bound 2d on the area comes form the continuity of the mapping $\tau \mapsto \operatorname{Area}(V)$. And the first statement of the lemma follows from this estimate.

Theorem 1.1 is simply a corollary of the previous lemma.

Lemma 2.12. Let $\tau \in [0, 1]$. Let U be a compact set with non empty interior included in the domain of attraction of ∞ of Q_1 and such that $U \subset S_1^r$. Then, there exists a sequence of compacts sets U^j with $U^0 = U$ and such that the following is true for all $j \geq 1$,

- 1. $\check{U}^j \neq \emptyset$,
- 2. $U^j \subset K_1$,
- 3. $L_{\tau}(U^j) \subset U^{j-1}$,

where $L_{\tau} = \varphi_1 \left(\Phi_1^a - \tau \right)$ is a Lavaurs map.

Proof. Let, for $j \ge 0$, $U^{j+1} = L_{\tau}^{-1}(U^j)$. By construction, $U^{j+1} \subset \mathring{K}_1$. Moreover, since L_{τ} is onto, U^{j+1} has non empty interior whenever U^j has non empty interior.

Lemma 2.13. Let $\tau \in [0, 1]$ and $\gamma \ge 0$.

Let $U \subset S_1^r$ be a compact set with non empty interior contained in the basin of attraction of ∞ of Q_1 .

Then, there exists $\alpha_0 > 0$ and a finite sequence $(U^j)_{j \leq \gamma}$ of compact sets $U^j \subset \mathring{K}_1$ with non empty interior, such that $U^0 = U$ and satisfying the following.

For all sequence of $\lambda_n = e^{2i\pi\alpha_n}$ with $\alpha_n \in [0, \alpha_0]$, such that $\{1/\alpha_n\}$ converges to τ , there exists a $n_0 \ge 0$ such that, for all $n \ge n_0$ and all $j \in \{1, \ldots, \lceil \gamma \rceil\}$,

$$Q_{\lambda_n}^{\circ p_n}(U^j) \subset U^{j-1}.$$

3 Convergence properties of the approximation of the area

Theorem 1.1 is a direct consequence of propositions 3.1 and 3.2 below.

Proposition 3.1. There exists a constant C > 0 satisfying the following.

Let $(\alpha_n)_n$ be a sequence of positive real numbers converging to 0 and $(N_n)_n$ be a sequence of whole numbers. Let $\lambda_n = e^{2i\pi\alpha_n}$.

Suppose that $N_n = o(2^{1/\alpha_n})$, then there exists $n_0 \in \mathbb{N}$ such that, for all $n \geq n_0$,

$$A(\lambda_n, 1, N_n) \ge C + \operatorname{Area} K_{\lambda_n}.$$

Proof. One must notice first that λ_n is such that $|\lambda_n| < 5$ and belongs to $\mathcal{N}_g = \{\lambda \in \mathbb{C}, G_\lambda(c_\lambda) \leq g\}$ for any $g \geq 0$.

For any subsequence $(\alpha_{n_j})_j$ such that $\left\{\frac{1}{\alpha_{n_j}}\right\}$ converges, let V and P be the open set and positive constant defined by lemma 2.11. Let R > 6. According lemma 2.11, V does not intersect the filled Julia set K_{λ_n} and there is a sequence $(p_n)_n$ of natural numbers converging to ∞ such that, for all $n \ge n_0$ big enough, $Q_{\lambda_n}^{\circ p_n}(V) \subset \mathbb{D}_R$.

For $n \ge n_0$, let $g_n = \left(\log \frac{11R}{6}\right) 2^{-p_n}$ and $r_n = e^{g_n}$. By lemma 2.4,

$$A(\lambda_n, 1, N_n) \ge \pi \left(1 - r_n^{2N_n+2}\right) + r_n^{2N_n} \left(\operatorname{Area} K_{\lambda_n} + \operatorname{Area} V\right).$$

By lemma 2.11, p_n is at a bounded distance from $\frac{1}{\alpha_n}$, and since $N_n = o\left(2^{1/\alpha_n}\right), r_n^{2p_n} = \exp\left(2^{-p_n+1}N_n\right)$ tends to 1. This implies the proposition.

Proposition 3.2. If $N_n = o(2^{1/\alpha_n})$, then

$$A(\lambda_n, 1, N_n) \to \operatorname{Area}(K_1).$$

Proof. For N fixed, $A(\lambda_n, 1, N)$ converges to A(1, 1, N) when $n \to \infty$. Moreover A(1, 1, N) converges to $\operatorname{Area}(K_1)$ when N grows, thus $\limsup A(\lambda_n, 1, N_n) \leq \operatorname{Area}(K_1)$.

Now we show that $\liminf_{n\to\infty} A(\lambda_n, 1, N_n) \geq \operatorname{Area}(K_1)$. Let, for $\varepsilon > 0$, $K_1^{\varepsilon} = \{z \in K_1 : d(z, \partial K_1) \geq \varepsilon\}$. Since, for ε small enough, K_1^{ε} is a compact subset of the interior of K_1 , there exists $m \in \mathbb{N}$ such that $Q_1^{\circ m}(K_1^{\varepsilon}) \subset \Omega_1^a$. Moreover there exists $\chi > 0$ such that

$$\max\left\{\left|\mathcal{R}e\left(\Phi_{\lambda}^{a}(z)\right)-\mathcal{R}e\left(\Phi_{\lambda}^{a}(z^{a})\right)\right|, z\in Q_{1}^{\circ m}(K_{1}^{\varepsilon})\right\}\leq\chi.$$

We may increase the value of m and χ so that the above is true also for Q_{λ_n} for all n big enough.

From point 7 of theorem 2.8, it follows that there exists a constant P independent of n such that for all $p \leq \frac{1}{\alpha_n} - P$, $Q_{\lambda_n}^{\circ p}(K_1^{\varepsilon}) \subset \mathbb{D}_6$.

As previously one concludes that, if $r_n = \left(\frac{11R}{6}\right)^{2^{-\frac{1}{\alpha_n}-P}}$, then

$$A(\lambda_n, 1, N_n) \ge \pi \left(1 - r_n^{2N_n+2}\right) + r_n^{2N_n} \operatorname{Area} K_1^{\varepsilon}.$$

Hence $\liminf A(\lambda_n, 1, N_n) \ge \operatorname{Area} K_1^{\varepsilon}$ for any $\varepsilon > 0$.

Proposition 3.3. Let $\gamma > 0$ and $\tau \in [0,1]$. Then there exists C > 0 satisfying the following.

Let $(\alpha_n)_n$ be a sequence of positive real numbers converging to 0 and such that $\left\lfloor \frac{1}{\alpha_n} \right\rfloor \to \tau$, and let $(N_n)_n$ be a sequence of whole numbers. Define $\lambda_n = e^{2i\pi\alpha_n}$. Suppose that $\log N_n \leq \frac{\gamma}{\alpha_n}$. Then, there exists $n_0 \in \mathbb{N}$ such that, for all $n \geq n_0$,

$$A(\lambda_n, 1, N_n) \ge C + \operatorname{Area} K_{\lambda_n}$$

Proof. Define $\gamma' = [\gamma] + 1$. According to lemma 2.13, there exists a finite sequence $(U^j)_{j < \gamma'}$ of compact sets $U^j \subset \check{K}_1$ with non empty interior, such that U^0 is a subset of \mathbb{D}_6 (see remark 2.9) and of the basin of attraction of ∞ of Q_{λ_n} , for all $n \ge n_0$ (for some n_0), and such that $Q_{\lambda_n}^{\circ p_n}(U^j) \subset U^{j-1}$, for $j = 1, \ldots, \gamma'$.

Then, from lemma 2.4, it follows that

$$A(\lambda_n, 1, N_n) \ge \pi \left(1 - r_n^{2N_n+2}\right) + r_n^{2N_n} \left(\operatorname{Area} K_{\lambda_n} + \operatorname{Area} U^{\gamma'}\right),$$

with $\log r_n = \log \frac{11R}{6} \cdot 2^{-p_n \gamma'}$ and $p_n = \left\lfloor \frac{1}{\alpha_n} \right\rfloor$.

Since

$$\log N_n - p_n \gamma' \le \gamma \left(\frac{1}{\alpha_n} - \left\lfloor \frac{1}{\alpha_n} \right\rfloor \right) - \left\lfloor \frac{1}{\alpha_n} \right\rfloor \to -\infty,$$

the above inequality implies the proposition.

References

- [1] Kari Astala. Area distortion of quasiconformal mappings. Acta Mathematica, 173(1):37-60, 1994.
- [2] Anna M. Benini, Aslı Deniz, Thomas Gauthier, and Tan Lei. Quo vadis? open problems related to MLC. 2012.
- [3] Xavier Buff and Arnaud Chéritat. Quadratic Julia sets with positive area. Ann. Math. (2), 176(2):673–746, 2012.
- [4] Adrien Douady. Does the julia set depend continuously on the polynomial? Proc.Symp.App.Math., (49), 1994.

- [5] Adrien Douady and John Hamal Hubbard. Étude dynamique des polynômes complexes., volume 84-85 of Publications Mathématiques d'Orsay [Mathematical Publications of Orsay]. Université de Paris-Sud, Département de Mathématiques, Orsay, 1984-85.
- [6] John H. Ewing and Glenn Schober. The area of the Mandelbrot set. Numer. Math., 61(1):59–72, 1992.
- [7] Pierre Fatou. Sur les équations fonctionnelles. Bull. Soc. Math. France, 47:161–271, 1919.
- [8] Pierre Fatou. Sur les équations fonctionnelles. Bull. Soc. Math. France, 48:33–94, 1920.
- [9] Jacek Graczyk and Grzegorz Światek. Generic hyperbolicity in the logistic family. Ann. of Math. (2), 146(1):1–52, 1997.
- [10] Jacek Graczyk and Grzegorz Świątek. The real Fatou conjecture, volume 144 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1998.
- [11] Thomas Hakon Gronwall. Some remarks on conformal representation. Annals of Mathematics, 16(1/4):pp. 72–76, 1914.
- [12] Inou Hiroyuki and Shishikura Mitsuhiro. The renormalization for parabolic fixed points and their perturbation. *preprint*, 2008.
- [13] Pierre Lavaurs. Systèmes dynamiques holomorphes: explosion de points périodiques paraboliques. PhD thesis, Université Paris-Sud, 1989.
- [14] Mikhail Yu Lyubich. On typical behavior of the trajectories of a rational mapping of the sphere. Soviet Math. Dokl., (27):22–25, 1983.
- [15] Mikhail Yu Lyubich. On the lebesgue measure of the julia set of a quadratic polynomial. *Preprint IMS at Stony Brook*, (10), 1991.
- [16] Curtis T. McMullen. Complex dynamics and renormalization, volume 135 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1994.
- [17] John Milnor. Dynamics in one complex variable, volume 160 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, third edition, 2006.
- [18] Carsten Lunde Petersen and Saeed Zakeri. On the julia set of a typical quadratic polynomial with a siegel disk. Annals of Mathematics, 159(1):pp. 1–52, 2004.
- [19] Mitsuhiro Shishikura. The hausdorff dimension of the boundary of the mandelbrot set and julia sets. Annals of Mathematics, 147(2):pp. 225–267, 1998.

- [20] Mitsuhiro Shishikura. Bifurcation of parabolic fixed points. In *The Mandelbrot set, theme and variations*, volume 274 of *London Math. Soc. Lecture Note Ser.*, pages 325–363. Cambridge Univ. Press, Cambridge, 2000.
- [21] Michel Zinsmeister. Basic parabolic implosion in five days (after A. Douady). 1997. Course notes.