

Irreducible decomposition and calculating of multiplicity of the symmetric and exterior powers representation of finite groups

Tomoyuki Tamura

May 2, 2019

Abstract

In this paper* we consider symmetric powers representation and exterior powers representation of finite groups, which generated by the representation which has finite dimension over the complex field. We calculate the multiplicity of irreducible component of two representations of some representation by using a character theory of representation and a pre- λ -ring, for example, the regular representation.

Introduction

Let G be a finite group. For any representation $\rho : G \rightarrow GL(V)$ where V is a finite dimension vector space over the complex field \mathbb{C} , we obtain the representation whose representation space is a i -th symmetric power space $S^i(V)$ (it is called to the i -th symmetric power representation) and the representation whose representation space is a i -th exterior power space $\Lambda^i(V)$ (it is called to the i -th exterior power representation) for any nonnegative integers $i \geq 0$ as follows.

$$\begin{aligned} S^i \rho & : G \rightarrow GL(S^i(V)), S^i(\rho)(g)(v_1 \cdots v_i) := (\rho(g)v_1) \cdots (\rho(g)v_i) \\ \Lambda^i \rho & : G \rightarrow GL(\Lambda^i(V)), \Lambda^i \rho(g)(v_1 \wedge \cdots \wedge v_i) := \rho(g)v_1 \wedge \cdots \wedge \rho(g)v_i. \end{aligned}$$

for any $g \in G$ and $v_1, \dots, v_i \in V$. Let V_1, V_2, \dots, V_k be all of irreducible representation spaces over \mathbb{C} of G . Since the representation $S^i(V)$ and $\Lambda^i(V)$ are the representation of G and over \mathbb{C} , so these representations are completely reducible from Maschkes Theorem and isomorphic to direct sum of some V_1, V_2, \dots, V_k as the representation of G . In addition, representations $S^i(V)$ and $\Lambda^i(V)$ have a finite dimension over \mathbb{C} too since V has a finite dimension over \mathbb{C} . Hence, these representation are isomorphic to direct sum of irreducible representation, and

*2010 Mathematics Subject Classification. Primary 20C15, Secondary 20C05; 13F99.
Keywords and Phrases; Symmetric powers representation; exterior powers representation; finite group; character table; λ -ring.

every irreducible component appears finitely, and uniquely. On the other words, we have

$$\begin{aligned}
 S^i(V) &\cong \overbrace{V_1 \oplus \cdots \oplus V_1}^{m_{i,1}} \oplus \overbrace{V_2 \oplus \cdots \oplus V_2}^{m_{i,2}} \oplus \cdots \oplus \overbrace{V_k \oplus \cdots \oplus V_k}^{m_{i,k}} \\
 \wedge^i(V) &\cong \overbrace{V_1 \oplus \cdots \oplus V_1}^{n_{i,1}} \oplus \overbrace{V_2 \oplus \cdots \oplus V_2}^{n_{i,2}} \oplus \cdots \oplus \overbrace{V_k \oplus \cdots \oplus V_k}^{n_{i,k}}
 \end{aligned}$$

and nonnegative integers $m_{i,1}, m_{i,2}, \dots, m_{i,k}, n_{i,1}, n_{i,2}, \dots, n_{i,k}$ are determined finitely and uniquely. The purpose of this paper is that calculating about it.

We consider about this character χ since V has a finite dimension. A set of all maps from G to \mathbb{C} , which is said to be $\text{Map}(G, \mathbb{C})$, becomes to pre- λ -ring (In fact, it is λ -ring). A character χ belongs to $\text{Map}(G, \mathbb{C})$. In addition, $S^i(\chi)$, where maps S^i are called to symmetric powers operation, implies that the character of the i -th symmetric powers representation, and $\lambda^i(\chi)$, where maps λ^i is called to λ -operation, implies that the character of the i -th exterior powers representation. we calculate $S^i(\chi)$ and $\lambda^i(\chi)$, and be expressed by sum of irreducible character.

Actually the general method of calculation of multiplicity using of character theory is known in the [2]. In this paper, we describe the how to calculate more efficient on the method of [2] about some finite groups and some representation.

The structure of the paper, §1 to §4 is a preparation, which is known as very well. It is discussed more detail than this paper in some reference. Main result of this paper describe in §5 and §6, however about §5.1 is described about the method which is written in [2]. §1 gives the definition and properties of a semiring and the ring completion. In §2, we provide a theory of symmetric polynomial and logarithmic derivation. In §3, we describe the definition and properties of pre- λ -ring. At last, we mix §1, §2 and §3 in §4, and consider the relation of characters and λ -operation.

In §5.2, we describe the method of generating function of multiplicity. In §5.3 to §5.5, we discuss about some representation space, which are one-dimension representation in §5.3, a regular representation and that generated by action G to the quotient G/N where N is a normal subgroup of G , in §5.4 and made by one-dimension representation of N that N is contained in a center of G in §5.5. In §5.6, we describe the relation of the irreducible and reducible representation. In §5.7, we introduce the method which is using the quotient group of G . At last we calculate the irreducible representation which has dimension more than 2 in some finite group in §6.

Contents

1 Semiring	1
1.1 Semiring	1
1.2 The ring completion	1

2	Polynomial and formal power series	2
2.1	Derivative of a formal power series	2
2.2	Symmetric polynomial	2
2.3	The inverse of multiplication and the logarithmic derivative	3
3	λ-operation	9
3.1	λ -operation	9
3.2	Adams operation	10
3.3	Symmetric powers operation	11
3.4	λ -operation and homomorphism	12
3.5	Binomial coefficient	14
4	Corresponding of character	16
5	Calculating method	20
5.1	Formulations of operations of a pre- λ -ring	20
5.2	The generating function of multiplicity	21
5.3	An one-dimension representation	23
5.4	A normal subgroup and the permutation representation	24
5.5	A normal subgroup and its one-dimension representation	30
5.6	Irreducible and reducible characters	34
5.7	Calculating that use the quotient group	35
6	Calculation of some finite groups	39
6.1	A symmetric group S_3	39
6.2	An alternating group A_4	40
6.3	A non-commutative group G_{21} which has order 21	41
6.4	A symmetric group S_4	42
6.5	An alternating group A_5	44
6.6	The dihedral group D_{2n}	47
6.7	The generalized quaternion group Q_{4n}	50
6.8	Heisenberg group modulo p H_p	52
7	Future work	54

1 Semiring

First, we describe about semiring, and the group and ring completion. we will be used it a set of isomorphism class of representation and its character. We can see more about semiring in §9.3 of [1] in detail.

In this paper , Assume that any ring has a unit element 1.

1.1 Semiring

Definition 1.1.1. We call that a semiring is a triple $(S, "+", "\cdot")$, where S is a set, "+" is the additive operation, "\cdot" is the multiplicative operation, and it is satisfied all the axioms of a ring except the existence of a negative or additive inverse.

Definition 1.1.2. A morphism from a semiring S to a semiring S' is the function $f : S \rightarrow S'$ such that $f(r + s) = f(r) + f(s)$, $f(rs) = f(r)f(s)$, $f(0) = 0$, $f(1) = 1$ for any elements r, s in S .

1.2 The ring completion

Next, we describe the ring completion of a semiring.

Definition 1.2.1. The ring completion of a semiring S is a pair (S^*, θ) , where S^* is a ring and θ is a morphism from a semiring S to a ring S^* such that if $f : S \rightarrow R$ is a morphism of semiring, there exists a ring morphism $g : S^* \rightarrow R$ such that $g \circ \theta = f$. Moreover, g is required to be unique.

In [1], Husemoller describe that there is a ring completion of any semiring S . We consider the set $S \times S$, and put the following equivalence relation. we define that two elements (a, b) and (a', b) in $S \times S$ are equivalent if there is a element $c \in S$ such that $a + b' + c = a' + b + c$ holds. Let $\langle a, b \rangle$ denote the equivalent class of (a, b) , and let S^* denote the set of all equivalent classes $\langle a, b \rangle$. S^* becomes a ring with operators $\langle a, b \rangle + \langle c, d \rangle := \langle a + c, b + d \rangle$, $\langle a, b \rangle \langle c, d \rangle = \langle ac + bd, ad + bc \rangle$, which is well-defined.

Proposition 1.2.2. (1) *The function $\theta : S \rightarrow S^*$ by $\theta(a) = \langle a, 0 \rangle$ is a morphism of a semiring.*

(2) *(The group completion) Let A be an abelian with the additive operator "+", and let $f : (S, "+") \rightarrow (A, "+")$ be a morphism of monoid. Then, there exists a group homomorphism uniquely $g : (S^*, "+") \rightarrow (A, "+")$ such that $g \circ \theta = f$.*

(3) *(The ring completion) Let R be a ring, and let $f : S \rightarrow R$ be a morphism of semiring. Then, there exists a ring homomorphism uniquely $g : S^* \rightarrow R$ such that $g \circ \theta = f$.*

A map g of this Proposition is constructed to $g(\langle a, b \rangle) = f(a) - f(b)$.

From this Proposition, we can see that there exists a ring completion of any semirings.

2 Polynomial and formal power series

The purpose of this chapter, we consider a polynomial, a formal power series, a symmetric polynomial, and those properties and logarithmic derivative, which will be used in next chapter, λ -ring. Let R be a commutative ring. We call $R[x_1, \dots, x_n]$ that a polynomial ring over R in n independent variables x_1, \dots, x_n , and we call $R[[x_1, \dots, x_n]]$ that a set of formal power series over R in n independent variables x_1, \dots, x_n , for each $n \in \mathbb{N}$. we can see about symmetric polynomial in [4], and we can see about logarithmic derivative in §3 of [3].

2.1 Derivative of a formal power series

Let $R[[t]]$ be a set of formal power series over R in variable t . we define a derivation of a formal power series $f(t) = \sum_{i=0}^{\infty} a_i t^i$ in $R[[t]]$ by

$$\frac{d}{dt}f(t) := \sum_{i=1}^{\infty} i a_i t^{i-1}.$$

This Definition is different of the Definition in the real functions, which is using a property of limit. However, we can also use next Proposition in this Definition.

Proposition 2.1.1. *The follow statements hold for elements a, b in R and f, g in $R[[t]]$.*

- (1) $\frac{d}{dt}(af + bg)(t) = a \frac{d}{dt}f(t) + b \frac{d}{dt}g(t).$
- (2) $\frac{d}{dt}(fg)(t) = f(t) \left(\frac{d}{dt}g(t) \right) + \left(\frac{d}{dt}f(t) \right) g(t).$
- (3) *If the constant term of g is 0, one has $\frac{d}{dt}f(g(t)) = \left(\frac{d}{dt}f \right)(g(t)) \frac{d}{dt}g(t).$*

2.2 Symmetric polynomial

Let S_n be a symmetric group each $n = 1, 2, 3, \dots$. For each $\sigma \in S_n$ and polynomial $f \in \mathbb{Z}[[x_1, \dots, x_n]]$, which coefficient are integer, we define $(\sigma f)(x_1, \dots, x_n)$ by $f(x_{\sigma^{-1}(1)}, \dots, x_{\sigma^{-1}(n)})$. Also, we call $\Lambda(X^n)$ as follows, and its element is called a symmetric polynomial.

$$\Lambda(X^n) := \{ f \in \mathbb{Z}[[x_1, \dots, x_n]] \mid \sigma f = f \text{ for } \forall \sigma \in S_n \}$$

For each $k = 1, \dots, n$, a k -th elementary symmetric polynomial e_k on x_1, \dots, x_n is denoted by the following statement.

$$e_k := \sum_{1 \leq i_1 < \dots < i_k \leq n} x_{i_1} \cdots x_{i_k}. \quad (2.1)$$

An elementary symmetric polynomial belongs to $\Lambda(X^n)$, and the elementary symmetric polynomials are algebraically independent. Hence, we can consider the following evaluation map of n elements a_1, \dots, a_n in R . It is a ring homomorphism.

$$F : \Lambda(X^n) = \mathbb{Z}[e_1, \dots, e_n] \rightarrow R, F(e_j) = a_j \quad (j = 1, \dots, n). \quad (2.2)$$

Next, we denote a symbol of complete symmetric polynomial and power-sum symmetric polynomial. For each $k \in \mathbb{N}$, $h_k(e_1, \dots, e_n)$ is a k -th complete symmetric polynomial on x_1, \dots, x_n , which is expressed by polynomial of e_1, \dots, e_n . Similarly, $Q_k(e_1, \dots, e_n)$ is a k -th power-sum symmetric polynomial on x_1, \dots, x_n , which is expressed by polynomial of e_1, \dots, e_n . On the other hand, the following statement holds.

$$h_k(e_1, \dots, e_n) := \sum_{1 \leq i_1 \leq \dots \leq i_k \leq n} x_{i_1} \cdots x_{i_k}, \quad (2.3)$$

$$Q_k(e_1, \dots, e_n) := x_1^k + \cdots + x_n^k. \quad (2.4)$$

2.3 The inverse of multiplication and the logarithmic derivative

Definition 2.3.1. For any ring homomorphism $F : R_1 \rightarrow R_2$, we define a map $F_\Lambda : R_1[[t]] \rightarrow R_2[[t]]$ by following equality for any formal power series $\sum_{i=0}^{\infty} a_i t^i$.

$$F_\Lambda \left(\sum_{i=0}^{\infty} a_i t^i \right) := \sum_{i=0}^{\infty} F(a_i) t^i. \quad (2.5)$$

F_Λ is a ring homomorphism from $R_1[[t]]$ to $R_2[[t]]$.

In the set of symmetric polynomial $\Lambda(X^n)[[t]]$, By using (2.1), (2.3) and the generating function of elementary and complete symmetric polynomials, we obtain the following equality.

$$\prod_{i=1}^n (1 + x_i t) = 1 + \sum_{i=1}^n e_i t^i, \quad (2.6)$$

$$\prod_{i=1}^n \left(\frac{1}{1 - x_i t} \right) = 1 + \sum_{i=1}^n h_i(e_1, \dots, e_n) t^i. \quad (2.7)$$

Hence, we can see the next Lemma from (2.2), (2.5) and (2.6).

Lemma 2.3.2. In any ring R and n element a_1, \dots, a_n in R , we define $F : \Lambda(X^n) \rightarrow R$ be a evaluation map by $F(e_j) = a_j$ ($j = 0, 1, \dots, n$), so we obtain

$$F_\Lambda \left(\prod_{i=1}^n (1 + x_i t) \right) = 1 + a_1 t + a_2 t^2 + \cdots + a_n t^n. \quad (2.8)$$

Let R be a commutative ring. We define the strict power series over R by the formal power series with coefficients in R and it has constant term 1. and $\Lambda(R)$ is defined by the set of all strict power series over R . For any ring homomorphism $F : R_1 \rightarrow R_2$. It is clear that $F_\Lambda(\Lambda(R_1)) \subset \Lambda(R_2)$ holds.

[2] and [3] called $\Lambda(R)$ the universal- λ -ring over R , which has a λ -ring structure as the name represents. However, we do not touch about it except of its addition, which is the multiplication of $R[[t]]$. About it, we have the next Proposition.

Proposition 2.3.3. $\Lambda(R)$ has a structure of group with the multiplication of formal power series over R . Also, let $f = 1 + \sum_{i=1}^{\infty} a_i t^i$ be a strict power series over R , and let inverse of f be $f^{-1} = 1 + \sum_{i=1}^{\infty} b_i t^i$. Then the coefficient b_i is a polynomial with integer coefficients in a_1, \dots, a_i . Moreover, that $b_i = (-1)^i h_i(a_1, \dots, a_i)$ are hold for each $i \geq 1$.

Proof. It is clear that $\Lambda(R)$ is closed about the multiplication, and has a unit element 1, we consider the inverse of multiplication. Let $f = 1 + \sum_{i=1}^{\infty} a_i t^i \in \Lambda(R)$ be a strict power series over R . Then we let b_n be the following equality by induction of $n \in \mathbb{N}$.

$$\begin{aligned} b_1 &:= -a_1, \\ b_n &:= -(b_{n-1}a_1 + b_{n-2}a_2 + \dots + b_1a_{n-1} + a_n) \quad (n \geq 2). \end{aligned} \tag{2.9}$$

It follows that $\sum_{p+q=n} b_p a_q = 0$, we have $(1 + \sum_{n=1}^{\infty} b_n t^n) f = 1$, which implies that we finish to proof of the existence of inverse $f^{-1} = 1 + \sum_{i=1}^{\infty} b_i t^i$. In particular, b_n is a polynomial with integer coefficients in a_1, \dots, a_n from (2.9), hence the coefficient of t^n in $g_n = 1 + a_1 t + \dots + a_n t^n$ is equal to b_n . Also, let $F_n : \Lambda(X^n) \rightarrow R$ be a evaluation map by setting $F_n(e_j) = a_j$ ($j = 1, 2, \dots, n$), we have

$$\begin{aligned} g_n^{-1} &= F_{n\Lambda} \left(\prod_{j=1}^n (1 + x_j t) \right)^{-1} \\ &= F_{n\Lambda} \left(\prod_{j=1}^n \frac{1}{1 - x_j(-t)} \right) \\ &= F_{n\Lambda} \left(1 + \sum_{j=1}^{\infty} h_j(e_1, \dots, e_n) (-t)^j \right) = 1 + \sum_{j=1}^{\infty} (-1)^j h_j(a_1, \dots, a_n) t^j \end{aligned}$$

from (2.7), (2.8). Comparing of coefficients of t^n in this equality, we have $b_n = (-1)^n h_n(a_1, \dots, a_n)$. \square

Definition 2.3.4. We define the map $\frac{d}{dt} \log : \Lambda(R) \rightarrow R[[t]]$ by $\frac{d}{dt} \log(f) := f' f^{-1}$ for any $f \in \Lambda(R)$.

Definition 2.3.4 is given the result of logarithmic derivation of the real function, then we call this map the logarithmic derivation too. We can define it since there is a inverse element for any elements of $\Lambda(R)$, which is domain of it.

From Proposition 2.1.1 (2), the follow statement hold for any $f, g \in \Lambda(R)$.

$$\frac{d}{dt} \log(fg) = \frac{d}{dt} \log(f) + \frac{d}{dt} \log(g). \quad (2.10)$$

It implies that the logarithmic derivation is a homomorphism from the multiplication of $\Lambda(R)$ to the addition of $R[[t]]$.

Proposition 2.3.5. *For any ring homomorphism $F : R_1 \rightarrow R_2$, the logarithmic derivation and a map $F_\Lambda : \Lambda(R_1) \rightarrow \Lambda(R_2)$ are commutative on $\Lambda(R_1)$. In the other words, The following formula is hold on $\Lambda(R_1)$.*

$$\frac{d}{dt} \log \circ F_\Lambda = F_\Lambda \circ \frac{d}{dt} \log.$$

Proof. From Proposition 2.3.3, we have

$$\begin{aligned} F_\Lambda \left(\frac{d}{dt} \log f(t) \right) &= F_\Lambda \left(\left(\sum_{i=1}^{\infty} i a_i t^{i-1} \right) \left(1 + \sum_{i=1}^{\infty} (-1)^i h_i(a_1, \dots, a_i) t^i \right) \right) \\ &= \left(\left(\sum_{i=1}^{\infty} i f(a_i) t^{i-1} \right) \left(1 + \sum_{i=1}^{\infty} (-1)^i h_i(f(a_1), \dots, f(a_i)) t^i \right) \right) \\ &= \frac{d}{dt} \log(F_\Lambda(f)). \end{aligned}$$

for all $f(t) = 1 + \sum_{i=1}^{\infty} a_i t^i \in \Lambda(R_1)$. □

Next, we say the relation about the strict power series and what is mapping by the logarithmic derivation.

Lemma 2.3.6. *Let a_1, a_2, \dots be the series of elements of R , and let f_n be equal to $1 + a_1 t + a_2 t^2 + \dots + a_n t^n \in \Lambda(R)$ for each $n \in \mathbb{N}$. Then, the followig statements hold.*

- (1) For integer $n \in \mathbb{N}$, we have $-t \frac{d}{dt} \log f_n(t) = \sum_{i=1}^{\infty} (-1)^i Q_i(a_1, \dots, a_n) t^i$.
- (2) For integers $k, n \in \mathbb{N}$ with $k \leq n$, we have $Q_k(a_1, \dots, a_n) = Q_k(a_1, \dots, a_k)$.
- (3) For integers $k, n \in \mathbb{N}$ with $k > n$, we have $Q_k(a_1, \dots, a_n) = Q_k(a_1, \dots, a_n, 0, \dots, 0)$, where the numbers of 0 of the right hand side is $k - n$.

Proof. (1) We consider a evaluation map $F_n : \Lambda(X^n) \rightarrow R$ setting by $F_n(e_j) =$

a_j ($j = 1, 2, \dots, n$). From (2.4), (2.8) and Proposition 2.3.5, we have

$$\begin{aligned}
-t \frac{d}{dt} \log f_n(t) &= -t \frac{d}{dt} \log F_{n\Lambda} \left(\prod_{j=1}^n (1 + x_j t) \right) \\
&= F_{n\Lambda} \left(-t \frac{d}{dt} \log \prod_{j=1}^n (1 + x_j t) \right) \\
&= F_{n\Lambda} \left(-t \sum_{j=1}^n \frac{d}{dt} \log(1 + x_j t) \right) \\
&= F_{n\Lambda} \left(-t \sum_{j=1}^n \sum_{i=0}^{\infty} x_j^{i+1} (-t)^i \right) \\
&= F_{n\Lambda} \left(\sum_{i=1}^{\infty} (-1)^i Q_i(e_1, \dots, e_n) t^i \right) = \sum_{i=1}^{\infty} (-1)^i Q_i(a_1, \dots, a_n) t^i.
\end{aligned}$$

(2) The coefficient of t^k in $-t \frac{d}{dt} \log f_n(t)$ is expressed by the polynomial with integer coefficients in a_1, \dots, a_k , so it is equal to the coefficient of t^k in $-t \frac{d}{dt} \log f_k(t)$. so, we have $Q_k(a_1, \dots, a_k) = Q_k(a_1, \dots, a_n)$. from (1).

(3) The polynomial f_n can be seen as $f_n = 1 + a_1 t + a_2 t^2 + \dots + a_n t^n + 0t^{n+1} + \dots + 0t^k$. Hence we have

$$-t \frac{d}{dt} \log f_n(t) = \sum_{i=1}^{\infty} (-1)^i Q_i(a_1, \dots, a_n, 0, \dots, 0) t^i \quad (2.11)$$

from (1). On the other hand, we have

$$-t \frac{d}{dt} \log f_n(t) = \sum_{i=1}^{\infty} (-1)^i Q_i(a_1, \dots, a_n) t^i \quad (2.12)$$

from (1). Therefore, The equalities $Q_k(a_1, \dots, a_n) = Q_k(a_1, \dots, a_n, 0, \dots, 0)$ is given by comparing of the coefficient of t^k in (2.11) and (2.12). \square

Proposition 2.3.7. *The following statements are equivalent for any series $f(t) = 1 + \sum_{i=1}^{\infty} a_i t^i$ and $g(t) = \sum_{i=1}^{\infty} b_i t^i$ in $R[[t]]$.*

- (1) $g(-t) = -t \frac{d}{dt} \log f(t)$.
- (2) $b_n = Q_n(a_1, \dots, a_n)$ for $n \geq 1$.
- (3) $\sum_{i=0}^{n-1} (-1)^i a_i b_{n-i} = (-1)^{n+1} n a_n$ for $n \geq 1$.

Proof. Let f_n be $1 + a_1t + a_2t^2 + \cdots + a_nt^n \in \Lambda(R)$ for each $n \in \mathbb{N}$. The coefficients of t^n in $-t \frac{d}{dt} \log f(t)$ is a polynomial with integer coefficients in a_1, \dots, a_n , then it is equal to the coefficients of t^n in $f_n(t)$. Then, we can see that the coefficient of t^n in $-t \frac{d}{dt} \log f(t)$ is equal to $(-1)^n Q_n(a_1, \dots, a_n)$ from Lemma 2.3.6 (1). Hence we finish to proof of that (1) and (2) are equivalent.

Next, we consider that (1) and (3) are equivalent. From (1), we have

$$\sum_{i=1}^{\infty} b_i (-t)^i = -t \frac{d}{dt} \log \left(1 + \sum_{i=1}^{\infty} a_i t^i \right) = \left(- \sum_{i=0}^{\infty} i a_i t^i \right) \left(1 + \sum_{i=1}^{\infty} a_i t^i \right)^{-1}$$

Then, we obtain

$$\left(\sum_{i=1}^{\infty} (-1)^i b_i t^i \right) \left(1 + \sum_{i=1}^{\infty} a_i t^i \right) = - \sum_{i=0}^{\infty} i a_i t^i$$

Now we compare the coefficients of t^n and multiple by $(-1)^n$, we have the equalities (3) for each $n \in \mathbb{N}$. we can also see the proof of (1) assumption (3) by following this reverse. \square

We define maps z_n and z_t , which is defined by the logarithmic derivation, and consider one of the condition of commutative ring, \mathbb{Z} -torsion-free and \mathbb{Q} -algebra.

Definition 2.3.8. A commutative ring R is said to be \mathbb{Z} -torsion-free if whenever $na = 0$ for some element $a \in R$ and integer $n \neq 0$, then $a = 0$. and a commutative ring R is said to be \mathbb{Q} -algebra if it is equipped with ring homomorphism from the rational numbers set \mathbb{Q} to R .

We define maps $f_n : R \rightarrow R$ by $f_n(r) = nr$ ($r \in R$) for each integer $n \neq 0$. Then, the map f_n is injective for any integer $n \neq 0$ if and only if R is \mathbb{Z} -torsion-free.

Moreover, If R is \mathbb{Q} -algebra, we can define $\frac{r}{n} := \frac{1}{n}r \in R$ for any $r \in R$ and integer $n \neq 0$. In particular, R is \mathbb{Z} -torsion-free.

Definition 2.3.9. For each $n \in \mathbb{N}$, we define maps $z_n : \Lambda(R) \rightarrow R$ by that $z_n(f)$ is the coefficient of t^n in $-t \frac{d}{dt} \log(f(t))$ multiplied $(-1)^n$. And, we define a map $z_t : \Lambda(R) \rightarrow \text{Map}(\mathbb{N}, R)$ by $z_t(f)(n) := z_n(f)$ for each strict power series $f \in \Lambda(R)$.

Since its Definition, we have

$$\sum_{i=1}^{\infty} z_n(f)(-t)^i = -t \frac{d}{dt} \log f(t) \tag{2.13}$$

for each $f \in \Lambda(R)$.

For a ring R , $\text{Map}(\mathbb{N}, R)$, which is used in next Proposition, is a commutative ring with the operation $(\alpha_1 + \alpha_2)(n) = \alpha_1(n) + \alpha_2(n)$, $(\alpha_1 \alpha_2)(n) = \alpha_1(n) \alpha_2(n)$.

Proposition 2.3.10. *The following statements hold about maps z_t and z_n .*

- (1) *We have $z_n(f) = Q_n(a_1, \dots, a_n)$ for any $f = 1 + \sum_{i=1}^{\infty} a_i t^i \in \Lambda(R)$*
- (2) *The map z_t is the homomorphism from the multiplication of $\Lambda(R)$ to the addition of $\text{Map}(\mathbb{N}, R)$. In particular, the map z_n is the homomorphism from the multiplication of $\Lambda(R)$ to the addition of R , for each $n \in \mathbb{N}$.*
- (3) *If R is \mathbb{Z} -torsion-free, then the map z_t is injective.*
- (4) *If R is \mathbb{Q} -algebra, then the map z_t is bijective.*

Proof. (1) It is clear from Proposition 2.3.7 (2) and (2.13).

(2) Since we have $z_t(fg) = z_t(f) + z_t(g)$ for each $f, g \in \Lambda(R)$ from (2.10) and (2.13), it is proved.

(3) Assume that R is \mathbb{Z} -torsion-free. Let h be a strict power series over R with $z_t(h) = 0$. we have

$$0 = -t \frac{d}{dt} \log(h) = h' h^{-1}$$

which imply $h' = 0$, and we have $h = 0$ by assumption that R is \mathbb{Z} -torsion-free.

(4) Assume that R is \mathbb{Q} -algebra, and let α be an element of $\text{Map}(\mathbb{N}, R)$. So, we let a_n be the following equality by induction of $n \in \mathbb{N}$.

$$a_1 := \alpha(1) \ (n = 1), \ a_n := \frac{(-1)^{n+1}}{n} \left(\sum_{i=0}^{n-1} (-1)^i a_i \alpha(n-i) \right) \ (n \geq 2).$$

Then, we have

$$\sum_{i=1}^{\infty} \alpha(i) (-t)^i = -t \frac{d}{dt} \log \left(1 + \sum_{i=1}^{\infty} a_i t^i \right)$$

from Proposition 2.3.7, and we obtain

$$z_t \left(1 + \sum_{i=1}^{\infty} a_i t^i \right) = \alpha,$$

which imply that the function z_t is surjective. Hence, we have that the function z_t is bijective with (3). \square

3 λ -operation

In this section, we describe the definition of a pre- λ -ring, which has axioms less than a λ -ring. we can see more in [1] §13, [2], [3], and [7].

3.1 λ -operation

Definition 3.1.1. A λ -semiring is a semiring S together with functions $\lambda^n : S \rightarrow S$ ($n = 0, 1, 2, \dots$) called λ -operation, such that for all $r, s \in S$, the following axioms are satisfied.

$$(L1) \quad \lambda^0(r) = 1.$$

$$(L2) \quad \lambda^1 = \text{id}_S.$$

$$(L3) \quad \lambda^n(r + s) = \sum_{i+j=n} \lambda^i(r)\lambda^j(s).$$

We call S a pre- λ -ring if S is a commutative ring.

Let R be a commutative ring, and let $\lambda^n : R \rightarrow R$ ($n = 0, 1, 2, \dots$) be maps. We define a map $\lambda_t : R \rightarrow R[[t]]$ by

$$\lambda_t(r) := \sum_{i=0}^{\infty} \lambda^i(r)t^i = \lambda^0(r) + \lambda^1(r)t + \lambda^2(r)t^2 + \dots \quad (r \in R) \quad (3.1)$$

which considered as a formal power series in t with coefficients in R . It is clear that axiom (L1) holds if and only if $\text{Im}(\lambda_t) \subset \Lambda(R)$, and axiom (L3) holds if and only if $\lambda_t(r + s) = \lambda_t(r)\lambda_t(s)$ for all $r, s \in R$. Now, we obtain the next lemma.

Lemma 3.1.2. *Let R be a commutative ring, and let $\lambda^n : R \rightarrow R$ ($n = 0, 1, 2, \dots$) be maps. We define the function $\lambda_t : R \rightarrow R[[t]]$ by (3.1), so the followings statements are equivalent.*

- (1) *A ring R is a pre- λ -ring together with functions $\{\lambda^n\}_{n=0,1,2,\dots}$*
- (2) *$\lambda_t(r + s) = \lambda_t(r)\lambda_t(s)$, and $\lambda_t(r) = 1 + rt + \dots$ for all $r, s \in R$.*

Example 3.1.3. The rings \mathbb{Z} of integers is a pre- λ -ring with $\lambda_t(n) = (1 + t)^n$ for $n \in \mathbb{Z}$. we will consider in detail with §3.5.

Example 3.1.4. For a finite group G , let $M_{\mathbb{C}}(G)$ denote the set of isomorphism classes $\{V\}$ of finite dimension representation of G over \mathbb{C} . The operation $\{V\} + \{W\} := \{V \oplus W\}$ and $\{V\}\{W\} := \{V \otimes_{\mathbb{C}} W\}$ make $M_{\mathbb{C}}(G)$ into a semiring. the functions $\lambda^n : M_{\mathbb{C}}(G) \rightarrow M_{\mathbb{C}}$, $\lambda^n(\{V\}) = \{\wedge^n(V)\}$ defines a λ -semiring structure on $M_{\mathbb{C}}(G)$. we will consider in detail with §4.

3.2 Adams operation

In this section, we define Adams operations in a pre- λ -ring and establish some of their properties. Moreover, we describe a pre- ψ -ring, which is a ring R together with ring endomorphisms of addition ψ^k that behave like the Adams operations in a pre- λ -ring, and it gives a pre- λ -ring structure on \mathbb{Q} -algebra.

Definition 3.2.1. For each $n = 1, 2, 3, \dots$ define the n -th Adams operation $\psi^n : R \rightarrow R$ by $\psi^n = z_n \circ \lambda_t$. In other words, we defined the Adams operation by the following equality holds

$$\psi_{-t}(r) = -t \frac{d}{dt} \log(\lambda_t(r))$$

for $r \in R$, where $\psi_t(r) = \sum_{i=1}^{\infty} \psi^i(r) t^i$.

Proposition 3.2.2. *The following statements hold in any pre- λ -ring R .*

- (1) $\psi^n(r) = Q_n(\lambda^1(r), \dots, \lambda^n(r))$.
- (2) $\psi^n(r) - \lambda^1(r)\psi^{n-1}(r) + \dots + \lambda^{n-1}(r)\psi^1(r) = (-1)^{n+1} n \lambda^n(r)$.

for all $n = 1, 2, 3, \dots, r \in R$.

Proof. Statement (1) follows from the definition and Proposition 2.3.10 (1), and statement (2) follows from the definition and Proposition 2.3.7.

Statement (2) of Proposition 3.2.2 is called Newton's formula since symmetric polynomial ring $\Lambda(X^n)$ is a λ -ring for all $n = 1, 2, 3, \dots$, and (2) gives a Newton's formula in symmetric polynomial. \square

Proposition 3.2.3. *Adams operation ψ^1 is the identity map of R . and each Adams operation is a homomorphism of the addition.*

Proof. We put $n = 1$ in statement (1) of Proposition 3.2.2. Since Lemma 3.1.2 and Proposition 2.3.10(2), ψ^n is additive. \square

Next, we define a pre- ψ -ring, and we describe its properties and \mathbb{Q} -algebra's properties, which produce a structure of pre- λ -ring.

Definition 3.2.4. A pre- ψ -ring is a commutative ring R together with homomorphism of the addition $\psi^n : R \rightarrow R$ ($n = 1, 2, \dots$) such that ψ^1 is the identity map of R .

Proposition 3.2.5. *Let S be a pre- ψ -ring and a \mathbb{Q} -algebra. Then, there exists a structure of pre- λ -ring on S such that Adams operation of this structure is ψ^n . Moreover, this structure is required to be unique.*

Proof. We define a map $\Psi : S \rightarrow \text{Map}(\mathbb{N}, S)$ by setting $\Psi(r)(n) = \psi^n(r)$ for all $r \in S$ and $n \in \mathbb{N}$. It is a homomorphism from the addition of S to the addition of $\text{Map}(\mathbb{N}, S)$. Furthermore, the map z_t is an isomorphism from the multiplication of $\Lambda(S)$ to the addition of $\text{Map}(\mathbb{N}, S)$ from Proposition 2.3.10.

Hence, we let λ_t be $\lambda_t := z_t^{-1} \circ \Psi$. it is a homomorphism from the addition of S to from the multiplication of $\Lambda(S)$, and

$$\lambda_t(r) = z_t^{-1} \circ \Psi(r) = 1 + rt + \dots$$

for any $r \in R$. so, λ_t gives a structure of pre- λ -ring to R , it's n -th Adams operation is ψ^n from Lemma 3.1.2. Uniqueness follows that a map $\frac{d}{dt} \log$ is bijective. \square

Example 3.2.6. For a finite group G , $\text{Map}(G, \mathbb{C})$ is a \mathbb{C} -algebra with the operation $(f_1 + f_2)(g) = f_1(g) + f_2(g)$, $(f_1 f_2)(g) = f_1(g) f_2(g)$, $(r f_1)(g) = r f_1(g)$ ($g \in G$). In particular, it is a \mathbb{Q} -algebra. We define the functions $\psi^n : \text{Map}(G, \mathbb{C}) \rightarrow \text{Map}(G, \mathbb{C})$ by

$$\psi^n(f)(g) := f(g^n) \quad (f \in \text{Map}(G, \mathbb{C}), g \in G) \quad (3.2)$$

for each $n=1,2,3,\dots$. Hence, $\text{Map}(G, \mathbb{C})$ becomes to a pre- ψ -ring, and be given a structure of pre- λ -ring such that Adams operation is (3.2).

3.3 Symmetric powers operation

In this section, we describe symmetric powers operation in a pre- λ -ring R , which is discussed in [7]. it will be used in symmetric powers representation.

Definition 3.3.1. We define the function $S_t : R \rightarrow R[[t]]$ by setting

$$S_t(r) := (\lambda_{-t}(r))^{-1} = \left(1 + \sum_{i=1}^{\infty} \lambda^i(r)(-t)^i\right)^{-1} \quad (r \in R).$$

and we define symmetric powers operation $S^n : R \rightarrow R$ by that $S^n(r)$ is coefficient of t^n in $S_t(r)$ for each $n \geq 0$.

It is clear that S^0 is a constant map of unit element of R .

Proposition 3.3.2. *The following statements hold in any pre- λ -ring R for any $n \geq 0$.*

$$S^n + (-1)\lambda^1 S^{n-1} + \dots + (-1)^{n-1} \lambda^{n-1} S^1 + \lambda^n = 0$$

Proof. It follows from definition $S_t \lambda_{-t} = 1$. \square

Proposition 3.3.3. *For any $r \in R$ and $n \in \mathbb{N}$, $S^n(r) = h_n(\lambda^1(r), \dots, \lambda^n(r))$ holds. In particular, S^1 is the identity map of R .*

Proof. By Proposition 2.3.3, we have

$$\begin{aligned} S_t(r) = (\lambda_{-t}(r))^{-1} &= 1 + \sum_{n=1}^{\infty} (-1)^n h_n(\lambda^1(r), \dots, \lambda^n(r)) (-t)^n \\ &= 1 + \sum_{n=1}^{\infty} h_n(\lambda^1(r), \dots, \lambda^n(r)) t^n. \end{aligned}$$

for any $r \in R$. Thus, comparing the coefficients of t^n , we obtain the first statement. \square

Proposition 3.3.4. *In any pre- λ -ring R , we have $S_t(r+s) = S_t(r)S_t(s)$, and $S^n(r+s) = \sum_{i+j=n} S^i(r)S^j(s)$ for $r, s \in R$, $n \in \mathbb{N}$.*

Proof. First statement follows from $S_t(r+s) = (\lambda_{-t}(r+s))^{-1} = (\lambda_{-t}(r))^{-1}(\lambda_{-t}(s))^{-1} = S_t(r)S_t(s)$. By comparing of coefficient of t^n in it, we have a second statement. \square

3.4 λ -operation and homomorphism

We discuss the morphism of λ -semiring and pre- λ -ring, and describe the relation of ring complement of semiring.

Definition 3.4.1. (1) Let S and S' be semirings. A semi- λ -homomorphism $f : S \rightarrow S'$ is a morphism of semiring such that $f \circ \lambda^n = \lambda^n \circ f$ for $n \geq 0$.
(2) Let R and R' be pre- λ -rings. A pre- λ -homomorphism $f : R \rightarrow R'$ is a ring homomorphism such that $f \circ \lambda^n = \lambda^n \circ f$ for $n \geq 0$.

About such morphism, we have next statements.

Proposition 3.4.2. (1) *The identity map of λ -semiring S is a semi- λ -homomorphism.*

(2) *Let S , S' and S'' be λ -semirings, and $f : S \rightarrow S'$ and $g : S' \rightarrow S''$ be semi- λ -homomorphisms. then $g \circ f : S \rightarrow S''$ is a semi- λ -homomorphism. In addition, These statements holds about pre- λ -ring and pre- λ -homomorphism.*

Thereafter, we call semi- λ -homomorphism and pre- λ -homomorphism to λ -homomorphism. And, If λ -homomorphism is injective, surjective, and bijective, we call it to λ -monomorphism, λ -epimorphism, λ -isomorphism each other.

Proposition 3.4.3. *Let R and R' be a pre- λ -rings, and let $f : R \rightarrow R'$ be a ring homomorphism.*

(1) *If $f : R \rightarrow R'$ is a λ -homomorphism, we have $\psi^n \circ f = f \circ \psi^n$, and $S^n \circ f = f \circ S^n$ for any $n \in \mathbb{N}$.*
(2) *Conversely, If f is satisfied $\psi^n \circ f = f \circ \psi^n$ for any $n \in \mathbb{N}$ and R' is a \mathbb{Z} -torsion-free, f is λ -homomorphism.*

Proof. (1) From (1) of Proposition 3.2.2, we have

$$\begin{aligned} \psi^n(f(r)) &= Q_n(\lambda^1(f(r)), \dots, \lambda^n(f(r))) \\ &= Q_n(f(\lambda^1(r)), \dots, f(\lambda^n(r))) \\ &= f(Q_n(\lambda^1(r), \dots, \lambda^n(r))) = f(\psi^n(r)) \end{aligned}$$

for any $r \in R$, $n \in \mathbb{N}$, Similarly, we obtain

$$\begin{aligned} S^n(f(r)) &= h_n(\lambda^1(f(r)), \dots, \lambda^n(f(r))) \\ &= h_n(f(\lambda^1(r)), \dots, f(\lambda^n(r))) \\ &= f(h_n(\lambda^1(r), \dots, \lambda^n(r))) = f(S^n(r)) \end{aligned}$$

from Proposition 3.3.3.

(2) We prove by induction of $n \in \mathbb{N}$. λ^1 is the identity map of R , so it is clear about $n = 1$. Suppose by induction that $f \circ \lambda^j = \lambda^j \circ f$ ($j = 1, \dots, n-1$). By the induction hypothesis and Newtons formula , we have

$$\begin{aligned} (-1)^{n+1}n\lambda^n(f(r)) &= \psi^n(f(r)) - \lambda^1(f(r))\psi^{n-1}(f(r)) + \dots + \lambda^{n-1}(f(r))\psi^1(f(r)) \\ &= f(\psi^n(r) - \lambda^1(r)\psi^{n-1}(r) + \dots + \lambda^{n-1}(r)\psi^1(r)) \\ &= f((-1)^{n+1}n\lambda^n(r)) = (-1)^{n+1}nf(\lambda^n(r)). \end{aligned}$$

Since R' is \mathbb{Z} -torsion-free by assumption, it follows that $\lambda^n(f(r)) = f(\lambda^n(r))$. This finishes the induction and the proof. \square

Proposition 3.4.4. *Let S be a λ -semiring, and commutative about its multiplication. and let (S^*, θ) be the ring completion of S . then, there exists a structure of pre- λ -ring in S^* such that θ is a λ -homomorphism. Moreover, this structure is required to be unique.*

Proof. Since S is commutative about it's multiplication, S^* is too. so, we can consider the formal power series over S^* . We define a map $G : S \rightarrow S^*[[t]]$ by setting

$$G(r) := \sum_{n=0}^{\infty} \theta \circ \lambda^n(r) t^n \quad (r \in S).$$

A map λ^0 in S is the constant map of unit element, so we have $\text{Im}(G) \subset \Lambda(S^*)$. Hence we have

$$\begin{aligned} G(r+s) &= \sum_{n=0}^{\infty} \theta \circ \lambda^n(r+s) t^n \\ &= \sum_{n=0}^{\infty} \theta \left(\sum_{i+j=n} \lambda^i(r) \lambda^j(s) \right) t^n \\ &= \sum_{n=0}^{\infty} \left(\sum_{i+j=n} \theta \circ \lambda^i(r) \theta \circ \lambda^j(s) \right) t^n = G(r)G(s) \end{aligned}$$

for all $r, s \in S$, which implies that G is a homomorphism from the addition of S to the multiplication of $\Lambda(S^*)$. Hence, There exists a map λ_t uniquely such that it is a homomorphism from the addition of S^* to the multiplication of $\Lambda(S^*)$, and it is satisfied $G = \lambda_t \circ \theta$, from Proposition 1.2.2(2). In addition, we have

$$\begin{aligned} \lambda_t(\langle a, b \rangle) = G(a)G(b)^{-1} &= (\theta \circ \lambda^0(a) + \theta \circ \lambda^1(a)t + \dots)(\theta \circ \lambda^0(b) - \theta \circ \lambda^1(b)t + \dots) \\ &= (1 + \langle a, 0 \rangle t + \dots)(1 - \langle b, 0 \rangle t + \dots) = 1 + \langle a, b \rangle t + \dots \end{aligned}$$

for all $\langle a, b \rangle \in S^*$. Therefore we obtain a pre- λ -ringstructure in S^* with λ_t . At last, we have

$$1 + \sum_{i=1}^{\infty} \theta \circ \lambda^i(r) t^i = G(r) = \lambda_t \circ \theta(r) = 1 + \sum_{i=1}^{\infty} \lambda^i \circ \theta(r) t^i$$

for all $r \in R$, hence comparing of coefficient of t^n gives that $\theta : S \rightarrow S^*$ is λ -homomorphsim. \square

3.5 Binomial coefficient

Let R be a \mathbb{Q} -algebra in this section, we prepare of a strict power series $(1+t)^r \in \Lambda(R)$ where $r \in R$, and describe its inverse of multiplication and derivative. We will be used it in §5.

Definition 3.5.1. We define a binomial symbols

$$\binom{r}{n} := \frac{r(r-1)\cdots(r-(n-1))}{n!} \quad (3.3)$$

actually lie in R for $r \in R$ and $n \in \mathbb{N}$. and we define

$$(1+t)^r := \sum_{i=1}^{\infty} \binom{r}{i} t^i. \quad (3.4)$$

Next, we describe inverse of multiplication of $(1+t)^r$ is $(1+t)^{-r}$.

Lemma 3.5.2. *The following statements hold for all $r \in R$, $n \in \mathbb{N}$.*

$$(-1)^{n+1} r \left(\sum_{i=0}^{n-1} (-1)^i \binom{r}{i} \right) = n \binom{r}{n}.$$

Proof. We prove by induction of $n \in \mathbb{N}$. If $n = 1$, it is clear. Suppose by induction that is a statement where $n - 1$. we have

$$(-1)^{n+1} r \left(\sum_{i=0}^{n-1} (-1)^i \binom{r}{i} \right) = -(n-1) \binom{r}{n-1} + r \binom{r}{n-1} = n \binom{r}{n}$$

for all $r \in R$, This finishes the induction and the proof. \square

Proposition 3.5.3. *The equalities $(1+t)^r(1+t)^s = (1+t)^{r+s}$ holds for any $r, s \in R$. In particular, $((1+t)^r)^{-1} = (1+t)^{-r}$ holds.*

Proof. We define $\psi^n : R \rightarrow R$ by $\psi^n := \text{id}_R$ for each $n \in \mathbb{N}$. then, R becomes to pre- ψ -ring together with the series $\{\psi^n\}_{n=1,2,3,\dots}$. Since Proposition 3.2.5, R is given a pre- λ -ringstructure uniquely such that ψ^n is n-th Adams operation. we show that it's λ -operation is satisfied

$$\lambda^n(r) = \binom{r}{n} \quad (3.5)$$

for all $r \in R$, $n = 0, 1, 2, \dots$ by induction of n . When $n = 0, 1$, it is clear since properties of λ -operation. Suppose by induction that $\lambda^j(r) = \binom{r}{j}$ ($j = 0, 1, 2, \dots, n-1$). By using Newtons formura(2) and Lemma 3.5.2, we compute

$$\begin{aligned} (-1)^{n+1} n \lambda^n(r) &= \psi^n(r) - \lambda^1(r) \psi^{n-1}(r) + \cdots + (-1)^{n-1} \lambda^{n-1}(r) \psi^1(r) \\ &= r \left(\sum_{i=0}^{n-1} (-1)^i \binom{r}{i} \right) = (-1)^{n+1} n \binom{r}{n} \end{aligned}$$

and divide by $(-1)^{n+1}n$, we obtain (3.5) for all $r \in R$. This finishes the induction. In particular, from property of λ_t , we have

$$(1+t)^{r+s} = \lambda_t(r+s) = \lambda_t(r)\lambda_t(s) = (1+t)^r(1+t)^s$$

for $r, s \in R$. □

Next, we describe the logarithmic derivative of $(1+t)^r$.

Lemma 3.5.4. *The following equality holds for all $a, r \in R, n \in \mathbb{N}$.*

$$-t \frac{d}{dt} \log(1 - a(-t)^n)^{\frac{r}{n}} = \sum_{i=1}^{\infty} r a^i (-t)^{in}$$

Proof. First, derivation of $(1+t)^r$ gives

$$\frac{d}{dt}(1+t)^r = r(1+t)^{r-1} \quad (3.6)$$

for $r \in R$. Since (3.6), Proposition 2.1.1 (3) Proposition 3.5.3, we have

$$\begin{aligned} -t \frac{d}{dt}(1 - a(-t)^n)^{\frac{r}{n}} &= (-t) \frac{r}{n} (1 - a(-t)^n)^{\frac{r}{n}-1} n a (-t)^{n-1} (1 - a(-t)^n)^{-\frac{r}{n}} \\ &= r a (-t)^n (1 - a(-t)^n)^{-1} = \sum_{i=1}^{\infty} r a_i (-t)^{in} \end{aligned}$$

for $r \in R, n \in \mathbb{N}$. □

Last, we describe the coefficients of t^n in $(1+t)^{-r}$ for any $r \in R$. In general, let f be a power series f over R , we have

$$f(t) = \sum_{i=0}^{\infty} \frac{f^{(i)}(0)}{i!} t^i$$

which $f^{(i)}(0)$ is a constant term of f that is differentiated i times. we put $f = (1+t)^{-r}$, then we have

$$\frac{f^{(i)}(0)}{i!} = \frac{(-r)(-r-1)\cdots(-r-(i-1))}{i!} = (-1)^i \binom{r+(i-1)}{i}$$

which imply that

$$(1+t)^{-r} = \sum_{n=0}^{\infty} \binom{r+(n-1)}{n} (-t)^n.$$

In particular, we put t to $-t$ in this statement, we get

$$(1-t)^{-r} = \sum_{n=0}^{\infty} \binom{r+(n-1)}{n} t^n \quad (3.7)$$

for $r \in R$.

4 Corresponding of character

In this chapter, we describe how to calculate a multiplicity of irreducible characters of the symmetric and exterior powers representation.

At first, we consider the one of Lemma. In general, let V be a vector space over \mathbb{C} which has a finite dimension $m = \dim V$, let $f : V \rightarrow V$ be a linear map, and we define $\text{Tr}(f)$ by the trace of the linear map f . About this values, $\text{Tr}(f)$ is equal to all eigenvalue of f , which is $\alpha_1, \dots, \alpha_n$.

Next, we define maps $f^n, S^n(f), \Lambda^n(f)$ by the following equalities hold for each $n \in \mathbb{N}$.

$$f^n : V \rightarrow V, f^n := \overbrace{f \circ \dots \circ f}^n \quad (4.1)$$

$$S^n(f) : S^n(V) \rightarrow S^n(V), \quad (4.2)$$

$$S^n(f)(v_1 \cdots v_n) := f(v_1) \cdots f(v_n),$$

$$\Lambda^n(f) : \Lambda^n(V) \rightarrow \Lambda^n(V), \quad (4.3)$$

$$\Lambda^n(f)(v_1 \wedge \cdots \wedge v_n) := f(v_1) \wedge \cdots \wedge f(v_n).$$

The vector spaces $V, S^n(V)$ and $\Lambda^n(V)$ have a finite dimension. About a trace of these linear map, we obtain the following equalities.

$$\text{Tr}(f^n) = \alpha_1^n + \cdots + \alpha_m^n = Q_n(\beta_1, \dots, \beta_m), \quad (4.4)$$

$$\text{Tr}(S^n(f)) = \sum_{1 \leq i_1 \leq \dots \leq i_n \leq m} \alpha_{i_1} \cdots \alpha_{i_n} = h_n(\beta_1, \dots, \beta_m), \quad (4.5)$$

$$\text{Tr}(\Lambda^n(f)) = \sum_{1 \leq i_1 < \dots < i_n \leq m} \alpha_{i_1} \cdots \alpha_{i_n} = \beta_n \quad (4.6)$$

where β_i be a elementary symmetric polynomial on $\alpha_1, \dots, \alpha_m$ for each $i \in \mathbb{N}$.

In this chapter afterward, we let G be an arbitrary finite multiplicative group with identity element, and we call the representation that has a finite dimension over \mathbb{C} . If we let $\rho : G \rightarrow GL(V)$ be the representation of G for each $i \geq 0$, we can give the i -th symmetric power representation $S^i(V)$ and the i -th exterior representation $\Lambda^i(V)$ that is defined by

$$S^i \rho : G \rightarrow GL(S^i(V)), (S^i \rho)(g) := S^i(\rho(g)), \quad (4.7)$$

$$\Lambda^i \rho : G \rightarrow GL(\Lambda^i(V)), (\Lambda^i \rho)(g) := \Lambda^i(\rho(g)). \quad (4.8)$$

from (4.2), (4.3), where $g \in G$. Remark that if the representation V is isomorphic to the representation W as the representation of G , then $S^i(V)$ is isomorphic to $S^i(W)$, and $\Lambda^i(V)$ is isomorphic to $\Lambda^i(W)$, as the representation of G , too.

As described under Example 3.1.4, the set of the all representation of G has a equivalence relation that is defined by the isomorphism as the representation of G . Let $M_{\mathbb{C}}(G)$ be the set of isomorphism classes $\{V\}$ of finite dimension representation of G over \mathbb{C} . The operation $\{V\} + \{W\} := \{V \oplus W\}$

and $\{V\}\{W\} := \{V \otimes_{\mathbb{C}} W\}$ make $M_{\mathbb{C}}(G)$ into a semiring. the functions $\lambda^n : M_{\mathbb{C}}(G) \rightarrow M_{\mathbb{C}}$, $\lambda^n(\{V\}) = \{\wedge^n(V)\}$ defines a λ -semiringstructure on $M_{\mathbb{C}}(G)$.

Now, let state once again the purpose of this paper here. Let V_1, V_2, \dots, V_k be all of representation spaces of the irreducible representation of G over \mathbb{C} . Since the representation $S^i(V)$ and $\wedge^i(V)$ are the representation of G and over \mathbb{C} , these representations are completely reducible from Maschkes Theorem, and isomorphic to direct sum of some V_1, V_2, \dots, V_k as the representation of G . In addition, representations $S^i(V)$ and $\wedge^i(V)$ have a finite dimension over \mathbb{C} too since V has a finite dimension over \mathbb{C} . Hence, these representation are isomorphic to direct sum of irreducible representation, and every irreducible component appears finitely, and uniquely. On the other words, we have

$$\begin{aligned} S^i(V) &\cong \overbrace{V_1 \oplus \dots \oplus V_1}^{m_{i,1}} \oplus \overbrace{V_2 \oplus \dots \oplus V_2}^{m_{i,2}} \oplus \dots \oplus \overbrace{V_k \oplus \dots \oplus V_k}^{m_{i,k}} \\ \wedge^i(V) &\cong \overbrace{V_1 \oplus \dots \oplus V_1}^{n_{i,1}} \oplus \overbrace{V_2 \oplus \dots \oplus V_2}^{n_{i,2}} \oplus \dots \oplus \overbrace{V_k \oplus \dots \oplus V_k}^{n_{i,k}} \end{aligned} \quad (4.9)$$

and the nonnegative numbers $m_{i,1}, \dots, m_{i,k}, n_{i,1}, \dots, n_{i,k}$ are determined finitely, and uniquely. The purpose of this paper is that calculating about it.

To achieve this purpose, we consider the character of the representation $\rho : G \rightarrow GL(V)$, which is a element of $\text{Map}(G, \mathbb{C})$, and is reffed that maps from $g \in G$ to the trace of $\rho(g)$. If the representation V is isomorphic to the representation W , the character of V is equal to the character of W . Hence, we can define a map $X : M_{\mathbb{C}}(G) \rightarrow \text{Map}(G, \mathbb{C})$ which maps from the representation V to the character of it. A map X is injective from $\text{char}(\mathbb{C})=0$ (from [6], Corollary 2.3.7).

In addition, as described under Example 3.2.6, $\text{Map}(G, \mathbb{C})$ is a \mathbb{C} -algebra with the operation $(f_1 + f_2)(g) = f_1(g) + f_2(g)$, $(f_1 f_2)(g) = f_1(g) f_2(g)$, $(r f_1)(g) = r f_1(g)$ ($g \in G$). In particular, it is a \mathbb{Q} -algebra. A map $X : M_{\mathbb{C}}(G) \rightarrow \text{Map}(G, \mathbb{C})$ is a ring monomorphism with these operator. Let X_1, \dots, X_k be characters of V_1, \dots, V_k , on the other hand, let χ_j be $X(\{V_j\})$ for each $j = 1, \dots, k$. In this case, the equalities (4.9) is equivalent to

$$\begin{aligned} X(\{S^i(V)\}) &= m_1 \chi_1 + \dots + m_k \chi_k, \\ X(\{\wedge^i(V)\}) &= n_1 \chi_1 + \dots + n_k \chi_k. \end{aligned} \quad (4.10)$$

At last, we consider about $X(\{S^i(V)\})$ and $X(\{\wedge^i(V)\})$, which are characters of the symmetric powers representation and the exterior powers representation. About these, we have the following Proposition.

The following equalities are hold for any $i = 0, 1, 2, \dots$, and for any representation of G .

$$\begin{aligned} (1) \quad X(\{\wedge^i(V)\}) &= \lambda^i(X(\{V\})), \\ (2) \quad X(\{S^i(V)\}) &= S^i(X(\{V\})). \end{aligned} \quad (4.11)$$

We describe the proof of this Proposition. About (1), See more in [2].

Proof. Let $(R_{\mathbb{C}}(G), \theta)$ be the ring completion of $M_{\mathbb{C}}(G)$. In this case, $R_{\mathbb{C}}(G)$ has a structure of a pre- λ -ring uniquely such that a map $\theta : M_{\mathbb{C}}(G) \rightarrow R_{\mathbb{C}}(G)$ is a λ -homomorphism from Proposition 3.1.2.

The claim of (1) is that a map $X : M_{\mathbb{C}}(G) \rightarrow \text{Map}(G, \mathbb{C})$ is a λ -homomorphism. At first, there is a ringhomomorphism $X' : R_{\mathbb{C}}(G) \rightarrow \text{Map}(G, \mathbb{C})$ uniquely such that the following equalities holds

$$X = X' \circ \theta \quad (4.12)$$

from Proposition 1.2.2. A λ -homomorphism θ can exchange the λ -operation, so If we have $X' \circ \lambda^i(\theta(\{V\})) = \lambda^i \circ X'(\theta(\{V\}))$ for any $\{V\} \in M_{\mathbb{C}}(G)$, then we can finish the proof of (1) from Proposition 3.4.2. Since $\text{Map}(G, \mathbb{C})$ is the \mathbb{Z} -torsion-free, it sufficed to show that

$$X' \circ \psi^i(\theta(\{V\})) = \psi^i \circ X'(\theta(\{V\})) \quad (4.13)$$

for any $i \in \mathbb{N}$, it is satisfied from Proposition 3.4.3.

Now, we prove (4.13). Let $\rho : G \rightarrow GL(V)$ be the representation of G with $m := \dim V$, let $\alpha_1, \dots, \alpha_m$ be eigenvalues of $\rho(g)$ for each $g \in G$, and let β_i be a i -th elementary symmetric polynomial on $\alpha_1, \dots, \alpha_m$. Since Proposition (3.2.2) (1), (4.12) and (4.6), we have

$$\begin{aligned} X' \circ \psi^i(\theta(\{V\}))(g) &= X' \left(Q_i(\lambda^1 \circ \theta(\{V\}), \dots, \lambda^i \circ \theta(\{V\})) \right)(g) \\ &= Q_i(X' \circ \theta \circ \lambda^1(\{V\})(g), \dots, X' \circ \theta \circ \lambda^i(\{V\})(g)) \quad (4.14) \\ &= Q_i(X \circ \lambda^1(\{V\})(g), \dots, X \circ \lambda^i(\{V\})(g)) = Q_i(\beta_1, \dots, \beta_i) \end{aligned}$$

and since (3.2), (4.12), (4.4), we have

$$\begin{aligned} \psi^i \circ X'(\theta(\{V\}))(g) &= X' \circ \theta(\{V\})(g^i) \\ &= X(\{V\})(g^i) \quad (4.15) \\ &= \alpha_1^i + \dots + \alpha_m^i = Q_i(\beta_1, \dots, \beta_m). \end{aligned}$$

To prove that (4.14) equals (4.15), we consider separately in the case of $i \leq m$ and $i > m$.

- If $i \leq m$, then we have $Q_i(\beta_1, \dots, \beta_i) = Q_i(\beta_1, \dots, \beta_m)$ from Lemma 2.3.6(2).
- Assume that $i > m$. If $k \in \mathbb{N}$ is satisfied $k \geq m$, then $\beta_k = 0$. hence, we have $Q_i(\beta_1, \dots, \beta_i) = Q_i(\beta_1, \dots, \beta_m, 0, \dots, 0) = Q_i(\beta_1, \dots, \beta_m)$ from Lemma 2.3.6(3).

Therefore we obtain (4.13), so we finish to prove (1).

Next, we consider about (2) with the same notation. Since the result of (1), Proposition 3.3.3 and (4.5), we have

$$\begin{aligned}
X(\{S^i(V)\})(g) &= h_i(\beta_1, \dots, \beta_i) \\
&= h_i(X(\{\Lambda^1(V)\})(g), \dots, X(\{\Lambda^i(V)\})(g)) \\
&= h_i((X \circ \lambda^1(\{V\}))(g), \dots, (X \circ \lambda^i(\{V\}))(g)) \\
&= h_i(\lambda^1 \circ X(\{V\}), \dots, \lambda^i \circ X(\{V\}))(g) = S^i(X(\{V\}))(g)
\end{aligned}$$

for any $g \in G$. so, we finish to prove (2) too. \square

Since (4.11), The equalities (4.10) is equivalence to

$$\begin{aligned}
S^i(X(\{V\})) &= X(\{S^i(V)\}) = m_1\chi_1 + \dots + m_k\chi_k, \\
\lambda^i(X(\{V\})) &= X(\{\Lambda^i(V)\}) = n_1\chi_1 + \dots + n_k\chi_k.
\end{aligned} \tag{4.16}$$

Hence, we obtain the following summary.

Let V be the representation of G . To calculate each multiplicity of irreducible representation of the symmetric powers representation $S^i(V)$ and the exterior powers representation $\Lambda^i(V)$, we calculate $S^i(X(\{V\}))$ and $\lambda^i(X(\{V\}))$, and express these character to sum of irreducible characters.

5 Calculating method

Now, we calculate the multiplicity of irreducible component of the symmetric and exterior powers representation of finite groups by some methods. Let G is a finite group, and let χ_1, \dots, χ_k be all of irreducible character of G . We denote χ by the character of the representation of G , then we calculate nonnegative integers $m_{i,j}$ and $n_{i,j}$ such that the following equalities hold.

$$\begin{aligned} S^i(\chi) &= m_{i,1}\chi_1 + \dots + m_{i,k}\chi_k, \\ \lambda^i(\chi) &= n_{i,1}\chi_1 + \dots + n_{i,k}\chi_k \end{aligned} \quad (5.1)$$

In addition, we define a character of the trivial representation which has 0-dimension by a zero element of a ring $\text{Map}(G, \mathbb{C})$.

By the property of λ -operation and symmetric powers operation, the map S^0 and λ^0 are constant maps to a unit element of $\text{Map}(G, \mathbb{C})$, and the map S^1 and λ^1 are identity map of $\text{Map}(G, \mathbb{C})$. Therefore, it is clear that the case of $i = 0, 1$. And, the case of $i > \chi(1)$, which is the dimension over of the representation space, $\lambda^i(\chi)$ is 0 from that dimension over \mathbb{C} of i -th exterior power representation is 0.

And, we define the inner product of $\text{Map}(G, \mathbb{C})$ by the following equality.

$$\langle f, f' \rangle_G := \frac{1}{|G|} \sum_{g \in G} f(g)f'(g^{-1}) \quad (f, f' \in \text{Map}(G, \mathbb{C})). \quad (5.2)$$

5.1 Formulations of operations of a pre- λ -ring

The statements about this section is described more detail in [2]. By using the inner product, we can write nonnegative integers $m_{i,j}, n_{i,j}$ in (5.1) as the following equalities.

$$m_{i,j} = \langle \chi_j, S^i(\chi) \rangle_G, \quad n_{i,j} = \langle \chi_j, \lambda^i(\chi) \rangle_G.$$

for any $i = 0, 1, 2, \dots$ and $j = 1, 2, \dots, k$. Hence the problem is the value of $S^i(\chi)(g)$ and $\lambda^i(\chi)(g)$ for $g \in G$, which is moved by an element of G the character $S^i(\chi)$ and $\lambda^i(\chi)$. The reason is that the value of $S^i(\chi)(g)$ and $\lambda^i(\chi)(g)$ is used in (5.2). Now, we describe the formula of operators of a pre- λ -ring, Newtons formula and Propositon 3.3.2 again. For all $n \in \mathbb{N}$, we have

$$\psi^n - \lambda^1 \psi^{n-1} + \dots + (-1)^{n-1} \lambda^{n-1} \psi^1 = (-1)^{n+1} n \lambda^n, \quad (5.3)$$

$$S^n + (-1) \lambda^1 S^{n-1} + \dots + (-1)^{n-1} \lambda^{n-1} S^1 + \lambda^n = 0. \quad (5.4)$$

We apply the character χ to these equation, and move an element $g \in G$. We describe Adams operation on $\text{Map}(G, \mathbb{C})$ as follow, in Example 3.2.6.

$$\psi^n(f)(g) = f(g^n) \quad (g \in G). \quad (5.5)$$

for $n \in \mathbb{N}$, $f \in \text{Map}(G, \mathbb{C})$. Therefore using (5.3), (5.4) and (5.5), we can calculate $S^i(\chi)(g)$ and $\lambda^i(\chi)(g)$.

In addition, remarks that the character $S^i(\chi)$ and $\lambda^i(\chi)$ is a class function, which is satisfied $S^i(\chi)(ghg^{-1}) = S^i(\chi)(h)$ and $\lambda^i(\chi)(ghg^{-1}) = \lambda^i(\chi)(h)$ for any elements $g, h \in G$.

Example 5.1.1 ([2], p.95). A symmetric group S_3 has only 2-dimension irreducible representation, so we consider about its 2nd exterior power representation.

At first, we describe conjugate classes and the character table of S_3 , which is the following statement and table.

$$C_1 = \{(1)\}, C_2 = \{(1\ 2), (1\ 3), (2\ 3)\}, C_3 = \{(1\ 2\ 3), (1\ 3\ 2)\}.$$

	C_1	C_2	C_3
χ_1	1	1	1
χ_2	1	-1	1
χ_3	2	0	-1

It is clear that $\lambda^0(\chi_3)$ equals χ_1 and $\lambda^1(\chi_3)$ equals χ_3 , we consider about $\lambda^2(\chi_3)$ by using Newton formula (5.3) when $n = 2$, we have

$$\lambda^2(\chi_3) = \frac{1}{2}(\chi_3^2 - \psi^2(\chi_3)).$$

Hence, (5.5) and the table that noted earlier give

$$\begin{aligned} \lambda^2(\chi_3)((1)) &= \frac{1}{2}(2^2 - 2) = 1, \\ \lambda^2(\chi_3)((1\ 2)) &= \frac{1}{2}(0^2 - 2) = -1, \\ \lambda^2(\chi_3)((1\ 2\ 3)) &= \frac{1}{2}((-1)^2 - (-1)) = 1. \end{aligned}$$

Then, we obtain $\lambda^2(\chi_3) = \chi_2$. This finish to calculate about $\lambda^2(\chi_3)$, however, is the inner product or the like, we may require further calculations actually.

5.2 The generating function of multiplicity

To calculate the character $S^i(\chi)$ and $\lambda^i(\chi)$ for any $i \geq 0$, we consider $S_t(\chi), \lambda_t(\chi)$. Expanding each of the power series, we obtain

$$\begin{aligned} S_t(\chi) &= \sum_{i=0}^{\infty} S^i(\chi)t^i = \sum_{i=0}^{\infty} \left(\sum_{j=1}^k m_{i,j} \chi_j \right) t^i = \sum_{j=1}^k \left(\sum_{i=0}^{\infty} \langle \chi_j, S^i(\chi) \rangle G^i t^i \right) \chi_j, \\ \lambda_t(\chi) &= \sum_{i=0}^{\infty} \lambda^i(\chi)t^i = \sum_{i=0}^{\infty} \left(\sum_{j=1}^k n_{i,j} \chi_j \right) t^i = \sum_{j=1}^k \left(\sum_{i=0}^{\infty} \langle \chi_j, \lambda^i(\chi) \rangle G^i t^i \right) \chi_j. \end{aligned} \tag{5.6}$$

In this section, we define the generating function of the multiplicity of irreducible component.

Definition 5.2.1. We define the formal power series $\langle \chi_j, S_t(\chi) \rangle_G$ and $\langle \chi_j, \lambda_t(\chi) \rangle_G$ with coefficient is nonnegative integer for $j = 1, \dots, k$ by the following equalities.

$$\begin{aligned}\langle \chi_j, S_t(\chi) \rangle_G &:= \sum_{i=0}^{\infty} \langle \chi_j, S^i(\chi) \rangle_G t^i = \frac{1}{|G|} \sum_{g \in G} \chi_j(g) S_t(\chi)(g^{-1}), \\ \langle \chi_j, \lambda_t(\chi) \rangle_G &:= \sum_{i=0}^{\infty} \langle \chi_j, \lambda^i(\chi) \rangle_G t^i = \frac{1}{|G|} \sum_{g \in G} \chi_j(g) \lambda_t(\chi)(g^{-1})\end{aligned}\tag{5.7}$$

By using (5.6), we obtain the following equalities.

$$S_t(\chi) = \sum_{j=1}^k \langle \chi_j, S_t(\chi) \rangle_G \chi_j, \quad \lambda_t(\chi) = \sum_{j=1}^k \langle \chi_j, \lambda_t(\chi) \rangle_G \chi_j\tag{5.8}$$

A formal power series $\langle \chi_j, S_t(\chi) \rangle_G$ and $\langle \chi_j, \lambda_t(\chi) \rangle_G$ imply that the generating function of the multiplicity about χ_j , hence it is believed that calculating such generating function is one means for computing the multiplicity. This method would be effective in calculating of the character of the symmetric powers representation especially of course in calculating of the character of the exterior powers representation.

Example 5.2.2. We calculate about power series $S_t(\chi_3)$, that χ_3 is 2-dimension irreducible representation of S_3 . From Example 5.1.1, we obtain

$$\lambda_t(\chi_3) = \chi_1 + \chi_3 t + \chi_2 t^2$$

which gives

$$\lambda_t(\chi_3)((1)) = (1+t)^2, \quad \lambda_t(\chi_3)((1\ 2)) = 1-t^2, \quad \lambda_t(\chi_3)((1\ 2\ 3)) = 1-t+t^2.$$

From Definition 3.3.1, which is Definition of Symmetric powers operation, we have

$$\begin{aligned}S_t(\chi_3)((1)) &= \frac{1}{(1-t)^2} = \sum_{i=0}^{\infty} \binom{i+1}{i} t^i, \\ S_t(\chi_3)((1\ 2)) &= \frac{1}{1-t^2} = \sum_{i=0}^{\infty} t^{2i}, \\ S_t(\chi_3)((1\ 2\ 3)) &= \frac{1}{1+t+t^2} = \frac{1-t}{1-t^3} = \sum_{i=0}^{\infty} t^{3i} - t^{3i+1}\end{aligned}\tag{5.9}$$

Putting these equalities into (5.7), we have

$$\begin{aligned}\langle \chi_1, S_t(\chi_3) \rangle_{S_3} &= \frac{1}{6} \left(\frac{1}{(1-t)^2} + \frac{3}{1-t^2} + \frac{2}{1+t+t^2} \right), \\ \langle \chi_2, S_t(\chi_3) \rangle_{S_3} &= \frac{1}{6} \left(\frac{1}{(1-t)^2} - \frac{3}{1-t^2} + \frac{2}{1+t+t^2} \right), \\ \langle \chi_3, S_t(\chi_3) \rangle_{S_3} &= \frac{1}{6} \left(\frac{2}{(1-t)^2} - \frac{2}{1+t+t^2} \right)\end{aligned}\tag{5.10}$$

More calculating, we have another expressing

$$\begin{aligned}\langle \chi_1, S_t(\chi_3) \rangle_{S_3} &= \frac{1}{(1-t^2)(1-t^3)}, \\ \langle \chi_2, S_t(\chi_3) \rangle_{S_3} &= \frac{t^3}{(1-t^2)(1-t^3)}, \\ \langle \chi_3, S_t(\chi_3) \rangle_{S_3} &= \frac{t+t^2}{(1-t^2)(1-t^3)} = \frac{t}{(1-t)(1-t^3)}.\end{aligned}\tag{5.11}$$

We have two form (5.10) and (5.11) of the generating function of multiplicity, (5.10) is oriented to see $S^n(\chi_3)$ for some $n \in \mathbb{N}$. For example, comparing the coefficients of t^6 in (5.9), (5.10), we have

$$S^6(\chi_3) = \frac{\binom{7}{6} + 3 + 2}{6} \chi_1 + \frac{\binom{7}{6} - 3 + 2}{6} \chi_2 + \frac{\binom{7}{6} - 1}{3} \chi_3 = 2\chi_1 + \chi_2 + 2\chi_3$$

On the other hand, (5.11) is more complexity than (5.10), however there is an advantage that if we have $\frac{1}{(1-t^2)(1-t^3)}$ in detail, we obtain the multiplicity for all $n \in \mathbb{N}$. It is believed that the change by the representation which one better in the calculation.

We describe coefficients of t^0, t^1, \dots, t^{10} in $\langle \chi_1, S_t(\chi_3) \rangle_{S_3}$, $\langle \chi_2, S_t(\chi_3) \rangle_{S_3}$ and $\langle \chi_3, S_t(\chi_3) \rangle_{S_3}$ below. We focus on that the coefficient of t^n in $\frac{1}{(1-t^2)(1-t^3)}$ is the number of the pair of nonnegative integer (x, y) such that $2x + 3y = n$.

	t^0	t^1	t^2	t^3	t^4	t^5	t^6	t^7	t^8	t^9	t^{10}	\dots
$\langle \chi_1, S_t(\chi_3) \rangle_{S_3}$	1	0	1	1	1	1	2	1	2	2	2	\dots
$\langle \chi_2, S_t(\chi_3) \rangle_{S_3}$	0	0	0	1	0	1	1	1	1	2	1	\dots
$\langle \chi_3, S_t(\chi_3) \rangle_{S_3}$	0	1	1	1	2	2	2	3	3	3	4	\dots

5.3 An one-dimension representation

Let χ_1 be a trivial character. In this section, we consider when χ has one dimension.

At first, if $i \geq 2$ then $\lambda^i(\chi) = 0$. hence we have

$$\lambda_t(\chi) = \chi_1 + \chi t.$$

Next, we obtain

$$S_t(\chi) = \frac{1}{\chi_1 - \chi t} = \sum_{i=0}^{\infty} \chi^i t^i$$

which implies that $S^i(\chi) = \chi^i$ for any $i \in \mathbb{N}$. A set of all character of one dimension representation, which is a subset of $\text{Map}(G, \mathbb{C})$, has a structure of a group with the multiplication, so let q is the order of χ about the multiplication, we have

$$\langle \chi_j, S_t(\chi) \rangle_G = \frac{1}{1-t^q} \sum_{i=0}^{q-1} \delta_{\chi_j, \chi^i} t^i.$$

Example 5.3.1. We consider about characters of one dimension representation of S_3 , which are χ_1 and χ_2 (See Example 5.1.1). The order of these character is 1 and 2 respectively, we obtain

$$\begin{aligned}\lambda_t(\chi_1) &= \chi_1 + \chi_1 t, & S_t(\chi_1) &= \frac{1}{1-t}\chi_1, \\ \lambda_t(\chi_2) &= \chi_1 + \chi_2 t, \\ S_t(\chi_2) &= \chi_1 + \chi_2 t + \chi_1 t^2 + \chi_2 t^3 + \cdots = \frac{1}{1-t^2}\chi_1 + \frac{t}{1-t^2}\chi_2.\end{aligned}$$

5.4 A normal subgroup and the permutation representation

In this section, we consider about the character which property is characterized by Adams operation, and contain the character of the permutation representation. And as a specific example, we describe the symmetric and exterior powers representation of a permutation representation which is generated by following action of G . G acts on the set of left cosets G/N of N in G by left multiplication $g(g'N) := (gg')N$, where N is a normal subgroup of G .

For any $g \in G$, we let $|g|$ be an order of an element g , and let (a, b) be a greatest common divisor of integers a and b . Since G is a finite group, we have $\psi^{m_1} = \psi^{m_2}$ for any $m_1, m_2 \geq 1$ with $m_1 \equiv m_2 \pmod{|G|}$. In addition, we define a function of G which has more strict condition that is characterized by $|G|$.

Definition 5.4.1. We define B by

$$B := \{f \in \text{Map}(G, \mathbb{C}) \mid \psi^n(f) = \psi^{(n, |G|)}(f) \text{ for all } n = 1, 2, \dots, |G|\}.$$

It is a subset of $\text{Map}(G, \mathbb{C})$,

A set B is closed by the addition from Proposition 3.2.3. Actually, a set B is a subring of $\text{Map}(G, \mathbb{C})$, and it is closed by λ -operation. However this proof use the properties of λ -ring, which is not described in this paper, and it is not use later. then, we do not describe this proof.

Example 5.4.2. Assume that a finite group G acts a finite set X . Then, let χ be the character of the representation, whose representation space is generated by elements of X as vector space. so, the character χ belongs to B .

Proof. For any $g \in G$, let X_g be a set of an element $x \in X$ such that $gx = x$. Therefore we have $\chi(g) = |X_g|$, hence it suffices to show that $X_{g^n} = X_{g^{(n, |G|)}}$ for any $n \geq 1$. Let x be an element of X_{g^n} . Then we have $g^n x = x$. Also, there are two integers a and b such that $an + b|G| = (n, |G|)$. We can choose an integer a to $a \in \mathbb{N}$, hence we have $g^{(n, |G|)} x = g^{an+b|G|} x = g^{an} x = x$. Therefore we obtain $x \in X_g^{(n, |G|)}$. It is clear that $X_g^{(n, |G|)} \subset X_{g^n}$. This finish to prove $X_{g^n} = X_{g^{(n, |G|)}}$ for any $g \in G$. \square

An element of B has a such property.

Theorem 5.4.3. *Let a_1, a_2, \dots, a_r be all of divisor of $|G|$ such that $1 = a_1 < a_2 < \dots < a_r = |G|$. For each $\chi \in B$, there are $b_1, \dots, b_r \in \text{Map}(G, \mathbb{C})$ uniquely, such that two following equalities hold.*

$$\lambda_t(\chi) = \prod_{i=1}^r (1 - (-t)^{a_i})^{\frac{b_i}{a_i}} \quad (5.12)$$

and

$$\psi^{a_l}(\chi) = \sum_{l'; a_l | a_{l'}} b_{l'} \quad (5.13)$$

for any $l = 1, \dots, r$.

Proof. Since G is a finite group, we have $\psi^{m_1}(\chi) = \psi^{m_2}(\chi)$ for any $m_1, m_2 \geq 1$ with $m_1 \equiv m_2 \pmod{|G|}$. Then, from the definition of Adams operation (Definition 3.2.1), we have

$$\begin{aligned} -t \frac{d}{dt} \log \lambda_t(\chi) &= \sum_{i=1}^r \psi^i(\chi) (-t)^i \\ &= \left(\frac{1}{1 - (-t)^{|G|}} \right) \left(\sum_{i=1}^{|G|} \psi^i(\chi) (-t)^i \right). \end{aligned} \quad (5.14)$$

And more, let T_l be the following equalities for any $l = 1, 2, \dots, r$.

$$T_l := \sum_{n=1, 2, \dots, |G|, (n, |G|)=a_l} (-t)^{a_n - 1}$$

Remark that T_l belongs to $\mathbb{Z}[t]$. Since (5.14) and $\chi \in B$, we obtain

$$\frac{d}{dt} \log \lambda_t(\chi) = \left(\frac{1}{1 - (-t)^{|G|}} \right) \left(\sum_{i=1}^r \psi^{a_i}(\chi) T_i \right). \quad (5.15)$$

Next, we let a polynomial F_l be the following equalities for any $l = 1, 2, \dots, r$.

$$F_l := \sum_{i=l}^r A_{l,i} T_i$$

whose coefficient of F_l is satisfied that following relations.

$$\begin{aligned} F_1 &:= \sum_{i=1}^r \psi^{a_i}(\chi) T_i \quad (A_{1,i} = \psi^{a_i}(\chi)), \\ F_l &:= (1 - (-t)^{|G|}) \frac{d}{dt} \log(1 - (-t)^{a_l})^{\frac{A_{l,l}}{a_l}} + F_{l+1}. \end{aligned} \quad (5.16)$$

About this definition we calculate both side of second equality, then we have

$$\begin{aligned}
F_l &= A_{l,l}T_l + \sum_{i=l+1}^r A_{l,i}T_i, \\
&(1 - (-t)^{|G|}) \frac{d}{dt} \log(1 - (-t)^{a_l})^{\frac{A_{l,l}}{a_l}} + F_{l+1} \\
&= A_{l,l} \frac{1 - (-t)^{|G|}}{1 - (-t)^{a_l}} + F_{l+1} \\
&= A_{l,l}((-t)^{a_l-1} + (-t)^{2a_l-1} + \dots + (-t)^{\frac{|G|}{a_l}a_l-1}) + \sum_{i=l+1}^r A_{l+1,i}T_i
\end{aligned}$$

which implies that

$$A_{l+1,i} = \begin{cases} A_{l,i} & (a_l \nmid a_i), \\ A_{l,i} - A_{l,l} & (a_l \mid a_i). \end{cases}$$

For each $A_{l,l}$ ($l = 1, 2, \dots, r$), let i_1, \dots, i_p be numbers such that $a_i \mid a_l$ and $1 = i_1 < \dots < i_p < a_l$. Then we have

$$\begin{aligned}
A_{l,l} = A_{l-1,l} = \dots = A_{i_p+1,l} &= A_{i_p,l} - A_{i_p,i_p} \\
&= \dots = A_{i_1,l} - A_{i_1,i_1} - \dots - A_{i_p,i_p} \\
&= \psi^{a_l}(\chi) - A_{i_1,i_1} - \dots - A_{i_p,i_p}
\end{aligned}$$

which imply that

$$\psi^{a_l}(\chi) = \sum_{l': a_l' \mid a_l} A_{l',l'}$$

In addition, calculating F_r we obtain

$$F_r = A_{r,r}T_r = A_{r,r}(-t)^{|G|-1} = (1 - (-t)^{|G|}) \frac{d}{dt} \log(1 - (-t)^{a_r})^{\frac{A_{r,r}}{a_r}}. \quad (5.17)$$

Therefore, by using (5.15), (5.16) and (5.17) we have

$$\begin{aligned}
\frac{d}{dt} \log \lambda_t(\chi) &= \frac{1}{1 - (-t)^{|G|}} F_1 \\
&= \frac{1}{1 - (-t)^{|G|}} \left((1 - (-t)^{|G|}) \frac{d}{dt} \log(1 - (-t)^{a_1})^{\frac{A_{1,1}}{a_1}} + F_2 \right) \\
&= \dots \\
&= \frac{1}{1 - (-t)^{|G|}} \sum_{i=1}^r (1 - (-t)^{|G|}) \frac{d}{dt} \log(1 - (-t)^{a_i})^{\frac{A_{i,i}}{a_i}} \\
&= \frac{d}{dt} \log \prod_{i=1}^r (1 - (-t)^{a_i})^{\frac{A_{i,i}}{a_i}}
\end{aligned}$$

which imply that

$$\lambda_t(\chi) = \prod_{i=1}^r (1 - (-t)^{a_i})^{\frac{A_{i,i}}{a_i}}$$

from the map $\frac{d}{dt} \log$ is injective. The uniqueness is proved by (5.13) and the induction of $l = 1, \dots, r$. □

We describe one of example which belongs to B . Let N be a normal subgroup of G . G acts on the quotient group G/N that the following statement holds

$$g \in G, \alpha = \pi(g') \implies g\alpha := \pi(gg')$$

where a map $\pi : G \rightarrow G/N$ is a natural projection. Let Π is a character of a representation which generalized by above action. Then, $m\Pi$ belongs to B for any $m \geq 1$ from Example 5.4.2. Now, we calculate $\langle \chi_j, S_t(m\Pi) \rangle_G$ and $\langle \chi_j, \lambda_t(m\Pi) \rangle_G$ for each $j = 1, \dots, k$.

A character Π is a induced representation from the trivial character from N to G . Hence the specific values is as follows.

$$\Pi(g) = \begin{cases} |G/N| & (g \in N) \\ 0 & (g \notin N) \end{cases} \quad (g \in G).$$

Now, we define the number $O_N(g)$ by the order of $\pi(g) \in G/N$.

Lemma 5.4.4. *We let a_1, \dots, a_r be all of divisor of $|G|$ such that $1 = a_1 < a_2 < \dots < a_r = |G|$. About elements $b_1, \dots, b_r \in \text{Map}(G, \mathbb{C})$ in Theorem 5.4.3, the following equality holds for any $i = 1, 2, \dots, r$ and $g \in G$.*

$$b_i(g) = \begin{cases} m |G/N| & (O_N(g) = a_i) \\ 0 & (O_N(g) \neq a_i) \end{cases} \quad (g \in G).$$

Proof. We prove it by induction of $i = 1, 2, \dots, r$. If $i = 1$, we have $b_1 = \psi^1(m\Pi) = m\Pi$ from (5.12), and N is equal to a set of all element $g \in G$ such that $O_N(g) = 1$. Hence we have this statement when $i = 1$ from the value of $m\Pi$.

Suppose by induction for $j < i$. By calculating b_i from (5.13), we have

$$b_i = \psi^{a_i}(m\Pi) - \sum_{l: a_l | a_i, l \neq i} b_l$$

Also, we obtain

$$\psi^{a_i}(m\Pi)(g) = \begin{cases} m |G/N| & (O_N(g) | a_i) \\ 0 & (O_N(g) \nmid a_i) \end{cases}$$

for all $g \in G$. Hence, we have

$$b_i(g) = \begin{cases} m |G/N| & (O_N(g) = a_i) \\ 0 & (O_N(g) \neq a_i) \end{cases}$$

by the induction hypothesis. This finishes the induction and the proof of the Corollary. □

From the above Proposition and Corollary, we have the following Theorem.

Theorem 5.4.5. *The following three statement holds about the character $m\Pi$.*

(1) *One has*

$$\begin{aligned}\lambda_t(m\Pi)(g) &= (1 - (-t)^{O_N(g)})^{\frac{m|G/N|}{O_N(g)}}, \\ S_t(m\Pi)(g) &= (1 - t)^{O_N(g)} - \frac{m|G/N|}{O_N(g)}.\end{aligned}\tag{5.18}$$

for any $g \in G$.

(2) *One has*

$$\begin{aligned}\langle \chi_j, S_t(m\Pi) \rangle_G &= \frac{1}{|G|} \sum_{g \in G} \chi_j(g) (1 - t^{O_N(g)})^{-\frac{m|G/N|}{O_N(g)}}, \\ \langle \chi_j, \lambda_t(m\Pi) \rangle_G &= \frac{1}{|G|} \sum_{g \in G} \chi_j(g) (1 - (-t)^{O_N(g)})^{\frac{m|G/N|}{O_N(g)}}\end{aligned}\tag{5.19}$$

for any $j = 1, \dots, k$.

(3) *In particular, if $n \geq 1$ is satisfied $(n, |G/N|) = 1$, we have*

$$S^n(m\Pi) = \frac{1}{|G/N|} \binom{m|G/N| + n - 1}{n} \Pi, \quad \lambda^n(m\Pi) = \frac{1}{|G/N|} \binom{m|G/N|}{n} \Pi$$

Proof. (1) It is clear by Lemma 5.4.4.

(2) We obtain (5.19) by apply (5.18) to (5.7).

(3) About (5.19), we collect the number $O_N(g)$, we have

$$\begin{aligned}\langle \chi_j, S_t(m\Pi) \rangle_G &= \frac{1}{|G|} \sum_{i=1}^r \left(\sum_{O_N(g)=a_i} \chi_j(g) \right) (1 - t^{a_i})^{-\frac{m|G/N|}{a_i}}, \\ \langle \chi_j, \lambda_t(m\Pi) \rangle_G &= \frac{1}{|G|} \sum_{i=1}^r \left(\sum_{O_N(g)=a_i} \chi_j(g) \right) (1 - (-t)^{a_i})^{\frac{m|G/N|}{a_i}}\end{aligned}\tag{5.20}$$

Also, we obtain

$$\frac{1}{|G|} \sum_{O_N(g)=1} \chi_j(g) = \frac{1}{|G/N|} \langle \chi_j, \Pi \rangle_G.$$

Assume that $n \geq 1$ is satisfied $(n, |G/N|) = 1$. The multiplication of χ_j in $S^n(m\Pi)$ and $\lambda^n(m\Pi)$ are the coefficient of t^n in each $\langle \chi_j, S_t(m\Pi) \rangle_G$ and $\langle \chi_j, \lambda_t(m\Pi) \rangle_G$. By $a_1 = 1$ and $O_N(g)$ is a divisor of $|G/N|$ for any $g \in G$, we

have

$$\begin{aligned}
\langle \chi_j, S^n(m\Pi) \rangle_G &= \frac{1}{|G|} \binom{m|G/N| + n - 1}{n} \left(\sum_{O_N(g)=a_i} \chi_j(g) \right) \\
&= \frac{1}{|G/N|} \binom{m|G/N| + n - 1}{n} \langle \chi_j, \Pi \rangle_G, \\
\langle \chi_j, \lambda^n(m\Pi) \rangle_G &= \frac{1}{|G|} \binom{m|G/N|}{n} \left(\sum_{O_N(g)=a_i} \chi_j(g) \right) \\
&= \frac{1}{|G/N|} \binom{m|G/N|}{n} \langle \chi_j, \Pi \rangle_G
\end{aligned}$$

from (5.20). Hence we have $S^n(m\Pi) = \frac{1}{|G/N|} \binom{m|G/N| + n - 1}{n} \Pi$ and $\lambda^n(m\Pi) = \frac{1}{|G/N|} \binom{m|G/N|}{n} \Pi$. \square

Example 5.4.6. We calculate a character of regular representation of S_3 , Π , which is obtained by putting $G = S_3$, $N = \{e\}$ and $m = 1$ in Theorem 5.4.5 About a character table of S^3 , see Example 5.1.1. About the generating function of a multiplicity of the exterior powers representation, we have

$$\begin{aligned}
\langle \chi_1, \lambda_t(\Pi) \rangle_{S_3} &= \frac{1}{6} \left((1+t)^6 + 3(1-t^2)^3 + 2(1+t^3)^2 \right), \\
\langle \chi_2, \lambda_t(\Pi) \rangle_{S_3} &= \frac{1}{6} \left((1+t)^6 - 3(1-t^2)^3 + 2(1+t^3)^2 \right), \\
\langle \chi_3, \lambda_t(\Pi) \rangle_{S_3} &= \frac{1}{6} \left(2(1+t)^6 - 2(1+t^3)^2 \right).
\end{aligned}$$

And about the generating function of a multiplicity of the symmetric powers representation, we have

$$\begin{aligned}
\langle \chi_1, S_t(\Pi) \rangle_{S_3} &= \frac{1}{6} \left(\frac{1}{(1-t)^6} + \frac{3}{(1-t^2)^3} + \frac{2}{(1-t^3)^2} \right), \\
\langle \chi_2, S_t(\Pi) \rangle_{S_3} &= \frac{1}{6} \left(\frac{1}{(1-t)^6} - \frac{3}{(1-t^2)^3} + \frac{2}{(1-t^3)^2} \right), \\
\langle \chi_3, S_t(\Pi) \rangle_{S_3} &= \frac{1}{6} \left(\frac{2}{(1-t)^6} - \frac{2}{(1-t^3)^2} \right),
\end{aligned}$$

In addition we describe coefficient of t^0, t^1, \dots, t^{10} in these generating function.

	t^0	t^1	t^2	t^3	t^4	t^5	t^6	t^7	t^8	t^9	t^{10}	...
$\langle \chi_1, \lambda_t(\Pi) \rangle_{S_3}$	1	1	1	4	4	1	0	0	0	0	0	...
$\langle \chi_2, \lambda_t(\Pi) \rangle_{S_3}$	0	1	4	4	1	1	1	0	0	0	0	...
$\langle \chi_3, \lambda_t(\Pi) \rangle_{S_3}$	0	2	5	6	5	2	0	0	0	0	0	...
$\langle \chi_1, S_t(\Pi) \rangle_{S_3}$	1	1	5	10	24	43	83	132	222	335	511	...
$\langle \chi_2, S_t(\Pi) \rangle_{S_3}$	0	1	2	10	18	43	73	132	207	335	490	...
$\langle \chi_3, S_t(\Pi) \rangle_{S_3}$	0	2	7	18	42	86	153	264	429	666	1001	...

5.5 A normal subgroup and its one-dimension representation

Let N be a normal subgroup of G that contains a central of G which is said to $Z(G)$. And let ζ be a character of a representation of N which has 1 dimension. Now, we define $\zeta_0 \in \text{Map}(G, \mathbb{C})$ as follows.

$$\zeta_0(g) = \begin{cases} \zeta(g) & (g \in N), \\ 0 & (g \notin N). \end{cases}$$

We let $m \in \mathbb{N}$ be that $m\zeta_0$ is a character of G , and in this section, we consider about symmetric and exterior powers representation which has character $m\zeta_0$. The notation $O_N(g)$ and (a, b) , we used as those in the previous section.

Remarks that there is a such $m \in \mathbb{N}$. For example, if $m = |G/N|$ then $m\zeta_0$ is a induced character of ζ to G . In addition if $|G/N|$ equals to square number, then putting $m = \sqrt{|G/N|}$ gives $\langle m\zeta_0, m\zeta_0 \rangle_G = 1$. However, $m\zeta_0$ is not necessarily the representation.

Theorem 5.5.1. *The following three statement holds about a character $m\zeta_0$.*

(1) *For any $g \in G$, we have*

$$\lambda_t(m\zeta_0)(g) = (1 - \zeta(g^h)(-t)^h)^{\frac{m}{h}} \quad (5.21)$$

where $h = O_N(g)$.

(2) *One has*

$$\begin{aligned} \langle \chi_j, S_t(m\zeta_0) \rangle_G &= \frac{1}{|G|} \sum_{g \in G} \chi_j(g) \left(1 - \zeta(g^{-O_N(g)})t^{O_N(g)}\right)^{-\frac{m}{O_N(g)}} \\ \langle \chi_j, \lambda_t(m\zeta_0) \rangle_G &= \frac{1}{|G|} \sum_{g \in G} \chi_j(g) \left(1 - \zeta(g^{-O_N(g)})(-t)^{O_N(g)}\right)^{\frac{m}{O_N(g)}} \end{aligned} \quad (5.22)$$

for any $j = 1, \dots, k$.

(3) *In particular, if $n \geq 1$ is satisfied $(n, |G/N|) = 1$, then we have*

$$S^n(m\zeta_0) = \binom{m+n-1}{n} \zeta_0^n, \quad \lambda^n(m\zeta_0) = \binom{m}{n} \zeta_0^n.$$

Proof. (1) By calculating Adams operation $\psi^n(m\zeta_0)(g)$ for any $n \geq 1$, we have

$$\psi^n(m\zeta_0)(g) = m\zeta_0(g^n) = \begin{cases} m\zeta(g^h)^i & (h \mid n, n = hi), \\ 0 & (h \nmid n) \end{cases}$$

since ζ has 1 dimension. Hence, we obtain

$$\sum_{i=1}^{\infty} m\zeta(g^h)^i (-t)^{hi} = -t \frac{d}{dt} \log \lambda_t(m\zeta_0)(g) \quad (5.23)$$

from the definition of Adams operation. Also, we have

$$-t \frac{d}{dt} \log(1 - \zeta(g^h)(-t)^h)^{\frac{m}{h}} = \sum_{i=1}^{\infty} m \zeta(g^h)^i (-t)^{hi} \quad (5.24)$$

from Lemma 3.5.4. By (5.23), (5.24) and that a map $\frac{d}{dt} \log$ is injective, this finish to prove (1).

(2) We obtain (5.22) by apply (5.21) to (5.7) for any $j = 1, \dots, k$.

(3) Let a_1, \dots, a_r be all of a divisor of $|G|$ such that $1 = a_1 < a_2 < \dots < a_r = |G|$. About (5.22) we collect the number $O_N(g)$, we have

$$\begin{aligned} \langle \chi_j, S_t(m\zeta_0) \rangle_G &= \frac{1}{|G|} \sum_{i=1}^r \left(\sum_{O_N(g)=a_i} \chi_j(g) (1 - \zeta(g^{-a_i}) t^{a_i})^{-\frac{m}{a_i}} \right), \\ \langle \chi_j, \lambda_t(m\zeta_0) \rangle_G &= \frac{1}{|G|} \sum_{i=1}^r \left(\sum_{O_N(g)=a_i} \chi_j(g) (1 - \zeta(g^{-a_i}) (-t)^{a_i})^{\frac{m}{a_i}} \right) \end{aligned} \quad (5.25)$$

Assume that $n \leq 1$ is satisfied $(n, |G/N|) = 1$. By $a_1 = 1$ and $O_N(g)$ is a divisor of $|G/N|$ for any $g \in G$, we have

$$\begin{aligned} \langle \chi_j, S^n(m\zeta_0) \rangle_G &= \frac{1}{|G|} \sum_{g \in N} \chi_j(g) \binom{m+n-1}{n} \zeta^n(g^{-1}) \\ &= \binom{m+n-1}{n} \langle \chi_j, \zeta_0^n \rangle_G, \\ \langle \chi_j, \lambda^n(m\zeta_0) \rangle_G &= \frac{1}{|G|} \sum_{g \in N} \chi_j(g) \binom{m}{n} \zeta^n(g^{-1}) \\ &= \binom{m}{n} \langle \chi_j, \zeta_0^n \rangle_G. \end{aligned}$$

from (5.25). Hence we have $S^n(m\zeta_0) = \binom{m+n-1}{n} \zeta_0^n$, $\lambda^n(m\zeta_0) = \binom{m}{n} \zeta_0^n$. \square

Example 5.5.2. Let ζ is a trivial character of N , and let m be the multiple of $|G/N|$, which is written by $m'|G/N|$. then $m\zeta_0$ is a m times the character of the representation which is generated by action G to G/N . If $n \leq 1$ is satisfied $(n, |G/N|) = 1$, we have

$$\begin{aligned} S^n(m'\Pi) &= S^n(m\zeta_0) = \binom{m+n-1}{n} \zeta_0^n = \frac{1}{|G/N|} \binom{m'|G/N|+n-1}{n} \Pi, \\ \lambda^n(m'\Pi) &= \lambda^n(m\zeta_0) = \binom{m}{n} \zeta_0^n = \frac{1}{|G/N|} \binom{m'|G/N|}{n} \Pi \end{aligned}$$

Calculating of this charcter Π had also done in Theorem 5.4.5, However If $N \subset Z(G)$, then we can calculate by this method.

Example 5.5.3. We let H_p be the following set.

$$H_p := \left\{ \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} \in GL(3, \mathbb{Z}/p\mathbb{Z}) \mid a, b, c \in \mathbb{Z}/p\mathbb{Z} \right\}$$

where p is a prime number such that $p \neq 2$. It is said to Heisenberg group modulo p . It is a non commutative group, has cardinality p^3 , and all of the element of H_p except unit element has the order p .

Conjugate classes and character table of H_p are the following statements and table where η is a primitive p -th root of unity.

$$C(h) := \left\{ \begin{pmatrix} 1 & 0 & h \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\} \quad (h \in \mathbb{Z}/p\mathbb{Z}),$$

$$C(e, f) := \left\{ \begin{pmatrix} 1 & e & k \\ 0 & 1 & f \\ 0 & 0 & 1 \end{pmatrix} \mid k \in \mathbb{Z}/p\mathbb{Z} \right\} \quad (e, f \in \mathbb{Z}/p\mathbb{Z}, (e, f) \neq (0, 0)).$$

	$C(h)$	$C(e, f)$
$\chi_{i,j} \ (i, j = 0, 1, \dots, p-1)$	1	η^{ei+fj}
$\tau_s \ (s = 1, 2, \dots, p-1)$	$p\eta^{sh}$	0

There are p character which have one-dimension. The reason is that the quotient group $H_p/D(H_p)$ is isomorphic to the direct sum of two cycle group which has order p where $D(H_p)$ is a commutator subgroup of H_p , which is the subgroup generated by the elements $xyx^{-1}y^{-1}$ for $x, y \in H_p$.

We obtain the irreducible character which has dimension p as follows. Let K be $K = \bigcup_{h=0}^{p-1} C(h) \cup \bigcup_{e=1}^{p-1} C(e, e)$, which is a subgroup of H_p and is a direct sum of two cycle group which have order p generated by two matrix

$$\alpha = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad \beta = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

The character $\tau_s \ (s = 1, 2, \dots, p-1)$ is given by the induced character of following character of K to H_p .

$$\gamma_s : K \rightarrow \mathbb{C}, \quad \gamma_s(\alpha^i \beta^j) = \eta^{js} \quad (i, j = 0, 1, \dots, p-1).$$

We consider the irreducible character τ_s for all $s = 1, 2, \dots, p-1$. At first, we define a map $\zeta_s : Z(H_p) \rightarrow \mathbb{C}$ as following while noted that $Z(H_p) = \bigcup_{h=0}^{p-1} C(h)$

$$\zeta_s \left(\begin{pmatrix} 1 & 0 & h \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) := \eta^{hs} \quad (h \in \mathbb{Z}/p\mathbb{Z}).$$

It is a character of $Z(H_p)$ which has one-dimension. In addition, we have $\tau_s = p(\zeta_s)_0$, hence let $n \geq 1$ be satisfied $(n, p) = 1$, since Theorem 5.5.1 we obtain

$$\lambda^n(\tau_s) = \binom{p}{n} (\zeta_s)_0^n = \frac{1}{p} \binom{p}{n} \tau_{n'}, \quad S^n(\tau_s) = \binom{p+n-1}{n} (\zeta_s)_0^n = \frac{1}{p} \binom{p+n-1}{n} \tau_{n'}$$

where n' is satisfied that $n' = 1, 2, \dots, p-1$ and $ns \equiv n' \pmod{p}$.

Next we calculate $S^{ip}(\tau_s)$ and $\lambda^{ip}(\tau_s)$ for $i \geq 1$. Since (5.21), we have the following table.

	$C(h)$	$C(e, f)$
$\lambda_t(\tau_s)$	$(1 + \eta^{sh}t)^p$	$1 + t^p$
$S_t(\tau_s)$	$(1 - \eta^{sh}t)^{-p}$	$(1 - t^p)^{-1}$

About $\lambda^p(\tau_s)$, we obtain $\lambda^p(\tau_s) = \chi_{0,0}$ since the coefficient of t^p in a polynomial which apply a element of $C(h)$ to $\lambda_t(\tau_s)$ is equal to $(\eta^{0h})(\eta^{1h}) \cdots (\eta^{ph}) = 1$. Next, we calculate about $S^{ip}(\tau_s)$. If we apply each irreducible character of H_p to (5.25), we obtain as follows about $\chi_{i',j'}$ ($i', j' = 0, 1, \dots, p-1$).

$$\begin{aligned} & \langle \chi_{i',j'}, S_t(\tau_s) \rangle_{H_p} \\ &= \frac{1}{p^3} \left(\sum_{h=0}^{p-1} (1 - \eta^{sh}t)^{-p} + p \sum_{e,f, (e,f) \neq (0,0)} \eta^{i'e+j'f} (1 - t^p)^{-1} \right) \\ &= \frac{1}{p^3} \left(\sum_{h=0}^{p-1} (1 - \eta^{sh}t)^{-p} - p(1 - t^p)^{-1} + p \left(\sum_{e=0}^p \eta^{i'e} \right) \left(\sum_{f=0}^p \eta^{j'f} \right) (1 - t^p)^{-1} \right) \\ &= \begin{cases} \frac{1}{p^3} \left(\sum_{h=0}^{p-1} (1 - \eta^{sh}t)^{-p} + p(p^2 - 1)(1 - t^p)^{-1} \right) & ((i', j') = (0, 0)) \\ \frac{1}{p^3} \left(\sum_{h=0}^{p-1} (1 - \eta^{sh}t)^{-p} - p(1 - t^p)^{-1} \right) & ((i', j') \neq (0, 0)) \end{cases} \end{aligned}$$

then, the coefficient of t^{ip} in $(1 - \eta^{sh}t)^{-p}$ is $\binom{p+ip-1}{ip} (\eta^{sh})^{ip} = \binom{p+ip-1}{p}$ since (3.7). hence we obtain

$$\begin{aligned} S^{ip}(\tau_s) &= \frac{1}{p^2} \left(\binom{p+ip-1}{ip} + (p^2 - 1) \right) \chi_{0,0} \\ &+ \sum_{i,j, (i,j) \neq (0,0)} \frac{1}{p^2} \left(\binom{p+ip-1}{ip} - 1 \right) \chi_{i,j} + \sum_{s'=1}^{p-1} a_{s'} \tau_{s'} \end{aligned} \quad (5.26)$$

About the multiplicity of τ'_s ($s' = 1, 2, \dots, p-1$) which is $a_{s'}$, we obtain 0 since calculating of dimension in the both side of (5.26).

To summarize the above, the following statement holds.

$$\lambda^n(\tau_s) = \begin{cases} \chi_{0,0} & (n = 0, p), \\ \frac{1}{p} \binom{p}{n} \tau_{n'} & (n = 1, \dots, p-1), \\ 0 & (n > p). \end{cases}$$

$$S^n(\tau_s) = \begin{cases} \frac{1}{p^2} \left(\binom{p+n-1}{n} + (p^2-1) \right) \chi_{0,0} \\ + \sum_{\substack{i,j \\ (i,j) \neq (0,0)}} \frac{1}{p^2} \left(\binom{p+n-1}{n} - 1 \right) \chi_{i,j} & (p \mid n), \\ \frac{1}{p} \binom{p+n-1}{n} \tau_{n'} & (p \nmid n). \end{cases}$$

for any $n \geq 1$ where $n' \geq 1$ is satisfied that $n' = 1, 2, \dots, p-1$ and $ns \equiv n' \pmod{p}$.

5.6 Irreducible and reducible characters

For any given character χ , we can write as follows.

$$\begin{aligned} S_t(\chi) &= S_t(\langle \chi_1, \chi \rangle_G \chi_1 + \langle \chi_2, \chi \rangle_G \chi_2 + \dots + \langle \chi_k, \chi \rangle_G \chi_k) \\ &= S_t(\chi_1)^{\langle \chi_1, \chi \rangle_G} \dots S_t(\chi_k)^{\langle \chi_k, \chi \rangle_G}, \\ \lambda_t(\chi) &= \lambda_t(\langle \chi_1, \chi \rangle_G \chi_1 + \langle \chi_2, \chi \rangle_G \chi_2 + \dots + \langle \chi_k, \chi \rangle_G \chi_k) \\ &= \lambda_t(\chi_1)^{\langle \chi_1, \chi \rangle_G} \dots \lambda_t(\chi_k)^{\langle \chi_k, \chi \rangle_G}. \end{aligned}$$

Therefore if we have the character of symmetric and exterior powers representation of all of the irreducible character, we obtain about the character of reducible representation too. Hence it will be important to calculate about the character of the irreducible representation. However, we have to know the result of the multiplication of the irreducible character with each other to calculating, and it is unlikely that it will become easier in most cases. Thus, the method of described in previous section may be simpler even though the character of reducible representation.

Example 5.6.1. We calculate about $\chi_1 + \chi$ for any character χ where χ_1 is a trivial character. We can apply about the natural representation of S_n or A_n which generated the action of S_n or A_n to a finite set $\{1, 2, \dots, n\}$. Thus, we have

$$S_t(\chi_1 + \chi) = \frac{1}{1-t} \chi_1 S_t(\chi) = \frac{1}{1-t} S_t(\chi)$$

which implies that $\langle \chi_j, S_t(\chi_1 + \chi) \rangle_G = \frac{1}{1-t} \langle \chi_j, S_t(\chi) \rangle_G$. And we have

$$\lambda_t(\chi_1 + \chi) = (\chi_1 + \chi_1 t) \left(\sum_{i=0} \lambda^i(\chi) t^i \right) = \chi_1 + \sum_{i=1} (\lambda^{i-1}(\chi) + \lambda^i(\chi)) t^i.$$

Example 5.6.2. Let χ be $\chi_1 + \chi_2 + \chi_3$ which is sum of all irreducible character of S_3 . One has

$$\begin{aligned}
\lambda_t(\chi) &= \lambda_t(\chi_1)\lambda_t(\chi_2)\lambda_t(\chi_3) \\
&= (\chi_1 + \chi_1 t)(\chi_1 + \chi_2 t)(\chi_1 + \chi_3 t + \chi_2 t^2) \\
&= \chi_1 + (\chi_1 + \chi_2 + \chi_3)t + (\chi_2 + \chi_2\chi_3 + \chi_1\chi_3 + \chi_1\chi_2)t^2 \\
&\quad + (\chi_1\chi_2\chi_3 + \chi_1\chi_2 + \chi_2\chi_2)t^3 + \chi_1\chi_2\chi_2 t^4 \\
&= \chi_1 + (\chi_1 + \chi_2 + \chi_3)t + (2\chi_2 + 2\chi_3)t^2 + (\chi_1 + \chi_2 + \chi_3)t^3 + \chi_1 t^4
\end{aligned}$$

5.7 Calculating that use the quotient group

Let N is a normal subgroup of G . There is a case that the multiplicity in symmetric and exterior powers representation can compute by the quotient group G/N . Also in this section, let a map $\pi_N : G \rightarrow G/N$ be a natural projection. In addition, let $\text{Cha}(G)$ and $\text{Cha}(G/N)$ be sets of all of character of G and G/N respectively. These sets are closed by the addition, the multiplication, λ -operation and symmetric powers operation. from §4.

Proposition 5.7.1. *We define a map $\pi_N^* : \text{Map}(G/N, \mathbb{C}) \rightarrow \text{Map}(G, \mathbb{C})$ as follows*

$$\pi_N^*(f) := f \circ \pi \quad (f \in \text{Map}(G/N, \mathbb{C})).$$

Then, the following statements hold.

- (1) *A map π_N^* is a λ -monomorphism.*
- (2) *We have $\langle f_1, f_2 \rangle_{G/N} = \langle \pi_N^*(f_1), \pi_N^*(f_2) \rangle_G$ for all $f_1, f_2 \in \text{Map}(G/N, \mathbb{C})$.*

Proof. (1) It is clear that a map π_N^* is a ring monomorphism, then we show that $\lambda^n \circ \pi_N^* = \pi_N^* \circ \lambda^n$ for any $n \geq 1$ $\text{Map}(G, \mathbb{C})$ which is a codomain of π_N^* , is the \mathbb{Z} -torsion-free, then it is sufficient to show that $\psi^n \circ \pi_N^* = \pi_N^* \circ \psi^n$ for any $n \geq 1$ from Proposition 3.4.3. For any $f \in \text{Map}(G/N, \mathbb{C})$, $n \geq 1$ and $g \in G$. we have

$$\psi^n(\pi_N^*(f))(g) = \pi_N^*(f)(g^n) = f \circ \pi(g^n) = f(\pi(g)^n) = \psi^n(f)(\pi(g)) = \pi_N^*(\psi^n(f))(g).$$

Hence, A ring monomorphism π_N^* is a λ -monomorphism.

- (2) For any $f_1, f_2 \in \text{Map}(G/N, \mathbb{C})$, we calculate the left hand side as follow.

$$\begin{aligned}
\langle \pi_N^*(f_1), \pi_N^*(f_2) \rangle_G &= \frac{1}{|G|} \sum_{g \in G} \pi_N^*(f_1)(g) \pi_N^*(f_2)(g^{-1}) \\
&= \frac{1}{|G|} \sum_{g \in G} f_1(\pi(g)) f_2(\pi(g)^{-1}) \\
&= \frac{|N|}{|G|} \sum_{\alpha \in G/N} f_1(\alpha) f_2(\alpha^{-1}) = \langle f_1, f_2 \rangle_{G/N}.
\end{aligned}$$

which imply to finish to prove (2). □

Lemma 5.7.2. *Let χ a character of a representation $\rho : G \rightarrow GL(V)$. we have following statements.*

- (1) $\text{Ker}(\rho) = \{g \in G \mid \chi(g) = \chi(e)\}$.
- (2) $N \subset \text{Ker}(\rho)$ holds if only and if $\chi(g) = \chi(e)$ holds for any $g \in N$.

Proof. (1) For any $g \in \text{Ker}(\rho)$, we have $\rho(g) = \text{id}_V$. hence $\chi(g) = \chi(e)$ holds.

Let n be $\dim V$, and let ω is a primitive $|G|$ -th root of unity. For each $g \in G$, since a representation ρ is a unitary, then there are integer numbers a_1, \dots, a_n such that $\chi(g) = \omega^{a_1} + \dots + \omega^{a_n}$. Assume that $\chi(g) = \chi(e) = n$ the we have $a_1, \dots, a_n = 0$ which imply that all eigenvalue of $\rho(g)$ are 1, then $\rho(g) = \text{id}_V$ holds. Hence, we have $g \in \text{Ker}(\rho)$. At last, it is clear that (2) holds from (1). \square

Proposition 5.7.3. *We define $F(N)$ as follows.*

$$F(N) := \{\chi \in \text{Cha}(G) \mid \chi(g) = \chi(e) \text{ for all } g \in N\}$$

This set is a character of a representation $\rho : G \rightarrow GL(V)$ which is satisfied $N \subset \text{Ker}(\rho)$, from Lemma 5.7.2 (2). About this, $F(N) = \pi_N^(\text{Cha}(G/N))$ holds.*

Proof. Let χ be the character of $F(N)$, and let ρ be the representation of G such that its character is χ . Since $N \subset \text{Ker}(\rho)$, There is a representation ρ' of G/N uniquely such that $\rho = \rho' \circ \pi$ holds from Lemma 5.7.2. Let χ' be a character of ρ' , we have $\chi = \chi' \circ \pi_N = \pi_N^*(\chi') \in \pi_N^*(\text{Cha}(G/N))$.

Conversely, assume that χ belongs to $\pi_N^*(\text{Cha}(G/N))$, and let χ' be $\chi = \pi_N^*(\chi')$. Then we have $\chi(g) = \pi_N^*(\chi')(g) = \chi'(e) = \chi(e)$ for any $g \in N$. Also we let $\rho' : G/N \rightarrow GL(V)$ the representation whose character is χ' . Then we have

$$\chi = \pi_N^*(\chi') = \chi' \circ \pi = \text{Tr} \circ (\rho' \circ \pi)$$

which implies that χ is a charcter of $\rho' \circ \pi$. Hence we have $\chi \in F(N)$. \square

Cororally 5.7.4. *The following statements holds for $f \in \text{Map}(G/N, \mathbb{C})$.*

- (1) *A map f belongs to $\text{Cha}(G/N)$ if and only if $\pi_N^*(f)$ belongs to $\text{Cha}(G)$.*
- (2) *A map f is a irreducible character of G/N if and only if $\pi_N^*(f)$ is a irreducible character of G .*

Proof. (1) is clear from Proposition 5.7.3, and (2) is given by (1) and Proposition 5.7.1 (2). \square

Let $\phi_1, \dots, \phi_{k'}$ be all of irreducible character of G/N . Since Corollary 5.7.4 (2), $\pi_N^*(\phi_1), \dots, \pi_N^*(\phi_{k'})$ are irreducible character of G Hence we can be written $\text{Cha}(G/N)$ and image of π_N^* , $F(N)$ as follows.

$$\begin{aligned} \text{Cha}(G/N) &= \{n_1\phi_1 + \dots + n_{k'}\phi_{k'} \mid n_1, \dots, n_{k'} \in \mathbb{Z}_+\} \\ F(N) &= \{n_1\pi_N^*(\phi_1) + \dots + n_{k'}\pi_N^*(\phi_{k'}) \mid n_1, \dots, n_{k'} \in \mathbb{Z}_+\}. \end{aligned}$$

Theorem 5.7.5. *Suppose that the character belongs to $F(N)$ and the character $\chi' \in \text{Cha}(G/N)$ is satisfied that $\chi = \pi_N^*(\chi')$ holds. Thus, the following statements holds.*

- (1) *Character $\lambda^i(\chi)$ and $S^i(\chi)$ belong to $F(N)$ for any $i \geq 1$.*
- (2) *For any $i = 0, 1, 2, \dots$ and $j = 1, 2, \dots, k'$, we have*

$$\langle \pi_N^*(\phi_j), S^i(\chi) \rangle_G = \langle \phi_j, S^i(\chi') \rangle_{G/N}, \quad \langle \pi_N^*(\phi_j), \lambda^i(\chi) \rangle_G = \langle \phi_j, \lambda^i(\chi') \rangle_{G/N}.$$

Proof. (1) In Proposition 5.7.1 (1), we proved that a map π_N^* can be exchanged λ -operation. From Proposition 3.4.3, it implies that π_N^* can be exchanged symmetric powers operation too. Therefore we have

$$\begin{aligned} \lambda^i(\chi) &= \lambda^i \circ \pi_N^*(\chi') = \pi_N^* \circ \lambda^i(\chi') \in F(N), \\ S^i(\chi) &= S^i \circ \pi_N^*(\chi') = \pi_N^* \circ S^i(\chi') \in F(N) \end{aligned}$$

for any $i \geq 0$.

(2) Since π_N^* can be exchanged λ -operation and symmetric powers operation we have the following equality from Proposition 5.7.1 (2) for any $i = 0, 1, 2, \dots$ and $j = 1, 2, \dots, k'$.

$$\begin{aligned} \langle \pi_N^*(\phi_j), \lambda^i(\chi) \rangle_G &= \langle \pi_N^*(\phi_j), \lambda^i \circ \pi_N^*(\chi') \rangle_G \\ &= \langle \pi_N^*(\phi_j), \pi_N^* \circ \lambda^i(\chi') \rangle_G = \langle \phi_j, \lambda^i(\chi') \rangle_{G/N}. \end{aligned}$$

Similarly, we have $\langle \pi_N^*(\chi_j), S^i(\chi) \rangle_G = \langle \chi_j, S^i(\chi') \rangle_{G/N}$. □

In summary, we can be calculated about a character χ which belongs to $F(N)$ by the character χ' of G/N . In particular, any character χ which is the character of the representation ρ belongs to $\text{Ker}(\rho)$, then we can consider on the representation of $G/\text{Ker}(\rho)$. In addition, this representation is faithful, hence we can assume the faithfully of the representation when we calculate the multiplicity.

Example 5.7.6. Let Π is a character which is generated by action G to G/N . Π belongs to $F(N)$, and correspond to a character of a regular representation of G/N .

Example 5.7.7. We consider irreducible characters of S_4 which has dimension 1 or 2. Conjugate classes and irreducible character which has dimension 1 or 2 is the following statement and table.

- (1) $\in C_1$, (1 2) $\in C_2$, (1 2 3) $\in C_3$, (1 2 3 4) $\in C_4$, (1 2)(3 4) $\in C_5$.

	C_1	C_2	C_3	C_4	C_5
ϕ_1	1	1	1	1	1
ϕ_2	1	-1	1	-1	1
ϕ_3	2	0	-1	0	2

Now, we let V is $C_1 \cup C_5$. Thus V is a normal subgroup of S_4 , and the quotient group S_4/V is isomorphic to S_3 . And $\phi_1, \phi_2, \phi_3 \in F(V)$ holds, which correspond to χ_1, χ_2, χ_3 by $\pi_V^* : \text{Map}(S_3, \mathbb{C}) \rightarrow F(V)$ where characters χ_1, χ_2, χ_3 are irreducible character of S_3 (See Example 5.1.1), Therefore the character sum of some ϕ_1, ϕ_2, ϕ_3 can be considered in S_3 . In particular, we can calculate about ϕ_3 from Example 5.1.1 and Example 5.2.2, then we have

$$\begin{aligned} \lambda_t(\phi_3) &= \phi_1 + \phi_3 t + \phi_2 t^2, \\ S_t(\phi_3) &= \frac{1}{(1-t^2)(1-t^3)} \phi_1 + \frac{t^3}{(1-t^2)(1-t^3)} \phi_2 + \frac{t+t^2}{(1-t^2)(1-t^3)} \phi_3. \end{aligned}$$

6 Calculation of some finite groups

In this chapter, we calculate characters of the symmetric powers representation and the exterior powers representation of some finite group, about symmetric group S_3 and S_4 , alternative group A_4 and A_5 , non-commutative group that has cardinality of 21 G_{21} , the dihedral groups D_{2n} , the generalized quaternion groups Q_{4n} and Heisenberg groups H_p .

At each some finite group, we give the conjugate class and character table. Next, we give the conjugate class where a inverse element of its belongs, a cardinality of conjugate class and a order of a element of conjugate class. These information of a finite group are used on calculate about inner product.

We calculate for only the irreducible representation with has dimension more than 2 is here. The reason is because if we have the result about irreducible representations, we can calculate about reducible representations from §5.6. In addition, The result of one dimension representation was given in 5.3, so we keep it to only give a order of the multiplication as a element of $\text{Map}(G, \mathbb{C})$. Some finite group and its character was described in §5, however we will write again.

Basically, the method of calculation about S_3 , A_4 , G_{21} , S_4 and A_5 , it will describe in the chapter §6.1 to §6.5, is the same as Example 5.1.1 and Example 5.2.2 At first, we calculate the character of exterior powers representation using by (5.3) and (5.5), next we calculate $\lambda_t(\chi)(g)$ and $S_t(\chi)(g)$ for each $g \in G$, and compute the generating function of multiplicity of irreducible component (5.7) for any irreducible character.

6.1 A symmetric group S_3

We have already said about S_3 in Example 5.1.1 and Example 5.2.2.

Conjugate classes and the character table of S_3 is the following statement and table.

$$C_1 = \{(1)\}, C_2 = \{(1\ 2), (1\ 3), (2\ 3)\}, C_3 = \{(1\ 2\ 3), (1\ 3\ 2)\}.$$

	C_1	C_2	C_3	Remarks
Inverse	C_1	C_2	C_3	
Cardinality	1	3	2	
Order	1	2	3	
χ_1	1	1	1	It is a trivial.
χ_2	1	-1	1	It has order 2.
χ_3	2	0	-1	

- About a character χ_3

	C_1	C_2	C_3
χ_3	2	0	-1
$\lambda_t(\chi_3)$	$(1+t)^2$	$1-t^2$	$1-t+t^2$
$S_t(\chi_3)$	$\frac{1}{(1-t)^2}$	$\frac{1}{1-t^2}$	$\frac{1}{1+t+t^2}$

$$\lambda_t(\chi_3) = \chi_1 + \chi_3 t + \chi_2 t^2.$$

$$\begin{aligned} S_t(\chi_3) &= \frac{1}{(1-t^2)(1-t^3)}\chi_1 + \frac{t^3}{(1-t^2)(1-t^3)}\chi_2 + \frac{t}{(1-t)(1-t^3)}\chi_3 \\ &= \frac{1}{(1-t^2)(1-t^3)}\left(\chi_1 + t^3\chi_2 + (t+t^2)\chi_3\right). \end{aligned}$$

6.2 An alternating group A_4

Conjugate classes and the character table of A_4 is the following statement and table where ω is a primitive 3-th root of unity.

$$(1) \in C_1, (1\ 2)(3\ 4) \in C_2, (1\ 2\ 3) \in C_3, (1\ 3\ 2) \in C_4.$$

	C_1	C_2	C_3	C_4	Remarks
Inverse	C_1	C_2	C_4	C_3	
Cardinality	1	3	4	4	
Order	1	2	3	3	
χ_1	1	1	1	1	It is a trivial.
χ_2	1	1	ω	ω^2	It has order 3.
χ_3	1	1	ω^2	ω	It has order 3.
χ_4	3	-1	0	0	

- About a character χ_4

	C_1	C_2	C_3	C_4
χ_4	3	-1	0	0
$\lambda_t(\chi_4)$	$(1+t)^3$	$(1-t)(1-t^2)$	$1+t^3$	$1+t^3$
$S_t(\chi_4)$	$\frac{1}{(1-t)^3}$	$\frac{1}{(1+t)(1-t^2)}$	$\frac{1}{1-t^3}$	$\frac{1}{1-t^3}$

$$\lambda_t(\chi_4) = \chi_1 + \chi_4 t + \chi_4 t^2 + \chi_1 t^3.$$

$$\begin{aligned} S_t(\chi_4) &= \frac{1-t^2+t^4}{(1-t^2)^2(1-t^3)}\chi_1 + \frac{t^2}{(1-t^2)^2(1-t^3)}\chi_2 \\ &+ \frac{t^2}{(1-t^2)^2(1-t^3)}\chi_3 + \frac{t}{(1+t)^2(1-t^3)}\chi_4 \\ &= \frac{1}{(1-t^2)^2(1-t^3)}\left((1-t^2+t^4)\chi_1 + t^2\chi_2 + t^2\chi_3 + (t+t^2+t^3)\chi_4\right). \end{aligned}$$

6.3 A non-commutative group G_{21} which has order 21

We let G_{21} be a finite group that is generated by two elements x and y which is satisfied relations $x^7 = y^3 = e$ and $y^{-1}xy = x^2$. This group has order 21, which has the most small of odd cardinality in the non-commutative group.

Conjugate classes and the character table of G_{21} is the following statement and table where ω is a primitive 3-th root of unity and η is a primitive 7-root of unity. In addition, we let the number a and b be satisfied equalities $a := \eta + \eta^2 + \eta^4 = (-1 + i\sqrt{7})/2$ and $b := \eta^3 + \eta^5 + \eta^6 = (-1 - i\sqrt{7})/2$. We have $a + b = -1$ and $ab = 2$.

$$C_1 = \{e\}, C_2 = \{x, x^2, x^4\}, C_3 = \{x^3, x^5, x^6\}, \\ C_4 = \{x^i y \mid i = 0, \dots, 5\}, C_5 = \{x^i y^2 \mid i = 0, \dots, 5\}.$$

	C_1	C_2	C_3	C_4	C_5	Remarks
Inverse	C_1	C_3	C_2	C_5	C_4	
Cardinality	1	3	3	7	7	
Order	1	7	7	3	3	
χ_1	1	1	1	1	1	It is a trivial, and has order 1.
χ_2	1	1	1	ω	ω^2	It has order 3.
χ_3	1	1	1	ω^2	ω	It has order 3.
χ_4	3	a	b	0	0	
χ_5	3	b	a	0	0	

- About a character χ_4

	C_1	C_2	C_3
χ_4	3	a	b
$\lambda_t(\chi_4)$	$(1+t)^3$	$1+at+bt^2+t^3$	$1+bt+at^2+t^3$
$S_t(\chi_4)$	$\frac{1}{(1-t)^3}$	$\frac{1}{1-at+bt^2-t^3}$	$\frac{1}{1-bt+at^2-t^3}$

	C_4	C_5
χ_4	0	0
$\lambda_t(\chi_4)$	$1+t^3$	$1+t^3$
$S_t(\chi_4)$	$\frac{1}{1-t^3}$	$\frac{1}{1-t^3}$

$$\begin{aligned}
\lambda_t(\chi_4) &= \chi_1 + \chi_4 t + \chi_5 t^2 + \chi_1 t^3. \\
S_t(\chi_4) &= \frac{1-t+t^4-t^7+t^8}{(1-t)(1-t^3)(1-t^7)} \chi_1 + \frac{t^4}{(1-t)(1-t^3)(1-t^7)} \chi_2 \\
&+ \frac{t^4}{(1-t)(1-t^3)(1-t^7)} \chi_3 + \frac{t-t^2+t^4}{(1-t)^2(1-t^7)} \chi_4 \\
&+ \frac{t^2-t^4+t^5}{(1-t)^2(1-t^7)} \chi_5 \\
&= \frac{1}{(1-t)(1-t^3)(1-t^7)} \left((1-t+t^4-t^7+t^8) \chi_1 + t^4 \chi_2 + t^4 \chi_3 \right. \\
&+ \left. (t+t^5+t^6) \chi_4 + (t^2+t^3+t^7) \chi_5 \right).
\end{aligned}$$

- About a character χ_5

	C_1	C_2	C_3
χ_4	3	b	a
$\lambda_t(\chi_4)$	$(1+t)^3$	$1+bt+at^2+t^3$	$1+at+bt^2+t^3$
$S_t(\chi_3)$	$\frac{1}{(1-t)^3}$	$\frac{1}{1-bt+at^2-t^3}$	$\frac{1}{1-at+bt^2-t^3}$

	C_4	C_5
χ_4	0	0
$\lambda_t(\chi_4)$	$1+t^3$	$1+t^3$
$S_t(\chi_3)$	$\frac{1}{1-t^3}$	$\frac{1}{1-t^3}$

$$\begin{aligned}
\lambda_t(\chi_5) &= \chi_1 + \chi_5 t + \chi_4 t^2 + \chi_1 t^3. \\
S_t(\chi_4) &= \frac{1-t+t^4-t^7+t^8}{(1-t)(1-t^3)(1-t^7)} \chi_1 + \frac{t^4}{(1-t)(1-t^3)(1-t^7)} \chi_2 \\
&+ \frac{t^4}{(1-t)(1-t^3)(1-t^7)} \chi_3 + \frac{t^2-t^4+t^5}{(1-t)^2(1-t^7)} \chi_4 \\
&+ \frac{t-t^2+t^4}{(1-t)^2(1-t^7)} \chi_5 \\
&= \frac{1}{(1-t)(1-t^3)(1-t^7)} \left((1-t+t^4-t^7+t^8) \chi_1 + t^4 \chi_2 + t^4 \chi_3 \right. \\
&+ \left. (t^2+t^3+t^7) \chi_4 + (t+t^5+t^6) \chi_5 \right).
\end{aligned}$$

6.4 A symmetric group S_4

Conjugate classes and the character table of S_4 is the following statement and table ([5] p.287).

$$(1) \in C_1, (1\ 2) \in C_2, (1\ 2\ 3) \in C_3, (1\ 2\ 3\ 4) \in C_4, (1\ 2)(3\ 4) \in C_5.$$

	C_1	C_2	C_3	C_4	C_5	Remarks
Inverse	C_1	C_2	C_3	C_4	C_5	
Cardinality	1	6	8	6	3	
Order	1	2	3	4	2	
χ_1	1	1	1	1	1	It is a trivial.
χ_2	1	-1	1	-1	1	It has order 2.
χ_3	3	1	0	-1	-1	
χ_4	3	-1	0	1	-1	$\chi_4 = \chi_2\chi_3$.
χ_5	2	0	-1	0	2	$\chi_5 \in F(C_1 \cup C_5)$.

- About a character χ_3

	C_1	C_2	C_3
χ_3	3	1	0
$\lambda_t(\chi_3)$	$(1+t)^3$	$(1+t)(1-t^2)$	$1+t^3$
$S_t(\chi_3)$	$\frac{1}{(1-t)^3}$	$\frac{1}{(1-t)(1-t^2)}$	$\frac{1}{1-t^3}$

	C_4	C_5
χ_3	-1	-1
$\lambda_t(\chi_3)$	$(1-t)(1+t^2)$	$(1-t)(1-t^2)$
$S_t(\chi_3)$	$\frac{1}{(1+t)(1+t^2)}$	$\frac{1}{(1+t)(1-t^2)}$

$$\begin{aligned}
\lambda_t(\chi_3) &= \chi_1 + \chi_3 t + \chi_4 t^2 + \chi_2 t^3. \\
S_t(\chi_3) &= \frac{1}{(1-t^2)(1-t^3)(1-t^4)} \chi_1 + \frac{t^6}{(1-t^2)(1-t^3)(1-t^4)} \chi_2 \\
&+ \frac{t}{(1-t)(1-t^2)(1-t^4)} \chi_3 + \frac{t^3}{(1-t)(1-t^2)(1-t^4)} \chi_4 \\
&+ \frac{t^2}{(1-t^2)^2(1-t^3)} \chi_5 \\
&= \frac{1}{(1-t^2)(1-t^3)(1-t^4)} \left(\chi_1 + t^6 \chi_2 + (t+t^2+t^3) \chi_3 \right. \\
&\left. + (t^3+t^4+t^5) \chi_4 + (t^2+t^4) \chi_5 \right).
\end{aligned}$$

- About a character χ_4

	C_1	C_2	C_3
χ_4	3	-1	0
$\lambda_t(\chi_4)$	$(1+t)^3$	$(1-t)(1-t^2)$	$1+t^3$
$S_t(\chi_4)$	$\frac{1}{(1-t)^3}$	$\frac{1}{(1+t)(1-t^2)}$	$\frac{1}{1-t^3}$

	C_4	C_5
χ_4	1	-1
$\lambda_t(\chi_4)$	$(1+t)(1+t^2)$	$(1-t)(1-t^2)$
$S_t(\chi_4)$	$\frac{1}{(1-t)(1+t^2)}$	$\frac{1}{(1+t)(1-t^2)}$

$$\begin{aligned}
\lambda_t(\chi_4) &= \chi_1 + \chi_4 t + \chi_4 t^2 + \chi_1 t^3. \\
S_t(\chi_4) &= \frac{1-t^3+t^6}{(1-t^2)(1-t^3)(1-t^4)} \chi_1 + \frac{t^3}{(1-t^2)(1-t^3)(1-t^4)} \chi_2 \\
&+ \frac{t^2}{(1-t)(1-t^2)(1-t^4)} \chi_3 + \frac{t-t^2+t^3}{(1-t)(1-t^2)(1-t^4)} \chi_4 \\
&+ \frac{t^2+t^4}{(1-t^2)^2(1-t^3)} \chi_5 \\
&= \frac{1}{(1-t^2)(1-t^3)(1-t^4)} \left((1-t^3+t^6) \chi_1 + t^3 \chi_2 \right. \\
&\left. + (t^2+t^3+t^4) \chi_3 + (t+t^3+t^5) \chi_4 + (t^2+t^4) \chi_5 \right).
\end{aligned}$$

- About a character χ_5 , which is described in Example 5.7.7, it corresponds to 2-dimension irreducible character of S_3 .

	C_1	C_2	C_3	C_4	C_5
χ_3	2	0	-1	0	2
$\lambda_t(\chi_5)$	$(1+t)^2$	$1-t^2$	$1-t+t^2$	$1-t^2$	$(1+t)^2$
$S_t(\chi_5)$	$\frac{1}{(1-t)^2}$	$\frac{1}{1-t^2}$	$\frac{1}{1+t+t^2}$	$\frac{1}{1-t^2}$	$\frac{1}{(1-t)^2}$

$$\begin{aligned}
\lambda_t(\chi_5) &= \chi_1 + \chi_5 t + \chi_2 t^2. \\
S_t(\chi_5) &= \frac{1}{(1-t^2)(1-t^3)} \chi_1 + \frac{t^3}{(1-t^2)(1-t^3)} \chi_2 + \frac{t}{(1-t)(1-t^3)} \chi_5 \\
&= \frac{1}{(1-t^2)(1-t^3)} \left(\chi_1 + t^3 \chi_2 + (t+t^2) \chi_5 \right)
\end{aligned}$$

6.5 An alternating group A_5

An alternating group A_5 is one of simple non-commutative group. Conjugate classes and the character table of A_5 is the following statement and table where η is the primitive 5-th root of unity. And we let numbers a and b be $a := \eta + \eta^{-1} = (-1 + \sqrt{5})/2$ and $b := \eta^2 + \eta^{-2} = (-1 - \sqrt{5})/2$. It is satisfied $a + b = ab = -1$ ([5] p.288).

$(1) \in C_1$, $(1\ 2)(3\ 4) \in C_2$, $(1\ 2\ 3) \in C_3$, $(1\ 2\ 3\ 4\ 5) \in C_4$, $(1\ 3\ 2\ 4\ 5) \in C_5$.

	C_1	C_2	C_3	C_4	C_5	Remarks
Inverse	C_1	C_2	C_3	C_4	C_5	
Cardinality	1	15	20	12	12	
Order	1	2	3	5	5	
χ_1	1	1	1	1	1	It is a trivial.
χ_2	4	0	1	-1	-1	
χ_3	5	1	-1	0	0	
χ_4	3	-1	0	-b	-a	
χ_5	3	-1	0	-a	-b	

- About χ_2

	C_1	C_2	C_3
χ_2	4	0	1
$\lambda_t(\chi_2)$	$(1+t)^4$	$(1-t^2)^2$	$(1+t)(1+t^3)$
$S_t(\chi_2)$	$\frac{1}{(1-t)^4}$	$\frac{1}{(1-t^2)^2}$	$\frac{1}{(1-t)(1-t^3)}$

	C_4	C_5
χ_2	-1	-1
$\lambda_t(\chi_2)$	$1-t+t^2-t^3+t^4$	$1-t+t^2-t^3+t^4$
$S_t(\chi_2)$	$\frac{1}{1+t+t^2+t^3+t^4}$	$\frac{1}{1+t+t^2+t^3+t^4}$

$$\begin{aligned}
\lambda_t(\chi_2) &= \chi_1 + \chi_2 t + (\chi_4 + \chi_5)t^2 + \chi_2 t^3 + \chi_1 t^4. \\
S_t(\chi_2) &= \frac{1-t^2+t^4-t^6+t^8}{(1-t^2)^2(1-t^3)(1-t^5)}\chi_1 \\
&+ \frac{t+t^6}{(1-t)(1-t^2)(1-t^3)(1-t^5)}\chi_2 + \frac{t^2}{(1-t)(1-t^2)^2(1-t^3)}\chi_3 \\
&+ \frac{t^3}{(1-t)(1-t^2)^2(1-t^5)}\chi_4 + \frac{t^3}{(1-t)(1-t^2)^2(1-t^5)}\chi_5 \\
&= \frac{1}{(1-t^2)^2(1-t^3)(1-t^5)} \left((1-t^2+t^4-t^6+t^8)\chi_1 \right. \\
&+ (t+t^2+t^6+t^7)\chi_2 + (t^2+t^3+t^4+t^5+t^6)\chi_3 \\
&+ (t^3+t^4+t^5)\chi_4 + (t^3+t^4+t^5)\chi_5 \left. \right).
\end{aligned}$$

- About χ_3

	C_1	C_2	C_3
χ_3	5	1	-1
$\lambda_t(\chi_3)$	$(1+t)^5$	$(1+t)(1-t^2)^2$	$(1-t+t^2)(1+t^3)$
$S_t(\chi_3)$	$\frac{1}{(1-t)^5}$	$\frac{1}{(1-t)(1-t^2)^2}$	$\frac{1}{(1+t+t^2)(1-t^3)}$

	C_4	C_5
χ_3	0	0
$\lambda_t(\chi_3)$	$1+t^5$	$1+t^5$
$S_t(\chi_3)$	$\frac{1}{1-t^5}$	$\frac{1}{1-t^5}$

$$\begin{aligned}
\lambda_t(\chi_3) &= \chi_1 + \chi_3 t + (\chi_2 + \chi_4 + \chi_5)t^2 + (\chi_2 + \chi_4 + \chi_5)t^3 + \chi_3 t^4 + \chi_1 t^5. \\
S_t(\chi_3) &= \frac{1-t^2+t^4+t^5+t^6-t^8+t^{10}}{(1-t^2)^2(1-t^3)^2(1-t^5)}\chi_1 + \frac{t^2+t^3-t^4+t^5+t^6}{(1-t)^2(1-t^3)^2(1-t^5)}\chi_2 \\
&+ \frac{t+t^2-t^3+t^4+t^5}{(1-t)(1-t^2)^2(1-t^3)^2}\chi_3 + \frac{t^3}{(1-t)^2(1-t^2)^2(1-t^5)}\chi_4 \\
&+ \frac{t^3}{(1-t)^2(1-t^2)^2(1-t^5)}\chi_5 \\
&= \frac{1}{(1-t^2)^2(1-t^3)^2(1-t^5)} \left((1-t^2+t^4+t^5+t^6-t^8+t^{10})\chi_1 \right. \\
&+ (t^2+3t^3+2t^4+2t^6+3t^7+t^8)\chi_2 \\
&+ (t+2t^2+t^3+2t^4+3t^5+2t^6+t^7+2t^8+t^9)\chi_3 \\
&+ (t^3+2t^4+3t^5+2t^6+t^7)\chi_4 \\
&\left. + (t^3+2t^4+3t^5+2t^6+t^7)\chi_5 \right).
\end{aligned}$$

- About χ_4

	C_1	C_2	C_3
χ_4	3	-1	0
$\lambda_t(\chi_4)$	$(1+t)^3$	$(1-t)(1-t^2)$	$1+t^3$
$S_t(\chi_4)$	$\frac{1}{(1-t)^3}$	$\frac{1}{(1+t)(1-t^2)}$	$\frac{1}{(1-t^3)}$

	C_4	C_5
χ_4	$-b$	$-a$
$\lambda_t(\chi_4)$	$1-bt-bt^2+t^3$	$1-at-at^2+t^3$
$S_t(\chi_4)$	$\frac{1}{1+bt-bt^2-t^3}$	$\frac{1}{1+at-at^2-t^3}$

$$\begin{aligned}
\lambda_t(\chi_4) &= \chi_1 + \chi_4 t + \chi_4 t^2 + \chi_1 t^3. \\
S_t(\chi_4) &= \frac{1+t-t^3-t^4-t^5+t^7+t^8}{(1+t)(1-t^2)(1-t^3)(1-t^5)}\chi_1 + \frac{t^3}{(1-t)(1-t^3)(1-t^5)}\chi_2 \\
&+ \frac{t^2}{(1-t^2)^2(1-t^3)}\chi_3 + \frac{t-t^3+t^5}{(1-t^2)^2(1-t^5)}\chi_4 + \frac{t^3}{(1-t^2)^2(1-t^5)}\chi_5 \\
&= \frac{1}{(1+t)(1-t^2)(1-t^3)(1-t^5)} \left((1+t-t^3-t^4-t^5+t^7+t^8)\chi_1 \right. \\
&+ (t^3+2t^4+t^5)\chi_2 + (t^2+t^3+t^4+t^5+t^6)\chi_3 \\
&\left. + (t+t^2-t^4+t^6)\chi_4 + (t^3+t^4+t^5)\chi_5 \right).
\end{aligned}$$

- About χ_5

	C_1	C_2	C_3
χ_5	3	-1	0
$\lambda_t(\chi_5)$	$(1+t)^3$	$(1+t)(1-t^2)$	$1+t^3$
$S_t(\chi_5)$	$\frac{1}{(1-t)^3}$	$\frac{1}{(1-t)(1-t^2)}$	$\frac{1}{(1-t^3)}$

	C_4	C_5
χ_5	$-a$	$-b$
$\lambda_t(\chi_5)$	$1-at-at^2+t^3$	$1-bt-bt^2+t^3$
$S_t(\chi_5)$	$\frac{1}{1+at-at^2-t^3}$	$\frac{1}{1+bt-bt^2-t^3}$

$$\begin{aligned}
\lambda_t(\chi_5) &= \chi_1 + \chi_5 t + \chi_5 t^2 + \chi_1 t^3. \\
S_t(\chi_5) &= \frac{1+t-t^3-t^4-t^5+t^7+t^8}{(1+t)(1-t^2)(1-t^3)(1-t^5)} \chi_1 + \frac{t^3}{(1-t)(1-t^3)(1-t^5)} \chi_2 \\
&+ \frac{t^2}{(1-t^2)^2(1-t^3)} \chi_3 + \frac{t^3}{(1-t^2)^2(1-t^5)} \chi_4 \\
&+ \frac{1-t^3+t^5}{(1-t^2)^2(1-t^5)} \chi_5 \\
&= \frac{1}{(1+t)(1-t^2)(1-t^3)(1-t^5)} \left((1+t-t^3-t^4-t^5+t^7+t^8) \chi_1 \right. \\
&+ (t^3+2t^4+t^5) \chi_2 + (t^2+t^3+t^4+t^5+t^6) \chi_3 \\
&+ \left. (t^3+t^4+t^5) \chi_4 + (t+t^2-t^4+t^6) \chi_5 \right).
\end{aligned}$$

6.6 The dihedral group D_{2n}

We define D_{2n} by the group that is generated by two elements a and b , which is satisfied the relation $a^n = b^2 = e$ and $bab^{-1} = a^{-1}$ for each $n \in \mathbb{N}$. It is a non-commutative group if $n \neq 1$. The group D_{2n} is called the dihedral group.

Conjugate classes and the character table of D_{2n} is different in the case of n is even and n is odd. Now, let η be the primitive n -th root of unity.

- If n is odd, conjugate classes and the character table of D_{2n} is the following statement and table.

$$C_i = \{a^i, a^{-i}\} \quad (i = 0, 1, \dots, \frac{n-1}{2}), \quad C' := \{ba^i \mid i = 0, 1, \dots, n-1\}.$$

	C_i	C'	Remarks
Inverse	C_i	C'	
Cardinality	2	n	The case of C_0 is 1.
Order	$\frac{n}{(i,n)}$	2	The case of C_0 is 1.
χ_1	1	1	It is a trivial.
χ_2	1	-1	It has order 2.
$\tau_k (k = 1, \dots, \frac{n-1}{2})$	$\eta^{ik} + \eta^{-ik}$	0	

- If n is even, conjugate classes and the character table of D_{2n} is the following statement and table.

$$\begin{aligned}
C_i &:= \{a^i, a^{-i}\} \quad (i = 0, 1, \dots, \frac{n}{2}), \\
C' &:= \{ba^{2i} \mid i = 0, 1, \dots, \frac{n}{2} - 1\}, \\
C'' &:= \{ba^{2i-1} \mid i = 0, 1, \dots, \frac{n}{2} - 1\}.
\end{aligned}$$

	C_i	C'	C''	Remarks
Inverse	C_i	C'	C''	
Cardinality	2	$\frac{n}{2}$	$\frac{n}{2}$	The case of C_0 or $C_{\frac{n}{2}}$ are 1.
Order	$\frac{n}{(i,n)}$	2	2	The case of C_0 is 1.
χ_1	1	1	1	It is a trivial.
χ_2	1	-1	-1	It has order 2.
χ_3	$(-1)^i$	1	-1	It has order 2.
χ_4	$(-1)^i$	-1	1	It has order 2.
$\tau_k (k = 1, \dots, \frac{n}{2} - 1)$	$\eta^{ik} + \eta^{-ik}$	0	0	

We calculate the character of symmetric and exterior powers representation of 2-dimension irreducible character $\tau_k (k = 1, 2, \dots, \frac{n-1}{2})$. It is different in the case of n . However, we define $\tau'_k \in \text{Map}(D_{2n}, \mathbb{C})$ by

$$\tau'_k(a^i) := \eta^{ik} + \eta^{-ik}, \quad \tau'_k(ba^i) := 0 \quad (i = 0, 1, \dots, n-1)$$

for each integer k . Hence we have $\tau_k = \tau'_k$ for any $k = 1, \dots, \frac{n-1}{2}$ if n is odd, and for any $k = 1, \dots, \frac{n}{2} - 1$ if n is even. Therefore, to consider τ'_k , we can also calculate in the case of both odd and even of n .

About any τ'_k , we have

- $\tau'_k \tau'_l = \tau'_{k+l} + \tau'_{k-l}$.
- $\tau'_0 = \chi_1 + \chi_2$, and If n is even, $\tau'_{\frac{n}{2}} = \chi_3 + \chi_4$.
- $\tau'_k = \tau'_{-k}$.

- If $k \equiv l \pmod{n}$, then $\tau'_k = \tau'_l$.
- $\chi_2 \tau'_k = \tau'_k$.

for any integers k and l . Using these properties, we obtain characters of symmetric and exterior powers representation of 2-dimension irreducible representation of D_{2n} from the next Proposition.

Proposition 6.6.1. *The following equalities hold for any integer k .*

$$(1) \lambda^2(\tau'_k) = \chi_2.$$

$$(2) S^{2m-1}(\tau'_k) = \sum_{i=1}^m \tau'_{(2i-1)k}, \quad S^{2m}(\tau'_k) = \sum_{i=1}^m \tau'_{2ik} + \chi_1 \quad (m \in \mathbb{N}).$$

Proof. (1) Since $\lambda^2(\tau'_k)$ has one-dimension, it suffices to calculate that $\lambda^2(\tau'_k)(ba^i)$ for any $i = 0, 1, \dots, n-1$. Using (5.3), we have

$$\lambda^2(\tau'_k)(ba^i) = \frac{1}{2}(\tau'_k(ba^i)^2 - \psi^2(\tau'_k)(ba^i)) = \frac{1}{2}(0^2 - 2) = -1. \quad (6.1)$$

Hence, we obtain $\lambda^2(\tau'_k) = \chi_2$ from character table.

(2) The result of (1) and (5.4) give the following equation.

$$S^m(\tau'_k) = \tau'_k S^{m-1}(\tau'_k) - \chi_2 S^{m-2}(\tau'_k) \quad (m \geq 2)$$

To prove (2), we use the induction by $m \in \mathbb{N}$. It is clear that $S^1(\tau'_k)$ equals τ'_k , and putting $m = 2$ in (6.1), we have

$$S^2(\tau'_k) = \tau'_k S^1(\tau'_k) - \chi_2 S^0(\tau'_k) = (\tau'_k)^2 - \chi_2 = \tau'_{2k} + \tau'_0 - \chi_2 = \tau'_{2k} + \chi_1$$

Hence, we prove (2) when $m = 1, 2$. Suppose by induction that (2) holds when $m \in \mathbb{N}$, we have

$$\begin{aligned} S^{2m+1}(\tau'_k) &= \tau'_k S^{2m}(\tau'_k) - \chi_2 S^{2m-1}(\tau'_k) \\ &= \tau'_k \left(\left(\sum_{i=1}^m \tau'_{2ik} \right) + \chi_1 \right) - \chi_2 \left(\sum_{i=1}^m \tau'_{(2i-1)k} \right) \\ &= \left(\sum_{i=1}^m (\tau'_{(2i+1)k} + \tau'_{(2i-1)k}) \right) + \tau'_k - \left(\sum_{i=1}^m \tau'_{(2i-1)k} \right) = \sum_{i=1}^{m+1} \tau'_{(2i-1)k} \\ S^{2m+2}(\tau'_k) &= \tau'_k S^{2m+1}(\tau'_k) - \chi_2 S^{2m}(\tau'_k) \\ &= \tau'_k \left(\sum_{i=1}^{m+1} \tau'_{(2i-1)k} \right) - \chi_2 \left(\left(\sum_{i=1}^m \tau'_{2ik} \right) + \chi_1 \right) \\ &= \left(\sum_{i=1}^{m+1} (\tau'_{2ik} + \tau'_{(2i-2)k}) \right) - \sum_{i=1}^m \tau'_{2ik} - \chi_2 = \sum_{i=1}^{m+1} \tau'_{2ik} + \chi_1. \end{aligned}$$

Hence, we have (2) when $m + 1$. This finishes the induction and the proof. \square

6.7 The generalized quaternion group Q_{4n}

We define Q_{4n} by the group that is generated by two elements a and b , which is satisfied the relation $a^{2n} = e, b^2 = a^n$ and $bab^{-1} = a^{-1}$ for each $n \geq 2$. The group Q_{4n} is called the generalized quaternion group.

Conjugate classes and the character table of Q_{2n} are following statements and table where η is a primitive $2n$ -th root of unity.

$$\begin{aligned} C_i &:= \{a^i, a^{-i}\} \quad (i = 0, 1, \dots, n), \\ C' &:= \{ba^{2i} \mid i = 0, 1, \dots, n-1\}, \\ C'' &:= \{ba^{2i-1} \mid i = 0, 1, \dots, n-1\}. \end{aligned}$$

- There are four character which have one-dimension. however, These two of the four depends on even or odd of n . Its commutator subgroup $D(Q_{4n})$, which is the subgroup generated by the elements $xyx^{-1}y^{-1}$ for $x, y \in Q_{4n}$, is the n -cycle group which is generated an element a^2 and the quotient group $Q_{4n}/D(Q_{4n})$ is a group with cardinality 4. The reason is that if n is even, then the quotient group $Q_{4n}/D(Q_{4n})$ is isomorphic to the direct sum of two 2-cycle group since an element $b^2 = a^n$ belongs to $D(Q_{4n})$, and if n is odd, then $Q_{4n}/D(Q_{4n})$ is isomorphic to the 4-cycle group since an element $b^2 = a^n$ do not belong to $D(Q_{4n})$.

	C_{2i}	C_{2i-1}	C'	C''	Remarks
Inverse	C_{2i}	C_{2i-1}	C'	C''	
Cardinality	2	2	$\frac{n}{2}$	$\frac{n}{2}$	The case of C_0 and C_n is 1.
Order	$\frac{n}{(2i, 2n)}$	$\frac{n}{(2i-1, 2n)}$	2	2	The case of C_0 is 1.
χ_1	1	1	1	1	It is a trivial.
χ_2	1	1	-1	-1	It has order 2
χ'_3	1	-1	1	-1	It is occurred when n is even. It has order 2.
χ'_4	1	-1	-1	1	It is occurred when n is even. It has order 2.
χ''_3	1	-1	i	$-i$	It is occurred when n is odd. It has order 4.
χ''_4	1	-1	$-i$	i	It is occurred when n is odd. It has order 4.

- In addition, characters of two-dimension representation, τ_k ($k = 1, 2, \dots, n-1$) are induced character of the following character

$$\zeta_k : \langle a \rangle \rightarrow \mathbb{C}, \quad \zeta_k(a^i) := \eta^{ik} \quad (i = 0, 1, \dots, 2n-1).$$

where $\langle a \rangle$ is a subgroup of Q_{4n} , which is a $2n$ -cycle group generated by an element a of Q_{4n} . The value of these character is the following table.

	C_i	C', C''
$\tau_k (k = 1, 2, \dots, n-1)$	$\eta^{ik} + \eta^{-ik}$	0

we calculate about the character τ_k . As in the case of the dihedral group, let $\tau'_k \in \text{Map}(Q_{4n}, \mathbb{C})$ be the following equalities for any integer k .

$$\tau'_k(a^i) := \eta^{ik} + \eta^{-ik}, \quad \tau'_k(ba^i) := 0 \quad (i = 0, 1, \dots, 2n-1).$$

Hence, we have $\tau_k = \tau'_k$ for any $k = 1, 2, \dots, n-1$.

About any τ'_k , we have

- $\tau'_k \tau'_l = \tau'_{k+l} + \tau'_{k-l}$.
- $\tau'_0 = \chi_1 + \chi_2, \quad \tau'_n = \begin{cases} \chi'_3 + \chi'_4 & (\text{if } n \text{ is odd.}), \\ \chi''_3 + \chi''_4 & (\text{if } n \text{ is even.}). \end{cases}$
- $\tau'_k = \tau'_{-k}$.
- $k \equiv l \pmod{2n} \implies \tau'_k = \tau'_l$.
- $\chi_2 \tau'_k = \tau'_k$.

for any integers k and l . Using these properties, we obtain characters of symmetric and exterior powers representation of 2-dimension irreducible representation of Q_{4n} from the next Proposition.

Proposition 6.7.1. *The following equalities about an element of $\text{Map}(Q_{4n}, \mathbb{C})$ hold for any integer k*

$$(1) \quad \lambda^2(\tau'_k) = \chi_2^{k+1}.$$

$$(2) \quad S^{2m-1}(\tau'_k) = \sum_{i=1}^m \tau'_{(2i-1)k}, \quad S^{2m}(\tau'_k) = \sum_{i=1}^m \tau'_{2ik} + (\chi_2)^{mk} \quad (m \in \mathbb{N}).$$

Proof. Plan of proof is almost the same as (1).

(1) Since $\lambda^2(\tau'_k)$ has one-dimension. Using by (5.3), we have

$$\begin{aligned} \lambda^2(\tau'_k)(a^i) &= \frac{1}{2}(\tau'_k(a^i)^2 - \psi^2(\tau'_k)(a^i)) = \frac{1}{2}(\eta^{2ki} + 2 + \eta^{-2ki} - \eta^{2ki} - \eta^{-2ki}) = 1, \\ \lambda^2(\tau'_k)(ba^i) &= \frac{1}{2}(\tau'_k(ba^i)^2 - \psi^2(\tau'_k)(ba^i)) = \frac{1}{2}(0^2 - \tau_k(a^n)) = (-1)^{k+1} \end{aligned}$$

Hence, we obtain $\lambda^2(\tau'_k) = \chi_2^{k+1}$ from character table.

(2) The result of (1) and (5.4) give the following equation.

$$S^m(\tau'_k) = \tau'_k S^{m-1}(\tau'_k) - \chi_2^{k+1} S^{m-2}(\tau'_k) \quad (m \geq 2) \quad (6.2)$$

To prove (2), we use the induction by $m \in \mathbb{N}$. It is clear that $S^1(\tau'_k)$ equals τ'_k , and putting $m=2$ in (6.2), we have

$$S^2(\tau'_k) = \tau'_k S^1(\tau'_k) - \chi_2^{k+1} S^0(\tau'_k) = (\tau'_k)^2 - \chi_2^{k+1} = \tau'_{2k} + \tau'_0 - \chi_2^{k+1} = \tau'_{2k} + \chi_2^{k+2} = \tau'_{2k} + \chi_2^k$$

Hence we prove (2) when $m = 1$. Suppose by induction that (2) holds when $m \in \mathbb{N}$, we have

$$\begin{aligned}
S^{2m+1}(\tau'_k) &= \tau'_k S^{2m}(\tau'_k) - \chi_2^{k+1} S^{2m-1}(\tau'_k) \\
&= \tau'_k \left(\left(\sum_{i=1}^m \tau'_{2ik} \right) + \chi_2^{mk} \right) - \chi_2^{k+1} \left(\sum_{i=1}^m \tau'_{(2i-1)k} \right) \\
&= \left(\sum_{i=1}^m (\tau'_{(2i+1)k} + \tau'_{(2i-1)k}) \right) + \tau'_k - \left(\sum_{i=1}^m \tau'_{(2i-1)k} \right) = \sum_{i=1}^{m+1} \tau'_{(2i-1)k} \\
S^{2m+2}(\tau'_k) &= \tau'_k S^{2m+1}(\tau'_k) - \chi_2^{k+1} S^{2m}(\tau'_k) \\
&= \tau'_k \left(\sum_{i=1}^{m+1} \tau'_{(2i-1)k} \right) - \chi_2^{k+1} \left(\left(\sum_{i=1}^m \tau'_{2ik} \right) + \chi_2^{mk} \right) \\
&= \left(\sum_{i=1}^{m+1} (\tau'_{2ik} + \tau'_{(2i-2)k}) \right) - \sum_{i=1}^m \tau'_{2ik} - \chi_2^{mk+k+1} \\
&= \sum_{i=1}^{m+1} \tau'_{2ik} + \chi_2^{mk+k+2} = \sum_{i=1}^{m+1} \tau'_{2ik} + \chi_2^{m(k+1)}
\end{aligned}$$

Hence, we have (2) when $m + 1$. This finishes the induction and the proof. \square

6.8 Heisenberg group modulo p H_p

We let H_p be the following sets.

$$H_p := \left\{ \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} \in GL(3, \mathbb{Z}/p\mathbb{Z}) \mid a, b, c \in \mathbb{Z}/p\mathbb{Z} \right\}$$

where p is a prime number such that $p \neq 2$. It is said to Heisenberg group modulo p .

Conjugate classes and character table of H_p are the following statements and table were η is a primitive p -th root of unity.

$$\begin{aligned}
C(h) &:= \left\{ \begin{pmatrix} 1 & 0 & h \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\} \quad (h \in \mathbb{Z}/p\mathbb{Z}), \\
C(e, f) &:= \left\{ \begin{pmatrix} 1 & e & k \\ 0 & 1 & f \\ 0 & 0 & 1 \end{pmatrix} \mid k \in \mathbb{Z}/p\mathbb{Z} \right\} \quad (e, f \in \mathbb{Z}/p\mathbb{Z}, (e, f) \neq (0, 0)).
\end{aligned}$$

	$C(h)$	$C(e, f)$	Remarks
Inverse	$C(-h)$	$C(-e, -f)$	
Cardinality	1	p	
Order	p	p	In the case of $C(0)$ is 1.
$\chi_{i,j}$ $(i, j = 0, 1, \dots, p-1)$	1	η^{ei+fj}	$\chi_{0,0}$ is a trivial and the other have order p .
τ_s ($s = 1, 2, \dots, p-1$)	$p\eta^{sh}$	0	

The calculate of τ_s had already described in Example 5.5.3, so we described the result again. we have

$$\lambda^n(\tau_s) = \begin{cases} \chi_{0,0} & (n = 0, p), \\ \frac{1}{p} \binom{p}{n} \tau_{n'} & (n = 1, \dots, p-1), \\ 0 & (n > p). \end{cases}$$

$$S^n(\tau_s) = \begin{cases} \frac{1}{p^2} \left(\binom{p+n-1}{n} + (p^2 - 1) \right) \chi_{0,0} \\ + \sum_{i,j \ (i,j) \neq (0,0)} \frac{1}{p^2} \left(\binom{p+n-1}{n} - 1 \right) \chi_{i,j} & (p \mid n), \\ \frac{1}{p} p + n - 1 n \tau_{n'} & (p \nmid n). \end{cases}$$

for any $n = 0, 1, 2, \dots$ where n' is a natural number such that $n' = 1, 2, \dots, p-1$ and $ns \equiv n' \pmod{p}$.

7 Future work

As we write this paper this time, there is the most important starting point which is [2] p.95 In there, Kuntson ,who is an author of [2]. calculate about a character of a 2-th exterior power representation of the irreducible representation of S_3 which has dimension 2 as a example (See Example 5.1.1). However, it is believed that there is not a literature or paper to refer the calculate of the multiplication of the symmetric and exterior representation of finite group since [2]. However, Seeing the result of the regular representation or irreducible representation of Heisenberg group modulo p , it is believed that this will help always somewhere. In the future works, in addition to calculate about other finite groups and representation, it is important to find the result of this paper.

As a final note, I would like to thank variety of people including Professor Hiroyuki Ochiai.

References

- [1] Dale Husemoller, Fiber Bundles Third Edition, Springer-Verlag, 1993
- [2] Donald Knutson, λ -Rings and the Representation Theory of the Symmetric Group, Springer-Verlag, 1973
- [3] Donald Yau, Lambda-rings, World Scientific, 2010
- [4] I.G. Macdonald, Symmetric Function and Hall Polynomials SECOND EDITION, Oxford science publications, 1995
- [5] Martin Isaacs I. , Character theory of finite groups, Academic press New York, 1976
- [6] Peter Schneider, Modular Representation Theory of Finite Groups, Springer-Verlag London, 2013
- [7] Ronco Maria, Free Lie algebra and lambda-ring structure, Bulletin of the Australian Mathematical Society. 11/1994; 50(03):373 - 382.

Graduate School of Mathematics, Kyushu-University,
Nishi-ku Fukuoka, 819-0395, Japan.
t-tamura@math.kyushu-u.ac.jp