

# Fixed points of local actions of nilpotent Lie groups on surfaces

Morris W. Hirsch  
 Mathematics Department  
 University of Wisconsin at Madison  
 University of California at Berkeley

September 14, 2021

## Abstract

Let  $G$  be connected nilpotent Lie group acting locally on a real surface  $M$ . Let  $\varphi$  be the local flow on  $M$  induced by a 1-parameter subgroup. Assume  $K$  is a compact set of fixed points of  $\varphi$  and  $U$  is a neighborhood of  $K$  containing no other fixed points.

*Theorem:* If the Dold fixed-point index of  $\varphi_t|_U$  is nonzero for sufficiently small  $t > 0$ , then  $\text{Fix}(G) \cap K \neq \emptyset$ .

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Local actions</b>	<b>5</b>
<b>3</b>	<b>The fixed-point index</b>	<b>6</b>
<b>4</b>	<b>Fixed point sets, stabilizers and ideals</b>	<b>9</b>
<b>5</b>	<b>Proof of Theorem 1</b>	<b>11</b>

## 1 Introduction

*Notation:*  $M$  is a manifold with tangent bundle  $TM$  and boundary  $\partial M$ . The set of vector fields on  $M$  is  $\mathfrak{v}(M)$ , and the set of vector fields that are  $C^1$  (continuously differentiable) is  $\mathfrak{v}^1(M)$ . The zero set of  $X \in \mathfrak{v}(M)$  is  $Z(X)$ .

$G$  always denotes a connected Lie group with unit element  $e$  and Lie algebra  $\mathfrak{g}$ . We usually treat  $\mathfrak{g}$  as the set of one-parameter subgroups  $X: \mathbb{R} \rightarrow G$ , but we also exploit the linear structure on  $\mathfrak{g}$  defined by its identification with the tangent space to  $G$  at  $e$ .

$\mathbb{R}$  denotes the vector space of real numbers,  $\mathbb{R}_+ = [0, \infty)$ ,  $\mathbb{R}^n$  is Euclidean  $n$ -space.  $\mathbb{Z}$  is the group of integers,  $\mathbb{N} := \{0, 1, \dots\}$  is the set of natural numbers, and  $\emptyset$  is the empty set. The frontier of a subset  $S$  of a topological space is denoted by  $\text{Fr}(S)$ . The fixed point set of  $f: A \rightarrow B$  is  $\text{Fix}(f) := \{a \in A: f(a) = a\}$ . Maps are continuous unless the contrary is indicated.

In 1885 Poincaré [25] published a seminal result on surface dynamics, extended to higher dimensions by Hopf [12] in 1925:

**Theorem (POINCARÉ-HOPF).** *Every vector field on a closed manifold  $M$  of nonzero Euler characteristic vanishes at some point.*

A smooth vector field induces a flow  $\Phi^X := \{\Phi_t^X\}_{t \geq 0}$  — a continuous action of the group  $\mathbb{R}$  — and the theorem implies  $\Phi$  has a fixed point. The same conclusion holds for semi-flows (continuous actions of  $\mathbb{R}_+$ ) on a broad class of spaces including all compact polyhedra and topological manifolds, thanks to Lefschetz’s Fixed Point Theorem [16]).

In his pioneering 1964 paper, E. Lima [18] generalized the Poincaré-Hopf Theorem to actions of connected abelian Lie groups on compact surfaces, allowing nonempty boundaries. This was extended to nilpotent groups in 1986 by my former student J. Plante [24] :

**Theorem (PLANTE).** *Every continuous action of a connected nilpotent Lie group on a compact surface of nonzero Euler characteristic has a fixed point.*

Our goal is a generalization of Plante’s Theorem to local actions (Section 2) on all surfaces. This necessitates replacement of the assumption  $\chi(M) \neq 0$ . The new hypothesis is based Dold’s fixed point index  $I(f) \in \mathbb{Z}$  for maps  $f: U \rightarrow M$ , where  $U \subset M$  is open and  $\text{Fix}(f)$  (see Section 3).

Let  $\Phi := \{\Phi_t\}_{t \in \mathbb{R}}$  be a local flow on  $M$ . A *block for  $\Phi$*  is a compact set

$$K \subset \text{Fix}(\Phi) := \bigcap_{t \in \mathbb{R}} \text{Fix}(\Phi_t)$$

having an *isolating neighborhood*  $U$ : a precompact open neighborhood of  $K$  such that  $K = \text{Fix}(\Phi) \cap \overline{U}$ . When  $\Phi := \Phi^X$ , the local flow generated by a vector field  $X$  on  $M$  then  $K \subset \text{Z}(X)$ , the *zero set* of  $X$ .

For sufficiently small  $t > 0$ , the *index of  $\Phi$  at  $K$*  is well defined by the formula

$$i_K(\Phi) := I(\Phi_t|U).$$

When  $\Phi$  comes from a vector field  $X$  we define

$$i_K(X) = i(X, U) := i_K(X).$$

In Theorem 1 and its corollaries  $G$  is nilpotent, and a local action  $\Phi$  of  $G$  on a surface  $M$  is postulated (see Section 2). Each  $X \in \mathfrak{g}$  is a one-parameter subgroup  $\mathbb{R} \rightarrow G$ , inducing a local flow  $\Phi^X$  on  $M$  whose fixed point set is denoted by  $\text{Fix}(X)$ . A block  $K$  for  $\Phi^X$  is an  $X$ -block.  $K$  is *essential* if  $i_K(X) \neq 0$ .

**Theorem 1.** *If  $K$  is an essential  $X$ -block, then  $\text{Fix}(G) \cap K \neq \emptyset$ .*

As distinct  $X$ -blocks are disjoint, we obtain:

**Corollary 2.** *If  $X$  has  $n$  essential blocks, then  $\text{Fix}(G)$  has at least  $n$  components. ■*

Let  $Z(\mathfrak{a})$  denote the set of common zeros of a set  $\mathfrak{a} \subset \mathfrak{v}(M)$ .

**Corollary 3.** *Let  $\mathfrak{a} \subset \mathfrak{v}^1(M)$  be a finite-dimensional linear subspace tangent to  $\partial M$  and forming a nilpotent Lie algebra under the Lie bracket operation. If  $K$  is an essential block of zeros for some  $X \in \mathfrak{g}$ , then  $Z(\mathfrak{a}) \neq \emptyset$ . ■*

The inspiration for Theorem 1 is a remarkable result of C. Bonatti [5]:

**Theorem (BONATTI).** *Assume  $\dim M \leq 4$ ,  $\partial M = \emptyset$ , and  $X, Y$  are commuting analytic vector fields  $M$ . If  $K$  is an essential block for the local flow generated by  $X$ , then  $Z(Y) \cap K \neq \emptyset$ .<sup>1</sup>*

This is one of the few fixed points theorem for noncompact Lie groups ( $\mathbb{R}^2$  in this case) acting on manifolds that are not compact, or have dimensions  $> 2$ , or have zero Euler characteristic. Another is Borel's Fixed Point Theorem, stated below.

When  $\partial M = \emptyset$ , our definition of the index of  $X$  in  $U$  extends Bonatti's definition for vector fields, which runs as follows. Let  $X$  be a  $C^1$  vector field on  $M$  generating the local flow  $\psi$ . Let  $U \subset M$  be an isolating neighborhood of an  $X$ -block  $K$ . Then  $i_K(X)$  equals the intersection number of  $X(U)$  and the image of the trivial field on  $U$  in the tangent bundle of  $M$ , which is Bonatti's definition. Equivalently: If a vector field  $Y$  is sufficiently close to  $X$  and  $Z(Y) \cap U$  is finite, then  $i_K(X)$  equals the the sum of the Poincaré-Hopf indices of the zeros of  $Y$  in  $U$ . This sum, the *Poincaré-Hopf index* of  $Y|U$ , is denoted by  $i_{PH}(Y|U)$ . See Proposition 9.

## Discussion

Plante's theorem does not extend to Lie groups that are solvable, or even supersolvable,<sup>2</sup> because he proved [24]:

---

<sup>1</sup>“The demonstration of this result involves a beautiful and quite difficult local study of the set of zeros of  $X$ , as an analytic  $Y$ -invariant set. Of course, analyticity is an essential tool in this study, and the validity of this type of result in the smooth case remains an open and apparently hard question.” —P. Molino [19]

<sup>2</sup>Supersolvable: All eigenvalues in the adjoint representation are real. This implies solvable.

- Every compact surface supports a fixed-point free  $C^\infty$  action by the group  $H$  of real matrices  $\begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix}$ , ( $a > 0$ ).

On the other hand, sufficiently strong assumptions imply fixed-point theorem for several natural classes of solvable group actions:

- There is a fixed point in every algebraic action of a solvable, linear, irreducible algebraic group on a complete algebraic variety over an algebraically closed field (Borel [3, 4]; see also Humphreys [17], Onishchik & Vinberg [22]). This celebrated result needs no assumptions on dimensions, Euler characteristics or compactness, and is valid for nonsmooth varieties.

- Borel's theorem extends to holomorphic actions of connected solvable Lie groups on compact Kähler manifolds  $M$  with  $H^1(M) = \{0\}$  (A. Sommese [26]).

- $\text{Fix}(G) \neq \emptyset$  when  $G$  is supersoluble and acts analytically on a compact surface  $M$  with  $\chi(M) \neq 0$  (A. Weinstein and M. W. Hirsch [15]). But this result fails for groups that are solvable but not supersoluble: The group of real matrices  $\begin{bmatrix} a & -b & x \\ b & a & y \\ 0 & 0 & 1 \end{bmatrix}$ , ( $a^2 + b^2 = 1$ ), acts without fixed point on the 2-sphere of oriented lines through the origin in  $\mathbb{R}^3$ . And it fails for  $C^\infty$  actions, by Plante's Theorem.

- The conclusion of Bonatti's theorem holds for analytic vector fields  $X, Y$  on a real or complex 2-manifold  $M$  without boundary, satisfying  $[X, Y] \wedge X = 0$  (Hirsch [11]). Two applications follow:

- Let  $\mathfrak{a}$  be a Lie algebra, perhaps infinite dimensional, of analytic vector fields on a real or complex 2-manifold with empty boundary. If  $X \in \mathfrak{a}$  spans a one-dimensional ideal,  $Z(\mathfrak{a})$  meets every essential  $X$ -block.

- Assume the center of  $G$  has positive dimension and  $M$  is a complex 2-manifold with empty boundary such that  $\chi(M) \neq 0$ . Then every holomorphic action of  $G$  on  $M$  has a fixed point.

M. Belliard [1] classified the pairs  $(G, M)$  where  $G$  has a continuous fixed-point free action on a compact surface  $M$ , relying on the classification of transitive surface actions in Mostow [21]. In particular:

- A solvable  $G$  acts without fixed point on the compact surface  $M$  iff  $G$  maps homomorphically onto  $\text{Aff}_+(\mathbb{R})$ .

For related work on the dynamics of Lie group actions see [2, 8, 9, 10, 13, 14].

**Open questions.** Is there an example of a connected nilpotent Lie group acting without fixed-point on a compact  $n$ -manifold,  $n > 2$ , having nonzero Euler characteristic? Does Bonatti's Theorem generalize to three or more vector fields, or to manifolds of dimensions  $> 4$ ?

## 2 Local actions

For a map  $f: A \rightarrow B$  we adopt the convention that notation such as  $f(\xi)$  presupposes that  $\xi$  is an element or subset of  $A$ . The domain  $A$  of  $f$  is denoted by  $\mathcal{D}f$  and the range of  $f$  by  $\mathcal{R}f := f(A)$ .

Let  $g, \phi$  denote maps. Regardless of the domains and ranges, the composition  $g \circ f$  is defined as the map (perhaps empty)  $g \circ f: x \mapsto g(f(x))$  whose domain is  $f^{-1}(\mathcal{D}g)$ .

The associative law holds for these compositions: The maps  $(h \circ g) \circ f$  and  $h \circ (g \circ f)$  have the same domain

$$D := \{x \in \mathcal{D}h: h(x) \in \mathcal{D}g, \quad g(h(x)) \in \mathcal{D}f\},$$

and

$$x \in D \implies (h \circ g)(f(x)) = h \circ ((g \circ f)(x)).$$

A *local homeomorphism* on a topological space  $Q$  is a homeomorphism between open subsets of  $Q$ . The set of these homeomorphisms is denoted by  $\text{LH}(Q)$ .

**Definition 4.** A *local action* of the connected Lie group  $G$  on a manifold  $M$  is a triple  $(G, M, \alpha)$  where  $\alpha: G \rightarrow \text{LH}(M)$  is a function having the following properties:

- The set  $\Omega_\alpha := \{(g, p) \in G \times M: p \in \mathcal{D}\alpha(g)\}$  is an open neighborhood of  $\{e\} \times M$ .
- The *evaluation map*

$$\text{ev}_\alpha: \Omega_\alpha \rightarrow M, \quad (g, p) \mapsto \alpha(g)(p)$$

is continuous.

- $\alpha(e)$  is the identity map of  $M$ .
- The maps  $\alpha(fg) \circ \alpha(h)$ ,  $\alpha(f) \circ \alpha(gh)$  agree on the intersection of their domains,  $(f, g, h \in G)$ .
- $\alpha(e)$  is the identity map of  $M$ .
- $\alpha(g^{-1}) = \alpha(g)^{-1}$ .

$\alpha$  is a *global action* provided  $\Omega_\alpha = G \times M$ . If  $G$  is connected and simply connected and  $M$  is compact, every local action extends to a unique global action.

In the rest of this section a local action  $(G, M, \alpha)$  is assumed. We sometimes omit the notation “ $\alpha$ ”, writing  $g$  for  $\alpha(g)$ ,  $\mathcal{D}g$  for  $\mathcal{D}\alpha(g)$ , and so forth.

A homomorphism  $\phi: H \rightarrow G$  of Lie groups induces the local action  $\alpha \circ \phi$  of  $H$  on  $M$  defined by

$$\alpha \circ \phi: h \rightarrow \alpha(\phi(h)),$$

called the *pullback* of  $\alpha$  by  $\phi$ . When  $\phi$  is an inclusion we set  $\alpha|_H := \alpha \circ \phi$ . In this way  $\alpha$  induces local actions of all Lie subgroups.

A *local flow* is a local action of the group  $\mathbb{R}$  of real numbers. If  $X: \mathbb{R} \rightarrow G$  is a one-parameter subgroup,  $\alpha \circ X$  is the local flow characterized by

$$(\alpha \circ X)(t): p \mapsto \alpha(X(t))(p), \quad (t \in \mathbb{R}, p \in \mathcal{D}\alpha(X(t))). \quad (1)$$

A Lie algebra homomorphism  $X \mapsto \hat{X}$  from  $\mathfrak{g}$  to  $C^\infty$  vector fields on  $M$  gives rise to a local action  $(G, M, \alpha)$  such that the maps  $t \mapsto (\alpha \circ X(t))$  are the integral curves of  $\hat{X}$ . (See Palais [23, Th. II.11], also Varadarajan [29, Th. 2.16.6]).

A set  $S \subset M$  is *invariant* provided  $g(S)$  is defined and in  $S$  for all  $g \in G$ , or more equivalently:

$$S \subset \mathcal{D}g \cap \mathcal{R}g.$$

The *orbit* of  $p \in M$  is the smallest invariant set containing  $p$ . The *fixed-point set* of  $\alpha$  is

$$\text{Fix}(G) := \bigcap_{g \in G} \text{Fix}(g).$$

**Proposition 5.**  $\text{Fix}(\mathfrak{g}) = \text{Fix}(G)$ .

*Proof.* Equation (1) implies  $\text{Fix}(\mathfrak{g}) \subset \text{Fix}(G)$ . A neighborhood of  $e$  is covered by one-parameter subgroups, and it generates  $G$  because  $G$  is connected. This implies  $\text{Fix}(G) \subset \text{Fix}(\mathfrak{g})$ . ■

If  $H \subset \mathfrak{g}$  is a connected Lie subgroup we set

$$\text{Fix}(H) = \text{Fix}(\alpha|_H).$$

**Proposition 6.** *Let  $G$  have a local action.*

- (i) *If  $p \in \text{Fix}(h) \cap \mathcal{D}g$ , then  $g(p) \in \text{Fix}(ghg^{-1})$ .*
- (ii) *If  $H \subset G$  is a connected normal Lie subgroup,  $\text{Fix}(H)$  is invariant under  $G$ .*

*Proof.* (i) is straightforward and implies (ii). ■

### 3 The fixed-point index

The late A. Dold [6, 7] defined a fixed-point index for a large class of maps having compact fixed-point sets. We use Dold's index to define an index for blocks in local flows. Dold's index  $I(f) \in \mathbb{Z}$  is defined for data  $f, V, M, \mathcal{S}$  where

- $V$  is an open set in a topological space  $\mathcal{S}$ ,
- $f: V \rightarrow \mathcal{S}$  is continuous with  $\text{Fix}(f)$  compact,
- $V$  is a *Euclidean neighborhood retract* (ENR): Some open set in a Euclidean space retracts onto a homeomorph of  $V$ .<sup>3</sup>

We will use the following properties of  $I(f)$ :

(D1)  $I(f) = I(f|V_0)$  if  $V_0 \subset V$  is an open neighborhood of  $\text{Fix}(f)$ .

(D2)  $I(f) = \begin{cases} 0 & \text{if } \text{Fix}(f) = \emptyset, \\ 1 & \text{if } f \text{ is constant.} \end{cases}$

(D3)  $I(f) = \sum_i I(f|V_i)$  if  $V$  is the union of finitely many disjoint open sets  $V_i$ .

(D4)  $I(f_0) = I(f_1)$  if there is a homotopy  $f_t: V \rightarrow \mathcal{S}$ , ( $0 \leq t \leq 1$ ) such that  $\bigcup_t \text{Fix}(f_t)$  is compact.

These correspond to (5.5.11), (5.5.12), (5.5.13) and (5.5.15) in Chapter VII of Dold's book [7].

(D5) Assume  $\mathcal{S}$  is a manifold,  $f$  is  $C^1$ ,  $\text{Fix}(f)$  is a singleton  $p \in M \setminus \partial M$ , and  $\text{Det } f'(p) \neq 0$ . Then  $I(f) = (-1)^\nu$ , where  $\nu$  is the number of distinct eigenvalues  $\lambda$  of  $f'(p)$  such that  $\lambda > 1$ . (See [7, VII.5.17, Ex. 3].)

(D6) Suppose  $\mathcal{S}$  is compact,  $V = \mathcal{S}$ , and  $f$  is homotopic to the identity. Then  $I(f) = \chi(\mathcal{S})$ . (See [7, VII.6.22].)

## The index for local flows

Let  $\varphi := \{\varphi_t\}_{t \in \mathbb{R}}$  be a local flow in a topological space  $\mathcal{S}$ .

A compact set  $K \subset \text{Fix}(\varphi)$  is a *block* for  $\varphi$ , or a  $\varphi$ -*block*, if it has an open, precompact ENR neighborhood  $V \subset \mathcal{S}$  such that  $\overline{V} \cap \text{Fix}(\varphi) = K$ . Such a  $V$  is said to be *isolating* for  $\varphi$ , and for  $(\varphi, K)$ . When  $\varphi$  is smooth this language agrees with the terminology for  $X$ -blocks in the Introduction, and  $\text{Fix}(\varphi) = Z(X)$ .

It turns out that the fixed-point index  $I(\varphi_t|V)$  for all sufficiently small  $t > 0$  depends only on  $\varphi$  and  $K$ :

**Proposition 7.** *If  $V$  is isolating for  $\varphi$ , there exists  $\tau > 0$  such that for all  $t \in (0, \tau]$ :*

(a)  $\text{Fix}(\varphi_t) \cap V$  is compact,

---

<sup>3</sup>The class of ENRs includes metrizable topological manifolds and triangulable subsets of Euclidean spaces.

(b)  $I(\varphi_t|V) = I(\varphi_\tau|V)$ .

*Proof.* If (a) fails there exist convergent sequences  $\{t_k\}$  in  $[0, \infty)$ , and  $\{p_k\}$  in  $V$ , such that

$$t_k \searrow 0, \quad p_k \in \text{Fix}(\varphi_{t_k}) \cap V, \quad p_k \rightarrow q \in \text{Fr}(V).$$

Joint continuity of  $(t, x) \mapsto \varphi_t(x)$  yields the contradiction  $q \in \text{Fix}(\varphi) \cap \text{Fr}(V)$ . Assertion (b) is a consequence of (a) and (D4). ■

**Definition 8.** Using the notation of Proposition 7 we define the *index* of  $\varphi$  in  $V$ , and at  $K$ , as:

$$i(\varphi, V) = i_K(\varphi) := I(\varphi_\tau|V).$$

$K$  and  $V$  are *essential* for  $\varphi$  if  $i_K(V) \neq 0$ . This implies  $K \neq \emptyset$  by (D2).

We say that  $\varphi$  is *smooth* and *generates*  $X \in \mathfrak{v}^1(M)$ , provided  $\mathcal{S}$  is a manifold  $M$  and

$$\left. \frac{\partial}{\partial t} \right|_{t=0} \varphi_t(p) = X_p, \quad (p \in M).$$

This implies  $X|_{\partial M}$  is tangent to  $\partial M$ , and  $Z(X) = \text{Fix}(\varphi)$ .

Recall that  $i_{PH}(X)$  denotes the Poincaré-Hopf index for vector fields  $X \in \mathfrak{v}(M)$  such that  $\partial M = \emptyset$  and  $Z(X)$  finite.

**Proposition 9.** Assume  $\partial M = \emptyset$ . Let  $\varphi$  be a smooth local flow on  $M$  generating  $X \in \mathfrak{v}(M)$ . Suppose  $V \subset M$  is isolating for  $\varphi$ . Let  $\{X^k\}$  be a sequence in  $\mathfrak{v}(M)$  converging to  $X$  such that each set  $Z(X^k) \cap \overline{V}$  is finite. Then  $i(\varphi, V) = i_{PH}(X^k|V)$  for sufficiently large  $k$ .

*Proof.* Choose a sequence  $\{Y^k\}$  in  $\mathfrak{v}^1(M)$  with  $Z(Y^k) \cap \overline{V}$  finite, with  $Y^k$  and so close to  $X^k$  that  $i_{PH}(Y^k|V) = i_{PH}(X^k|V)$  and  $\lim_k Y^k = X$ . Let  $\psi^k$  denote the local flow of  $Y^k$ . There exists  $\rho > 0$  such that

$$\lim_{k \rightarrow \infty} \psi_t^k(x) = \varphi_t(x) \quad \text{uniformly for } (t, x) \in [0, \rho] \times \overline{V}.$$

The conclusion follows by applying (D4) to  $f_0 := \varphi_t|V$  and  $f_1 := \psi_t^k|V$  for sufficiently small  $t > 0$ . ■

This result can be used to show that the Dold index and the Bonatti index coincide in situations where both are defined.

Now assume  $G$  has a local action on  $M$ . Every  $X \in \mathfrak{g}$  generates a local flow  $\varphi^X$  on  $M$ . A block  $K \subset \text{Fix}(\varphi^X)$  is called an  $X$ -block. When  $U \subset M$  is  $U$  is isolating for  $\alpha \circ X$  we say  $U$  is isolating for  $X$ , and set

$$i(X, U) = i_K(X) := i(\varphi^X, U).$$

$K$  is *essential* for  $X$  provided  $i_K(X) \neq 0$ .



**Proposition 10.** *Assume  $V \subset M$  is isolating for  $X$ .*

(a) *The set*

$$\mathfrak{N}(X, V, \mathfrak{g}) := \{Y \in \mathfrak{g} : V \text{ is isolating for } Y \text{ and } i(Y, V) = i(X, V)\}$$

*is an open neighborhood of  $X$  in  $\mathfrak{g}$ .*

(b) *If  $\bar{V}$  is a compact invariant manifold,  $i(Y, V) = \chi(\bar{V})$  for all  $Y \in \mathfrak{N}(X, V, \mathfrak{g})$ .*

*Proof.* (a) Compactness of  $\text{Fr}(V)$  implies that the set  $\mathfrak{N}(\mathfrak{g}) := \{Y \in \mathfrak{v}(\mathfrak{g}) : \text{Fix}(Y) \cap \text{Fr}(V) = \emptyset\}$  is an open neighborhood of  $X$ , and  $V$  is isolating for every  $Y \in \mathfrak{N}(\mathfrak{g})$ . If  $Y \in \mathfrak{N}(\mathfrak{g})$  is sufficiently close to  $X$  and  $0 \leq s \leq 1$  then  $Y_s := (1-s)X + sY$  also lies in  $\mathfrak{N}(\mathfrak{g})$ , and therefore  $i(Y, V) = i(X, V)$  by (D4).

(b) Follows from (D6). ■

## 4 Fixed point sets, stabilizers and ideals

As usual,  $G$  denotes a connected Lie group with Lie algebra  $\mathfrak{g}$ . When  $G$  is nilpotent, its exponential map  $\mathfrak{g} \rightarrow G$  is an analytic diffeomorphism sending subalgebras onto closed subgroups, and ideals onto normal subgroups. In some situations  $\mathfrak{g}$  is more convenient than  $G$  because it has a natural linear structure.

A local action of  $G$  on a surface  $M$  is assumed. Note that  $\text{Fix}(G) = \text{Fix}(\mathfrak{g})$  because  $G$  is connected. The *isotropy group* of  $p \in M$  is the subgroup  $G_p \subset G$  generated by

$$\{g \in G : g(p) = p\}.$$

The *stabilizer* of  $p \in M$  is the subalgebra  $\mathfrak{g}_p \subset \mathfrak{g}$  generated by

$$\{Y \in \mathfrak{g} : p \in \text{Fix}(Y)\}.$$

The *stabilizer* of  $S \subset M$  is  $\mathfrak{g}_S := \bigcap_{p \in S} \mathfrak{g}_p$ . Evidently a one-parameter subgroup  $Y : \mathbb{R} \rightarrow G$  belongs to  $\mathfrak{g}_p$  iff  $Y(\mathbb{R}) \subset G_p$ .

**Lemma 11.** *If  $\mathfrak{g}$  is nilpotent and  $\dim(\mathfrak{g}) \geq 2$ , every element of  $\mathfrak{g}$  lies in an ideal of codimension one.*

*Proof.* If  $\dim(\mathfrak{g}) = 2$  the conclusion is trivial because  $\mathfrak{g}$  is abelian. Assume inductively:  $\dim(\mathfrak{g}) = d \geq 3$  and the lemma holds for Lie algebras of lower dimension. Let  $Y \in \mathfrak{g}$  be arbitrary. Fix a 1-dimensional central ideal  $\mathfrak{j}$  and a surjective Lie algebra homomorphism

$$\pi : \mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{j}.$$

By the inductive assumption  $\pi(Y)$  belongs to a codimension-one ideal  $\mathfrak{f} \subset \mathfrak{g}/\mathfrak{j}$ , whence  $Y$  belongs to the codimension-one ideal  $\pi^{-1}(\mathfrak{f}) \subset \mathfrak{g}$ . ■

The following simple result is useful:

**Proposition 12.** *If  $p \in M$  and  $u, w \subset \mathfrak{g}_p$  are linear subspaces such that  $u + w = \mathfrak{g}$ , then  $p \in \text{Fix}(\mathfrak{g})$ . ■*

The set  $C(\mathfrak{g})$  of codimension-one ideals has a natural structure as a projective variety in the real projective space  $\mathbf{P}^{d-1}$ ,  $d = \dim(\mathfrak{g})$ , and is given the corresponding metrizable topology.

**Proposition 13** (PLANTE [24]). *Assume  $\mathfrak{g}$  is nilpotent.*

- (i) *Every component of  $C(\mathfrak{g})$  has positive dimension.*
- (ii) *Every codimension-one subalgebra of  $\mathfrak{g}$  is an ideal.*
- (iii) *If  $O \subset M$  is a one-dimensional orbit of the local action, then*

$$p \in O \implies \mathfrak{g}_p = \mathfrak{g}_O \in C(\mathfrak{g}).$$

## Minimal sets

A *minimal set* (for the local action of  $G$  on  $M$ ) is a nonempty compact invariant set containing no smaller such set. Compact orbits are minimal sets; all other minimal sets are *exceptional*.

An orbit homeomorphic to the unit circle  $\mathbf{S}^1$  is a *circle orbit*. A circle orbit is *isolated* if it has a neighborhood containing no other circle orbit; otherwise it is *nonisolated*.

The following proposition is adapted from Plante [24]:

**Proposition 14.** *Let  $M_1 \subset M$  be a compact surface.*

- (i) *The number of exceptional minimal sets in  $M_1$  is at most half the genus of  $M_1$ .*
- (ii) *The union of the minimal sets in  $M_1$  is compact.*
- (iii) *The union of the nonisolated circle orbits in  $M_1$  is compact.*
- (iv) *If  $C \subset M$  is a nonisolated circle orbit, every neighborhood of  $C$  contains a compact invariant surface  $Q$  such that:*
  - *each component  $P$  of  $Q$  is either an annulus or a Möbius band,*
  - *$\Lambda \setminus P$  contains at most finitely many minimal sets.*

*Proof.* (i) is a generalization of [24, Lemma 2.3], with the same proof. Slight revisions of arguments on [24, page 155] prove the other assertions. ■

## 5 Proof of Theorem 1

Recall that  $G$  is a connected nilpotent Lie group with a local action on a surface  $M$ , the Lie algebra of  $G$  is  $\mathfrak{g}$ , and  $K \subset M$  is an essential block of fixed points for the induced local flow of a one-parameter subgroup  $X: \mathbb{R} \rightarrow G$ .

The theorem states that  $\text{Fix}(G) \cap K \neq \emptyset$ , which is trivial if  $\dim G \leq 1$ . Assume inductively:  $\dim G > 1$  and the conclusion holds for groups of lower dimension.

Every neighborhood of  $K$  in  $M$  contains an isolating neighborhood  $U$  for  $(X, K)$  such that  $\overline{U}$  is a compact surface. It suffices to prove for all such  $U$  that

$$\text{Fix}(G) \cap U \neq \emptyset. \quad (2)$$

**Lemma 15.** *If  $U$  contains only finitely many minimal sets, Equation (2) holds.*

*Proof.* Since  $\dim(\mathfrak{g}) > 1$  and  $\mathfrak{g}$  is covered by codimension-one ideals (Lemma 11),  $\mathfrak{g}$  contains a set  $\{Y_k\}_{k \in \mathbb{N}}$  converging to  $X$ , and a sequence  $\{\mathfrak{h}_k\}$  of pairwise distinct, codimension-one ideals, such that:  $Y_k \in \mathfrak{h}_k$ ,  $U$  is isolating for  $Y_k$ , and  $i(Y, U) \neq 0$  (Proposition 10). As the set  $K_k := \text{Fix}(\mathfrak{h}_k) \cap U$  is compact, nonempty by the induction hypothesis, and invariant by Proposition 6(ii), there is a minimal set  $L_k \subset K_k \subset U$ . The hypothesis of the Lemma implies there exist indices  $i, j$  such that  $\mathfrak{h}_i \neq \mathfrak{h}_j$  and  $L_i = L_j$ . Equation (2) now follows from Proposition 12. ■

In verifying Equation (2) we can assume  $U$  contains infinitely many minimal sets, thanks to Lemma 15. Setting  $M_1 := \overline{U}$  in Proposition 14, we see that all but finitely many of these are circle orbits, and there is a nonempty, compact, invariant surface  $P \subset \overline{U}$  such that:

$$\chi(P) = 0 \text{ and } \overline{U} \setminus P \text{ contains only finitely many minimal sets.} \quad (3)$$

If  $\text{Fix}(G) \cap P \neq \emptyset$ , Equation (2) holds and the proof is complete. Henceforth assume:

$$\text{Fix}(G) \cap P = \emptyset \quad (4)$$

**Lemma 16.** *There exists  $Z \in \mathfrak{g}$  such that:*

- (a)  $U$  is isolating for  $Z$ ,
- (b)  $i(Z, U) = i(X, U) \neq 0$ ,
- (c)  $\text{Fix}(Z) \cap \partial P = \emptyset$ .

*Proof.* Each of the finitely many components  $C_i \subset \partial P$  contains no fixed point by (4), hence it is a circle orbit. Proposition 13(iii) shows that property (c) holds for all  $Z$  in the dense open set  $\mathfrak{g} \setminus \bigcup_i \mathfrak{g}_{C_i}$ , while (a) and (b) hold if  $Z$  is in the nonempty open set  $\mathfrak{R}(X, U, \mathfrak{g})$  (see Proposition 10). Thus the Lemma is satisfied by all  $Z$  in the nonempty set  $\mathfrak{R}(X, U, \mathfrak{g}) \setminus \bigcup \mathfrak{g}_{C_i}$ . ■

Fix  $Z$  as in Lemma 16. Then

$$i(Z, U) \neq 0. \quad (5)$$

Since  $\text{Fix}(Z) \cap \partial P = \emptyset$  by Lemma 16(c), the sets  $U \setminus P$  and  $P \setminus \partial P$  are isolating for  $Z$ . Therefore

$$i(Z, U) = i(Z, U \setminus P) + i(Z, P \setminus \partial P). \quad (6)$$

by (D3) in Section 3. Now  $i(Z, P \setminus \partial P) = 0$ , by Theorem 10(b) with  $V := P \setminus \partial P$ . Consequently

$$i(Z, U \setminus \partial P) \neq 0 \quad (7)$$

by (5) and (6). Equations (3) and (7) show that  $U \setminus P$  contains only finitely many minimal sets. Therefore Lemma 15 yields

$$\text{Fix}(G) \cap (U \setminus P) \neq \emptyset,$$

implying (2). This completes the proof of Theorem 1. ■

## References

- [1] M. Belliard, *Actions sans points fixes sur les surfaces compactes*, Math. Zeit. **225** (1997), 453–465
- [2] M. Belliard & I. Liousse, *Actions affines sur les surfaces*, Publications IRMA, Universite de Lille, **38** exposé X (1996)
- [3] A. Borel, *Groupes linéaires algébriques*, Annals Math. **64** (1956), 20–80
- [4] A. Borel, “Linear algebraic groups,” notes taken by Hyman Bass. W. A. Benjamin, Inc., New York-Amsterdam, 1969
- [5] C. Bonatti, *Champs de vecteurs analytiques commutants, en dimension 3 ou 4: existence de zéros communs*, Bol. Soc. Brasil. Mat. (N.S.) **22** (1992), 215–247
- [6] A. Dold, *Fixed point index and fixed point theorem for Euclidean neighborhood retracts*, Topology **4** (1965), 1–8
- [7] A. Dold, “Lectures on Algebraic Topology,” Die Grundlehren der mathematischen Wissenschaften Bd. 52, second edition. Springer-Verlag New York, 1972
- [8] D. B. A. Epstein & W. P. Thurston, *Transformation groups and natural bundles*, Proc. London Math. Soc. **38** (1979), 219–236

- [9] M. W. Hirsch, *Actions of Lie groups and Lie algebras on manifolds*, “A Celebration of the Mathematical Legacy of Raoul Bott,” Centre de Recherches Mathématiques, U. de Montréal. Proceedings & Lecture Notes vol. 50 (P. R. Kotiuga, ed.), Amer. Math. Soc. Providence RI (2010), 69-78
- [10] M. W. Hirsch, *Smooth actions of Lie groups and Lie algebras on manifolds*, J. Fixed Point Th. App. **10** (2011), 219–232
- [11] M. W. Hirsch, *Zero sets of Lie algebras of analytic vector fields on real and complex 2-manifolds*, in preparation (2014)
- [12] H. Hopf, *Vektorfelder in Mannifaltigkeiten*, Math. Annalen **95** (1925), 340–367
- [13] J. Horne, *A locally compact connected group acting on the plane has a closed orbit*, Illinois J. Math. **9** (1965) 644–650
- [14] J. Hounie, *Minimal sets of families of vector fields on compact surfaces*, J. Diff. Geom. **16** (1981), 739–744
- [15] M. W. Hirsch & A. Weinstein, *Fixed points of analytic actions of super-soluble Lie groups on compact surfaces*, Ergod. Th. Dyn. Sys. **21** (2001), 1783–1787
- [16] S. Lefschetz *Intersections and transformations of complexes and manifolds*, Trans. Amer. Math. Soc. **28** (1926), 1–49.
- [17] J. Humphreys, “Linear Algebraic Groups.” Springer-Verlag, New York 1975
- [18] E. Lima, *Common singularities of commuting vector fields on 2-manifolds*, Comment. Math. Helv. **39** (1964) 97–110
- [19] P. Molino, Review of Bonatti [5], Math. Reviews MR1179486 (93h:57044) (1993)
- [20] P. Molino & F.-J. Turiel, *Une observation sur les actions de  $\mathbf{R}^p$  sur les variétés compactes de caractéristique non nulle*, Comment. Math. Helv. **61** (1986), 370-375
- [21] G. Mostow, *The extensibility of local Lie groups of transformations and groups on surfaces*, Annals of Math. **52** (1950), 606–636.
- [22] A. Onishchik & É. Vinberg, “Lie Groups and Algebraic Groups,” Springer-Verlag, Berlin 199)
- [23] R. S. Palais, “A Global Formulation of the Lie Theory of Transformation Groups.” Memoirs Amer. Math. Soc. no. 22. American Mathematical Society, Providence 1957

- [24] J. Plante, *Fixed points of Lie group actions on surfaces*, Erg. Th. Dyn. Sys. **6** (1986), 149–161
- [25] H. Poincaré, *Sur les courbes définies par une équation différentielle*, J. Math. Pures Appl. **1** (1885) 167–244
- [26] A. Sommese, *Borel's fixed point theorem for Kaehler manifolds and an application*, Proc. Amer. Math. Soc. **41** (1973), 51–54.
- [27] F.-J. Turiel, *An elementary proof of a Lima's theorem for surfaces*, Publ. Mat. **3** (1989), 555–557
- [28] F.-J. Turiel, *Analytic actions on compact surfaces and fixed points*, Manuscripta Mathematica **110** (2003) 195–201
- [29] V. Varadarajan, “Lie Groups, Lie Algebras, and Their Representations.” Springer-Verlag, New York 1976