# Representation theory of logics: a categorial approach

Darllan Conceição Pinto \* Hugo Luiz Mariano †

December 2013

#### Abstract

The major concern in the study of categories of logics is to describe condition for preservation, under the a method of combination of logics, of meta-logical properties. Our complementary approach to this field is study the "global" aspects of categories of logics in the vein of the categories  $S_s$ ,  $L_s$ ,  $L_s$ , studied in [AFLM3]. All these categories have good properties however the category of logics L does not allow a good treatment of the "identity problem" for logics ([Bez]): for instance, the presentations of "classical logics" (e.g., in the signature  $\{\neg, \lor\}$  and  $\{\neg', \rightarrow'\}$ ) are not  $L_s$ -isomorphic. In this work, we sketch a possible way to overcome this "defect" (and anothers) by a mathematical device: a representation theory of logics obtained from category theoretic aspects on (Blok-Pigozzi) algebraizable logics. In this setting we propose the study of (left and right) "Morita equivalence" of logics and variants. We introduce the concepts of logics (left/right)-(stably) -Morita-equivalent and show that the presentations of classical logics are stably Morita equivalent but classical logics and intuitionist logics are not stably-Morita-equivalent: they are only stably-Morita-adjointly related.

# 1 Introduction

In the 1990's rise many methods of combinations of logics ([CC3]). They appear in dual aspects: as processes of decomposition or analysis of logics (e.g., the "Possible Translation Semantics" of W. Carnielli, , [Car]) or as a processes of composition or synthesis of logics (e.g., the "Fibrings" of D. Gabbay, [Ga]). This was the main motivation of categories of logics. The major concern in the study of categories of logics (CLE-UNICAMP, IST-Lisboa) is to describe condition for preservation, under the combination method, of meta-logical properties ([CCCSS], [ZSS]). Our complementary approach to this field is study the "global" aspects of categories of logics ([AFLM1], [AFLM2], [AFLM3], [MaMe]).

The initial steps on "global" approach to categories of logics are given in the sequence of papers [AFLM1], [AFLM2] and [AFLM3]: they present very simple but too strict notions of logics and morphisms, with "good" categorial properties ([AR]) but unsatisfactory treatment of the "identity problem" of logics ([Bez]). More flexible notions of morphisms between logics are considered in [FC], [BCC1], [BCC2], [CG]: this alternative notion allows better approach to the identity problem however has many categorial "defects". A "refinement" of those ideas is provided in [MaMe]: are considered categories of logics satisfying simultaneously certain natural conditions: (i) represent the major part of logical systems; (ii) have good categorial properties; (iii) allow a natural notion of algebraizable logical system ([BP], [Cze]); (iv) allow satisfactory treatment of the "identity problem" of logics.

In the present work we present and alternative approach to overcome the problems above through a *mathematical device*: a representation theory of logics obtained from category theoretic aspects on (Blok-Pigozzi) algebraizable logics:

Motivation 1: analogy: logics ← rings

- "Representation theory of rings":
- $-R \in obj(Ring) \iff R Mod \text{ (respec., } Mod R) \in CAT;$
- (left/right) Morita equivalence of rings:

 $R \equiv R' \Leftrightarrow R - Mod \simeq R' - Mod \text{ (respec., } Mod - R \simeq Mod - R').$ 

- "Representation theory of propositional logics":
- $-l \in obj(Log) \iff l Mod \text{ (respec., } Mod l): \text{ diagrams of categories and functors (respec.: diagrams of categories, functors and natural transformations);}$

<sup>\*</sup>Instituto de Matemática e Estatística, University of São Paulo, Brazil. Emails: darllan\_math@hotmail.com, hugomar@ime.usp.br

 $<sup>^{\</sup>dagger}$ Research supported by FAPESP, under the Thematic Project LOGCONS: Logical consequence, reasoning and computation (number 2010/51038-0).

- (left/right) Morita equivalence of logics and variants;
- left and right are conceptually and technically distincts.

Motivation 2: analogy: logics ← topology

- "Algebraic Topology": (objects: topological spaces or logics)
- define a general theory of "mathematical invariants" to mesure the degree of distinctions of arbitrary logics;
- develope general methods of calculation of "invariants" (in some sense);
- introduce new forms of comparation of objects.

# 2 Preliminaries

# 2.1 Categories of Signatures and Categories of Logics

If we want define and study categories of logics, we must provide answers to the two natural questions: (i) how to represent a logical system? (ii) what are the relevant notions de morphisms? ([CC1], [CC2], [CCRS], [SSC]). In the following we adopt a simple –and sintatical– approach to this theme.

**2.1.**  $S_s$ , the category of signature and (strict or simple) signature morphisms:

A (propositional, finitary) signature is a sequence of pairwise disjoint sets  $\Sigma = (\Sigma_n)_{n \in \mathbb{N}}$ . In what follows,  $X = \{x_0, x_1, \ldots, x_n, \ldots\}$  will denote a fixed enumerable set (written in a fixed order). Denote  $F(\Sigma)$  (repectively  $F(\Sigma)[n]$ ), the set of  $\Sigma$ -formulas over build from X (respec.  $\{x_0, \ldots, x_{n-1}\}$ ). A (strict) morphism,  $f: \Sigma \to \Sigma'$ , is a sequence of functions:  $f = (f_n)_{n \in \mathbb{N}} : (\Sigma_n)_{n \in \mathbb{N}} \to (\Sigma'_n)_{n \in \mathbb{N}}$ ; this induces a function between formula algebras  $\hat{f}: F(\Sigma) \to F(\Sigma')$ .  $\square$ 

**2.2.**  $S_s \simeq Set^{\mathbb{N}}$ , is a finitely locally presentable category and the fp signatures are the "finite support" signatures.  $\square$ 

Recall:

- (i) locally presentable (= accessible + complete/cocomplete) ([AR], [MP]);
- (ii) a category is  $\kappa$ -accessible if it has  $\kappa$ -filtered colimits and a set of  $\kappa$ -presentable objects such that every object is a  $\kappa$ -filtered colimit of these objects
- **2.3.**  $\mathcal{L}_s$ , the category of (strict) logics over  $\mathcal{S}_s$ :

A logic is a pair  $l = (\Sigma, \vdash)$ , where  $\Sigma$  is a signature and  $\vdash$  is a tarskian consequence operator. A  $\mathcal{L}_s$ -morphism,  $f: (\Sigma, \vdash) \to (\Sigma', \vdash')$ , is a (strict) signature morphism  $f \in \mathcal{S}_s(\Sigma, \Sigma')$  such that  $\hat{f}: F(\Sigma) \to F(\Sigma')$  is a  $(\vdash, \vdash')$ -translation:  $\Gamma \vdash \psi \Rightarrow \hat{f}[\Gamma] \vdash' \hat{f}(\psi)$ , for all  $\Gamma \cup \{\psi\} \subseteq F(\Sigma)$  (i.e., it is "continuous").

**2.4.**  $\mathcal{L}_s$  is a  $\omega$ -locally presentable category and the fp logics are given by a finite set of "axioms" and "inference rules" over a fp signature.

- **2.5.**  $A_s$ , the category (strict) of BP-algebraizable logics (see [**BP**]):
- objects: logic  $l = (\Sigma, \vdash)$ , that has some algebraizing pair  $((\delta \equiv \epsilon), \Delta)$ ;
- morphisms:  $f: l \to l': f \in \mathcal{L}_s(l, l')$  and "preserves algebraizing pair" (well defined).


#### Recall that:

#### **2.6.** Functors:

- Forgetful functors:  $U: \mathcal{A} \to \mathcal{L}_s$ ;  $U': \mathcal{A} \to \mathcal{S}_s$ ;
- If  $f \in \mathcal{A}(l_0, l_1)$  and  $\mathcal{K}^i \subseteq \Sigma^i Str$  is the quasivariety equivalent algebraic semantic of  $l_i$ , then  $f^* : \Sigma^1 Str \to \Sigma^0 Str$   $(M_1 \mapsto (M_1)^f)$  restricts to  $f^* : \mathcal{K}^1 \to \mathcal{K}^0$ .

<b>2.7.</b> Limits and colimits in $\mathcal{A}_s$ : $U$ creates products over "bounded" diagrams. $U'$ creates colimits over non empty diagrams. $U$ creates filtered colimits over if $(l, (\gamma_i)_{i \in (I, \leq)})$ is a colimit cocone, then given $M \in \Sigma - Str$ , $M \in \mathcal{K} \Leftrightarrow M^{\gamma_i} \in \mathcal{K}_i$ , $\forall i \in I$ .	vits,
<b>2.8.</b> $\mathcal{A}_s$ is a finitely accessible category (but not complete/cocomplete). Moreover $U: \mathcal{A} \to \mathcal{L}_s$ is a $\omega$ -access functor.	ible
<b>2.9.</b> Remote algebrization revisited (LFIs):  • $F_i: \mathcal{X}_i \to \mathcal{Y}, i = 0, 1$ , accessible functors $\Rightarrow (F_0 \to F_1)$ is an accessible category;  • accessible categories have a <i>small</i> weakly initial family;  Proposition: For each $l \in Obj(\mathcal{L}_s)$ , there is a <i>small</i> family of $\mathcal{L}_s$ -morphisms $(\eta_i: l \to U(l_i))_{i \in I}$ such that for e $l' \in Obj(\mathcal{A})$ and $f \in \mathcal{L}_s(l, U(l'))$ , there are $i \in I$ and $f_i \in \mathcal{A}(l_i, l')$ such that $U(f_i) \circ \eta_i = f$ .  Corollary: A weak universal property of $\eta: l \to \prod_{i \in I} U(l_i)$ .  Questions: Describe conditions on $l$ such that: $\{l_i: i \in I\} \subseteq (\mathcal{A})_{fp}$ .	ach
$ -\{l_i: i \in I\} \text{ be bounded and } U(\prod_{i \in I} l_i) \xrightarrow{\cong} \prod_{i \in I} U(l_i). $ $ -\text{ we can replace } \exists \iff \exists!. $ Then the ingredients are "canonical" $(I \cong I', l_i \cong l_{i'})$ and allows us to define "the algebraizable spectrum of the logic (analogogy with rings: $R \in Obj(cRing_1) \leadsto (\alpha_P : R \to Frac(R/P))_{P \in Spec(R)}). $	c <i>l</i> "
<b>2.10.</b> But they does not allow a good treatment of the "identity problem" for logics: for instance, the presentation of "classical logics" (e.g., in the signature $\{\neg, \lor\}$ and $\{\neg', \rightarrow'\}$ ) are not $\mathcal{L}_s$ -isomorphic.	ons
In this work, we sketch a possible way to overcome this "defect", by a mathematical device.	
2.11. Other categories of logics	
• $\mathcal{L}_f$ : logical translations with "flexible" signature morphisms $c_n \in \Sigma_n \mapsto \varphi'_n \in F(\Sigma')[n]$ ([FC])	
• $Q\mathcal{L}_f$ : "quotient" category: $f \sim g$ iff $\check{f}(\varphi) \dashv \vdash \check{g}(\varphi)$ . The logics $l$ and $l'$ are equipollent ([ <b>CG</b> ]) iff $l$ and $l'$ are $Q\mathcal{L}_f$ -isomorphic.	
• $\mathcal{L}_f^c \subseteq \mathcal{L}_f$ : "congruential" logics: $\varphi_0 \dashv \vdash \psi_0, \dots, \varphi_{n-1} \dashv \vdash \psi_{n-1} \Rightarrow c_n(\varphi_0, \dots, \varphi_{n-1}) \dashv \vdash c_n(\psi_0, \dots, \psi_{n-1})$ . The inclusion functor $\mathcal{L}_f^c \hookrightarrow \mathcal{L}_f$ has a left adjoint.	
• $Lind(\mathcal{A}_f)\subseteq \mathcal{A}_f$ : "Lindenbaum algebraizable" logics: $\varphi \dashv \vdash \psi \Leftrightarrow \vdash \varphi \Delta \psi$ (well defined). $Lind(\mathcal{A}_f)\subseteq \mathcal{L}_f^c$ and the inclusion functor $Lind(\mathcal{A}_f)\hookrightarrow \mathcal{A}_f$ has a left adjoint.	
• $Q\mathcal{L}_f^c$ (or simply $\mathcal{Q}_f^c$ ): "good" category of logics: represents the major part of logics; has good categorial proper (is an accessible category complete/cocopmplete); solves the identity problem for the presentations of classical logic terms of isomorphism; allows a good notion of algebraizable logic ([MaMe]).	ties ogic
2.12. Dense morphism	
• $f: l \to l' \in \mathcal{L}_f$ is dense iff $\forall \varphi'_n \in F(\Sigma')[n] \ \exists \varphi_n \in F(\Sigma)[n]$ such that $\varphi'_n \dashv' \vdash \check{f}(\varphi_n)$ .	
• $f: l \to l'$ is a $\mathcal{L}$ -epimorphism (= surjective at each level $n \in \mathbb{N}$ ), thus it is a dense $\mathcal{L}$ -morphism.	
• $l' \in \mathcal{L}_f^c \Rightarrow f$ is dense iff $\forall c'_n \in \Sigma'_n \ \exists \varphi_n \in F(\Sigma)[n]$ such that $c'_n(x_0, \dots, x_{n-1}) \dashv \vdash \check{f}(\varphi_n)$ .	

- **2.13.**  $Q_f^c$ -isos: For  $h \in \mathcal{L}_f^c(l, l')$ , are equivalent:
  - $[h] \in \mathcal{Q}^{c}_{f}(l, l')$  is  $\mathcal{Q}^{c}_{f}$ -isomorphism;
  - h is a dense morphism and h is a conservative translation (i.e.,  $\Gamma \vdash \psi \Leftrightarrow \check{h}[\Gamma] \vdash' \check{h}(\psi)$ , for all  $\Gamma \cup \{\psi\} \subseteq F(\Sigma)$ ).  $\square$
- 2.14. Quotient categories of (Lindenbaum) algebraizable logics
- $QLind(\mathcal{A}_f) \hookrightarrow Q\mathcal{A}_f$ :
- closed under directed colimits
- reflective subcategory
- ullet both have non-empty colimits



# 2.2 Algebraizable Logics and Categories


**2.15.** Recall that in the theory of Blok-Pigozzi, to each algebraizable logic  $a = (\Sigma, \vdash)$  is canonically associated a unique quasivariety QV(a) in the same signature  $\Theta$  (its "algebraic codification").

**Lemma 2.16.** The inclusion functor has a left adjoint  $(L, I): QV \rightleftharpoons \alpha - Str:$  given by  $M \mapsto M/\theta_M$  where  $\theta_M$  is the least  $\Sigma$ -congruence in M such that  $M/\theta_M \in QV$ . Moreover, the unity of the adjunction (L, I) has components  $(q_M)_{M \in \Sigma - Str}$ , where  $q_M : M \twoheadrightarrow M/\theta_M$  is the quotient homomorphism.

**Remark 2.17.** The (forgetful) functor  $(QV \xrightarrow{I} \Sigma - Str \xrightarrow{U} Set)$  has the (free) functor  $(Set \xrightarrow{F} \Sigma - Str \xrightarrow{L} QV)$ ,  $Y \mapsto F(Y)/\theta_{F(Y)}$ , as left adjoint. Moreover, if  $\sigma_Y : Y \to U \circ F(Y)$  is the Y-component of the unity of the adjunction (F,U), then  $(Y \xrightarrow{t_Y} UILF(Y)) := (Y \xrightarrow{\sigma_Y} UF(Y) \xrightarrow{U(q_{F(Y)})} UILF(Y))$  is the Y-component of the adjunction  $(L \circ F, U \circ I)$ .

**Theorem 2.18.** Let  $h \in \mathcal{A}_f(a, a')$ , then the induced functor  $h^* : \Sigma' - Str \to \Sigma - Str \ (M' \mapsto (M')^h)$ , "commutes over Set" (i.e.,  $U \circ h^* = U'$ ) and has the following additional properties:

- (a) it has restriction  $h^* : QV(a') \to QV(a)$  (i.e.  $I \circ h^* = h^* \circ I'$ );
- (b) there is a natural epimorphism  $\tilde{h}: L \circ h^{\star} \to h^{\star} \cap cL'$ , that restricts to  $L \circ h^{\star} \circ I' = h^{\star} \cap cL' \circ I'$



Good representation theory of  $Lind(A_f)$ 

**Proposition 2.19.** Let  $g_0, g_1 : l \to a \in \mathcal{L}_f$ , with  $a \in Lind(\mathcal{A}_f)$ .

- (a)  $g_0$  is dense  $\Rightarrow g^* \upharpoonright : QV(a) \to \Sigma str$  is full, faitful and injective on objects.
- (b)  $[g_0] = [g_1] \in \mathcal{Q}_f \Rightarrow g_0^{\star} \models g_1^{\star} \models QV(a) \rightarrow \Sigma str.$

**Proposition 2.20.** Let  $a = (\Sigma, \vdash)$  be a Lindenbaum algebraizable then:

- (a)  $F(\Sigma)/\Delta = F(\Sigma)/(\dashv \vdash)$  is a  $\Sigma$ -structure.
- (b)  $F(\Sigma)/\Delta \in QV(a)$ .
- (c)  $F(\Sigma)/\Delta$  is the free QV(a)-object over the set  $X = \{x_0, \ldots, x_n, \ldots\}$ .

**Proposition 2.21.** Let a and a' be Lindenbaum algebraizable logics. If  $a \overset{[h']}{\rightleftharpoons} a'$  is a pair of inverse  $QLind(\mathcal{A}_f)$ -

isomorphisms (i.e., are  $Q_f^c$ -isomorphisms that preserve algebraizing pair) then:  $QV(a) \stackrel{h' \uparrow}{\underset{h \uparrow}{\rightleftharpoons}} QV(a')$  is an isomorphism of categories.

**Lemma 2.22.**Let  $\Sigma, \Sigma' \in Obj(S_f)$ . Consider  $H: \Sigma' - Str \to \Sigma - Str$  a functor that "commutes over Set" (i.e.  $U \circ H = U'$ ) and, for each set Y, let  $\eta_H(Y): F(Y) \to H(F'(Y))$  be the unique  $\Sigma$ -morphism such that  $(Y \xrightarrow{\sigma_Y} UF(Y) \xrightarrow{U(\eta_Y)} UHF'(Y)) = (Y \xrightarrow{\sigma_Y'} U'F'(Y))$ . Then:

- (a) For each set Y and each  $\psi \in F(Y)$ ,  $Var(\eta_H(Y)(\psi)) \subseteq Var(\psi)$ ;
- (b)  $(\eta_H(Y))_{Y \in Set}$  is a natural transformation  $\eta_H : F \to H \circ F'$ ;
- (c) If H is an isomorphism of categories, then  $\eta_H(Y)$  "preserves variables" (i.e.,  $\forall \psi \in F(Y)$ ,  $Var(\eta_H(Y)(\psi)) = Var(\psi)$ ) and H preserves (strictly) products and substructures.
- (d) For each  $n \in \mathbb{N}$ , let  $X_n := \{x_0, \dots, x_{n-1}\} \subseteq X$ , if  $\eta_H(X_n)$  "preserves variables", then the mapping  $c_n \in \Sigma_n \mapsto \eta_H(X_n)(c_n(x_0, \dots, x_{n-1})) \in F'(X_n)$  determines a  $\mathcal{S}_f$ -morphism  $m_H : \Sigma \to \Sigma'$ .

### 2.23.

- (a) Let  $\Sigma, \Sigma' \in Obj(\mathcal{S}_f)$ . Let  $H: \Sigma' Str \to \Sigma Str$  be a "signature" functor, i.e. a functor satisfying (s1), (s2), (s3): (s1) H "commutes over Set" (s2)  $\eta_H$  "preserves variables" (s3) H preserves (strictly) products and substructures.
- (b) Denote  $S_f^{\dagger}$  the subcategory of the category of diagrams (i.e., the category whose objects are categories and the arrows are change of base morphisms (i.e., some pairs (functors, natural transformations)), given by all the categories  $\Sigma str$  and morphisms  $(H, \eta_H)$  where H is a signature functor.
- (c) Let  $a, a' \in Obj(Lind(A_f))$ . Let  $H : \Sigma' Str \to \Sigma Str$  be a "Lindenbaum" functor, i.e. a signature functor also satisfying (l1), (l2), (l3):

(l1) H has a (unique) restriction to the quasivarieties  $H \upharpoonright : QV(a') \to QV(a)$ (l2)  $\check{m}_H(\Delta) \dashv \vdash \Delta'$  (l3)  $\check{m}_H(\delta) = \check{m}_H(\varepsilon) \vDash_{QV(a')} \vDash \delta' =$ 

 $(l2)\ \check{m}_H(\Delta) \dashv' \vdash \Delta' \\ (d)\ \mathrm{Let}\ a, a' \in Obj(Lind(\mathcal{A}_f)) \ \mathrm{and}\ H : \Sigma' - Str \rightarrow \Sigma - Str \ \mathrm{be}\ \mathrm{a}\ \mathrm{``Lindenbaum''}\ \mathrm{functor.}\ \mathrm{For\ each\ set}\ Y,\ \mathrm{let}\ \bar{\eta}_H(Y) :$ 

 $LF(Y) \to H \upharpoonright (L'F'(Y))$  be the unique QV(a)-morphism such that  $(Y \xrightarrow{t_Y} UILF(Y) \xrightarrow{UI(\bar{\eta}_Y)} UIH \upharpoonright L'F'(Y)) =$ 

 $(Y \xrightarrow{t'_Y} U'I'L'F'(Y))$ . Then  $(\bar{\eta}_H(Y))_{Y \in Set}$  is a natural transformation  $\bar{\eta}_H : L \circ F \to H \upharpoonright \circ L' \circ F'$ .

(d) Denote  $Lind(\mathcal{A}_f)^{\dagger}$  the subcategory of the category of diagrams, given by all the subcategories  $QV(a) \hookrightarrow \Sigma - str$  and morphisms  $(H \uparrow, \bar{\eta}_H)$  where H is a Lindenbaum functor.

### Theorem 2.24.

- (a) The categories  $S_f$  and  $S_f^{\dagger}$  are anti-isomorphic. More precisely, given  $\Sigma, \Sigma' \in S_f$ , the mappings  $h \in S_f(\Sigma, \Sigma') \mapsto (h^{\star}, \eta_{h^{\star}}) \in S_f(\Sigma' str, \Sigma str)^{\dagger}$  and  $(H, \eta_H) \in S_f(\Sigma' str, \Sigma str)^{\dagger} \mapsto m_H \in S_f(\Sigma, \Sigma')$  are inverse bijections.
- (b) The pair of inverse anti-isomorphisms above restricts to a pair of inverse anti-isomorphisms between the categories  $Lind(\mathcal{A}_f)$  and  $Lind(\mathcal{A}_f)^{\dagger}$ .

Moreover, the inverse isomorphisms establish a correspondence:

- (c) If  $h \in Lind(\mathcal{A}_f)(a, a')$  and  $H \in Lind(\mathcal{A}_f)^{\dagger}$  are in correspondence, then also are in correspondence the equivalence class  $\{h' \in Lind(\mathcal{A}_f)(a, a') : [h] = [h'] \in QLind(\mathcal{A}_f)(a, a')\}$  and the equivalence class  $\{H' \in Lind(\mathcal{A}_f)^{\dagger} : H' \models H \uparrow, \bar{\eta}_{H'} = \bar{\eta}_H\}$ .
- (d) If  $h \in Lind(\mathcal{A}_f)(a,a')$  and  $H \in Lind(\mathcal{A}_f)^{\dagger}$  are in correspondence, then [h] is a  $QLind(\mathcal{A}_f)$ -isomorphism (i.e., h is an equipolence of logics)  $\Leftrightarrow$   $(H \upharpoonright, \bar{\eta}_H)$  is an isomorphism of change of bases.

# 3 Representation Theory of Logics

### 3.1 General Logics and Categories

Let  $U: \mathcal{L}_s \to Lind(\mathcal{A}_s)$  denote the forgetful functor.

- Objects: To each logic  $l = (\Sigma, \vdash)$ , are associated two pairs (left and right) of data:
- (I) two comma categories (over  $Lind(A_s)$ ):
- $(l \to U)$ , the "left algebrizable spectrum of l" (analysis process);
- $(U \to l)$ , the "right algebrizable spectrum of l" (synthesis process).




(II) two diagrams (left and right "representation diagram"):

- l- $Mod \iff (l \to U, I);$
- $Mod-l \iff (U \to l, L)$ .


$$l\text{-}Mod: (l \to U)^{op} \longrightarrow (\Sigma - str \leftarrow CAT)$$

$$(a_0, f_0) \mapsto (\Sigma - str \leftarrow QV(a_0))$$

$$(a_0, f_0) \overset{h}{\mapsto} (a_1, f_1) \quad \mapsto \quad ((QV(a_1), f_1^{\star} \upharpoonright I_1) \overset{h^{\star}}{\to} (QV(a_0), f_0^{\star} \upharpoonright I_0))$$



$$\begin{array}{cccc} Mod\text{-}l: & (U \to l) \longrightarrow (2 - CAT \leftarrow \Sigma - str) \\ (a_0, f_0) & \mapsto & (QV(a_0) \stackrel{L_0f_0^{\uparrow}}{\leftarrow} \Sigma - str) \\ (a_0, f_0) & \mapsto & (a_1, f_1) & \mapsto & ((QV(a_0), L_0f_0^{\star} \upharpoonright) \stackrel{(h^{\uparrow}, \tilde{h})}{\rightarrow} (QV(a_1), L_1f_1^{\star} \upharpoonright)) \end{array}$$



$$L_0 f_0^\star \overset{\tilde{h}f_1^\star}{\Rightarrow} h^\star \upharpoonright L_1 f_1^\star \quad \text{where} \quad L_0 h^\star \overset{\tilde{h}}{\Rightarrow} h^\star \upharpoonright L_1$$

Since: 
$$(L_0 h^* g^* \overset{\tilde{h}g^*}{\Rightarrow} h^* \upharpoonright L_1 g^* \overset{h^* \tilde{g}}{\Rightarrow} h^* \upharpoonright g^* \upharpoonright L_2) = L_0(gh)^* \overset{\widetilde{gh}}{\Rightarrow} (gh)^* \upharpoonright L_2$$
  
then:  $(L_0 f_0^* \overset{\tilde{h}f_1^*}{\Rightarrow} h^* \upharpoonright L_1 f_1^* \overset{h^* \upharpoonright (\tilde{g}f_2^*)}{\Rightarrow} h^* \upharpoonright g^* \upharpoonright L_2 f_2^*) = L_0(gh)^* \overset{\widetilde{gh}f_2^*}{\Rightarrow} (gh)^* \upharpoonright L_2 f_2^*$ 

- Arrows: Morphisms between logics  $t: l \to l'$  induce two pairs (left and right) of data:
- (I) a left/right "spectral" functor:


$$(l \to U) \stackrel{\neg \circ t}{\leftarrow} (l' \to U);$$

$$(U \to l) \stackrel{t \circ -}{\to} (U \to l').$$

(II) a left/right "representation diagram" morphism:


$$(l\text{-}Mod) \overset{t^{\star} \circ -}{\leftarrow} (l'\text{-}Mod);$$

$$(Mod\text{-}l) \stackrel{-\circ t^{\star}}{\rightarrow} (Mod\text{-}l').$$



### 3.1. The category of all left modules: LM

objects: a left module for a logic l, i.e. the functor  $left(l): (l \to U)^{op} \stackrel{l-Mod}{\longrightarrow} (\Sigma - str \leftarrow CAT) \hookrightarrow (CAT \leftarrow CAT)$ arrows: a pair  $(B,\tau): left(l') \to left(l)$ , where  $B: (l' \to U) \to (l \to U)$  is a "change of bases" functor, and  $\tau: left(l') \Rightarrow left(l) \circ B$  is a natural transformation with the additional compatibility condition: for each  $(a'_0, f'_0), (a'_1, f'_1) \in (l' \to U) \implies Proj(\tau_{(a'_0, f'_0)}) = Proj(\tau_{(a'_1, f'_1)}) : \Sigma' - str \to \Sigma - str$ 



### Proposition 3.2.

(a) 
$$t: l \stackrel{\cong}{\to} l' \in \mathcal{L}_s \Rightarrow (-\circ t, (t^*, id)) : left(l') \stackrel{\cong}{\to} left(l) \in \mathbf{LM}.$$


(b)  $can_l: l \to l^{(c)}$  induces a LM-isomorphism:  $left(l^{(c)}) \stackrel{\cong}{\to} left(l)$ 

#### 3.2Morita equivalence of logics and variants

• (left/right) Morita equivalence of rings: an equivalence relation coarser than isomorphism (Ex.: For rings,  $R \equiv Mat_{n \times n}(R)$ )

#### **Definition 3.3.** The logics l and l' are left Morita equivalent when:

- (a) Let S be a full subcategory of  $(l \to U)$ . S is called generic if  $S^{op} \hookrightarrow (l \to U)^{op} \stackrel{l-Mod}{\to} (CAT \to \Sigma str)$  is "relatively cofinal" (in the image...)
  - (b) The logics l and l' are left Morita equivalent when there are:
- generic subcategories  $S \hookrightarrow (l \to U)$  and  $S' \hookrightarrow (l' \to U)$ ;
- functors  $B: S' \to S$  and  $B': S \to S'$ ;
- "natural comparations":  $(T, \tau)$  and  $(T', \tau')$
- \*  $T: \Sigma' str \to \Sigma str$  is a functor and for each  $(a', f') \in S'$ ,  $\tau_{f'}: QVcod(f') \stackrel{\cong}{\to} QVcod(B(f'))$  is a isomorphism of categories such that  $B(f')^* \circ \tau_{f'} = T \circ f'^*$  and for each  $(a'_0, f'_0) \xrightarrow{g} (a'_1, f'_1) \in S'$



**Theorem 3.4.** If  $left(l) \cong left(l')$  then l and l' are left Morita equivalent. In particular:

- (a) If  $l \cong l'$ , then l and l' are left Morita equivalent.
- (b) l and  $l^{(c)}$  are left Morita equivalent.

**Theorem 3.5.** If l and l' are equipollent, then they are left Morita equivalent.

**Lemma 3.6.** Let  $l \stackrel{[t]}{\rightarrow} l'$  be a  $Q_f$ -isomorphism. Then:

(a) For each Lindenbaum algebraizale logic  $a' = (\alpha', \vdash')$  and any dense  $\mathcal{L}_s$ -morphism  $f' : l' \to a'$ , consider:

- $\alpha = (\Sigma_n / \equiv_n)_{n \in \mathbb{N}}$ , where  $c_n \equiv_n d_n$  iff  $\hat{f}(\check{t}(c_n)) = \hat{f}(\check{t}(d_n))$ ;
- $\bullet$  f = auoc.
- h be the unique  $S_f$ -morphim such that  $h \bullet f = f' \bullet t$  in  $S_f$  (thus  $\check{h} \circ \hat{f} = \hat{f}' \circ \check{t}$ )
- $a = (\alpha, h^*(\vdash_{a'}));$

Then:


- a is a Lindenbaum algebraizable logic;
- $h: a \to a'$  is a  $\mathcal{L}_f^c$ -morphism preserves algebraizable pair and is an weak equivalence i.e. it induces a  $\mathcal{Q}_f^c$ -isomorphism;
- $f: l \to a$  is a  $\mathcal{L}_s$ -epimorphism (= surjective at each level  $n \in \mathbb{N}$ ), thus it is a dense  $\mathcal{L}_s$ -morphism.

(b) If  $a_0' = (\alpha_0', \vdash_0')$ ,  $a_1' = (\alpha_1', \vdash_1')$  are Lindenbaum algebraizate logic;  $f_0' : l' \to a_0'$ ,  $f_1' : l' \to a_1'$  are dense  $\mathcal{L}_s$ -morphisms and  $g' : a_0' \to a_1'$  is a  $\mathcal{L}_s$ -morphism such that  $g'f_0' = f_1'$ , then:

- there is a unique  $S_s$ -morphism  $g: \alpha_0 \to \alpha_1$  such that  $gf_0 = f_1$ ; Moreover:
- $h_1g = g'h_0$ ;
- g is a  $A_s$ -morphism.

**Definition 3.7.** The logics l and l' are left-stably Morita equivalent when:

- there are functors:  $\Sigma' str \underset{F'}{\rightleftharpoons} \Sigma str;$
- there are functors:  $colim_{f' \in (l' \to U)} \ QV cod(f') \stackrel{E}{\underset{E'}{\rightleftharpoons}} colim_{f \in (l \to U)} \ QV cod(f);$ such that:
- E and E' are quasi-inverse equivalence functors;
- the diagram below commutes:



**Proposition 3.8.** l and l' are left Morita equivalent  $\Rightarrow l$  and l' are left-stably Morita equivalent.

Corollary 3.9.  $left(l) \cong left(l') \implies l$  and l' are left-stably Morita equivalent.


**Proposition 3.10.**  $a \cong a' \in QLind(A_f) \Rightarrow a \text{ and } a' \text{ are left stably Morita equivalent.}$ 

Corollary 3.11. The presentations of classical logics are left stably Morita equivalent.

**Proposition 3.12.** Concerning the relations between Classical logics and Intuitionist logics:

- (a) They are not stably-Morita-equivalent.
- (b) But they are only stably-Morita-adjointly related:

 $L: HA \to BA: H \mapsto H/F_H, \text{ where } F_H = \langle \{a \leftrightarrow \neg \neg a : a \in H\} \rangle$ 



### 4 Final Remarks and Future Works

- Present the adequade definitions "on the right side" that allow get basic results analogous to "left side". Note that the considerations "on left" and "on right" are distincts conceptually ("left" is adequate for analysis of logics; "right" is related to synthesis process) and technically (a 2-categorial aspect is needed "on right").
- Describe necessary/sufficient conditions for Morita equivalence of logics (and variants).
- Induce new (functorial) morphisms between logics from the representation diagrams left(l) and right(l).
- Analise categories of fractions of categories of logics.
- Define a general theory of "mathematical invariants" to mesure the degree of distinctions of arbitrary logics and develope general methods of calculation of invariants (in some sense).
- Understand new notions of identity of logics.
- Describe similar construction on alternative base categories ([MaMe]). (Example: the study of LFIs by Possible Translations Semantics.)

# References

- [AFLM1] P. Arndt, R. A. Freire, O. O. Luciano, H. L. Mariano, Fibring and Sheaves, Proceedings of IICAI-05, Special Session at the 2nd Indian International Conference on Artificial Intelligence, Pune, India, (2005), 1679-1698.
- [AFLM2] P. Arndt, R. A. Freire, O. O. Luciano, H. L. Mariano, On the category of algebraizable logics, CLE e-Prints 6(1) (2006), 24 pages.
- [AFLM3] P. Arndt, R. A. Freire, O. O. Luciano, H. L. Mariano, A global glance on categories in Logic, Logica Universalis 1 (2007), 3-39.
- [AR] J. Adámek, J. Rosický, Locally Presentable and Accessible Categories, Lecture Notes Series of the LMS 189, Cambridge University Press, Cambridge, Great Britain, 1994.

- [Bez] J.Y. Béziau, From Consequence Operator to Universal Logic: A Survey of General Abstract Logic in Logica Universalis: Towards a General Theory of Logic (J.-Y. Beziau, ed.), Birkhaeuser, Basel, 2005.
- [BC] J. Bueno-Soler, W.A. Carnielli, *Possible-translations algebraization for paraconsistent logics*, Bulletin of the Section of Logic **34(2)** (2005), 77-92, University of Lodz, Polônia; CLE e-Prints **5(6)** (2005), 13 páginas.
- [BCC1] J. Bueno, M.E. Coniglio, W.A. Carnielli, Finite algebraizability via possible-translations semantics, Proceedings of CombLog 04 Workshop on Combination of Logics: Theory and Applications, (editors: W.A. Carnielli, F.M. Dionísio e P. Mateus), (2004), 79-86.
- [BCC2] J. Bueno-Soler, M.E. Coniglio, W.A. Carnielli, *Possible-Translations Algebraizability*, Paraconsistency with no Frontiers (2006) (editors: J.-Y. Beziau, W. Carnielli), North-Holland.
- [BP] W. J. Blok, D. Pigozzi, Algebraizable logics, Memoirs of the AMS 396, American Mathematical Society, Providence, USA, 1989.
- [Car] W. A. Carnielli, Many-valued logics and plausible reasoning, Proceedings of the XX International Congress on Many-Valued Logics, IEEE Computer Society, University of Charlotte, USA, (1990), 328-335.
- [Con] M. E. Coniglio, The Meta-Fibring environment: Preservation of meta-properties by fibring, CLE e-Prints 5(4) (2005), 36 páginas.
- [Cze] J. Czelakowski, Protoalgebraic logic, Trends in Logic, Studia Logica Library, Kluwer Academic Publishers, 2001.
- [CC1] W. A. Carnielli, M. E. Coniglio, A categorial approach to the combination of logics, Manuscrito 22 (1999), 64-94.
- [CC2] W. A. Carnielli, M. E. Coniglio, Transfers between logics and their applications, Studia Logica 72 (2002); CLE e-Prints 1(4) (2001), 31 pages.
- [CC3] W. A. Carnielli, M. E. Coniglio, Combining Logics, Stanford Encyclopedia of Philosophy, http://plato.stanford.edu/entries/logic-combining/
- [CC4] W. A. Carnielli, M. E. Coniglio, Splitting logics, in We Will Show Them: Essays in Honour of Dov Gabbay, volume 1, (2005), 389414, College Publications, (Artemov, S., Barringer, H., d'Avila Garcez, A. S., Lamb, L. C., and Woods, J., editors).
- [CCGGS] W. A. Carnielli, M. Coniglio, D. Gabbay, P. Gouveia, C. Sernadas, Analysis and Synthesis of Logics, volume 35 of Applied Logic Series, (2008), Springer.
- [CCRS] C. Caleiro, W. Carnielli, J. Rasga, C. Sernadas, Fibring of Logics as a Universal Construction, Handbook of Philosophical Logic 13 (2005) (editors: D. Gabbay, F. Guenthner), Kluwer Academic Publishers.
- [CCCSS] C. Caleiro, W. Carnielli, M. E. Coniglio, A. Sernadas, C. Sernadas, Fibring Non-Truth-Functional Logics: Complete-ness Preservation, Journal of Logic, Language and Information 12(2) (2003), 183-211; CLE e-Prints 1(1) (2001), 34 pages.
- [CG] C. Caleiro, R. Gonçalves, Equipollent logical systems, Logica Universalis: Towards a General Theory of Logic (Editor J.-Y. Beziau) (2007), 97-110.
- [CR] C. Caleiro, J. Ramos, (2004), *Cryptofibring*, Proceedings of CombLog 04 Workshop on Combination of Logics: Theory and Applications, Lisboa, Portugal (editores: W. A. Carnielli, F. M. Dionísio, P. Mateus) (2004), 87-92.
- [CSS2] M. E. Coniglio, A. Sernadas, C. Sernadas, Fibring logics with topos semantics, Journal of Logic and Computation 13(4) (2003), 595-624.
- [FC] V. L. Fernández, M. E. Coniglio, Fibring algebraizable consequence systems, Proceedings of CombLog 04 Workshop on Combination of Logics: Theory and Applications, (editors: W.A. Carnielli, F.M. Dionísio and P. Mateus), (2004), 93-98.
- [Ga] D. Gabbay, Fibred Semantics and the weaving of logics: part 1, Journal of Symbolic Logic 61(4) (1996), 1057-1120.
- [MaMe] H. L. Mariano, C. A. Mendes, Towards a good notion of categories of logics, 5°-TACL, Marseilles-France, july 2011, 207-210.
- [MP] M. Makkai, R. Paré, Accessible categories: The Foundations of Categorical Model Theory, Contemporary Mathematics 104, American Mathematical Society, Providence, USA, 1989.
- [SSC] A. Sernadas, C. Sernadas, C. Caleiro, Fibring of logics as a categorial construction, Journal of logic and computation, 9(2) (1999), 149-179.
- [ZSS] A. Zanardo, A. Sernadas, C. Sernadas, Fibring: Completeness preservation, The Journal of Symbolic Logic 66(1) (2001), 414-439.