Anti-norms on finite von Neumann algebras

Jean-Christophe Bourin^{*} and Fumio Hiai[†]

Abstract

As the reversed version of usual symmetric norms, we introduce the notion of symmetric anti-norms $\|\cdot\|_{!}$ defined on the positive operators affiliated with a finite von Neumann algebra with a finite normal trace. Related to symmetric anti-norms, we develop majorization theory and superadditivity inequalities of the form $\|\psi(A+B)\|_{!} \geq \|\psi(A)\|_{!} + \|\psi(B)\|_{!}$ for a wide class of functions ψ .

Keywords: finite von Neumann algebra, symmetric norm, symmetric anti-norm, τ -measurable operator, majorization, unitary orbit, spectral dominance, Fuglede-Kadison determinant.

AMS subjects classification 2010: 47A30, 46L52, 46L51

1 Introduction

In Functional Analysis, symmetrically normed Banach functions spaces are classical objects as well as their non-commutative generalizations in the setting of τ -measurable operators affiliated with a von Neumann algebra with a faithful normal finite trace τ . Symmetric norms are homogeneous convex functional completely determined by their values on the positive cone of the function space or the operator algebra. This point of view may motivate the study of concave, homogeneous functionals on positive operators. It is our concern in this article.

A part of our work could fit in a very general setting, for instance, in the C^* -algebra framework. However, we confine to finite von Neumann algebras for two reasons. First, dealing with a von Neumann algebra equipped with a normal finite trace allows to consider unbounded operators, and hence to develop a theory parallel to most of the usual non-commutative Banach function spaces. Secondly, the finiteness assumption allows to consider many functionals which would not make sense in the non-finite case, such as the Fuglede-Kadison determinant. The assumption is also essential for some technical reasons (for instance, a unitary operator can be taken as a phase of the polar decomposition). Moreover, this setup of a finite von Neumann algebra naturally extends the matrix approach in our previous work.

We call our functionals, defined on the positive part \mathcal{N}^+ of a finite von Neumann algebra \mathcal{N} , symmetric anti-norms on \mathcal{N}^+ , as the triangle inequality for norms is then

^{*}Supported by ANR 2011-BS01-008-01.

[†]Supported by Grant-in-Aid for Scientific Research (C)26400103.

reversed. Section 2 gives the precise definition and exhibits an important family of such anti-norms which are derived from symmetric norms. For the convenience of the reader and to fix terminologies which may have some variants in the literature, our discussion also covers basic facts on symmetric norms. Our approach of symmetric norms may have its own interest and originality. Section 4 is devoted to the (non-obvious) extension of these anti-norms to the whole set of densely-defined positive operators affiliated with \mathcal{N} . Here we consider the more classical case of symmetric norms as well. Section 5 presents a superadditivity inequality for convex functions which is a far extension of a classical trace inequality of Rotfel'd. Several norm inequalities follow from this antinorm inequality. In Sections 6 and 7, we focus on a special class of symmetric norms and anti-norms which corresponds, in the commutative case, to the class of rearrangement invariant function spaces. The theory is then related to majorization relations.

Most of the results, norm and anti-norm inequalities given in Sections 4–7, are based on operator inequalities via unitary orbits. These essential operator inequalities are established in Section 3. The idea of the proofs consists in combining a unitary orbit technique for spectral dominance in a finite factor with the disintegration of \mathcal{N} into its factorial components. These results nicely extend the scope of some well-known matrix inequalities to the general finite von Neumann algebra setting.

2 Symmetric norms and symmetric anti-norms

Let \mathcal{N} be a finite von Neumann algebra acting on a separable Hilbert space \mathcal{H} with a faithful normal finite trace τ , and \mathcal{N}^+ the set of positive operators in \mathcal{N} . Let $\overline{\mathcal{N}}$ denote the set of τ -measurable operators affiliated with \mathcal{N} (see [13] for details), and $\overline{\mathcal{N}}^+$ the positive cone of $\overline{\mathcal{N}}$. Since τ is finite, $\overline{\mathcal{N}}$ is the set of all densely-defined closed operators affiliated with \mathcal{N} .

In this article, a symmetric norm $\|\cdot\|$ on \mathcal{N} means a norm satisfying $\|UXV\| = \|X\|$ for all $X \in \mathcal{N}$ and all unitaries $U, V \in \mathcal{N}$. The monotonicity of such a norm in the next lemma is a well-known simple fact [14, Lemma 3.2, Corollary 3.3]. We give an alternative proof. The letter I stands for the identity (of any algebra).

Lemma 2.1. Any symmetric norm $\|\cdot\|$ on \mathcal{N} is monotone, i.e., $\|A\| \leq \|B\|$ if $A, B \in \mathcal{N}^+$ and $A \leq B$.

Proof. Let $T \in \mathcal{N}$ be a contraction. Note that $|T| = (V_1 + V_2)/2$ where $V_1 := |T| + i\sqrt{I - |T|^2}$ and $V_2 := |T| - i\sqrt{I - |T|^2}$ are unitaries. As T = U|T| for some unitary $U \in \mathcal{N}$ (since \mathcal{N} is a finite von Neumann algebra), we have $T = (U_1 + U_2)/2$ with two unitaries $U_1, U_2 \in \mathcal{N}$. Therefore $||TXT|| \leq ||X||$ for all $X \in \mathcal{N}$. Now, assume that $0 \leq A \leq B$ in \mathcal{N} . Then there exists a contraction $C \in \mathcal{N}^+$ such that $A = B^{1/2}CB^{1/2} = VC^{1/2}BC^{1/2}V^*$ for some unitary $V \in \mathcal{N}$. Hence $||A|| = ||C^{1/2}BC^{1/2}|| \leq ||B||$.

Consequently, any symmetric norm on \mathcal{N} is continuous with respect to the operator norm $\|\cdot\|_{\infty}$. In fact, since $|X| \leq \|X\|_{\infty}I$, we have

$$||X|| \le ||X||_{\infty} ||I||, \quad X \in \mathcal{N}.$$
 (2.1)

A symmetric norm $\|\cdot\|$ on \mathcal{N} depends only on its values on positive operators via the polar decomposition, and its restriction to \mathcal{N}^+ satisfies

- (1) $\|\alpha A\| = \alpha \|A\|$ for all $A \in \mathcal{N}^+$ and all scalars $\alpha \ge 0$,
- (2) $||A|| = ||UAU^*||$ for all $A \in \mathcal{N}^+$ and all unitaries $U \in \mathcal{N}$,
- (3) $||A|| \le ||A + B|| \le ||A|| + ||B||$ for all $A, B \in \mathcal{N}^+$.

The first inequality in (3) follows from Lemma 2.1. Conversely, if $\|\cdot\|$ is a non-negative functional on \mathcal{N}^+ satisfying (1)–(3), then $\|X\| := \||X|\|$ for $X \in \mathcal{N}$ becomes a symmetric norm (more precisely, semi-norm) on \mathcal{N} , as immediately shown by a triangle inequality in [1] or by Proposition 3.4 below.

We introduce the notion of symmetric anti-norms on the positive cone \mathcal{N}^+ , by replacing the convexity/subadditivity of symmetric norms with concavity/superadditivity.

Definition 2.2. A symmetric anti-norm $\|\cdot\|_{!}$ on \mathcal{N}^{+} is a functional taking values in $[0, \infty)$ satisfying the following properties:

- (1): $\|\alpha A\|_{!} = \alpha \|A\|_{!}$ for all $A \in \mathcal{N}^{+}$ and all scalars $\alpha \geq 0$,
- (2)! $||A||_! = ||UAU^*||_!$ for all $A \in \mathcal{N}^+$ and all unitaries $U \in \mathcal{N}$,
- $(3)_{!} ||A + B||_{!} \ge ||A||_{!} + ||B||_{!} \text{ for all } A, B \in \mathcal{N}^{+}.$
- $(4)_{!} ||A + \varepsilon I||_{!} \searrow ||A||_{!} \text{ as } \varepsilon \searrow 0 \text{ for all } A \in \mathcal{N}^{+}.$

This definition was first introduced in [6, 7] for the matrix algebra \mathbb{M}_n . In the matrix case, (4)! is equivalent to the usual continuity with respect to the operator norm.

Typical example of symmetric anti-norms are $A \mapsto {\tau(A^q)}^{1/q}$, $0 < q \leq 1$, and $A \mapsto {\tau(A^{-1/p})}^{-1/p}$, $0 . The latter is first defined on the invertible part of <math>\mathcal{N}^+$, and understood for non-invertible operators as

$$\{\tau(A^{-p})\}^{-1/p} := \lim_{\varepsilon \searrow 0} \{\tau((A + \varepsilon I)^{-p})\}^{-1/p},$$

where the finiteness assumption $\tau(I) < \infty$ is essential to have non-trivial functionals on \mathcal{N}^+ . These Schatten like functionals with negative exponents are a special case of a more general family.

Definition 2.3. Fix a symmetric norm $\|\cdot\|$ on \mathcal{N} and p > 0. For each $A \in \mathcal{N}^+$, since $\|(A + \varepsilon I)^{-p}\|^{-1/p}$ decreases as $\varepsilon \searrow 0$ by Lemma 2.1, we can define

$$||A||_{!} := \lim_{\varepsilon \searrow 0} ||(A + \varepsilon I)^{-p}||^{-1/p}.$$

Note that if A is invertible, then the above $||A||_!$ is equal to $||A^{-p}||^{-1/p}$, i.e.,

$$||A^{-p}||^{-1/p} = \lim_{\varepsilon \searrow 0} ||(A + \varepsilon I)^{-p}||^{-1/p}.$$
(2.2)

We call this functional $A \in \mathcal{N}^+ \mapsto ||A||_!$ a *derived anti-norm* and say that it is derived from $|| \cdot ||$ and p.

A derived anti-norm is indeed a symmetric anti-norm as claimed in the next statement.

Theorem 2.4. The above functional $\|\cdot\|_1$ derived from a symmetric norm $\|\cdot\|$ on \mathcal{N} and a p > 0 satisfies

$$||A + B||_! \ge ||A||_! + ||B||_!$$

for every $A, B \in \mathcal{N}^+$. Hence $\|\cdot\|_!$ is a symmetric anti-norm on \mathcal{N}^+ .

To prove this result, we begin with an operator arithmetic-geometric mean inequality.

Lemma 2.5. Let $A, B \in \mathcal{N}^+$. Then, there exists a unitary $V \in \mathcal{N}$ such that

$$|BA| \le \frac{A^2 + VB^2V^*}{2}$$

Proof. Consider the operator on $\mathcal{H} \oplus \mathcal{H}$,

$$\begin{bmatrix} A^2 & AB \\ BA & B^2 \end{bmatrix}$$

which is a positive operator. Thus, for any $V \in \mathcal{N}$ so is

$$\begin{bmatrix} I & -V \end{bmatrix} \begin{bmatrix} A^2 & AB \\ BA & B^2 \end{bmatrix} \begin{bmatrix} I \\ -V^* \end{bmatrix}$$

Letting V^* be the unitary factor in the polar decomposition $BA = V^*|BA|$ yields the inequality.

Combining Lemmas 2.1 and 2.5 yields $||AB|| \leq (s||A^2||+s^{-1}||B^2||)/2$ for all symmetric norms, $A, B \in \mathcal{N}^+$, and s > 0. Thus, minimizing over s and considering operators $X, Y \in \mathcal{N}$ with $|X^*| = A$, $|Y^*| = B$, we obtain the Cauchy-Schwarz inequality for symmetric norms.

Corollary 2.6. Let $X, Y \in \mathcal{N}$. Then, for any symmetric norm on \mathcal{N} ,

$$||X^*Y|| \le ||X^*X||^{1/2} ||Y^*Y||^{1/2}$$

As a byproduct of this inequality we get from $\|\cdot\|$ another symmetric norm.

Corollary 2.7. If $\|\cdot\|$ is a symmetric norm on \mathcal{N} , then $X \mapsto \|X^*X\|^{1/2}$ is a symmetric norm on \mathcal{N} too.

For symmetric anti-norms, the following proposition is known in the matrix case [6].

Lemma 2.8. If $\|\cdot\|_!$ is a symmetric anti-norm on \mathcal{N}^+ and 0 < q < 1, then $A \mapsto \|A^q\|_!^{1/q}$ is a symmetric anti-norm on \mathcal{N}^+ too.

Proof. The same proof as in the matrix case [6] shows that it is a homogeneous, unitarily invariant and concave functional. The continuity property (4)_!, i.e., $||A^q||_!^{1/q} = \lim_{\varepsilon \searrow 0} ||(A + \varepsilon)^q||_!^{1/q}$ is obvious from the monotonicity of $|| \cdot ||_!$ since $A^q \leq (A + \varepsilon I)^q \leq A^q + \varepsilon^q I$. We are now in a position to prove the theorem.

Proof. (Theorem 2.4) Let $\|\cdot\|$ be a symmetric norm on \mathcal{N} . Let $A, B \in \mathcal{N}^+$ be invertible and assume that $\|A^{-1}\| = \|B^{-1}\| = 1$. As $t \mapsto t^{-1}$ is operator convex on $(0, \infty)$, we have, for 0 < s < 1,

$$\|(sA + (1-s)B)^{-1}\| \le \|sA^{-1} + (1-s)B^{-1}\| \le s + (1-s) = 1$$

and so

$$||(sA + (1 - s)B)^{-1}||^{-1} \ge 1.$$

For general invertible $S, T \in \mathcal{N}^+$, taking in this estimate $A = ||S^{-1}||S$, $B = ||T^{-1}||T$, and $s = ||S^{-1}||^{-1}/(||S^{-1}||^{-1} + ||T^{-1}||^{-1})$ yields

$$||(S+T)^{-1}||^{-1} \ge ||S^{-1}||^{-1} + ||T^{-1}||^{-1}.$$

Therefore, $A \mapsto ||A||_! := ||A^{-1}||^{-1}$ is a homogeneous and concave/superadditive functional on the invertible part of \mathcal{N}^+ . It can be extended with the same properties to the whole of \mathcal{N}^+ by the limit formula $||A||_! := \lim_{\varepsilon \searrow 0} ||A + \varepsilon I||_!$. Hence, this functional derived from a symmetric norm $|| \cdot ||$ and p = 1 is a symmetric anti-norm on \mathcal{N}^+ .

Next we consider a functional derived from a symmetric norm $\|\cdot\|$ and an arbitrary p > 0. We have $p = 2^n q$ where n is a positive integer and 0 < q < 1. By Corollary 2.7 applied n times and the first step of the proof,

$$||A||_{!}^{(n)} := \lim_{\varepsilon \searrow 0} ||(A + \varepsilon I)^{-2^{n}}||^{-1/2^{n}}$$

is a symmetric anti-norm on \mathcal{N}^+ . Applying Lemma 2.8 shows that $A \mapsto \left(\|A^q\|_{!}^{(n)} \right)^{1/q}$ is a symmetric anti-norm too, which is readily verified to be the functional derived from $\|\cdot\|$ and p.

3 Inequalities via unitary orbits

This section is a main technical part of this article. The next theorems give superadditive or subadditive operator inequalities via unitary orbits for convex or concave functions, which will be of essential use in Sections 4 and 5.

Theorem 3.1. Let g(t) be a non-negative convex function on $[0, \infty)$ with g(0) = 0. Then, for every $A, B \in \overline{\mathcal{N}}^+$ and every $\varepsilon > 0$, there exist unitaries $U, V \in \mathcal{N}$ such that

$$g(A+B) + \varepsilon I \ge Ug(A)U^* + Vg(B)V^*.$$

Theorem 3.2. Let f(t) be a non-negative concave function on $[0, \infty)$. Then, for every $A, B \in \overline{\mathcal{N}}^+$ and every $\varepsilon > 0$, there exist unitaries $U, V \in \mathcal{N}$ such that

$$f(A+B) \le Uf(A)U^* + Vf(B)V^* + \varepsilon I.$$

Before proving the theorems we recall the notion of the spectral scale [26]. The spectral scale of $A \in \overline{\mathcal{N}}^+$ is defined as

$$\lambda_t(A) := \inf\{s \in \mathbb{R} : \tau(\mathbf{1}_{(s,\infty)}(A)) \le t\}, \qquad t \in (0,\tau(I)), \tag{3.1}$$

where $\mathbf{1}_{(s,\infty)}(A)$ is the spectral projection of A corresponding to (s,∞) . We write $\lambda(A)$ for the function $t \mapsto \lambda_t(A)$ on $(0,\tau(I))$, which is non-increasing and right-continuous. Furthermore, we write $\lambda_0(A)$ and $\lambda_{\tau(I)}(A)$ for $\lim_{t \searrow 0} \lambda_t(A)$ and $\lim_{t \nearrow \tau(I)} \lambda_t(A)$, respectively, which are the maximal and minimal spectra of A (when A is bounded). The generalized s-numbers [13] of $X \in \overline{\mathcal{N}}$ is $\mu_t(X) := \lambda_t(|X|), t \in (0, \tau(I))$.

What we will use to prove the theorems is the following lemma. The lemma is rather well-known but we give the proof for the convenience of the reader.

Lemma 3.3. Let \mathcal{N} be a finite factor and $A, B \in \overline{\mathcal{N}}^+$. If B spectrally dominates A, *i.e.*, $\lambda_t(A) \leq \lambda_t(B)$ for all $t \in (0, \tau(I))$, then for every $\varepsilon > 0$ there exists a unitary $U \in \mathcal{N}$ such that $UAU^* \leq B + \varepsilon I$.

Proof. Since the matrix case is obvious without εI in the right-hand side, we may assume that \mathcal{N} is a type II₁ factor with the normalized trace τ . Choose an increasing family $\{F_t\}_{0 \le t \le 1}$ of projections in \mathcal{N} such that $\tau(F_t) = t$ for all $t \in [0, 1]$. Define

$$\tilde{A} := \int_0^1 \lambda_t(A) \, dF_t, \qquad \tilde{B} := \int_0^1 \lambda_t(B) \, dF_t.$$

We then have $\lambda(\tilde{A}) = \lambda(A), \lambda(\tilde{B}) = \lambda(B)$ and $\tilde{A} \leq \tilde{B}$. Hence the assertion follows since $||A - V\tilde{A}V^*||_{\infty} < \varepsilon/2$ and $||B - W\tilde{B}W^*||_{\infty} < \varepsilon/2$ for some unitaries $V, W \in \mathcal{N}$ by [18, Lemma 4.1].

We now turn to the proofs of the theorems, which are based on the spectral dominance theorem [9] and the central direct decomposition.

Proof. (Theorem 3.1) First assume that \mathcal{N} is a finite factor. Then the matrix case is [2, Theorem 2.1] (without εI in the left-hand side). The proof in the type II₁ factor case is similar based on [9]. For any contraction $Z \in \mathcal{N}$ and any $T \in \overline{\mathcal{N}}^+$ it is known [9, Lemma 10 (ii)] that $Z^*g(T)Z$ spectrally dominates $g(Z^*TZ)$. Hence, by Lemma 3.3,

$$Z^*g(T)Z + \varepsilon I \ge Wg(Z^*TZ)W^* \tag{3.2}$$

for some unitary $W \in \mathcal{N}$. Then, by arguing as in the proof of [2, Theorem 2.1] or [8, Corollary 3.2], one can see that the claimed inequality holds for some unitaries $U, V \in \mathcal{N}$.

For the non-factor case, as in [16], we take the central direct integral decomposition into factors (see [27]) as

$$\{\mathcal{N},\mathcal{H}\} = \int_{\Omega}^{\oplus} \{\mathcal{N}_{\omega},\mathcal{H}_{\omega}\} \, d\nu(\omega), \qquad \tau = \int_{\Omega}^{\oplus} \tau_{\omega} \, d\nu(\omega) \tag{3.3}$$

over a finite measure space $(\Omega, \mathcal{B}, \nu)$ that may be assumed to be complete. Then A, B are represented as

$$A = \int_{\Omega}^{\oplus} A_{\omega} \, d\nu(\omega), \qquad B = \int_{\Omega}^{\oplus} B_{\omega} \, d\nu(\omega) \tag{3.4}$$

with unique (a.e.) measurable fields $\omega \mapsto A_{\omega}, B_{\omega}$ of τ_{ω} -measurable positive operators affiliated with \mathcal{N}_{ω} . For each $\omega \in \Omega$, from the first step of the proof, there are unitaries $U, V \in \mathcal{N}_{\omega}$ such that

$$g(A_{\omega} + B_{\omega}) + \varepsilon I_{\omega} \ge Ug(A_{\omega})U^* + Vg(B_{\omega})V^*.$$
(3.5)

Now, define $F(\omega)$ to be the set of pairs (U, V) of unitaries in \mathcal{N}_{ω} satisfying (3.5), and prove that there are measurable fields $\omega \mapsto U_{\omega}$ and $\omega \mapsto V_{\omega}$ such that $(U_{\omega}, V_{\omega}) \in F(\omega)$ for all $\omega \in \Omega$. For this, as in [16], we may assume that $\omega \mapsto \mathcal{H}_{\omega}$ is a constant field \mathcal{H}_0 . Then $F(\cdot)$ is a multifunction whose values are non-empty closed subsets of a Polish space $B(\mathcal{H}_0)_1 \times B(\mathcal{H}_0)_1$, where $B(\mathcal{H}_0)_1$ is the closed unit ball of $B(\mathcal{H}_0)$ with the strong^{*} topology. By using [20, Theorem 6.1] and [16, Lemma 3.2(1)] as well as the fact that $\overline{\mathcal{N}}$ with the τ -measure topology is a Polish space, we infer that the graph

$$\{(\omega, U, V) \in \Omega \times B(\mathcal{H}_0)_1 \times B(\mathcal{H}_0)_1 : (U, V) \in F(\omega)\}$$

of $F(\cdot)$ belongs to $\mathcal{B} \otimes \mathcal{B}(B(\mathcal{H}_0)_1 \times B(\mathcal{H}_0)_1)$, where $\mathcal{B}(B(\mathcal{H}_0)_1 \times B(\mathcal{H}_0)_1)$ is the Borel σ -field of $B(\mathcal{H}_0)_1 \times B(\mathcal{H}_0)_1$. Hence, as in [16] the measurable selection theorem (e.g., [20]) yields measurable fields $\omega \mapsto U_\omega$ and $\omega \mapsto V_\omega$ as desired, so we obtain the claimed inequality with the unitaries $U = \int_{\Omega}^{\oplus} U_\omega \, d\nu(\omega)$ and $V = \int_{\Omega}^{\oplus} V_\omega \, d\nu(\omega)$ in \mathcal{N} .

Proof. (Theorem 3.2) The matrix case is in [2, Theorem 2.1]. In the type II₁ factor case, inequality (3.2) for a contraction $Z \in \mathcal{N}$ and an $T \in \overline{\mathcal{N}}^+$ is, in turn, reversed as

$$Z^*f(T)Z \le Wf(Z^*TZ)W^* + \varepsilon I$$

by [9, Lemma 10 (i)] and Lemma 3.3 similarly. (Here, note that although our assumption on f is slightly weaker than that in [9], the proof of [9, Lemma 10 (i)] can easily be modified to show that $f(Z^*TZ)$ spectrally dominates $Z^*f(T)Z$.) Hence the desired assertion follows in the factor case. Now, the proof for the non-factor case is the same as above.

An idea of the above proofs is to combine a unitary orbit technique with the measurable selection theorem. We end the section with another illustration of the idea, along the lines of [8, Proposition 2.11] for the matrix case.

Proposition 3.4. Let g(t) be a non-decreasing convex function on $[0, \infty)$. Then, for every $X, Y \in \overline{\mathcal{N}}$ and every $\varepsilon > 0$, there exist unitaries $U, V \in \mathcal{N}$ such that

$$g(|X+Y|) \le \frac{Ug(|X|+|Y|)U^* + Vg(|X^*|+|Y^*|)V^*}{2} + \varepsilon I.$$

Proof. For any $X \in \overline{\mathcal{N}}$, the decomposition $X = |X^*|^{1/2} V |X|^{1/2}$ with a unitary V shows that

$$\begin{bmatrix} |X^*| & X\\ X^* & |X| \end{bmatrix}$$

is a positive τ -measurable operator affiliated with $\mathbb{M}_2(\mathcal{N}) = \mathcal{N} \otimes \mathbb{M}_2$. Consequently,

$$\begin{bmatrix} |X^*| + |Y^*| & X + Y \\ X^* + Y^* & |X| + |Y| \end{bmatrix}$$

also belongs to $\overline{\mathbb{M}_2(\mathcal{N})}^+$. Let W be the unitary part in X + Y = W|X + Y|. Then

$$\begin{bmatrix} -W^* & I \end{bmatrix} \begin{bmatrix} |X^*| + |Y^*| & X + Y \\ X^* + Y^* & |X| + |Y| \end{bmatrix} \begin{bmatrix} -W \\ I \end{bmatrix}$$

is in $\overline{\mathcal{N}}^+$, so that

$$|X+Y| \le \frac{|X|+|Y|+W^*(|X^*|+|Y^*|)W}{2}.$$
(3.6)

As g(t) is non-decreasing and convex, in the factor case, (3.6) combined with [13, Proposition 4.6 (ii)] shows that g(|X+Y|) is spectrally dominated by $\{g(|X|+|Y|)+W^*g(|X^*|+|Y^*|)W\}/2$. Using Lemma 3.3 completes the proof of the proposition when \mathcal{N} is a factor. The general case follows by using the measurable selection method as in the previous proofs.

4 Extension of symmetric norms and anti-norms

The aim of this section is to show that a symmetric norm on \mathcal{N} and a symmetric antinorm on \mathcal{N}^+ can naturally be extended, respectively, to $\overline{\mathcal{N}}$ and to $\overline{\mathcal{N}}^+$. First, let $\|\cdot\|$ be a symmetric norm on \mathcal{N} . For each $X \in \overline{\mathcal{N}}$ and s > 0, the function $t \mapsto \beta_s(t) := \min\{s, t\}$ is used to define $|X| \wedge s := \beta_s(|X|)$. Since Lemma 2.1 implies that $\||X| \wedge s\|$ is increasing as $s \nearrow \infty$, a natural extension of $\|\cdot\|$ to $\overline{\mathcal{N}}$ is given as

$$||X|| := \lim_{s \nearrow \infty} ||X| \land s|| = \sup_{s > 0} ||X| \land s|| \in [0, \infty].$$

Proposition 4.1. The above extension of $\|\cdot\|$ becomes a symmetric norm on $\overline{\mathcal{N}}$ (permitting value ∞).

Proof. It is immediate to see that the extended $\|\cdot\|$ on $\overline{\mathcal{N}}$ satisfies $\|\alpha X\| = |\alpha| \|X\|$ for all $\alpha \in \mathbb{C}$ and $\|UXV\| = \|X\|$ for all unitaries $U, V \in \mathcal{N}$. For every $A, B \in \overline{\mathcal{N}}^+$, s > 0 and $\varepsilon > 0$, since β_s is concave on $[0, \infty)$, by Theorem 3.2 there are unitaries $U, V \in \mathcal{N}$ such that

 $(A+B) \wedge s \le U(A \wedge s)U^* + V(B \wedge s)V^* + \varepsilon I.$

By Lemma 2.1 this implies that

$$\|(A+B) \wedge s\| \le \|A \wedge s\| + \|B \wedge s\| + \varepsilon \|I\|.$$

Letting $\varepsilon \searrow 0$ and $s \nearrow \infty$ gives $||A+B|| \le ||A||+||B||$. Next we extend the monotonicity of $||\cdot||$ to $A \le B$ in $\overline{\mathcal{N}}^+$. For every s > 0 and $\varepsilon > 0$ there exists a unitary $U \in \mathcal{N}$ such that $U(A \land s)U^* \le B \land s + \varepsilon I$. Since $\lambda_t(A \land s) \le \lambda_t(B \land s)$ for all $t \in (0, \tau(I))$, this follows from Lemma 3.3 when \mathcal{N} is a factor. For the non-factor case, we can use the measurable selection method under the central direct decomposition as in the previous section. Therefore, $||A \land s|| \le ||B \land s|| + \varepsilon ||I||$, which implies that $||A|| \le ||B||$. Now, the subadditivity $||X + Y|| \le ||X|| + ||Y||$ in the whole $\overline{\mathcal{N}}$ follows by the triangle inequality

$$|X+Y| \le U|X|U^* + V|Y|V^*$$

for some unitaries $U, V \in \mathcal{N}$ ([13, Lemma 4.3], [21]), or else by use of (3.6).

Secondly, let $\|\cdot\|_{!}$ be a symmetric anti-norm on \mathcal{N}^+ . For every $A \in \overline{\mathcal{N}}^+$ and s > 0, since $\|A \wedge s\|_{!}$ increasing as $s \nearrow \infty$, we can extend $\|\cdot\|_{!}$ to $\overline{\mathcal{N}}^+$ as

$$\|A\|_{!} := \lim_{s \nearrow \infty} \|A \wedge s\|_{!} \in [0, \infty].$$

Proposition 4.2. The above extension of $\|\cdot\|_1$ to $\overline{\mathcal{N}}^+$ still satisfies the three conditions $(1)_1, (2)_1$ and $(3)_1$.

Proof. Since $(1)_!$ and $(2)_!$ are immediate by definition, we may prove $(3)_!$. Let $A, B \in \overline{\mathcal{N}}^+$, s > 0 and $\varepsilon > 0$ be arbitrary. We show that there exists a unitary $U \in \mathcal{N}$ such that

 $(A+B) \wedge 2s + \varepsilon I \ge U(A \wedge s + B \wedge s)U^*.$ (4.1)

When \mathcal{N} is a factor, this follows from Lemma 3.3 since we have, for $t \in (0, \tau(I))$,

$$\lambda_t((A+B) \wedge 2s) = \lambda_t(A+B) \wedge 2s \ge \lambda_t(A \wedge s + B \wedge s) \wedge 2s = \lambda_t(A \wedge s + B \wedge s)$$

due to $A \wedge s + B \wedge s \leq 2sI$. For the non-factor case, under the decompositions (3.3) and (3.4) we have $A \wedge s = \int_{\Omega}^{\oplus} A_{\omega} \wedge s \, d\nu(\omega)$ and similarly for $B \wedge s$ and $(A + B) \wedge 2s$. For each $\omega \in \Omega$, from the above factor case, there is a unitary $V \in \mathcal{N}_{\omega}$ such that

$$(A_{\omega} + B_{\omega}) \wedge 2s + \varepsilon I_{\omega} \ge V(A_{\omega} \wedge s + B_{\omega} \wedge s)V^*.$$

Now, define $F(\omega)$ to be the set of unitaries $V \in \mathcal{N}_{\omega}$ satisfying (4.1), and use the measurable selection method as before to obtain a measurable field $\omega \mapsto U_{\omega}$ such that $U_{\omega} \in F(\omega)$ for all $\omega \in \Omega$. Therefore, we have (4.1) with the unitary $U := \int_{\Omega}^{\oplus} U_{\omega} d\nu(\omega)$ in \mathcal{N} , so that $\|(A + B) \wedge 2s + \varepsilon I\|_{!} \geq \|A \wedge s\|_{!} + \|B \wedge s\|_{!}$. Letting $\varepsilon \searrow 0$ and $s \nearrow \infty$ gives $\|A + B\|_{!} \geq \|A\|_{!} + \|B\|_{!}$.

Remark 4.3. Let $\mathcal{N} = \bigoplus_{n=1}^{\infty} \mathcal{N}_n$ where $\mathcal{N}_n = \mathbb{M}_2$ for all n. We equip \mathcal{N} with the trace $\tau = \sum_{n\geq 1} 2^{-n} \tau_n$ where τ_n is the standard trace on \mathcal{N}_n . The function $t \mapsto \beta_n(t) = \min\{t, n\}$ is concave and there exist $A_n, B_n \in \mathbb{M}_2^+$ such that $\operatorname{Tr} \beta_n(A_n + B_n) < \operatorname{Tr} \beta_n(A_n) + \operatorname{Tr} \beta_n(B_n)$. Hence, there exist $A, B \in \overline{\mathcal{N}}^+$ such that, for all $n \in \mathbb{N}, \tau((A + B) \wedge n)) < \tau(A \wedge n) + \tau(A \wedge n)$. Such a phenomenon for τ (regarded as an anti-norm) explains why the proof of the superadditivity part (3)! in Proposition 4.2 is non-trivial. Derived anti-norms extended to $\overline{\mathcal{N}}^+$ have the following simple properties; (a) and (b) may be used to check that a symmetric anti-norm is not a derived one.

Proposition 4.4. Let $\|\cdot\|_!$ be a derived anti-norm on \mathcal{N}^+ , which is derived from a symmetric norm $\|\cdot\|$ on \mathcal{N} and a p > 0. The following hold for the extensions of $\|\cdot\|_!$ to $\overline{\mathcal{N}}^+$ and $\|\cdot\|$ to $\overline{\mathcal{N}}$.

(a)
$$||A||_{!} < \infty$$
 for all $A \in \overline{\mathcal{N}}^{+}$.

- (b) If $A \in \overline{\mathcal{N}}^+$ is singular, i.e., the kernel of A is non-trivial, then $||A||_{!} = 0$.
- (c) If $A \in \overline{\mathcal{N}}^+$ is nonsingular, then $||A||_! = ||A^{-p}||^{-1/p}$ (with convention $\infty^{-1/p} = 0$).

Proof. The proof of (a) is easy and left to the reader.

(b) Assume that ker $A \neq \{0\}$, and let P be the projection onto ker A. For each s > 0 and $\varepsilon > 0$ we have $(A \wedge s + \varepsilon I)^{-p} \ge \varepsilon^{-p}P$ and so $||(A \wedge s + \varepsilon I)^{-p}|| \ge \varepsilon^{-p}||P||$. By definition (2.2),

$$\|A \wedge s\|_{!} = \lim_{\varepsilon \searrow 0} \|(A \wedge s + \varepsilon I)^{-p}\|^{-1/p} \le \lim_{\varepsilon \searrow 0} \varepsilon \|P\|^{-1/p} = 0.$$

Therefore, $||A||_{!} = 0.$

(c) First we extend (2.2) to a nonsingular $A \in \overline{\mathcal{N}}^+$. Let $A \in \overline{\mathcal{N}}^+$ with ker $A = \{0\}$; hence $A^{-p} \in \overline{\mathcal{N}}^+$. For every s > 0 and $\varepsilon > 0$, set

$$\phi_s(\varepsilon) := \sup_{x \ge 0} \left\{ x^{-p} \wedge s - (x + \varepsilon)^{-p} \right\}.$$

Then it is clear that $\phi_s(\varepsilon) > 0$ and $\lim_{\varepsilon \searrow 0} \phi_s(\varepsilon) = 0$ for each s > 0. Since $||(A + \varepsilon I)^{-p}||$ increases as $\varepsilon \searrow 0$ by Lemma 2.1 and

$$A^{-p} \wedge s \le (A + \varepsilon I)^{-p} + \phi_s(\varepsilon)I,$$

we have $||A^{-p} \wedge s|| \leq \lim_{\varepsilon \searrow 0} ||(A + \varepsilon I)^{-p}||$. Hence $||A^{-p}|| \leq \lim_{\varepsilon \searrow 0} ||(A + \varepsilon I)^{-p}||$. On the other hand, since $(A + \varepsilon I)^{-p} \leq A^{-p} \wedge \varepsilon^{-p}$, we have $||(A + \varepsilon I)^{-p}|| \leq ||A^{-p}||$ for every $\varepsilon > 0$. Therefore,

$$||A^{-p}||^{-1/p} = \lim_{\varepsilon \searrow 0} ||(A + \varepsilon I)^{-p}||^{-1/p}.$$
(4.2)

Now, looking at the function $x \ge 0 \mapsto (x \land s + \varepsilon)^{-p}$, one can easily see that

$$(A + \varepsilon I)^{-p} \le (A \wedge s + \varepsilon I)^{-p} \le (A + \varepsilon I)^{-p} + s^{-p}I$$

and hence

$$\|(A + \varepsilon I)^{-p}\|^{-1/p} \ge \|(A \wedge s + \varepsilon I)^{-p}\|^{-1/p} \ge (\|(A + \varepsilon I)^{-p}\| + s^{-p}\|I\|)^{-1/p}.$$

Thanks to (4.2), letting $\varepsilon \searrow 0$ gives

$$||A^{-p}||^{-1/p} \ge ||A \land s||_! \ge (||A^{-p}|| + s^{-p}||I||)^{-1/p}$$

so that $||A||_! = ||A^{-p}||^{-1/p}$ follows.

5 Superadditivity for convex functions

The aim of this section is to prove the next superadditivity theorem for a symmetric anti-norm involving a convex function g(t) on $[0, \infty)$. Note that a non-negative convex function g(t) on $[0, \infty)$ is superadditive if and only if g(0) = 0. Also, note that the assumption on g(t) in the theorem is best possible; indeed, the assumption is necessary even for the classical Rotfel'd trace inequality for matrices.

Theorem 5.1. Let $A, B \in \overline{\mathcal{N}}^+$ and let g(t) be a non-negative convex function on $[0, \infty)$ with g(0) = 0. Then, for all symmetric anti-norms on \mathcal{N}^+ ,

$$||g(A+B)||_! \ge ||g(A)||_! + ||g(B)||_!.$$

Here, and in the whole sequel, we consider that symmetric anti-norms on \mathcal{N}^+ are automatically defined on the full cone $\overline{\mathcal{N}}^+$ as in Proposition 4.2, and similarly symmetric norms on \mathcal{N} are defined on the whole $\overline{\mathcal{N}}$ as in Proposition 4.1.

Theorem 5.1 claims a numerical inequality; however, its proof relies on an operator inequality presented in Section 3. Indeed, when $A, B \in \mathcal{N}^+$, it is a straightforward consequence of Theorem 3.1. In the unbounded case, we cannot argue by letting $\varepsilon \searrow 0$ in Theorem 3.1 because (4)! may not hold in $\overline{\mathcal{N}}^+$; for instance, when $\|\cdot\|_!$ is the Fuglede-Kadison determinant (see the end of Section 6), there is an $A \in \overline{\mathcal{N}}^+$ which has the non-trivial kernel (hence $\|A\|_! = 0$) but satisfies $\|A + \varepsilon I\|_! = \infty$ for all $\varepsilon > 0$.

Proof. Let $A, B \in \overline{\mathcal{N}}^+$ and s > 0 be arbitrary. Since g(t) is continuous and nondecreasing on $[0, \infty)$, we have thanks to [13, Lemma 2.5 (iv)]

$$\lambda_t(g((A+B) \land 2s)) = g(\lambda_t(A+B) \land 2s)$$

$$\geq g(\lambda_t(A \land s+B \land s))$$

$$= \lambda_t(g(A \land s+B \land s)), \qquad t \in (0, \tau(I)).$$

Given $\varepsilon > 0$, when \mathcal{N} is a finite factor, Lemma 3.3 then entails

$$g((A+B) \wedge 2s) + \varepsilon I \ge Wg(A \wedge s + B \wedge s)W^*$$

for some unitary $W \in \mathcal{N}$. This inequality can be extended to the non-factor case, by using the measurable selection method under the central direct decomposition as in Section 3, while full details are left to the reader. Hence, Theorem 3.1 applied to $A \wedge s$, $B \wedge s$ in place of A, B shows that

$$g((A+B) \wedge 2s) + 2\varepsilon I \ge Ug(A \wedge s)U^* + Vg(B \wedge s)V^*$$

for some unitaries $U, V \in \mathcal{N}$. Therefore,

$$||g((A+B) \wedge 2s) + 2\varepsilon I||_! \ge ||g(A \wedge s)||_! + ||g(B \wedge s)||_!$$

so that $\|g((A+B) \wedge 2s)\|_{!} \geq \|g(A \wedge s)\|_{!} + \|g(B \wedge s)\|_{!}$. Since a simple estimation gives

$$\|g(A)\|_{!} = \lim_{s \nearrow \infty} \|g(A) \wedge s\|_{!} = \lim_{s \nearrow \infty} \|g(A \wedge s)\|_{!},$$

the claimed inequality follows.

In the rest of the section we collect a few special illustrations of Theorem 5.1.

Corollary 5.2. Let $A, B \in \overline{\mathcal{N}}^+$ be nonsingular and p_1, \ldots, p_m be positive scalars such that $\sum_{i=1}^m p_i \ge 1$. Then, for all symmetric norms on \mathcal{N} ,

$$\prod_{i=1}^{m} \left\| (A+B)^{-p_i} \right\|^{-1} \ge \prod_{i=1}^{m} \left\| A^{-p_i} \right\|^{-1} + \prod_{i=1}^{m} \left\| B^{-p_i} \right\|^{-1}$$

Proof. Let $g_i(t)$ be strictly increasing convex functions on $[0, \infty)$ with $g_i(t) = 0$, and let $q_i > 0, 1 \le i \le m$. Theorem 5.1 applied to the derived anti-norms $A \mapsto ||A^{-q_i}||^{-1/q_i}$ yields, in view of Proposition 4.4 (c),

$$\left\|g_{i}^{-q_{i}}(A+B)\right\|^{-1/q_{i}} \ge \left\|g_{i}^{-q_{i}}(A)\right\|^{-1/q_{i}} + \left\|g_{i}^{-q_{i}}(B)\right\|^{-1/q_{i}}, \quad 1 \le i \le m$$

Now, assume that $\sum_{i=1}^{m} q_i = 1$. By the elementary inequality from the concavity of the weighted geometric mean,

$$\prod_{i=1}^{m} (a_i + b_i)^{q_i} \ge \prod_{i=1}^{m} a_i^{q_i} + \prod_{i=1}^{m} b_i^{q_i}$$

with $a_i = \|g_i^{-q_i}(A)\|^{-1/q_i}$ and $b_i = \|g_i^{-q_i}(B)\|^{-1/q_i}$, we have

$$\prod_{i=1}^{m} \left\| g_i^{-q_i}(A+B) \right\|^{-1} \ge \prod_{i=1}^{m} \left\| g_i^{-q_i}(A) \right\|^{-1} + \prod_{i=1}^{m} \left\| g_i^{-q_i}(B) \right\|^{-1}.$$
(5.1)

Let $p = \sum_{i=1}^{m} p_i$. Take in (5.1) $g_i(t) = t^p$ and $q_i = p_i/p$, $1 \le i \le m$, to obtain the required estimate.

Corollary 5.3. Let $A, B \in \overline{\mathcal{N}}^+$ be nonsingular. Then, for all symmetric norms on \mathcal{N} , and m = 1, 2, ...,

$$\left\|\sum_{k=1}^{m} (A+B)^{-k}\right\|^{-1} \ge \left\|\sum_{k=1}^{m} A^{-k}\right\|^{-1} + \left\|\sum_{k=1}^{m} B^{-k}\right\|^{-1}$$

Proof. Let $m \ge 2$ and let

$$g(t) := \frac{t^m}{1 + t + \dots + t^{m-1}} = \frac{t^{m+1} - t^m}{t^m - 1}.$$

Then g(0) = 0 and for t > 0,

$$g''(t) = \frac{mt^{m-2}\left\{(m-1)t^{m+1} - (m+1)t^m + (m+1)t - (m-1)\right\}}{(t^m - 1)^3}$$

For
$$\phi(t) := (m-1)t^{m+1} - (m+1)t^m + (m+1)t - (m-1)$$
 compute
 $\phi'(t) = (m+1)\{(m-1)t^m - mt^{m-1} + 1\},\$
 $\phi''(t) = (m+1)m(m-1)t^{m-2}(t-1).$

Since $\phi(1) = \phi'(1) = \phi''(1) = 0$, we see that $\phi(t) \leq 0$ for $0 \leq t \leq 1$ and $\phi(t) \geq 0$ for $t \geq 1$. Hence $g''(t) \geq 0$ for all t > 0, so g(t) is convex on $(0, \infty)$. Note that $g(t) = \left(\sum_{k=1}^{m} t^{-k}\right)^{-1}$ for t > 0. So, applying Theorem 5.1 to this function g(t) and the derived anti-norm $A \mapsto ||A^{-1}||^{-1}$ proves the corollary.

The next corollary involves an anti-norm specific to the matrix algebra \mathbb{M}_n . Here $\wedge^m A$ denotes the *m*th antisymmetric tensor power of a matrix A (see [4]).

Corollary 5.4. Let $A, B \in \mathbb{M}_n^+$ be nonsingular. Let $g : [0, \infty) \to [0, \infty)$ be a strictly increasing convex function with g(0) = 0. Then, for all $0 < q \le 1$ and all m = 1, 2, ...,

$$\left\{\frac{\operatorname{Tr}\,\wedge^m g^q(A+B)}{\operatorname{Tr}\,\wedge^{m-1} g^q(A+B)}\right\}^{1/q} \ge \left\{\frac{\operatorname{Tr}\,\wedge^m g^q(A)}{\operatorname{Tr}\,\wedge^{m-1} g^q(A)}\right\}^{1/q} + \left\{\frac{\operatorname{Tr}\,\wedge^m g^q(B)}{\operatorname{Tr}\,\wedge^{m-1} g^q(B)}\right\}^{1/q}.$$

Note that letting m = q = 1 we recapture the Rotfel'd trace inequality.

Proof. By a theorem of Marcus and Lopes [24] (also [25, p. 116]), the functional on positive nonsingular matrices

$$A \mapsto \frac{\operatorname{Tr} \wedge^m A}{\operatorname{Tr} \wedge^{m-1} A}$$

is superadditive. This can be extended as an anti-norm on the whole of \mathbb{M}_n^+ by using condition (4). The corollary then follows from Lemma 2.8 combined with Theorem 5.1.

6 Full symmetry and majorization

In this section we consider a stronger symmetry property of norms and anti-norms in connection with majorization relations. We will focus on the case of diffuse algebras. Indeed, the case of \mathbb{M}_n is simpler as well as classical. Meanwhile, the setting of a general finite von Neumann algebra \mathcal{N} is inappropriate to apply the majorization technique. This issue may be justified by the fact [14, Theorem 3.27] that (\mathcal{N}, τ) with $\tau(I) = 1$ satisfies the *weak Dixmier property* (i.e., $\tau(A)$ is in the $\|\cdot\|_{\infty}$ -closure of the convex hull of $\{B \in \mathcal{N}^+ : \lambda(B) = \lambda(A)\}$ for every $A \in \mathcal{N}^+$) if and only if either (\mathcal{N}, τ) is a subalgebra of $(\mathbb{M}_n, n^{-1}\mathrm{Tr})$ containing all diagonal matrices, or \mathcal{N} is diffuse.

Thus, in Sections 6 and 7, we shall always write \mathcal{M} (differently from \mathcal{N}) to denote a diffuse finite von Neumann algebra with a faithful normal trace τ such that $\tau(I) = 1$.

Definition 6.1. A symmetric norm $\|\cdot\|$ on \mathcal{M} is said to be *fully symmetric* (or *re-arrangement invariant*) if $\lambda(A) = \lambda(B)$ implies $\|A\| = \|B\|$ for $A, B \in \mathcal{M}^+$. Also, a symmetric anti-norm $\|\cdot\|_!$ on \mathcal{M}^+ is said to be *fully symmetric* if the same property holds for $\|\cdot\|_!$.

The next proposition says that the symmetry and the full symmetry properties are equivalent when \mathcal{M} is a II₁ factor (this is also true and well-known for \mathbb{M}_n).

Proposition 6.2. If \mathcal{M} is a factor, then any symmetric norm and any symmetric anti-norm are fully symmetric.

Proof. Let $A, B \in \mathcal{M}^+$ and assume that $\lambda(A) = \lambda(B)$. For a symmetric norm $\|\cdot\|$, by Lemmas 2.1 and 3.3 we have $\|A\| \leq \|B\| + \varepsilon \|I\|$ for every $\varepsilon > 0$, so $\|A\| \leq \|B\|$ and the reverse inequality is similar. The proof is similar for a symmetric anti-norm by using condition (4)₁.

Recall some notions of majorization relevant to our discussions below. For $A, B \in \overline{\mathcal{M}}^+$, the submajorization $A \prec_w B$ is defined as $\lambda(A) \prec_w \lambda(B)$, i.e.,

$$\int_0^t \lambda_s(A) \, ds \le \int_0^t \lambda_s(B) \, ds, \qquad t \in (0,1),$$

and the majorization $A \prec B$ means that $A \prec_w B$ and $\tau(A) = \tau(B) < \infty$. The supermajorization $A \prec^w B$ is defined as

$$\int_t^1 \lambda_s(A) \, ds \ge \int_t^1 \lambda_s(B) \, ds, \qquad t \in (0,1).$$

The log-supermajorization $A \prec^{w(\log)} B$ is defined as

$$\int_t^1 \log \lambda_s(A) \, ds \ge \int_t^1 \log \lambda_s(B) \, ds, \qquad t \in (0,1).$$

These definitions make sense since the integrals always exist permitting $\pm \infty$. (The finiteness assumption $\tau(I) < \infty$ is essential to introduce the supermajorization and the log-supermajorization.)

Example 6.3. (1) For each $t \in (0, 1]$ the functional

$$\|X\|_{(t)} := \int_0^t \mu_s(X) \, ds, \qquad X \in \mathcal{M}$$

is a fully symmetric norm on \mathcal{M} , that is the continuous version of the Ky Fan k-norm for matrices. The triangle inequality of $\|\cdot\|_{(t)}$ is a consequence of the submajorization $\mu(X+Y) \prec_w \mu(X) + \mu(Y)$ for $X, Y \in \overline{\mathcal{M}}$ (see [17]).

(2) For each $t \in [0, 1)$ the functional

$$||A||_{\{t\}} := \int_{1-t}^{1} \lambda_s(A) \, ds, \qquad A \in \mathcal{M}^+$$

is a fully symmetric anti-norm on \mathcal{M}^+ . The superadditivity of $\|\cdot\|_{\{t\}}$ is a consequence of the majorization $\lambda(A+B) \prec \lambda(A) + \lambda(B)$ for $A, B \in \mathcal{M}^+$ (see [17]). This anti-norm is not a derived anti-norm. (3) For each $t \in [0, 1)$ and p > 0, the derived anti-norm from the above $\|\cdot\|_{(t)}$ and p is written as

$$||A||_{!} := \lim_{\varepsilon \searrow 0} ||(A + \varepsilon I)^{-p}||_{(t)}^{-1/p} = \left(\int_{1-t}^{1} \lambda_{s}(A)^{-p} \, ds\right)^{-1/p}, \qquad A \in \mathcal{M}^{+}$$
(6.1)

with the usual convention $0^{-p} = \infty$ and $\infty^{-1/p} = 0$. Obviously, this derived anti-norm is fully symmetric. One can easily find a sequence $A_n, A \in \mathcal{M}^+$ such that $||A_n - A||_{\infty} \to 0$ and $||A_n||_1 = 0$ for all n, but $||A||_1 > 0$. Therefore, this $|| \cdot ||_1$ is not $|| \cdot ||_{\infty}$ -continuous on \mathcal{M}^+ . On the other hand, the derived anti-norm from $|| \cdot ||_{\infty}$ (and any p > 0) is $\lambda_1(A)$, which is $|| \cdot ||_{\infty}$ -continuous on \mathcal{M}^+ . Thus, the continuity behavior with respect to the operator norm in the diffuse case is subtler than the matrix case.

Since \mathcal{M} is diffuse, we can choose a family $\{F_t\}_{0 \le t \le 1}$ of projections as in the proof of Lemma 3.3. In our discussions below we will use such a family $\{F_t\}$ without mentioning explicitly.

6.1 Fully symmetric norms

The following properties of fully symmetric norms were more or less discussed in study of non-commutative Banach function spaces (e.g., [11, 12, 28]), usually as working assumptions rather than results. The book [23] contains a nice discussion on this topic.

Proposition 6.4. Let $\|\cdot\|$ be a fully symmetric norm on \mathcal{M} .

- (a) For every $A, B \in \overline{\mathcal{M}}^+$, $A \prec_w B$ implies $||A|| \leq ||B||$.
- (b) If $A, A_n \in \overline{\mathcal{M}}^+$ and $A_n \nearrow A$ in the τ -measure topology (or more weakly $\lambda_t(A_n) \nearrow \lambda_t(A)$ for a.e. $t \in (0,1)$), then $||A_n|| \nearrow ||A||$.
- (c) $||X||_1 ||I|| \leq ||X|| \leq ||X||_\infty ||I||$ for all $X \in \overline{\mathcal{M}}$, where $||X||_1 := \tau(|X|)$ and $||X||_\infty := \lim_{t \searrow 0} \lambda_t(|X|)$, the operator norm.

Proof. (a) (for bounded operators) Assume that $A, B \in \mathcal{M}^+$ and $A \prec_w B$. Since $f := \lambda(A) \prec_w g := \lambda(B)$ as functions in $L^{\infty}(0,1)^+$, by [10, Theorem 1.1] there is an $h \in L^{\infty}(0,1)^+$ such that $f \leq h \prec g$. Let Γ denote the set of bijective Borel transformations on (0,1) preserving the Lebesgue measure. In the same way as in the proof of [19, Theorem 2.1, Lemma 2.2] we can see that the decreasing rearrangement h^* of h is in the $\|\cdot\|_{\infty}$ -closure of the convex hull of $\{\gamma f : \gamma \in \Gamma\}$, where $(\gamma f)(t) := f(\gamma^{-1}t)$. Thus, for every $\varepsilon > 0$ there are $\gamma_i \in \Gamma$ and $\lambda_i > 0, 1 \leq i \leq k$, such that $\sum_{i=1}^k \lambda_i = 1$ and $h^* \leq \sum_{i=1}^k \lambda_i \gamma_i g + \varepsilon 1$. Let $C := \int_0^1 h^*(t) dF_t$ and $B_i := \int_0^1 (\gamma_i g)(t) dF_t$. Since $\lambda(B_i) = \lambda(B)$, the monotonicity and the full symmetry of $\|\cdot\|$ yield

$$||A|| \le ||C|| \le \sum_{i=1}^{k} \lambda_i ||B_i|| + \varepsilon ||I|| = ||B|| + \varepsilon ||I||.$$

Hence $||A|| \leq ||B||$.

(b) First we prove that if $A, A_n \in \mathcal{M}^+$ and $\lambda_t(A_n) \nearrow \lambda_t(A)$ a.e., then $||A_n|| \nearrow ||A||$. By (a), $||A_n||$ is increasing in n. For every $\varepsilon > 0$, since $\lambda_0(A_n) = ||A_n||_{\infty} \nearrow \lambda_0(A) = ||A||_{\infty}$ and

$$\lim_{t \searrow 0} \frac{1}{t} \int_0^t (\lambda_s(A) - \lambda_s(A_n)) \, ds = \lambda_0(A) - \lambda_0(A_n),$$

one can choose $n_0 \in \mathbb{N}$ and $\delta > 0$ so that $t^{-1} \int_0^t (\lambda_s(A) - \lambda_s(A_{n_0})) ds < \varepsilon$ for all $t \in (0, \delta)$, and hence

$$\frac{1}{t} \int_0^t (\lambda_s(A) - \lambda_s(A_n)) \, ds < \varepsilon, \qquad t \in (0, \delta), \ n \ge n_0.$$

Furthermore, for any $t \in [\delta, 1)$ one has

$$\frac{1}{t} \int_0^t (\lambda_s(A) - \lambda_s(A_n)) \, ds \le \frac{1}{\delta} \int_0^1 (\lambda_s(A) - \lambda_s(A_n)) \, ds \searrow 0 \quad \text{as } n \to \infty$$

by the dominated convergence theorem. Hence there exists an $n_1 \in \mathbb{N}$ such that

$$\frac{1}{t} \int_0^t (\lambda_s(A) - \lambda_s(A_n)) \, ds < \varepsilon, \qquad t \in [\delta, 1), \ n \ge n_1.$$

If $n \ge \max\{n_0, n_1\}$, then the above estimates imply that $A \prec_w A_n + \varepsilon I$ so that $||A|| \le ||A_n|| + \varepsilon ||I||$ by (a). Thus $||A_n|| \nearrow ||A||$.

Next assume that $A, A_n \in \overline{\mathcal{M}}^+$ and $\lambda_t(A_n) \nearrow \lambda_t(A)$ a.e. For every s > 0, since $\lambda_t(A_n \land s) \nearrow \lambda_t(A \land s)$ a.e., we have $||A_n \land s|| \nearrow ||A \land s||$ from the first step. Hence $||A_n|| = \sup_{s>0} ||A_n \land s||$ is increasing in n and

$$||A|| = \sup_{s>0} ||A \wedge s|| = \sup_{s>0, n \in \mathbb{N}} ||A_n \wedge s|| = \sup_{n \in \mathbb{N}} ||A_n||.$$

Therefore, $||A_n|| \nearrow ||A||$.

(c) For $X \in \mathcal{M}$, $||X|| \leq ||X||_{\infty} ||I||$ was given in (2.1). Since $\tau(|X|)I \prec |X|$, $||X||_1 ||I|| \leq ||X||$ by (a). These can easily extend to all $X \in \overline{\mathcal{M}}$ by (b).

(a) (for unbounded operators) Let $A, B \in \overline{\mathcal{M}}^+$ and assume $A \prec_w B$. We may assume that $||B|| < \infty$ and so $||A||_1 \le ||B||_1 < \infty$ by (c). Fix $0 < \rho < 1$. Then, for each $n \in \mathbb{N}$, since $\int_0^t \lambda_s(B) \wedge m \, ds \nearrow \int_0^t \lambda_s(B) \, ds$ as $m \to \infty$ uniformly in $t \in (0, 1)$, one can choose an $m \in \mathbb{N}$ such that $\rho A \wedge n \prec_w B \wedge m$ and hence $\|\rho A \wedge n\| \le \|B\|$ by (a). Hence $\|A\| \le \|B\|$ follows by letting $n \to \infty$ and then $\rho \nearrow 1$.

In view of Example 6.3(1) we have

Corollary 6.5. Let $X, Y \in \overline{\mathcal{M}}$. Then $|X| \prec_w |Y|$ (i.e., $\mu(X) \prec_w \mu(Y)$) if and only if $||X|| \leq ||Y||$ for all fully symmetric norms on \mathcal{M} (extended to $\overline{\mathcal{M}}$).

6.2 Fully symmetric anti-norms

Fully symmetric anti-norms have the following properties. It would be worthwhile to consider these properties in parallel to those in Proposition 6.4.

Proposition 6.6. Let $\|\cdot\|_{!}$ be a fully symmetric anti-norm on \mathcal{M}^+ .

- (a) For every $A, B \in \overline{\mathcal{M}}^+$, $A \prec^w B$ implies $||A||_! \ge ||B||_!$.
- (b) If $A, A_n \in \mathcal{M}^+$ and $A_n \searrow A$ (or more weakly $\lambda_t(A_n) \searrow \lambda_t(A)$ for a.e. $t \in (0,1)$), then $||A_n||_! \searrow ||A||_!$.
- (c) $\lambda_1(A) \|I\|_! \leq \|A\|_! \leq \tau(A) \|I\|_!$ for all $A \in \overline{\mathcal{M}}$, where $\lambda_1(A) := \lim_{t \nearrow 1} \lambda_t(A)$.

Proof. (a) Since $A \prec^w B$ implies $A \wedge s \prec^w B \wedge s$ for any s > 0 (seen as in the discrete case [25, p. 167]), we may consider the case $A, B \in \mathcal{M}^+$. Since $f := \lambda(A) \prec^w g := \lambda(B)$ as functions in $L^{\infty}(0, 1)^+$, there is an $h \in L^{\infty}(0, 1)^+$ such that $f \geq h \prec g$. The remaining proof being similar to that of Proposition 6.4 (a), we omit the details.

(b) Assume that $A, A_n \in \mathcal{M}$ and $\lambda_t(A_n) \searrow \lambda_t(A)$ for a.e. $t \in (0, 1)$. By (a), $||A_n||_1$ is decreasing in n. For every $\varepsilon > 0$, since $\lambda_1(A_n) \searrow \lambda_1(A)$ and

$$\lim_{t \searrow 0} \frac{1}{t} \int_{1-t}^{1} (\lambda_s(A_n) - \lambda_s(A)) \, ds = \lambda_1(A_n) - \lambda_1(A),$$

one can show that $A + \varepsilon I \prec^w A_n$ for all sufficiently large n, as in the proof of Proposition 6.4 (b) by replacing $t^{-1} \int_0^t \text{ with } t^{-1} \int_{1-t}^1$. By (a) this implies that $||A + \varepsilon I||_! \ge \lim_n ||A_n||_!$. Letting $\varepsilon \searrow 0$ gives $||A||_! \ge \lim_n ||A_n||_!$ and so $||A_n||_! \searrow ||A||_!$.

(c) For $A \in \overline{\mathcal{M}}^+$, $\lambda_1(A)I \leq A$ implies $\lambda_1(A)||I||_! \leq ||A||_!$. Assuming $\tau(A) < \infty$, we have $\tau(A)||I||_! \geq ||A||_!$ from $\tau(A)I \prec A$.

In view of Example 6.3(2) we have

Corollary 6.7. Let $A, B \in \overline{\mathcal{M}}^+$. Then $A \prec^w B$ if and only if $||A||_! \leq ||B||_!$ for all fully symmetric anti-norms on \mathcal{M}^+ (extended to $\overline{\mathcal{M}}^+$).

Remark 6.8. Proposition 6.4 (b) means that a fully symmetric norm extended to $\overline{\mathcal{M}}$ satisfies the *Fatou property* (see [12]). Proposition 6.6 (b) is considered as the "anti-Fatou property". Even though (4)_! may not hold in $\overline{\mathcal{M}}^+$ as noted in Section 5, it is not known whether the anti-Fatou property holds for $A, A_n \in \overline{\mathcal{M}}^+$ when $||A_n||_! < \infty$ and $A_n \searrow A$ in the τ -measure topology. For fully symmetric derived anti-norms, this property will be shown in the next subsection.

6.3 Fully symmetric derived anti-norms

In the rest of the section we will consider fully symmetric derived anti-norms.

Lemma 6.9. Let $\|\cdot\|_{!}$ be a derived anti-norm on \mathcal{M}^+ , which is derived from a symmetric norm $\|\cdot\|$ on \mathcal{M} and a p > 0. Then $\|\cdot\|_{!}$ is fully symmetric if and only if so is $\|\cdot\|$.

Proof. From (2.1), a symmetric norm on \mathcal{M} is fully symmetric if the condition in Definition 6.1 holds for invertible $A, B \in \mathcal{M}^+$. From condition (4)_!, the same is true for a symmetric anti-norm. Hence the assertion follows from the relations $||A||_! = ||A^{-p}||^{-1/p}$ and $||A|| = ||A^{-1/p}||_!^{-p}$ for invertible $A \in \mathcal{M}^+$.

Properties (a) and (b) of Proposition 6.6 are strengthened for fully symmetric derived anti-norms as follows. It is worth noting that $A \prec^{w(\log)} B$ is weaker than $A \prec^{w} B$ for $A, B \in \overline{\mathcal{M}}^+$.

Proposition 6.10. Let $\|\cdot\|_{!}$ be a fully symmetric derived anti-norm on \mathcal{M}^+ .

- (a) For every $A, B \in \overline{\mathcal{M}}^+$, $A \prec^{w(\log)} B$ implies $||A||_! \ge ||B||_!$.
- (b) If $A, A_n \in \overline{\mathcal{M}}^+$ and $A_n \searrow A$ in the τ -measure topology (or more weakly $\lambda_t(A_n) \searrow \lambda_t(A)$ for a.e. $t \in (0,1)$), then $||A_n||_! \searrow ||A||_!$. In particular, (4)! holds in $\overline{\mathcal{M}}^+$.

Proof. From Lemma 6.9, let $\|\cdot\|_{!}$ be derived from a fully symmetric norm $\|\cdot\|$ and a p > 0.

(a) Assume that $A \prec^{w(\log)} B$. Since this implies that $A \wedge s \prec^{w(\log)} B \wedge s$ for all s > 0 (similarly to the assertion for \prec^w in the proof of Proposition 6.6, by considering the function $\log(e^x \wedge s)$), it is enough to assume that $A, B \in \mathcal{M}^+$. Furthermore, by replacing A with $A + \varepsilon I$ for any $\varepsilon > 0$, A may be assumed invertible. First, assume that $\int_0^1 \log \lambda_s(B) \, ds = -\infty$, and we prove that $\|B\|_! = 0$ for every derived anti-norm $\|\cdot\|_!$. By Proposition 6.4 (c) it suffices to prove this for the derived anti-norm from $\|\cdot\|_1$ and any p > 0. So we may show that $\lim_{\varepsilon \searrow 0} \|(B + \varepsilon I)^{-p}\|_1 = \infty$. Since $\int_0^1 \log \lambda_s(B) \, ds = -\infty$ implies that $\lambda_1(B) = 0$, we have $-\log \lambda_s(B) \leq \lambda_s(B)^{-p}$ for all s sufficiently near 1. Hence by (6.1) for t = 1,

$$\lim_{\varepsilon \searrow 0} \|(B + \varepsilon I)^{-p}\|_1 = \int_0^1 \lambda_s(B)^{-p} \, ds = \infty.$$

Next assume that B as well as A is invertible. Then $A \prec^{w(\log)} B$ means $\log \lambda(A) \prec^{w} \log \lambda(B)$. For every p > 0, since $\log \lambda(A^{-p}) \prec_{w} \log \lambda(B^{-p})$, we have $A^{-p} \prec_{w} B^{-p}$ by [17, Proposition 1.2]. Hence by Proposition 6.4 (a), $||A^{-p}|| \leq ||B^{-p}||$ for any fully symmetric norm $|| \cdot ||$, implying the assertion.

Finally, assume that B is not invertible but $\int_0^1 \log \lambda_s(B) \, ds > -\infty$, so $\lambda_s(B) > 0$ for all $s \in (0, 1)$ and $\log \lambda(B)$ is integrable on (0, 1). One can fix an $r_0 \in (0, 1)$ such that $\lambda_{r_0}(B) \leq \lambda_1(A)$ (since A is assumed invertible while B is not). For every $\delta > 0$ there exists an $r \in (r_0, 1)$ such that

$$\int_{r}^{1} (\log \lambda_{r}(B) - \log \lambda_{s}(B)) \, ds \le \delta(1 - r_{0}),$$

so we define

$$\hat{B} := \int_0^r \lambda_s(B) \, dF_s + \int_r^1 \lambda_r(B) \, dF_s,$$

which is invertible. If $t \in (r_0, 1)$, then

$$\int_{t}^{1} (\log \lambda_{s}(A) + \delta) \, ds \ge \int_{t}^{1} \log \lambda_{s}(A) \, ds \ge \int_{t}^{1} \log \lambda_{r_{0}}(B) \, ds \ge \int_{t}^{1} \log \lambda_{s}(\hat{B}) \, ds.$$

If $t \in (0, r_0]$, then

$$\begin{split} \int_{t}^{1} (\log \lambda_{s}(A) + \delta) \, ds &\geq \int_{t}^{1} \log \lambda_{s}(B) \, ds + \delta(1 - r_{0}) \\ &= \int_{t}^{1} \log \lambda_{s}(\hat{B}) \, ds - \int_{r}^{1} (\log \lambda_{r}(B) - \log \lambda_{s}(B)) \, ds + \delta(1 - r_{0}) \\ &\geq \int_{t}^{1} \log \lambda_{s}(\hat{B}) \, ds. \end{split}$$

The above estimates imply that $e^{\delta}A \prec^{w(\log)} \hat{B}$. Since \hat{B} is invertible, $||(e^{\delta}A)^{-p}|| \leq ||\hat{B}^{-p}||$ as in the previous case. Therefore,

$$e^{\delta} \|A^{-p}\|^{-1/p} \ge \|\hat{B}^{-p}\|^{-1/p} = \|\hat{B}\|_{!} \ge \|B\|_{!}$$

by Proposition 6.6 (a) since $\lambda(\hat{B}) \geq \lambda(B)$. Letting $\delta \searrow 0$ gives $||A||_! \geq ||B||_!$.

(b) If ker $A \neq \{0\}$, then there is a $\delta \in (0,1)$ such that $\lambda_{1-\delta}(A) = 0$. Since $\lambda_{1-\delta/2}(A_n) \searrow 0$, for every $\varepsilon > 0$ there exists an $n_0 \in \mathbb{N}$ such that $\lambda_{1-\delta/2}(A_n) < \varepsilon$ for all $n \ge n_0$. For each $n \ge n_0$, letting $P_n := \mathbf{1}_{[0,\varepsilon]}(A_n)$ we have $\tau(P_n) \ge \delta/2$ and $(A_n + \varepsilon I)^{-p} \ge (2\varepsilon)^{-p}P_n$. Therefore,

$$\|(A_n + \varepsilon I)^{-p}\| \ge (2\varepsilon)^{-p} \|P_n\| \ge (2\varepsilon)^{-p} \|I\| \delta/2$$

so that $||A_n||_1 \leq 2\varepsilon(||I||\delta/2)^{-1/p}$ for all $n \geq n_0$. Hence $||A_n||_1 \searrow 0$.

Next assume that ker $A = \{0\}$, so $A^{-p} \in \overline{\mathcal{M}}^+$. Since

$$\lambda_t(A_n^{-p}) = \lambda_{1-t}(A_n)^{-p} \nearrow \lambda_{1-t}(A)^{-p} = \lambda_t(A^{-p})$$

for a.e. $t \in (0, 1)$, we have $||A_n^{-p}|| \nearrow ||A^{-p}||$ by Proposition 6.4 (b). Therefore, $||A_n||_! \searrow ||A||_!$ by Proposition 4.4 (c).

The next theorem is concerned with the converse implication of Proposition 6.10(a).

Theorem 6.11. Let $A, B \in \overline{\mathcal{M}}^+$ and assume that $\int_0^1 \lambda_s(B)^{-p} ds < \infty$ for some p > 0 (in particular, this is the case if $B \ge \delta I$ for some $\delta > 0$). Then the following two conditions are equivalent:

(i) $A \prec^{w(\log)} B;$

(ii) $||A||_{!} \geq ||B||_{!}$ for every fully symmetric derived anti-norm $||\cdot||_{!}$ on \mathcal{M}^{+} .

To prove the theorem, we first give a lemma. When A is invertible, the lemma is [3, Lemma 4.3.6] with a simpler proof.

Lemma 6.12. Let $A \in \mathcal{M}^+$ and assume that $\int_0^1 \lambda_s(A)^{-p} ds < \infty$ for some p > 0. Then, for every $t \in (0, 1]$,

$$\exp\left(\frac{1}{t}\int_{1-t}^{1}\log\lambda_s(A)\,ds\right) = \lim_{p\searrow 0}\left(\frac{1}{t}\int_{1-t}^{1}\lambda_s(A)^{-p}\,ds\right)^{-1/p} = \lim_{p\searrow 0}t^{1/p}\|A^{-p}\|_{(t)}^{-1/p},$$

where $\|\cdot\|_{(t)}$ is in Example 6.3 (1).

Proof. Replacing A with αA for some $\alpha > 0$, we may suppose that $A \leq I$. Assume that $\int_0^1 \lambda_s(A)^{-p_0} ds < \infty$ for some $p_0 > 0$; then $0 < \lambda_s(A) \le 1$ for all $s \in (0,1)$ and $-\lambda_s(A)^{-p}\log\lambda_s(A)$ is integrable on (0,1) for every $p \in (0,p_0)$. Write $\varphi(s,p) := \lambda_s(A)^{-p}$ for $s \in (0, 1)$ and $p \in (0, p_0)$. Since

$$\frac{\varphi(s,p) - \varphi(s,0)}{p} = \partial_p \varphi(s,\theta p) = -\lambda_s(A)^{-\theta p} \log \lambda_s(A) \le -\lambda_s(A)^{-p} \log \lambda_s(A),$$

where $\theta \in (0, 1)$ (depending on s, p) and

$$\lim_{p \searrow 0} \frac{\varphi(s,p) - \varphi(s,0)}{p} = \partial_p \varphi(s,0) = -\log \lambda_s(A),$$

it follows from the Lebesgue convergence theorem that, for every $t \in (0, 1]$,

$$\frac{d}{dp} \int_{1-t}^{1} \varphi(s,p) \, ds \Big|_{p=+0} = \lim_{p \searrow 0} \int_{1-t}^{1} \frac{\varphi(s,p) - \varphi(s,0)}{p} \, ds = -\int_{1-t}^{1} \log \lambda_s(A) \, ds,$$

where $\frac{d}{dp}(\cdot)|_{p=+0}$ means the right derivative at p=0. Therefore,

$$\lim_{p \searrow 0} \left[-\frac{1}{p} \log \left(\frac{1}{t} \int_{1-t}^{1} \lambda_s(A)^{-p} \, ds \right) \right] = -\frac{d}{dp} \log \left(\frac{1}{t} \int_{1-t}^{1} \lambda_s(A)^{-p} \, ds \right) \Big|_{p=+0}$$
$$= \frac{1}{t} \int_{1-t}^{1} \log \lambda_s(A) \, ds,$$

which is equivalent to the desired limit formula.

We now prove the theorem.

Proof. (Theorem 6.11) By Proposition 6.10 (a) we may prove that (ii) \Rightarrow (i), so assume that $\int_0^1 \lambda_s(B)^{-p} ds < \infty$ for some p > 0 and $||A||_1 \ge ||B||_1$ for all fully symmetric derived anti-norms. It suffices to show that, for each $t \in (0,1)$ fixed, if $||A^{-p}||_{(t)}^{-1/p} \ge ||B^{-p}||_{(t)}^{-1/p}$ for all p > 0, then $\int_{1-t}^{1} \log \lambda_s(A) ds \ge \int_{1-t}^{1} \log \lambda_s(B) ds$ holds. Since all the relevant quantities depend on $\lambda(A), \lambda(B)$ restricted on (1-t, 1), we may assume that $A, B \in \mathcal{M}^+$, by replacing A, B with $A \wedge \alpha, B \wedge \beta$ where $\alpha := \lambda_{1-t}(A), \beta := \lambda_{1-t}(B)$, respectively. Then for every $\delta > 0$ we have $\|(A + \delta I)^{-p}\|_{(t)}^{-1/p} \ge \|A^{-p}\|_{(t)}^{-1/p} \ge \|B^{-p}\|_{(t)}^{-1/p}$. Applying Lemma 6.12 to $A + \delta I$ and B yields

$$\int_{1-t}^{1} \log(\lambda_s(A) + \delta) \, ds \ge \int_{1-t}^{1} \log \lambda_s(B) \, ds.$$

Letting $\delta \searrow 0$ gives the desired inequality.

Remark 6.13. Note that there is a $B \in \mathcal{M}^+$ such that $\int_0^1 \log \lambda_s(B) \, ds > -\infty$ but $\int_0^1 \lambda_s(B)^{-p} ds = \infty$ for all p > 0. For instance, this is the case when $\lambda_s(B) =$

٦	1	-	
T	J	J	

 $\exp(-1/\sqrt{1-s})$. For such a $B \in \mathcal{M}^+$ and every p > 0, if $\|\cdot\|$ is a fully symmetric norm on \mathcal{M}^+ , then by Proposition 6.4 (c) we have

$$\lim_{\varepsilon \searrow 0} \|(B + \varepsilon I)^{-p}\| \ge \lim_{\varepsilon \searrow 0} \|(B + \varepsilon I)^{-p}\|_1 \|I\| = \int_0^1 \lambda_s(B)^{-p} \, ds \, \|I\| = \infty.$$

This means that $||B||_{!} = 0$ for every fully symmetric derived anti-norm, so (ii) of Theorem 6.11 is satisfied for any $A \in \mathcal{M}^+$. Therefore, (ii) \Rightarrow (i) does not hold for general $A, B \in \mathcal{M}^+$. Such a subtle difference between the two conditions (i) and (ii) never occurs in the matrix case: In the matrix algebra \mathbb{M}_n , the conditions (i) and (ii) in Theorem 6.11 are equivalent; (i) \Rightarrow (ii) is shown in [7, Lemma 4.10], and (ii) \Rightarrow (i) is implicit in [7, Example 4.5], the discrete version of the anti-norms in (6.2).

The following is a consequence of Lemma 6.12.

Corollary 6.14. For every $t \in (0, 1]$ the functional

$$\Delta_t(A) := \exp\left(\frac{1}{t} \int_{1-t}^1 \log \lambda_s(A) \, ds\right), \qquad A \in \mathcal{M}^+ \tag{6.2}$$

is a symmetric anti-norm on \mathcal{M}^+ .

Proof. The properties $(1)_!$ and $(2)_!$ of Definition 2.2 for Δ_t are clear and $(4)_!$ is immediate from the monotone convergence theorem. To show $(3)_!$, we may assume in view of $(4)_!$ that $A, B \in \mathcal{M}^+$ are invertible. Then Lemma 6.12 yields

$$\Delta_t(A) = \lim_{p \searrow 0} t^{-1/p} \|A^{-p}\|_{(t)}^{-1/p}$$
(6.3)

and the same expressions for $\Delta_t(B)$ and $\Delta_t(A+B)$. Hence (3)! for Δ_t follows from that of the derived anti-norms $||A^{-p}||_{(t)}^{-1/p}$.

The symmetric anti-norms Δ_t are not derived ones, but (6.3) says that they are in the boundary of the derived anti-norms. In particular, when t = 1,

$$\Delta(X) := \exp\left(\int_0^1 \log \lambda_s(|X|) \, ds\right) = \Delta_1(|X|), \qquad X \in \mathcal{M}$$

is the Fuglede-Kadison determinant [15]. This is extended to $\overline{\mathcal{M}}$ by Proposition 4.2 and the above expression holds whenever $\int_0^1 \log \lambda_s(|X|) ds$ makes sense permitting $\pm \infty$. The determinant Δ has been useful in the non-commutative H^{∞} theory (e.g., [3, 5]).

7 Superadditivity with more functions

For a fully symmetric derived anti-norm, Theorem 5.1 for convex functions can be extended to a considerably larger class of superadditive functions given as follows:

Let S be the class of functions $\psi : [0, \infty) \to [0, \infty)$ such that $\psi(t) = f \circ g(t)$ for some superadditive log-concave function $f : [0, \infty) \to [0, \infty)$ and some superadditive convex function $g : [0, \infty) \to [0, \infty)$.

Recall that $f : [0, \infty) \to [0, \infty)$ is log-concave if $f((a+b)/2) \ge \sqrt{f(a)f(b)}$ for all $a, b \ge 0$, i.e., if $\log f : [0, \infty) \to [-\infty, \infty)$ is concave. A convex function $g : [0, \infty) \to [0, \infty)$ is superadditive if and only if g(0) = 0. Note that any function $\psi(t)$ in \mathcal{S} is superadditive and non-decreasing on $[0, \infty)$ with $\psi(0) = 0$.

Any superadditive log-concave function $\psi : [0, \infty) \to [0, \infty)$ is in S. Any convex function $\psi : [0, \infty) \to [0, \infty)$ with $\psi(0) = 0$ is in S. The next examples point out some functions in S which are in the intersection of these two subclasses, or only in one subclass, or in none of them. We implicitly assume that superadditive functions are defined on $[0, \infty)$.

- **Example 7.1.** The power functions $t \mapsto t^p$, $p \ge 1$, and the angle function at any $\alpha > 0, t \mapsto (t \alpha)_+ := \max\{t \alpha, 0\}$, are superadditive, convex and log-concave. The function $t \mapsto t \arctan t$ is also superadditive, convex and log-concave.
 - For any $\gamma > 1$, the functions $t \mapsto \sinh t^{\gamma}$ and $t \mapsto t \exp t^{\gamma}$ are superadditive and convex, but not log-concave.
 - When $1 \leq \alpha < \beta$, the function $t \mapsto \min\{t^{\alpha}, t^{\beta}\}$ is superadditive and log-concave, but not convex. The function $t \mapsto t^{\alpha} \exp(-1/t^{\beta})$ is the same whenever $\alpha \geq 1$ and $\beta > 2\alpha - 1 + 2\sqrt{\alpha(\alpha - 1)}$. When 0 < a < b, the function $t \mapsto (t - a)\mathbf{1}_{[b,\infty)}(t)$ is also the same though not continuous.
 - For $f(t) = \min\{t^{\alpha}, t^{\beta}\}$ with $1 \leq \alpha < \beta$ and $g(t) = \sinh t^{\gamma}$ or $t \exp t^{\gamma}$ with $\gamma > 1$, $f \circ g(t)$ is a function in \mathcal{S} , but neither log-concave nor convex.

Recall that \mathcal{M} stands for a (finite) diffuse algebra. The superadditivity results in this section also hold with \mathbb{M}_n in place of \mathcal{M} with similar though simpler proofs. The next theorem is the main result of this section.

Theorem 7.2. Let $A, B \in \overline{\mathcal{M}}^+$ and let $\psi(t)$ be a function in S. Then, for any fully symmetric derived anti-norm on $\overline{\mathcal{M}}^+$,

$$\|\psi(A+B)\|_{!} \ge \|\psi(A)\|_{!} + \|\psi(B)\|_{!}.$$

The proof is based on Theorem 3.1 and the next lemma.

Lemma 7.3. Let $\|\cdot\|_{!}$ be a fully symmetric derived anti-norm on \mathcal{M}^+ . Let $A, B \in \overline{\mathcal{M}}^+$. If $f : [0, \infty) \to [0, \infty)$ is superadditive and log-concave, then

$$||f(A+B)||_! \ge ||f(A)||_! + ||f(B)||_!.$$

Proof. For each s > 0, since $\lambda(f(A+B)) \ge \lambda(f(A \land s + B \land s))$, by Proposition 6.10 (a) we have $||f(A+B)||_! \ge ||f(A \land s + B \land s)||_!$. Since $||f(A)||_! = \lim_{s \nearrow \infty} ||f(A \land s)||_!$ and similarly for $||f(B)||_!$ as in the proof of Theorem 5.1, we may assume that $A, B \in \mathcal{M}^+$. We prove that

$$f(A+B) \prec^{w(\log)} \int_0^1 f(\lambda_t(A) + \lambda_t(B)) \, dF_t.$$
(7.1)

As f(t) is non-decreasing, we have $\lambda(f(A + B)) \geq \lambda(f(A))$ so that $||f(A + B)||_! \geq ||f(A)||_!$, and similarly $||f(A+B)||_! \geq ||f(B)||_!$. Hence, the claimed inequality is obvious if $||f(A)||_! = 0$ or $||f(B)||_! = 0$. So assume that $||f(A)||_! > 0$ and $||f(B)||_! > 0$. This implies by Proposition 4.4 (b) that f(A) and f(B) are nonsingular, so $f(\lambda_t(A)) > 0$ and $f(\lambda_t(B)) > 0$ for all $t \in (0, 1)$. Hence f(t) > 0 for all $t > t_0 := \min\{\lambda_1(A), \lambda_1(B)\}$. Furthermore, thanks to Proposition 6.10 (b), it suffices to prove the claimed inequality for $A + \varepsilon I$ and $B + \varepsilon I$ for any $\varepsilon > 0$. Thus we can assume that f(t) > 0 for all $t \geq t_0$.

Then, from the majorization $\lambda(A+B) \prec \lambda(A) + \lambda(B)$ and the concavity of log f, we have

$$\log \lambda(f(A+B)) = \log f(\lambda(A+B)) \prec^w \log f(\lambda(A) + \lambda(B)),$$

which means that the log-supermajorization (7.1) holds. As f(t) is further superadditive, (7.1) entails

$$f(A+B) \prec^{w(\log)} \int_0^1 (f(\lambda_t(A)) + f(\lambda_t(B))) dF_t.$$

Therefore, Proposition 6.10(a) implies that

$$\|f(A+B)\|_{!} \ge \left\|\int_{0}^{1} f(\lambda_{t}(A)) \, dF_{t}\right\|_{!} + \left\|\int_{0}^{1} f(\lambda_{t}(B)) \, dF_{t}\right\|_{!}$$
$$= \|f(A)\|_{!} + \|f(B)\|_{!}.$$

Proof. (Theorem 7.2) As in the proof of the previous lemma, we may and do assume
that $A, B \in \mathcal{M}^+$. Let $\psi(t)$ in \mathcal{S} be written as $\psi(t) = f \circ g(t)$, with $f(t)$ superadditive
log-concave and $g(t)$ superadditive convex. If $\ \psi(A)\ _{!} = 0$ or $\ \psi(B)\ _{!} = 0$, then the
claimed inequality follows as in the proof of the previous lemma. So we may assume that
$\ \psi(A)\ _{!} > 0$ and $\ \psi(B)\ _{!} > 0$. This implies that $f(\lambda_t(g(A))) > 0$ and $f(\lambda_t(g(B))) > 0$
for all $t \in (0,1)$. Hence, $f(t) > 0$ for all $t > t_0 := \min\{\lambda_1(g(A)), \lambda_1(g(B))\}$, so that
$f(t)$ is continuous on (t_0,∞) . For every $\varepsilon > 0$ let $U, V \in \mathcal{M}$ be unitaries as given in
Theorem 3.1. Then we have

$$\lambda_t(f(g(A+B)+\varepsilon I)) \ge \lambda_t(f(Ug(A)U^*+Vg(B)V^*))$$

for all $t \in (0, 1)$. By Proposition 6.10 (a) and Lemma 7.3, this implies that

$$\|f(g(A+B)+\varepsilon I)\|_{!} \ge \|f(Ug(A)U^{*}+Vg(B)V^{*})\|_{!}$$
$$\ge \|f(g(A))\|_{!}+\|f(g(B))\|_{!}.$$

Letting $\varepsilon \searrow 0$ yields the claimed inequality thanks to Proposition 6.10 (b).

Finally, we return to a general finite von Neumann algebra \mathcal{N} with a faithful normal trace τ , $\tau(I) = 1$, and extend Theorem 7.2 to $\overline{\mathcal{N}}^+$ with a restriction on derived antinorms. For this, we start with a fully symmetric norm ρ on the commutative von Neumann algebra $L^{\infty}(0,1)$ with the trace $\int_{0}^{1} \cdot dt$ (expectation). Define a fully symmetric norm $\|\cdot\|_{\rho}$ on \mathcal{N} as

$$||X||_{\rho} := \rho(\mu(X)), \qquad X \in \mathcal{N},$$

which we call a ρ -symmetric norm. This way of construction of symmetric norms is common in the theory of non-commutative Banach function spaces (e.g., [11, 12]). Let $\|\cdot\|_{!}$ be the (fully symmetric) derived anti-norm on \mathcal{N}^+ that is derived from $\|\cdot\|_{\rho}$ and a p > 0. In case of a diffuse \mathcal{M} , any fully symmetric norm on \mathcal{M} is a ρ -symmetric norm with

$$\rho(h) := \left\| \int_0^1 h(t) \, dF_t \right\|, \qquad h \in L^{\infty}(0,1),$$

so the restriction here on fully symmetric derived anti-norms is indeed no restriction in the diffuse case. However, on a general \mathcal{N} we have a fully symmetric norm which is not written as a ρ -symmetric norm.

Consider the tensor product (diffuse) von Neumann algebra

$$\mathcal{M} := \mathcal{N} \,\overline{\otimes} \, L^{\infty}(0,1)$$

with the tensor product trace $\tau \otimes \int_0^1 \cdot dt$, and define the ρ -symmetric norm $\|\cdot\|_{\rho}$ on \mathcal{M} and the corresponding derived anti-norm $\|\cdot\|_!$ on \mathcal{M}^+ in the same way as above. Then the following equations are obvious:

$$||X||_{\rho} = ||X \otimes 1||_{\rho}, \quad X \in \overline{\mathcal{N}} ; \qquad ||A||_{!} = ||A \otimes 1||_{!}, \quad A \in \overline{\mathcal{N}}^{+}.$$

Therefore, all the results concerning fully symmetric derived anti-norms on \mathcal{M}^+ in Section 6 and in this section remain true for the derived anti-norm $\|\cdot\|_!$ on \mathcal{N}^+ from $\|\cdot\|_{\rho}$ as above. In particular, we have

Corollary 7.4. Let $\|\cdot\|_{!}$ be a derived anti-norm on \mathcal{N}^{+} that is derived from a ρ -symmetric norm $\|\cdot\|_{\rho}$ on \mathcal{N} and a p > 0. Then, for every $A, B \in \overline{\mathcal{N}}^{+}$ and every function $\psi(t)$ in \mathcal{S} ,

$$\|\psi(A+B)\|_{!} \ge \|\psi(A)\|_{!} + \|\psi(B)\|_{!}.$$

Corollary 7.5. Let $g : [0, \infty) \to [0, \infty)$ be a convex function with g(0) = 0, and let $\psi(t)$ be a strictly increasing function in S. Then, for all nonsingular $A, B \in \mathcal{N}^+$ and all 0 ,

$$\frac{\tau(g^p(A+B))}{\tau(\psi^{p-1}(A+B))} \ge \frac{\tau(g^p(A))}{\tau(\psi^{p-1}(A))} + \frac{\tau(g^p(B))}{\tau(\psi^{p-1}(B))}$$

Proof. By Theorem 5.1 the functional $A \mapsto \{\tau(g^p(A))\}^{1/p}$ is superadditive with finite values on \mathcal{N}^+ . By the previous corollary, $A \mapsto \{\tau(\psi^{p-1}(A))\}^{1/(p-1)}$ is superadditive with strictly positive finite values on the nonsingular part of \mathcal{N}^+ . Hence, their *p*-weighted geometric mean is again superadditive on the nonsingular part of \mathcal{N}^+ . \Box

Note that $\|\cdot\|_{(t)}$ is a ρ -symmetric norm and (6.3) is valid in \mathcal{N}^+ . Hence, through Corollary 7.4 applied to the derived anti-norm from $\|\cdot\|_{(t)}$, we have

$$\Delta_t(\psi(A+B)) \ge \Delta_t(\psi(A)) + \Delta_t(\psi(B)), \qquad A, B \in \overline{\mathcal{N}}^+$$

for all $t \in (0, 1]$ and all functions $\psi(t)$ in \mathcal{S} . For the Fuglede-Kadison determinant Δ , since $\Delta(\sqrt{\psi\omega}(A)) = {\Delta(\psi(A))\Delta(\omega(A))}^{1/2}$ for $\psi, \omega : [0, \infty) \to [0, \infty)$ and $A \in \overline{\mathcal{N}}^+$, we furthermore have

Corollary 7.6. Let $\psi(t)$ and $\omega(t)$ be two functions in S. Then, for all $A, B \in \overline{\mathcal{N}}^+$,

$$\Delta\left(\sqrt{\psi\omega}(A+B)\right) \ge \Delta\left(\sqrt{\psi\omega}(A)\right) + \Delta\left(\sqrt{\psi\omega}(B)\right).$$

This is a substantial generalization of the Minkowski inequality for $\Delta(A)$ on \mathcal{M}^+ given in [3] as a consequence of a variational expression of Δ . In addition, it is worth noting that the concavity of $A \mapsto \Delta(f(A))$ on \mathcal{M}^{sa} for a positive concave function fwas shown in [22] (a similar result for matrices is in [6]).

References

- C. A. Akemann, J. Anderson and G. K. Pedersen, Triangle inequalities in operator algebras, *Linear Multilinear Algebra* 11 (1982), 167–178.
- [2] J. S. Aujla and J.-C. Bourin, Eigenvalues inequalities for convex and log-convex functions, *Linear Algebra Appl.* 424 (2007), 25–35.
- [3] W. B. Arveson, Analyticity in operator algebras, Amer. J. Math. 89 (1967), 578–642.
- [4] R. Bhatia, *Matrix Analysis*, Springer, New-York, 1997.
- [5] D. P. Blecher and L. E. Labuschagne, Von Neumann algebraic H[∞] theory, in Function Spaces, Contemp. Math., 435, 2007, Amer. Math. Soc., pp. 89–114.
- [6] J.-C. Bourin and F. Hiai, Norm and anti-norm inequalities for positive semi-definite matrices, Internat. J. Math. 22 (2011), 1121–1138.
- [7] J.-C. Bourin and F. Hiai, Jensen and Minkowski inequalities for operator means and anti-norms, *Linear Algebra Appl.* 456 (2014), 22–53.
- [8] J.-C. Bourin and E.-Y. Lee, Unitary orbits of Hermitian operators with convex or concave functions, Bull. London Math. Soc. 44 (2012), 1085–1102.
- [9] L. G. Brown and H. Kosaki, Jensen's inequality in semi-finite von Neumann algebras, J. Operator theory 23 (1990), 3–19.
- [10] K. M. Chong, Doubly stochastic operators and rearrangement theorem, J. Math. Anal. Appl. 56 (1976), 309–316.
- [11] P. G. Dodds, T. K.-Y. Dodds and B. de Pagter, Non-Commutative Banach Function Spaces, Math. Z. 201 (1989), 583–597.

- [12] P. G. Dodds, T. K.-Y. Dodds and B. de Pagter, Noncommutative Köthe duality, Trans. Amer. Math. Soc. 339 (1993), 717–750.
- [13] T. Fack and H. Kosaki, Generalized s-numbers of τ-measurable operators, Pacific J. Math. 123 (1986), 269–300.
- [14] J. Fang, D. Hadwin, E. Nordgren and J. Shen, Tracial gauge norms on finite von Neumann algebras satisfying the weak Dixmier property, J. Funct. Anal. 255 (2008), 142–183.
- [15] B. Fuglede and R. Kadison, Determinant theory in finite factors, Ann. of Math. 55 (1952), 520–530.
- [16] F. Hiai, Spectral relations and unitary mixing in semifinite von Neumann algebras, Hokkaido Math. J. 17 (1988), 117–137.
- [17] F. Hiai and Y. Nakamura, Majorizations for generalized s-numbers in semifinite von Neumann algebras, Math. Z. 195 (1987), 17–27.
- [18] F. Hiai and Y. Nakamura, Distance between unitary orbits in von Neumann algebras, *Pacific J. Math.* 138 (1989), 259–294.
- [19] F. Hiai and Y. Nakamura, Closed convex hulls of unitary orbits in von Neumann algebras, Trans. Amer. Math. Soc. 323 (1991), 1–38.
- [20] C. J. Himmelberg, Measurable relations, Fund. Math. 87 (1975), 53–72.
- [21] H. Kosaki, On the continuity of the map $\varphi \to |\varphi|$ from the predual of a W*-algebra, J. Funct. Anal. **59** (1984), 123–131.
- [22] E. H. Lieb and G. K. Pedersen, Convex multivariable trace functions, *Rev. Math. Phys.* 14 (2002), 631–648.
- [23] S. Lord, F. Sukochev and D. Zanin, Singular Traces: Theory and Applications, De Gruyter Studies in Mathematical Physics 46, De Gruyter, Berlin, 2013.
- [24] M. Marcus and L. Lopes, Inequalities for symmetric functions and Hermitian matrices, Canad. J. Math. 9 (1957), 305–312.
- [25] A. W. Marshall, I. Olkin and B. C. Arnold, *Inequalities: Theory of Majorization and Its Applications*, Springer, New York, second edition, 2011.
- [26] D. Petz, Spectral scale of self-adjoint operators and trace inequalities, J. Math. Anal. Appl. 109 (1985), 74–82.
- [27] M. Takesaki, Theory of Operator Algebras. I, Springer-Verlag, New York-Heidelberg, 1979.
- [28] K. Watanabe, Some results on non-commutative Banach function spaces, Math. Z. 210 (1992), 555–572.

Jean-Christophe Bourin Laboratoire de mathématiques, Université de Franche Comté, 25030 Besançon, France jcbourin@univ-fcomte.fr

Fumio Hiai Tohoku University (Emeritus), Hakusan 3-8-16-303, Abiko 270-1154, Japan hiai.fumio@gmail.com