

ISOGENIES OF JACOBIANS

V. MARCUCCI, J.C. NARANJO, AND G.P. PIROLA

ABSTRACT. We prove by means of the study of the infinitesimal variation of Hodge structure and a generalization of the classical Babbage-Enriques-Petri theorem that the Jacobian variety of a generic element of a codimension k subvariety of \mathcal{M}_g is not isogenous to a distinct Jacobian if $g > 3k + 4$. We extend this result to $k = 1, g \geq 5$ by using degeneration methods.

1. INTRODUCTION

Let \mathcal{Z} be a subvariety of the moduli space \mathcal{M}_g of complex smooth curves of genus g of codimension $k > 0$. We want to show that under some numerical restrictions, the Jacobian of a generic element of \mathcal{Z} is not isogenous to a distinct Jacobian. In other words, all the curves of genus g contained in $J\mathcal{C}$, with C generic in \mathcal{Z} , are birationally equivalent. This is an extension of the Theorem proved by Bardelli and Pirola (see [BP]) for the whole \mathcal{M}_g and can be seen as a Noether-Lefschetz locus problem for surfaces which are the product of two curves of the same genus (see Corollary (6.2)). More precisely, our result is as follows:

Theorem 1.1. *Let $\mathcal{Z} \subset \mathcal{M}_g$ a codimension $k > 0$ subvariety. Assume that $g > 3k + 4$ (in particular $g > 7$), then the Jacobian of a generic curve C of \mathcal{Z} is not isogenous to another Jacobian. The same is true for $k = 1$ and $g \geq 5$.*

Observe that the Theorem fails for $g = 4$ and $k = 1$: in this case \mathcal{M}_4 is a divisor in \mathcal{A}_4 , therefore intersecting in the Siegel upper space \mathcal{H}_4 the Jacobian locus \mathcal{J}_4 with the image $j(\mathcal{J}_4)$ by the action of a fixed isogeny j , we get a divisor in \mathcal{M}_4 where the Jacobian of a generic element is isogenous to a different Jacobian.

For $g > 3k + 4$, our strategy is as follows: after a base change we have two families of smooth complex curves of genus g on a base variety W , $\pi : \mathcal{C} \rightarrow W$ and $\pi' : \mathcal{C}' \rightarrow W$, and a family of isogenies of the associated family of Jacobians, that is

$$\chi : J(\mathcal{C}') \rightarrow J(\mathcal{C}).$$

This means that for $t \in W$ the map $\chi_t : J(C'_t) \rightarrow J(C_t)$ is an isogeny, here $C'_t = \pi'^{-1}(t)$ and $C_t = \pi^{-1}(t)$.

It follows that the associated rational Hodge structures are isomorphic. Consider the local (polarized) systems $\Lambda_{\mathbb{Z}} = R^1\pi_*\mathbb{Z}$ and $\Lambda'_{\mathbb{Z}} = R^1\pi'_*\mathbb{Z}$ and tensoring by \mathbb{C} $\Lambda_{\mathbb{C}} = R^1\pi_*\mathbb{C}$ and $\Lambda'_{\mathbb{C}} = R^1\pi'_*\mathbb{C}$.

In particular the infinitesimal variation of Hodge structure associated to the Hodge filtration of $\Lambda^{1,0} \subset \Lambda_{\mathbb{C}}$ and $\Lambda'^{1,0} \subset \Lambda'_{\mathbb{C}}$ are isomorphic. We borrowed this basic observation from Claire Voisin (see Remark (4.2.5) in [BP]). It is well known

Naranjo has been partially supported by the Proyecto de Investigación MTM2012-38122-C03-02; Pirola has been partially supported by Gnsaga and by MIUR PRIN 2012: Moduli, strutture geometriche e loro applicazioni. Part of this research has been done during the visit of Pirola to the IMUB in the spring of 2013.

that the infinitesimal invariant of Hodge structure of curves determines the quadrics that contain a canonical curve (see [CGGH]). This allows to translate our problem to a geometric one. Let $I(2)$ and $J(2)$ be the space of quadrics that contain the canonical curve associated to C_t and D_t . It follows that under a choice of a suitable canonical embedding $I(2) \cap J(2)$ has codimension $\geq k$, where k is the codimension of $m(W)$ in \mathcal{M}_g and m is the modular mapping $m : W \rightarrow \mathcal{M}_g$. We can bound the codimension k by using the Clifford index. For this we prove a result that gives an interesting (at least in our opinion) reconstruction result of the curve from a *partial system of quadrics*. It is a generalization of the Babbage-Enriques-Petri theorem (see e.g. chapter 3, section 3 in [ACGH]).

Theorem 1.2. *Let C be a curve of genus g and Clifford index c . Let $I_2 \subset \text{Sym}^2 H^0(C, \omega_C)$ be the vector space of the equations of the quadrics containing C and let $K \subset I_2$ be a linear subspace of codimension k . If $g > 2k + 5$ and $c > k + 1$, then C is the only irreducible non-degenerate curve contained in the intersection of the quadrics of K .*

Corollary 1.3. *Let C be a generic curve in a codimension k subvariety \mathcal{Z} of \mathcal{M}_g . Let $I_2 \subset \text{Sym}^2 H^0(C, \omega_C)$ be the vector space of the equations of the quadrics containing C . Let $K \subset I_2$ be a linear subspace of codimension k . If $g > 3k + 4$, then C is the only irreducible non-degenerate curve contained in the intersection of the quadrics of K .*

The Corollary is a consequence of the Theorem (1.2). Indeed, let c be the Clifford index of a generic element of \mathcal{Z} . The locus of curves with given Clifford index can have several components and the minimal codimension is attained when c is realized by a g_d^1 linear series, with $c = d - 2$. This follows easily by a parameter count from a result in [CM] where it is proved that a curve C with Clifford index c is either $(c + 2)$ -gonal or the dimension of the Brill-Noether locus $W_{c+3}^1(C)$ is at least 1.

Then, by Riemann-Hurwitz, the codimension of the component of the curves with a g_{c+2}^1 linear series is $3g - 3 - (2g - 2 + 2(c + 2) - 3) = g - 2c - 2$. Hence, since we assume $g > 3k + 4$, we have

$$k \geq g - 2c - 2 > 3k + 4 - 2c - 2 = 3k - 2c + 2.$$

Therefore $c > k + 1$ and the result follows from (1.2) since $g > 3k + 4$ implies $g > 2k + 5$ for $k > 0$.

In section 2 we start the proof of Theorem (1.1) under the hypothesis $g > 3k + 4$ by reducing it to Corollary (1.3) following Voisin's observation indicated above. Theorem (1.2) will be proved in Section 3. The idea of the proof is as follows: assuming the existence of a second non-degenerate curve in the intersection of the quadrics we select linearly independent points x_i in this curve. Then, by constructing a suitable rank 2 vector bundle on C , we are able to find points $p_j \in C$ such that the linear span of the points x_i is contained in the linear span of the points p_j . From this it is easy to obtain a contradiction by using a theorem of Ran [R].

To prove the divisorial case of the main theorem, we use the original approach in [BP], based on the analysis of the map

$$\chi_{\mathcal{Z}} : \Lambda_{\mathcal{Z}} \rightarrow \Lambda'_{\mathcal{Z}}$$

(in fact we will work with the dual lattices, that is, with the homology groups). If we can prove that $\chi_{\mathbb{Z}}(\Lambda_{\mathbb{Z}}) = n\Lambda'_{\mathbb{Z}}$, $n \in \mathbb{Z}$ we will get that \mathcal{C}'_t is isomorphic to \mathcal{C}_t and χ_t is given by multiplication by n . We use degeneration to Δ_0 , and study the monodromy action on $\Lambda_{\mathbb{Z}}$. The basic geometric information is now encoded on the generalized Jacobians. Roughly speaking one has to prove that part of the limit map χ_0 is multiplication by n . This gives that the map χ_t is multiplication by n on *a part of the invariant cycles*. We need finally to have degenerations with independent monodromy to complete the proof. It is clear that to follow this strategy one needs to control the degeneration type. Using the theory of divisors on \mathcal{M}_g and following a valuable suggestion of Gavril Farkas, we realize the above program when $c = 1$. The degeneration procedure is performed in sections 4, 5 and 6. We will prove the existence of a type of degeneration (if $c = 1$) to the union of a curve of genus $g - 2$ and two generic elliptic tails. The *independent* degenerations to Δ_0 are obtained by letting the elliptic tails becomes singular. To extract more information from the degeneration we analyze the type of monodromy involved that we classify in three cases (a,b,c of section 5). Then we analyze the geometry of the generalized Jacobians by comparing their extension classes. In section 6 we complete the proof by comparing the invariants of the two degenerations.

Acknowledgements: We thank Gavril Farkas for useful and stimulating conversations on the moduli space of curves.

2. REDUCTION TO A PROBLEM ON QUADRICS THROUGH THE CANONICAL CURVE

The aim of this section is to prove that Corollary (1.3) implies Theorem (1.1) under the hypothesis $g > 3k + 4$ and $k > 0$.

Remember that an isogeny $\chi : A' \rightarrow A$ between principally polarized Abelian varieties $(A', L_{A'})$ and (A, L_A) such that $\chi^* L_A \cong L_{A'}^{\otimes m}$ is determined by a subgroup H of the group of m -torsion points A'_m totally isotropic with respect to the Riemann bilinear form

$$e_m : A'_m \times A'_m \rightarrow \mu_m$$

(being $A = A'/H$) and a level subgroup \tilde{H} of the theta group $\mathcal{G}(L_{A'}^{\otimes m})$, see ([M], chapter 23). Then the moduli space of those isogenies can be rewritten as

$$\tilde{\mathcal{A}}_g^m = \{\chi : A' \rightarrow A, \chi^*(L_A) \cong L_{A'}^{\otimes m}\}_{/\cong} = \{(A', L_{A'}; H, \tilde{H})\}_{/\cong}$$

and the forgetful map is a finite covering $\varphi : \tilde{\mathcal{A}}_g^m \rightarrow \mathcal{A}_g$. Moreover the map $\psi : \tilde{\mathcal{A}}_g^m \rightarrow \mathcal{A}_g$ sending $\chi : A' \rightarrow A$ to (A, L_A) is another covering space.

Given a generic isogeny $\chi : A' \rightarrow A$ we consider tangent spaces in the following diagram:

$$\begin{array}{ccc} \tilde{\mathcal{A}}_g^m & \xrightarrow{\psi} & \mathcal{A}_g \\ \varphi \downarrow & & \\ \mathcal{A}_g & & \end{array}$$

and we get an isomorphism λ as follows:

$$\begin{array}{ccc}
T_{\widetilde{\mathcal{A}}_g^m, \chi} & \xrightarrow[\cong]{d\psi} & T_{\mathcal{A}_g, A} = \text{Sym}^2 H^0(A, T_A) \\
d\varphi \downarrow \cong & \nearrow \lambda & \\
T_{\mathcal{A}_g, A'} & & \\
\parallel & & \\
\text{Sym}^2 H^0(A', T_{A'}) & &
\end{array}$$

Coming back to our problem let us assume that the locus of curves in \mathcal{M}_g with Jacobian isogenous to the Jacobian of some curve in \mathcal{Z} contains a codimension k component $\mathcal{Z}' \subset \mathcal{M}_g$. Our hypothesis on k implies that a generic element $C' \in \mathcal{Z}'$ satisfies $\text{End}(JC) \cong \mathbb{Z}$ (see [CGT] or [P]). Therefore an isogeny $\chi : JC' \rightarrow JC$ must satisfy that the pull-back of the principal polarization in JC is a multiple of the principal polarization in JC' . Hence there exists an integer m and an irreducible variety $\mathcal{R} \subset \widetilde{\mathcal{A}}_g^m$ dominating \mathcal{Z}' and \mathcal{Z} through φ and ψ respectively. Set $\mathcal{M} := \varphi^{-1}(\mathcal{M}_g)$ and $\mathcal{M}' := \psi^{-1}(\mathcal{M}_g)$. Then $\mathcal{R} \subset \mathcal{M} \cap \mathcal{M}'$. Fix a generic element $\chi : JC' \rightarrow JC$ in \mathcal{R} . In the following diagram we consider in the first row the natural inclusions of tangent spaces at χ and we put in the second row its image by $d\varphi$:

$$\begin{array}{ccccccc}
T_{\mathcal{R}, \chi} & \hookrightarrow & T_{\mathcal{M}, \chi} & \hookrightarrow & T_{\mathcal{M}, \chi} + T_{\mathcal{M}', \chi} & \hookrightarrow & T_{\widetilde{\mathcal{A}}_g^m, \chi} \\
\downarrow \cong & & \downarrow \cong & & \parallel & & \downarrow \cong \\
T_{\mathcal{Z}, JC} & \hookrightarrow & T_{\mathcal{M}_g, JC} = H^0(C, \omega_C^{\otimes 2})^* & \hookrightarrow & \bar{T} & \hookrightarrow & \text{Sym}^2 H^0(C, \omega_C)^*
\end{array}$$

Observe that, by the Grassmann formula, the dimension of \bar{T} is at most $3g-3+k$. Set $K(C) := \text{Kernel}(\text{Sym}^2 H^0(C, \omega_C) \rightarrow \bar{T}^*)$, this is a subspace of the vector space $I_2(C)$ of the quadrics containing the image of C by the canonical map. The codimension of $K(C)$ in $I_2(C)$ is at most k . By using ψ instead of φ we get the corresponding vector space $K(C') \subset I_2(C')$ and we obtain a canonical isomorphism $K(C) \cong K(C')$. Then the Corollary (1.3) implies that C and C' are isomorphic and, since $\text{End}(JC) = \mathbb{Z}$, the isogeny is a multiple of the identity.

3. HIGH CODIMENSION FAMILY OF QUADRICS THROUGH THE CANONICAL CURVE

This section is devoted to the proof of Theorem (1.2). We fix the notation $K \subset I_2$ of the statement. We assume that the intersection of all the quadrics of K contains an irreducible non-degenerate curve different from C . In particular we can select $k+1$ linearly independent points $x_i \in \bigcap_{Q \in K} Q \subset \mathbb{P}H^0(C, \omega_C)^*$ such that $x_i \notin C$. We choose a representative of x_i in $H^0(C, \omega_C)^*$ and we denote it with the same symbol. Then $x_i \otimes x_i \in \text{Sym}^2 H^0(C, \omega_C)^*$. We denote by L the linear variety spanned by these points.

Let R, R' be the quotients I_2/K and $Sym^2 H^0(C, \omega_C)/K$ respectively. Then we have the diagram of vector spaces

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & K & \longrightarrow & I_2 & \longrightarrow & R \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \\
& & Sym^2 H^0(C, \omega_C) & = & Sym^2 H^0(C, \omega_C) & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & R' & \longrightarrow & H^0(C, \omega_C^{\otimes 2}) & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \\
& & 0 & & 0 & &
\end{array}$$

and its dual

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \uparrow & & \uparrow & & \\
0 & \longrightarrow & I_2^* & \longrightarrow & K^* & \longrightarrow & 0 \\
& & \uparrow & & \uparrow & & \\
& & Sym^2 H^0(C, \omega_C)^* & = & Sym^2 H^0(C, \omega_C)^* & & \\
& & \uparrow & & \uparrow & & \\
0 & \longrightarrow & H^1(C, T_C) = H^0(C, \omega_C^{\otimes 2})^* & \longrightarrow & R'^* & \longrightarrow & R^* \longrightarrow 0 \\
& & \uparrow & & \uparrow & & \\
& & 0 & & 0 & &
\end{array}$$

Since all the quadrics of K vanish on x_i then the image of L in K^* is zero, hence $L \subset R'^*$. Since L has dimension $k+1$ and, by the hypothesis on K , $\dim R = k$ we have $H^1(C, T_C) \cap L \neq (0)$. Let α be a non-trivial element in this intersection. Looking at $H^1(C, T_C) = Ext^1(\omega_C, \mathcal{O}_C)$ as classes of extensions we associate to α a rank 2 vector bundle E_α and a short exact sequence:

$$0 \longrightarrow \mathcal{O}_C \longrightarrow E_\alpha \longrightarrow \omega_C \longrightarrow 0.$$

The coboundary map $H^0(C, \omega_C) \longrightarrow H^1(C, \mathcal{O}_C)$ is the cup-product with α . Since $\alpha \in L$, then $\alpha = \sum_{i=1}^{k+1} a_i x_i \otimes x_i$. Therefore, denoting by H_i the kernel of the form $x_i : H^0(C, \omega_C) \longrightarrow \mathbb{C}$, the intersection $H_1 \cap \dots \cap H_{k+1}$ is contained in $Ker(\cdot \cup \alpha)$, in fact

$$(1) \quad Ker(\cdot \cup \alpha) = \bigcap_{i \text{ with } a_i \neq 0} H_i.$$

We can assume that $x_1, \dots, x_{k'}, k' \leq k+1$, are the points such that $a_i \neq 0$. Then there are $g - k'$ sections of $H^0(C, \omega_C)$ lifting to E_α . Let $W \subset H^0(C, E_\alpha)$ be the vector space generated by these sections. We consider the wedge product of

sections:

$$\psi : \Lambda^2 W \hookrightarrow \Lambda^2 H^0(C, E_\alpha) \longrightarrow H^0(C, \det E_\alpha) = H^0(C, \omega_C).$$

The hypothesis $g > 2k + 5$ implies that the projectivization of the kernel of ψ (which has codimension at most g) intersects in $\mathbb{P}(\Lambda^2 W)$ the Grassmannian of the decomposable elements. Hence there are two sections $s_1, s_2 \in W \subset H^0(C, E_\alpha)$ such that $s_1 \wedge s_2 = 0$. This means that they generate a rank 1 torsion free sheaf $M_\alpha \subset E_\alpha$, hence a line bundle. By construction $h^0(C, M_\alpha) \geq 2$. Let us consider the quotient of sheaves $Q_\alpha = E_\alpha/M_\alpha$ and let Q_α^0 be the quotient of Q_α by its torsion subsheaf. The kernel L_α of the natural map $E_\alpha \mapsto Q_\alpha^0$ is a line bundle that by construction contains M_α . In particular $h^0(C, L_\alpha) \geq 2$. Observe that $Q_\alpha^0 \cong \omega_C \otimes L_\alpha^{-1}$. We get a diagram:

$$(2) \quad \begin{array}{ccccccc} & & & 0 & & & \\ & & & \downarrow & & & \\ & & & L_\alpha & & & \\ & & & \downarrow & & & \\ 0 & \rightarrow & \mathcal{O}_C & \xrightarrow{\quad} & E_\alpha & \xrightarrow{\quad} & \omega_C \rightarrow 0 \\ & & \searrow \rho & & \downarrow & & \\ & & & & \omega_C \otimes L_\alpha^{-1} & & \\ & & & & \downarrow & & \\ & & & & 0 & & \end{array}$$

Note that $\rho \neq 0$, otherwise the section of E_α represented by the horizontal arrow $\mathcal{O} \mapsto E_\alpha$ would belong to W which contradicts the definition of W .

Observe that the existence of the map ρ implies that $h^0(C, \omega_C \otimes L_\alpha^{-1})$ is positive. We distinguish two cases:

Case 1: $h^0(C, \omega_C \otimes L_\alpha^{-1}) \geq 2$. Then we can use L_α to compute the Clifford index of the curve. We have:

$$h^0(C, L_\alpha) + h^0(C, \omega_C \otimes L_\alpha^{-1}) \geq h^0(C, E_\alpha) \geq g - k' + 1$$

that combined with Riemann-Roch gives $2h^0(C, L_\alpha) \geq \deg(L_\alpha) + 2 - k'$. Therefore

$$\deg(L_\alpha) - 2h^0(C, L_\alpha) + 2 \leq k' \leq k + 1$$

which is contradiction, since $k + 1 < c$ by hypothesis.

Case 2: $h^0(C, \omega_C \otimes L_\alpha^{-1}) = 1$. Then $h^0(C, L_\alpha) \geq g - k'$. Let e be the degree of $\omega_C \otimes L_\alpha^{-1} \cong \mathcal{O}_C(p_1 + \dots + p_e)$.

Claim: we have that $e \leq k'$.

Indeed, since ρ induces an isomorphism $H^0(C, \mathcal{O}_C) \cong H^0(C, \omega_C \otimes L_\alpha^{-1})$, then the map $H^0(C, E_\alpha) \longrightarrow H^0(C, \omega_C \otimes L_\alpha^{-1})$ is surjective and we get:

$$g - k' \leq h^0(C, L_\alpha) = h^0(C, \omega_C \otimes L_\alpha^{-1}) + 2g - 2 - e + 1 - g = g - e,$$

the claim follows.

Coming back to diagram (2) we obtain

$$H^0(C, L_\alpha) = H^0(C, \omega_C(-p_1 - \dots - p_e)) \subset \text{Ker}(\cdot \cup \alpha) = \bigcap_{i=1, \dots, k'} H_i.$$

By dualizing, we obtain the inclusion of linear spans

$$\langle x_1, \dots, x_{k'} \rangle \subset \langle p_1, \dots, p_e \rangle.$$

We denote by \tilde{C}_0 a non-degenerate irreducible curve, $\tilde{C}_0 \neq C$, contained in all the quadrics parametrized by K . Let \tilde{C} be the normalization of \tilde{C}_0 and let γ be the normalization map. By choosing generically the $k+1$ points $x_i \in \tilde{C}$, we can assume that k' and e are constant, so the correspondence:

$\Gamma = \{(x_1 + \dots + x_{k'}, p_1 + \dots + p_e) \mid \langle \gamma(x_1), \dots, \gamma(x_{k'}) \rangle \subset \langle p_1, \dots, p_e \rangle\} \subset \tilde{C}^{(k')} \times C^{(e)}$ dominates $\tilde{C}^{(k')}$. Moreover, since \tilde{C} is non-degenerate, the fibers of $\pi_2 : \Gamma \rightarrow C^{(e)}$ must be finite. Since $e \leq k' \leq \dim \Gamma = \dim \pi_2(\Gamma) \leq e$ we obtain that $e = k'$. On the other hand the natural rational maps

$$\begin{aligned} C^{(e)} &\dashrightarrow \text{Sec}^e(C) \subset \text{Grass}(e-1, \mathbb{P}^{g-1}) \\ \tilde{C}^{(e)} &\dashrightarrow \text{Sec}^e(\tilde{C}_0) \subset \text{Grass}(e-1, \mathbb{P}^{g-1}) \end{aligned}$$

are generically injective by the uniform position theorem (remember $e = k' \leq k+1$ and $2k+5 < g$), hence the correspondence Γ is of bidegree $(1, 1)$ and therefore $\tilde{C}^{(e)}$ and $C^{(e)}$ are birational. In particular $g(\tilde{C}) = g(C) = g$ (the induced map on Jacobians $J\tilde{C} \rightarrow JC$ has to be dominant since the image generates and the same in the opposite direction). By Ran's Theorem on symmetric products, see [R], we get $C \cong \tilde{C}$. Observe that both curves have to be canonical, hence \tilde{C}_0 is smooth and there is a linear projective transformation $\varphi : \tilde{C} \rightarrow C$. Coming back to our argument and choosing k generic points $x_i \in \tilde{C}$ we have that $\langle x_1, \dots, x_k \rangle = \langle \varphi(x_1), \dots, \varphi(x_k) \rangle$, hence φ leaves $\text{Sec}^k(C)$ invariant and must be the identity, so $\tilde{C} = C$ which is a contradiction.

4. DIVISOR CASE, INTERSECTION WITH THE BOUNDARY

Now we start the proof of the codimension 1 case of the main Theorem assuming $g \geq 5$. We put now \mathcal{D} instead of \mathcal{Z} . The initial step of our degeneration procedure is to show that the intersection of \mathcal{D} with the boundary contains appropriate stable curves. These curves have to contain enough information to deduce from them the main result for the general smooth curve. The goal of this section is to define a family of convenient reducible curves and to prove that they appear in the closure of \mathcal{D} .

We start by recalling the following well-known facts on the rational Picard group of the compactified moduli space $\overline{\mathcal{M}}_g$ of stable curves (see for instance [ACG]):

$$\text{Pic}_{\mathbb{Q}} \mathcal{M}_g = \lambda \mathbb{Q},$$

where λ is the Hodge class. Moreover:

$$\overline{\mathcal{M}}_g \setminus \mathcal{M}_g = \bigcup_{i=0}^{\lfloor \frac{g-1}{2} \rfloor} \Delta_i$$

and

$$Pic_{\mathbb{Q}}(\overline{\mathcal{M}}_g) = \langle \lambda, \delta_0, \delta_1, \dots, \delta_{\lfloor \frac{g-1}{2} \rfloor} \rangle \mathbb{Q}$$


where $\delta_i, i > 0$ is the class of the divisor Δ_i whose general point represents a nodal curve $C_1 \cup C_2$, C_1, C_2 being integral, smooth curves of genus i and $g - i$ intersecting in one point. And δ_0 is the class of Δ_0 whose general point represents an irreducible curve with exactly one node.

We denote by d the class of $\overline{\mathcal{D}}$ in the rational Picard group. Then we can write:

$$(3) \quad d = a\lambda + \sum_{i \geq 0} a_i \delta_i.$$

Remark 4.1. We note that a must be different from zero. Otherwise the class of \mathcal{D} in the rational Picard group of \mathcal{M}_g would be zero. Since the Satake compactification $\overline{\mathcal{M}}_g^s$ of \mathcal{M}_g is a projective variety and the boundary $\overline{\mathcal{M}}_g^s - \mathcal{M}_g$ has codimension 2, given a smooth point p of \mathcal{D} , there exists a complete curve C in \mathcal{M}_g going through p and cutting \mathcal{D} transversally. Hence $C \cdot \mathcal{D} \neq 0$ which is a contradiction.


Now we consider a complete integral curve B in \mathcal{M}_{g-2} (it exists because $g-2 \geq 3$) and we fix two elliptic curves E_1, E_2 with arbitrary j -invariants $j_1, j_2 \in \mathcal{M}_1$. Denote by Γ_b the smooth curve of genus $g - 2$ corresponding to b . We consider the set of the stable curves obtained by glueing to Γ_b the two elliptic curves in two distinct points p_1 and p_2 of Γ_b . This does not depend on the choice of the points on the elliptic curves. This family is parametrized by the symmetric product $\Gamma_b^{(2)} \setminus \Delta_{\Gamma_b}$ minus the diagonal.



So we have a well-defined map

$$\begin{aligned} \Gamma_b^{(2)} \setminus \Delta_{\Gamma_b} &\longrightarrow \Delta_1 \subset \overline{\mathcal{M}}_g \\ p_1 + p_2 &\mapsto E_1 \cup_{p_1} \Gamma \cup_{p_2} E_2, \end{aligned}$$

which extends to the whole symmetric product by sending $2p$ to the following curve: glue the infinity point of a \mathbb{P}^1 with the point p and then glue E_1, E_2 to other two points in the line.



We note that these curves also belong to Δ_2 .

Finally, by moving b in the curve B we obtain a complete threefold $T \subset \Delta_1$. In other words, this threefold can be seen as the image in $\overline{\mathcal{M}}_g$ of the relative symmetric product over B :

$$T = \bigcup_{b \in B} \Gamma_b^{(2)}.$$

Our aim is to study the restriction of the divisor $\overline{\mathcal{D}}$ to T . To do this we make a computation in $\text{Pic}_{\mathbb{Q}}(T)$. Denote by S the surface in T obtained as the union of all the diagonals:

$$S = \bigcup_{b \in B} \Delta_{\Gamma_b}.$$

We will need the following vanishing results.

Lemma 4.2. *The restriction of the class δ_1 to S is zero: $\delta_1|_S = 0$.*

Proof. We fix a smooth curve C of genus 2 with a marked point x . We glue C with Γ_b identifying x with $p \in \Gamma_b$. Then, by moving p in Γ_b and b in B , we construct an algebraic surface S_C such that $\Delta_1 \cap S_C = \emptyset$. Therefore $\delta_1 \cdot S_C = 0$. Now we degenerate C to a genus 2 curve with a marked point consisting in the two elliptic curves E_1, E_2 glued to a \mathbb{P}^1 in 0 and 1 respectively and infinity being the marked point. Therefore by adding the curve Γ_b identifying ∞ with p we get our surface S as a limit of a family of algebraic surfaces S_C as above. We get that $\delta_1 \cdot S = 0$. \square

Lemma 4.3. *For each $b \in B$, $\lambda|_{\Delta_{\Gamma_b}} = 0$.*

Proof. The Hodge structure is constant along the diagonal. \square

We also will use the following basic observation:

Lemma 4.4. *Let N be a complete curve in $\overline{\mathcal{M}}_g$. Then $\rho|_N \neq 0$ for at least one class $\rho \in \{\lambda, \delta_0, \dots, \delta_{\lfloor \frac{g-1}{2} \rfloor}\}$.*

The main result of this section is the following

Proposition 4.5. *The restriction $d|_T$ is not a multiple of the class of S in $\text{Pic}_{\mathbb{Q}}(T)$, i.e. $d|_T \neq mS$ for all $m \in \mathbb{Q}$. In particular $\overline{\mathcal{D}} \cap T \neq \emptyset$ and this intersection contains elements outside S .*

Proof. We use the notation introduced in (3). By contradiction, assume that $d|_T = mS$. Notice that Δ_i does not intersect T for $i = 0$ and $i \geq 3$ and that $\Delta_2 \cap T = S$, so we get:

$$d|_T = mS = a\lambda|_T + a_1\delta_1|_T + a_2kS,$$

for some k . Therefore

$$(m - a_2k)S = a\lambda|_T + a_1\delta_1|_T.$$

Restricting to one diagonal Δ_{Γ_b} and using Lemmas 4.2 and 4.3 we deduce that $m - a_2k = 0$. Restricting now to S we get $a\lambda|_T = 0$. Since λ is not trivial on T we obtain that $a = 0$ which contradicts Remark (4.1). \square

Remark 4.6. Observe that the isomorphism classes of E_1 and E_2 are arbitrary, hence they could represent the ∞ class. Then the limit curves we were looking for are:

$$\mathcal{L} = \{E_1 \cup_{p_1} \Gamma \cup_{p_2} E_2 \in \overline{\mathcal{D}} \mid \Gamma \in \mathcal{M}_{g-2}, p_1 \neq p_2, E_1, E_2 \in \overline{\mathcal{M}}_1\} \subset \overline{\mathcal{D}}.$$

We recall that a generic point of M_g , $g \geq 3$ is contained in a complete curve (see Remark (4.1)). Then a consequence of the proposition (4.5) is

Corollary 4.7. *There is a subvariety $\mathcal{R} \subset \mathcal{M}_{g-2}$ of codimension at most 1 such that for any $\Gamma \in \mathcal{R}$ and for any $E_1, E_2 \in \overline{\mathcal{M}}_1$, there are curves $E_1 \cup_{p_1} \Gamma \cup_{p_2} E_2$ in \mathcal{L} .*


Observe that all these elements belong to Δ_1 and they belong to Δ_0 if and only if at least one of the elliptic curves represents the infinity class.

Remark 4.8. With the same techniques one can prove that the divisor $\overline{\mathcal{D}}$ contains irreducible curves with only one node. Moreover, by considering a surface in Δ_i , $i > 0$ given by fixed smooth curves of genus $g - i$ and i and moving the intersection point, one also shows with the same procedure that the divisor contain “generic” elements of Δ_i . We do not use these facts in the rest of the paper.

5. LIMITS OF ISOGENIES

As in section 2 we assume the existence of isogenies $JC' \rightarrow JC$ for generic elements $C \in \mathcal{D}$. In order to glue all these maps together to provide a family $\chi : \mathcal{J}' \rightarrow \mathcal{J}$ we need to pull-back to a suitable branched cover of our divisor. Since our calculations will be mainly of local nature we still denote this space of parameters as \mathcal{D} . Our goal is to get as much information as possible from the specialization of the family of isogenies to the curves of \mathcal{L} .

Let $L = E_1 \cup_{p_1} \Gamma \cup_{p_2} E_2 \in \mathcal{L} \subset \overline{\mathcal{D}}$ a fixed limit curve (see Remark (4.6) and Corollary (4.7)). We assume that one of the elliptic curves has a node.



Observe that $L \in \Delta_0 \cap \Delta_1$. Let $\overline{\mathcal{D}}_0$ be a component of $\overline{\mathcal{D}} \cap \Delta_0$ containing L . Since we can assume that the generic element of this component does not belong to Δ_i , $i \geq 2$, we obtain the following four cases:

- a) The generic element of $\overline{\mathcal{D}}_0$ is an irreducible curve with only one node.
- b) The generic element of $\overline{\mathcal{D}}_0$ consists of an irreducible curve with only one node with an elliptic curve attached in a smooth point.

- c) The generic element of $\overline{\mathcal{D}}_0$ consists of a smooth irreducible curve with an elliptic nodal curve attached in a smooth point.
- d) The generic element of $\overline{\mathcal{D}}_0$ consists of two irreducible smooth curves glued in two different points.
- e) The generic element of $\overline{\mathcal{D}}_0$ is an irreducible curve with two nodes.

For topological reasons the cases d) and e) can not occur. Indeed, the number of nodal points in a stable curve such that the curve remains connected when we remove the point can not decrease under specialization. Since there are two such points in a generic point of type d) and e) and only one in our limit curve L we can ignore these cases.

In this section we will specialize the isogeny χ to a generic curve of the component $\overline{\mathcal{D}}_0$ that contains our limit curve L .

Remark 5.1. We will use several times that the limit of a family of isogenies of Jacobians is also an isogeny, that is, an étale surjective map between the generalized Jacobians. We briefly give an argument to justify this. Consider two semistable flat families of curves parametrized by a disk $\pi' : \mathcal{C}' \rightarrow \mathbb{D}$ and $\pi : \mathcal{C} \rightarrow \mathbb{D}$ such that the curves are smooth away from the central fibres $\mathcal{C}'_0, \mathcal{C}_0$. Given a family of isogenies $\varphi : \mathcal{J}\mathcal{C}' \rightarrow \mathcal{J}\mathcal{C}$ over the punctured disk, the existence of a map $\varphi_0 : \mathcal{J}\mathcal{C}'_0 \rightarrow \mathcal{J}\mathcal{C}_0$ of semiabelian varieties is given in [BP], p. 267. We want to prove that this map is also an isogeny. Recall that the family of Jacobians is given by

$$\mathcal{J}\mathcal{C}' = R^1\pi'_*\mathcal{O}/R^1\pi'_*\mathbb{Z}$$

and analogously for $\mathcal{J}\mathcal{C}$. The fact that φ_t is an isogeny for $t \neq 0$ implies that there is a sequence over the punctured disk

$$0 \rightarrow R^1\pi'_*\mathbb{Z} \rightarrow R^1\pi_*\mathbb{Z} \rightarrow \tau \rightarrow 0,$$

where τ is a torsion local system. Hence tensoring with \mathbb{Q} we obtain an isomorphism

$$R^1\pi'_*\mathbb{Q} \cong R^1\pi_*\mathbb{Q}.$$

Since the rational cohomology of the central fibre can be identified with the invariant subspace by the action of the monodromy we have that:

$$H^1(\mathcal{C}'_0, \mathbb{Q}) \cong H^1(\mathcal{C}'_{|\mathbb{D}^*}, \mathbb{Q})^{\text{inv}} \cong H^0(\mathcal{C}'_{|\mathbb{D}^*}, R^1\pi'_*\mathbb{Q}),$$

and therefore we conclude that $H^1(\mathcal{C}'_0, \mathbb{Q}) \cong H^1(\mathcal{C}_0, \mathbb{Q})$. So the map φ_0 is an isogeny between the generalized Jacobians.

When we go to the limit the information we obtain is different according to the three cases explained above.

Case a). We consider the normalization map $p : \Delta_0 \dashrightarrow \mathcal{M}_{g-1}$ restricted to $\overline{\mathcal{D}}_0$. There are two possibilities according to the dimension of the generic fiber of $p_0 := p|_{\overline{\mathcal{D}}_0}$.

Case a.1): Assume that p_0 is dominant, therefore the generic fibre has dimension $\dim \overline{\mathcal{D}}_0 - \dim \mathcal{M}_{g-1} = 3g - 5 - 3g + 6 = 1$. For a generic element $t_0 \in \overline{\mathcal{D}}_0$ the limit map $\chi_{t_0} : \mathcal{J}\mathcal{C}'_{t_0} \rightarrow \mathcal{J}\mathcal{C}_{t_0}$ gives a diagram of extensions:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{C}^{*r} & \rightarrow & \mathcal{J}\mathcal{C}'_{t_0} & \rightarrow & \mathcal{J}\tilde{\mathcal{C}}'_{t_0} \rightarrow 0 \\ & & \downarrow \gamma & & \downarrow \chi_{t_0} & & \downarrow \tilde{\chi}_{t_0} \\ 0 & \rightarrow & \mathbb{C}^* & \rightarrow & \mathcal{J}\mathcal{C}_{t_0} & \rightarrow & \mathcal{J}\tilde{\mathcal{C}}_{t_0} \rightarrow 0 \end{array}$$

where \tilde{C}'_{t_0} and \tilde{C}_{t_0} stand for the normalizations of C'_{t_0} and C_{t_0} respectively. Since χ_{t_0} has finite kernel, r must be one and $\gamma(z) = z^m$ for some non-zero integer m . Since $g - 1 \geq 4$ we can apply the main result in [BP] and we get that $\tilde{C}'_{t_0} = \tilde{C}_{t_0}$ and the isogeny $\tilde{\chi}_{t_0}$ is n times the identity. Assume that C_{t_0} (resp. C'_{t_0}) is obtained from \tilde{C}_{t_0} by pinching two distinct points p, q (resp. p', q'). As in [BP], section 2, to compare the extension classes of each horizontal short exact sequence, we decompose the last diagram into

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{C}^* & \longrightarrow & JC'_{t_0} & \longrightarrow & J\tilde{C}'_{t_0} \longrightarrow 0 \\ & & \downarrow \gamma & & \downarrow \chi_{t_0} & & \parallel \\ 0 & \longrightarrow & \mathbb{C}^* & \longrightarrow & E & \longrightarrow & J\tilde{C}_{t_0} \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \tilde{\chi}_{t_0} \\ 0 & \longrightarrow & \mathbb{C}^* & \longrightarrow & JC_{t_0} & \longrightarrow & J\tilde{C}_{t_0} \longrightarrow 0. \end{array}$$

We identify (up to sign) the extension class

$$[JC'_{t_0}] \in Ext(J\tilde{C}_{t_0}, \mathbb{C}^*) \cong Pic^0(J\tilde{C}_{t_0}) \cong J\tilde{C}_{t_0}$$

with $p' - q'$ and analogously $[JC_{t_0}]$ with $p - q$. Then the equality $[E] = \gamma_*([JC'_{t_0}]) = \tilde{\chi}_{t_0}^*([JC_{t_0}])$ provides the following relation in $J\tilde{C}_{t_0}$:

$$n(p - q) = \pm m(p' - q').$$

Hence we can assume that $np + mq' = mp' + nq$ in $Pic(\tilde{C}_{t_0})$. We assume also that the points are different. Since the dimension of the generic fibre of p_0 is 1 we have a one dimensional family of maps $\tilde{C}_{t_0} \rightarrow \mathbb{P}^1$ of degree $n + m$ with two fibers as above. The Riemann-Hurwitz Theorem implies that

$$2g(\tilde{C}_{t_0}) - 2 = 2g - 4 = (n + m)(2g(\mathbb{P}^1) - 2) + 2(n - 1) + 2(m - 1) + r = -4 + r,$$

so the number r of the ramification points out of the special fibers $np + mq'$ and $mp' + nq$ is $r = 2g$. Then the Hurwitz scheme of maps of degree $n + m$ into \mathbb{P}^1 with $r + 2 = 2g + 2$ discriminant points must cover \mathcal{M}_{g-1} with generic fibers of dimension 1. Comparing dimensions:

$$2g + 2 - \dim \text{Aut}(\mathbb{P}^1) - \dim \text{generic fiber} = 2g + 2 - 4 = 2g - 2 \geq \dim \mathcal{M}_{g-1} = 3g - 6,$$

which contradicts the hypothesis $g \geq 5$. Hence we get that the extension is the same and $n = m$.

Case a.2): Assume that the generic fiber of p_0 has dimension 2. As before we get a diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{C}^* & \longrightarrow & JC'_{t_0} & \longrightarrow & J\tilde{C}'_{t_0} \longrightarrow 0 \\ & & \downarrow m & & \downarrow \chi_{t_0} & & \downarrow \tilde{\chi}_{t_0} \\ 0 & \longrightarrow & \mathbb{C}^* & \longrightarrow & JC_{t_0} & \longrightarrow & J\tilde{C}_{t_0} \longrightarrow 0 \end{array}$$

but now we do not have the genericity of \tilde{C}_{t_0} so we can not directly apply the main result in [BP]. The relation between extension classes is in this case

$$m(p' - q') = \tilde{\chi}_{t_0}^*(p - q),$$

in $J\tilde{C}'_{t_0}$. In other words, the isogeny $\tilde{\chi}_{t_0}$ induces a map between the surfaces $\tilde{C}'_{t_0} - \tilde{C}_{t_0}$ and $m(\tilde{C}'_{t_0} - \tilde{C}'_{t_0})$. By using the arguments of the section 3 in [BP] one easily checks that, as before, the curves are the same and the map is a multiplication by an integer. So we have proved the following result:

Proposition 5.2. *Let $\chi : \mathcal{J}' \rightarrow \mathcal{J}$ be a family of isogenies parametrized by \mathcal{D} and let $\chi_{t_0} : JC'_{t_0} \rightarrow JC_{t_0}$ be a specialization to a generic point t_0 of a component of the boundary $\overline{\mathcal{D}} \cap \Delta_0$, where the curve C_{t_0} is an irreducible curve with only one node. Then $C'_{t_0} \cong C_{t_0}$ and χ_{t_0} is the multiplication by a non-zero integer.*

Case b): We assume now that the limit curve L belongs to an irreducible component $\overline{\mathcal{D}}_0$ of $\overline{\mathcal{D}} \cap \Delta_0$ whose generic element consists in a smooth curve Γ of genus $g-1$ and a nodal elliptic curve E_∞ (i.e. a \mathbb{P}^1 with the points 0 and 1 identified) glued to Γ in a point $p \in \Gamma$. We denote as above by $C_{t_0} = \Gamma \cup_p E_\infty$ a generic curve in $\overline{\mathcal{D}}_0$. Observe that the natural map $\overline{\mathcal{D}}_0 \dashrightarrow \mathcal{M}_{g-1}$ must be dominant by a count of dimensions (the fibre has dimension at most 1), therefore we can assume that Γ is generic in \mathcal{M}_{g-1} .

We consider the specialization of the family of isogenies to our curve $\chi_{t_0} : JC'_{t_0} \rightarrow JC_{t_0} = J\Gamma \times \mathbb{C}^*$ which fits in a diagram of extensions:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{C}^{*r} & \rightarrow & JC'_{t_0} & \rightarrow & J\tilde{C}'_{t_0} \rightarrow 0 \\ & & \downarrow \gamma & & \downarrow \chi_{t_0} & & \downarrow \tilde{\chi}_{t_0} \\ 0 & \rightarrow & \mathbb{C}^* & \rightarrow & J\Gamma \times \mathbb{C}^* & \rightarrow & J\Gamma \rightarrow 0 \end{array}$$

where \tilde{C}'_{t_0} stands for the normalization of C'_{t_0} . Since χ_{t_0} has finite kernel, r must be one and $\gamma(z) = z^m$ for some non-zero integer m . Since $g-1 \geq 4$ we can apply the main result in [BP] and we get that $\tilde{C}'_{t_0} = \Gamma$ and the isogeny $\tilde{\chi}_{t_0}$ is n times the identity. So the diagram above becomes:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{C}^* & \rightarrow & JC'_{t_0} & \rightarrow & J\Gamma \rightarrow 0 \\ & & \downarrow m & & \downarrow \chi_{t_0} & & \downarrow n \\ 0 & \rightarrow & \mathbb{C}^* & \rightarrow & J\Gamma \times \mathbb{C}^* & \rightarrow & J\Gamma \rightarrow 0 \end{array}$$

Since the extension class of the first row corresponds to a generalized Jacobian, there exist points $q_1, q_2 \in \Gamma$ such that this class corresponds (up to sign) to $q_1 - q_2 \in J\Gamma$. Therefore, since the class of the second row is zero we get that $m(q_1 - q_2) = 0$.

Moving the point p in Γ we have a positive dimensional family of pairs of points $(q_1, q_2) \in C \times C$ with this property. This family has to be the diagonal since the map $\Gamma \times \Gamma \rightarrow J\Gamma$ has degree 1 and finite fibres out of the diagonal. Hence we obtain $q_1 = q_2$.

Hence the extension given by the first row is also trivial: $JC'_{t_0} \cong J\Gamma \times \mathbb{C}^*$ and $\chi_{t_0} = \begin{pmatrix} n & 0 \\ 0 & m \end{pmatrix}$. We get the following result:

Proposition 5.3. *Let $\chi : \mathcal{J}' \rightarrow \mathcal{J}$ be a family of isogenies parametrized by \mathcal{D} and let $\chi_{t_0} : JC'_{t_0} \rightarrow JC_{t_0}$ be a specialization to a generic point t_0 of a component of the boundary $\overline{\mathcal{D}} \cap \Delta_0$, where the curve C_{t_0} is a reducible curve consisting in a smooth curve of genus $g-1$ with a nodal elliptic curve E_∞ attached in one point. Then $C'_{t_0} \cong C_{t_0}$ and the isogeny in the compact part is the multiplication by a non zero integer.*

Case c): Finally we assume that the limit curve L belongs to an irreducible component $\overline{\mathcal{D}}_0$ of $\overline{\mathcal{D}} \cap \Delta_0$ whose generic element consists in a nodal curve Γ_0 of genus $g - 1$ and an elliptic curve E glued to Γ_0 in a smooth point $p \in \Gamma_0$. We denote as above by $C_{t_0} = \Gamma_0 \cup_p E$ a generic curve in $\overline{\mathcal{D}}_0$. Observe that the natural map $\overline{\mathcal{D}}_0 \dashrightarrow \Delta_0(\mathcal{M}_{g-1})$ must be dominant (here $\Delta_0(\mathcal{M}_{g-1})$ denotes the Δ_0 divisor in the moduli space \mathcal{M}_{g-1}). Indeed the generic fibre of this map has dimension at most 2 parametrized by the smooth point p in Γ_0 and the moduli of the elliptic curves. Since $\dim \overline{\mathcal{D}}_0 = 3g - 5$ and $\dim \Delta_0(\mathcal{M}_{g-1}) = 3g - 7$, the dominance follows. Therefore we can assume that Γ_0 is generic in $\Delta_0(\mathcal{M}_{g-1})$, and also its normalization $\tilde{\Gamma}_0$ is generic in \mathcal{M}_{g-2} . Moreover all the curves $\Gamma_0 \cup_p E$ are contained in $\overline{\mathcal{D}}_0$ for a generic Γ_0 .

As in the other cases we consider the specialization of the family of isogenies to our curve $\chi_{t_0} : JC'_{t_0} \rightarrow JC_{t_0} = J\Gamma_0 \times E$ which fits in a diagram of extensions:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{C}^{*r} & \rightarrow & JC'_{t_0} & \rightarrow & J\tilde{C}'_{t_0} \rightarrow 0 \\ & & \downarrow \gamma & & \downarrow \chi_{t_0} & & \downarrow \tilde{\chi}_{t_0} \\ 0 & \rightarrow & \mathbb{C}^* & \rightarrow & J\Gamma_0 \times E & \rightarrow & J\tilde{\Gamma}_0 \times E \rightarrow 0 \end{array}$$

where \tilde{C}'_{t_0} stands for the normalization of C'_{t_0} . Since χ_{t_0} has finite kernel, r must be one and $\gamma(z) = z^m$ for some non-zero integer m .

We claim that $J\tilde{C}'_{t_0}$ must be a product of Jacobians (in other words, the smooth curve \tilde{C}'_{t_0} is reducible). We prove this by contradiction, assume that \tilde{C}'_{t_0} is irreducible and compare the extension classes. This gives (up to sign) a relation $m(p' - q') = \tilde{\chi}_{t_0}(p - q)$ in $J\tilde{C}'_{t_0}$. Moving the points p and q in the fixed curve $\tilde{\Gamma}_0$ we get that the image of $\tilde{\Gamma}_0 - \tilde{\Gamma}_0$ by the isogeny is the surface $\tilde{C}'_{t_0} - \tilde{C}'_{t_0}$ which is impossible since it is contained in the proper abelian subvariety $\tilde{\chi}_{t_0}^*(J\tilde{\Gamma}_0)$.

By the genericity of $\tilde{\Gamma}_0$ we can assume that $J\tilde{\Gamma}_0$ is simple and then $J\tilde{C}'_{t_0} \cong J\tilde{C}''_{t_0} \times E'$, where \tilde{C}''_{t_0} is an irreducible curve of genus $g - 2$ and E' stands for a smooth elliptic curve. So C'_{t_0} is a nodal curve C''_{t_0} with the elliptic curve attached in a smooth point. The extension class is the difference of two points $p'', q'' \in \tilde{C}''_{t_0}$. The diagram above becomes:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{C}^* & \rightarrow & JC''_{t_0} \times E' & \rightarrow & J\tilde{C}''_{t_0} \times E' \rightarrow 0 \\ & & \downarrow m & & \downarrow \chi_{t_0} & & \downarrow \tilde{\chi}_{t_0} = \tilde{\chi}''_{t_0} \times \varphi \\ 0 & \rightarrow & \mathbb{C}^* & \rightarrow & J\Gamma_0 \times E & \rightarrow & J\tilde{\Gamma}_0 \times E \rightarrow 0, \end{array}$$

where $\varphi : E' \rightarrow E$ is a non-constant map of elliptic curves. The relation between extension classes is in this case (up to sign):

$$m(p'' - q'') = (\tilde{\chi}''_{t_0})^*(p - q),$$

in $J\tilde{C}''_{t_0}$. In other words, the isogeny $\tilde{\chi}''_{t_0}$ induces a map between the surfaces $\tilde{\Gamma}_0 - \tilde{\Gamma}_0$ and $m(\tilde{C}''_{t_0} - \tilde{C}''_{t_0})$. By using the arguments of Section 3 in [BP] one easily checks the following facts: the curves C''_{t_0} and Γ_0 are isomorphic, the isogeny $\tilde{\chi}''_{t_0}$ is a non-zero multiple n of the identity, $n = m$ and then $\chi_{t_0} = n \times \varphi$.

Proposition 5.4. *Let $\chi : \mathcal{J}' \rightarrow \mathcal{J}$ be a family of isogenies parametrized by \mathcal{D} and let $\chi_{t_0} : JC'_{t_0} \rightarrow JC_{t_0}$ be a specialization to a generic point t_0 of a component of*

the boundary $\overline{\mathcal{D}} \cap \Delta_0$, where the curve C_{t_0} is a reducible curve consisting in a nodal curve Γ_0 of genus $g-1$ with an elliptic curve E attached in one smooth point, then also C'_{t_0} is of the form $\Gamma_0 \cup_p E'$, for some elliptic curve E' , and the isogeny induces the multiplication by n on $J\Gamma_0$.

6. END OF THE PROOF

Let us go back to our family of isogenies $\chi : \mathcal{J}' \rightarrow \mathcal{J}$ parametrized by (some covering of) the divisor \mathcal{D} .

Consider a generic point $t \in \mathcal{D}$ corresponding to smooth curves C'_t and C_t . Observe that for all t the isogeny is determined by the map at the level of homology groups

$$\chi_{t,\mathbb{Z}} : H_1(C'_t, \mathbb{Z}) \rightarrow H_1(C_t, \mathbb{Z})$$

which we still denote by χ_t . We set $\Lambda_t \subset H_1(C_t, \mathbb{Z})$ for the image of χ_t . This is a sublattice of maximal rank $2g$. We first note that the proof of Proposition (4.2.1) in [BP] applies verbatim to obtain the following result:

Proposition 6.1. *Assume that $\Lambda_t = nH_1(C_t, \mathbb{Z})$ for some positive integer n . Then $C'_t \cong C_t$ and χ_t is multiplication by n .*

Therefore to finish the proof of the Theorem we have to show the equality $\Lambda_t = nH_1(C_t, \mathbb{Z})$. To do this, the main idea is to get information on Λ_t from the homology groups of some convenient limits C_0 . In the previous sections we have shown the existence of certain limits and we have seen in Propositions (5.2), (5.3) and (5.4) how is the image of the limit of χ_t when t goes to one of these degenerations.

To pass information from some C_0 to the smooth point we use the following principle: We can assume that there exists a disc $\mathbb{D} \subset \overline{\mathcal{D}} \subset \overline{\mathcal{M}}_g$ centered at the class of the curve C_0 such that the curves C_t, C'_t corresponding to $\mathbb{D} \setminus \{0\}$ are smooth. After performing a base change (that we skip to simplify the notation) we can assume that there is a family of isogenies $\chi_{\mathbb{D}} : \mathcal{J}'_{\mathbb{D}} \rightarrow \mathcal{J}_{\mathbb{D}}$ that coincides with the original isogeny χ_t for a generic t . We denote $\chi_0 : JC'_0 \rightarrow JC_0$ the limit isogeny. We call $\mathcal{C}'_{\mathbb{D}}$ and $\mathcal{C}_{\mathbb{D}}$ the corresponding families of curves. Since the central fibres C'_0 and C_0 are retracts of $\mathcal{C}'_{\mathbb{D}}$ and $\mathcal{C}_{\mathbb{D}}$ respectively, we have a diagram as follows:

$$\begin{array}{ccc} H_1(C'_t, \mathbb{Z}) & \twoheadrightarrow & H_1(\mathcal{C}'_{\mathbb{D}}, \mathbb{Z}) = H_1(C'_0, \mathbb{Z}) \\ \downarrow \chi_t & & \downarrow \chi_0 \\ H_1(C_t, \mathbb{Z}) & \twoheadrightarrow & H_1(\mathcal{C}_{\mathbb{D}}, \mathbb{Z}) = H_1(C_0, \mathbb{Z}). \end{array}$$

Let us consider first the simplest case: assume that we are in the case a) and that C_0 is a generic element of $\overline{\mathcal{D}} \cap \Delta_0$, so it is an irreducible curve with only one node. Remember that C_0 can be degenerated in this component to a curve L consisting of a curve of genus $g-2$ with two attached curves, one elliptic (called E_2) and the other nodal and rational (E_{∞}). By Proposition (5.2) $C'_0 \cong C_0$ and $\chi_0 = n \cdot \text{Id}$. The kernel of the horizontal maps are generated by the vanishing cycles a'_1 and a_1 respectively, therefore $\chi_t(a'_1) = la_1$ for some l . On the other hand we can lift a basis in $H_1(C_0, \mathbb{Z})$ and construct symplectic bases $a'_1, b'_1, \dots, a'_g, b'_g$ in $H_1(C'_t, \mathbb{Z})$ and $a_1, b_1, \dots, a_g, b_g$ in $H_1(C_t, \mathbb{Z})$ in such a way that:

$$\chi_s(a'_i) = na_i + s_i a_1, \quad \chi_t(b'_1) = nb_1 + t_1 a_1, \quad \chi_t(b'_i) = nb_i + t_i a_1$$

for some integers s_i and t_i and $i \geq 2$. We also can assume that a_g, b_g correspond to cycles which become a basis of the homology of $E_2 \subset L$.

By the genericity of the curve C_t in a divisor of the moduli space the pull-back of the theta divisor is a multiple of the theta divisor in JC'_t . This translates into the existence of a non-zero integer m such that the cup-product satisfies

$$\chi_t(x) \cup \chi_t(y) = mx \cup y.$$

Then we obtain

$$\begin{aligned} m &= \chi_t(a'_1) \cup \chi_t(b'_1) = la_1 \cup (nb_1 + t_1a_1) = ln \\ m &= \chi_t(a'_2) \cup \chi_t(b'_2) = n^2, \end{aligned}$$

so $n = l$. With similar computations it is easy to prove that $s_i = t_i = 0$ for $i \geq 2$, hence:

$$(4) \quad \chi_t(a'_1) = na_1, \quad \chi_t(a'_i) = na_i, \quad \chi_t(b'_i) = nb_i \quad \text{for } i \geq 2.$$

To get the piece of information which is still unknown we need to consider a second limit. We have seen in Corollary (4.7) that fixing Γ we can move freely the two elliptic curves E_1, E_2 , that is $E_1 \cup \Gamma \cup E_2 \in \overline{\mathcal{D}}$ for all E_1, E_2 . We select a second limit curve $\hat{L} = E_1 \cup \Gamma \cup E_\infty$ in such a way that the corresponding vanishing cycle is now a_g . Again, to simplify, we assume that \hat{L} belongs to the case a) of section 5. Then, with the same argument, we get that χ_t satisfies (for the same symplectic basis):

$$\chi_t(a'_i) = \hat{n}a_i, \quad \chi_t(b'_i) = \hat{n}b_i \quad \text{for } i \leq g-1$$

and $\chi_t(a'_g) = \hat{n}a_g$. Therefore $\hat{n} = n$ and $\Lambda_t = nH_1(C_t, \mathbb{Z})$ for all $s \neq 0$. Hence we have finished (under the assumption that L and \hat{L} are limit curves of nodal curves). The rest of the proof consists in the description of the small modifications that have to be done to take care of the rest of the cases. Observe that the information on the limit given in the case b) (see Proposition (5.3)) is the same as that given in the case a), that is we have again the relations (4). So we only have to take care of the situation when at least one of the limit curves L, \hat{L} belong to the case c). Assume for example that L does. Using the first limit as above we know that

$$\chi_t(a'_1) = na_1, \quad \chi_t(a'_i) = na_i, \quad \chi_t(b'_i) = nb_i \quad \text{for } 2 \leq i \leq g-1.$$

The difference with the previous cases is that we have no control on $\chi_t(b'_1)$. Remember that L is the limit of a curve $\Gamma_0 \cup E$ where Γ_0 is a nodal curve intersecting the elliptic curve E in a smooth point. Denoting $\tilde{\Gamma}_0$ the normalization of Γ_0 we note that all the nodal curves with this normalization belong to the divisor $\overline{\mathcal{D}}_0$. So we change freely the node in such a way that the vanishing cycle of the node becomes a_2 (instead of a_1). By using the two vanishing cycles we get that $\chi_t(b'_1) = nb'_1$ so we recover again the relations (4). This finishes the proof of the theorem. \square

Our theorem can be interpreted as a type of Noether-Lefschetz problem in the following way: consider in $\mathcal{M}_g \times \mathcal{M}_g$ the set

$$\mathcal{NL}_g = \{(C', C) \mid \text{rank } NS(C \times C') \geq 3\}.$$

A consequence of what we have proved is the following result:

Corollary 6.2. *For $g \geq 5$ all the components of \mathcal{NL}_g outside the diagonal have dimension less than or equal to $3g - 5$.*

The first natural problem one could face in this context is to investigate the existence of dimension 10 components in \mathcal{NL}_5 . Similar problems on isogenies can be considered for other families of abelian varieties (see for example [NP]).

REFERENCES

- [ACGH] E. Arbarello, M. Cornalba, P. A. Griffiths, J. Harris: Geometry of algebraic curves. Vol. I. Grundlehren der Mathematischen Wissenschaften, 267. Springer-Verlag, New York, 1985.
- [ACG] E. Arbarello, M. Cornalba, P. A. Griffiths: Geometry of algebraic curves. Vol. II. Grundlehren der Mathematischen Wissenschaften, 268. Springer-Verlag, New York, 2011.
- [BP] F. Bardelli, G.P. Pirola: Curves of genus g lying on a g -dimensional Jacobian variety, Invent. math., 95 (1989), pp. 263–276.
- [CM] M. Coppens, G. Martens: Secant spaces and Clifford’s theorem, Compositio Mathematica, 78 (1991), pp. 193–212.
- [CGGH] J. Carlson, M. Green, P. Griffiths, M. Cornalba: Infinitesimal variations of Hodge structure. I, Compositio Math. 50 (1983), 109205.
- [CGT] C. Ciliberto, G. van der Geer, M. Teixidor: On the number of parameters of curves whose Jacobians possess nontrivial endomorphisms, J. Algebraic Geom. 1 (1992), 215–229.
- [M] D. Mumford: Abelian Varieties. Oxford University Press, Bombay, 1974.
- [NP] J.C. Naranjo and G.P. Pirola: On the genus of curves in the generic Prym variety, Indag. Math. (N.S.) 5 (1994), 101–105.
- [P] G.P. Pirola: Base number theorem for abelian varieties. An infinitesimal approach, Math. Ann. 282 (1988), 361–368.
- [R] Z. Ran: On a Theorem of Martens, Rend. Sem. Mat. Univ. Politec. Torino, 44 (1986), pp. 287–291.

VALERIA MARCUCCI, UNIVERSITÀ DEGLI STUDI DI PAVIA, DIPARTIMENTO DI MATEMATICA, VIA FERRATA, 1, 27100 PAVIA, ITALY
E-mail address: `valeria.marcucci@unipv.it`

JUAN CARLOS NARANJO, UNIVERSITAT DE BARCELONA, FACULTAT DE MATEMÀTIQUES, GRAN VIA 585, 08007 BARCELONA, SPAIN
E-mail address: `jcnaranjo@ub.edu`

GIAN PIETRO PIROLA, UNIVERSITÀ DEGLI STUDI DI PAVIA, DIPARTIMENTO DI MATEMATICA, VIA FERRATA, 1, 27100 PAVIA, ITALY
E-mail address: `gianpietro.pirola@unipv.it`

This figure "geogebra_thumbnail.png" is available in "png" format from:

<http://arxiv.org/ps/1405.2545v2>