

ON SMALL ABSTRACT QUOTIENTS OF LIE GROUPS AND LOCALLY COMPACT GROUPS

LINUS KRAMER

Dem Andenken von Günter Pickert gewidmet.

1. INTRODUCTION

Suppose that G is a locally compact group, that Γ is a discrete, finitely generated group, and that

$$\varphi : G \longrightarrow \Gamma$$

is an 'abstract' surjective homomorphism. We are interested in conditions which imply that φ is automatically continuous. We obtain a complete answer to this question in the case where G is a topologically finitely generated locally compact abelian group or an almost connected Lie group. In these two cases the well-known structure theory for such groups G leads quickly to a solution. The question becomes much more difficult if one assumes only that G is a locally compact group. This leads to interesting questions about normal subgroups in infinite products and in ultraproducts. Los' Theorem, the solution of the 5th Hilbert Problem, and recent results by Nikolov–Segal can be combined to answer the question.

The topic of this article is thus of a rather group-theoretic nature. However, locally compact transformation groups are ubiquitous in geometry. Conversely, several proofs in this article rely on geometry, at different levels of abstraction. The 'bootstrap Lemma' 6.3 is based on a little exercise in spherical geometry, which was still taught at German high schools in Günter Pickert's early years. But we also use some observations about euclidean and spherical buildings.

The article is organized as follows. In Section 2 we consider small abstract quotients of locally compact abelian groups. In Section 3 we do the same for Lie groups. Section 4 then extends these results to compact and to locally compact groups. In Section 5 we make some observations about normal subgroups in infinite products. This leads ultimately to questions about normal subgroups in ultraproducts. We adopt a sheaf-theoretic viewpoint which is elementary but useful. Section 6 gives a detailed proof for the nonexistence of countable quotients of compact connected perfect groups, following Nikolov–Segal.

1991 *Mathematics Subject Classification.* 51H, 22D.
Supported by SFB 878.

The necessary background material on Lie groups and locally compact groups can be found in the excellent books by Hilgert–Neeb [13], Stroppel [31], and Hofmann–Morris [17], [16]. As the referee pointed out, it is likely that many of our results can be extended to Pro-Lie groups, using the structure theory developed in [16]. At some places, this is straight-forward, as in Corollary 2.3, or in Lemma 4.1, which holds verbatim for Pro-Lie groups. The generalization of other results in this article would seem to require new ideas.

1.1. Notation and conventions. The cardinality of a set S is written as $\#S$. We call a set S *countable* if $\#S \leq \aleph_0$. Throughout this article, all topological groups and spaces are assumed to be Hausdorff, unless stated otherwise. The identity component in a topological group G is denoted by G° . This is a closed normal subgroup of G . We say that a topological group G is *topologically finitely generated* if G has a finitely generated dense subgroup.

We call a quotient G/N of a topological group G *abstract* if no topological assumptions like closedness of the subgroup N in G are made. Similarly, we call a homomorphism between topological groups *abstract* if no continuity assumptions are made.

We frequently use the fact that a subgroup of finite index in a finitely generated group is also finitely generated. It suffices to prove this for finitely generated free groups, where it follows from the Nielsen–Schreier Theorem [27, 6.1.1].

A group Γ is called *residually finite* if the intersection of all normal subgroups of finite index in Γ is trivial. Equivalently, Γ embeds in a product of finite groups. Every finitely generated abelian group is residually finite, and subgroups of residually finite groups are again residually finite.

A group G is *virtually 'X'* (where 'X' is a group-theoretic property) if G has a finite index subgroup which has property 'X'.¹

A group element g is called *divisible* if the equation $x^n = g$ has a solution in the given group for every integer $n \geq 1$. A group is called *divisible* if it consists of divisible elements, and *divisibly generated* if it is generated by a set of divisible elements. Divisibility will play an important role in some proofs. The following group-theoretic facts illustrate some of the consequences and non-consequences of divisibility.

1.2. Some facts about divisibility in groups.

- (i) *Homomorphisms preserve divisibility.*
- (ii) *A group of finite exponent (e.g. a finite group) contains no divisible elements besides the identity. Hence a residually finite group contains no divisible elements besides the identity. In particular, a finitely generated abelian group contains no divisible elements besides the identity.*
- (iii) *There exist finitely generated divisible groups. Moreover, every countable group (e.g. the additive group \mathbb{Q}) is isomorphic to a subgroup of some 2-generated group.*

¹Finite group theory is therefore concerned with virtually trivial groups—this indicates the drawbacks of unrestricted virtualization.

(iv) *A connected locally compact group is divisibly generated.*

Proof. Claim (i) is obvious. For (ii) we note that if a group H has finite exponent $m > 1$ and if $h \in H$ is different from the identity, then the equation $x^m = h$ cannot be solved in H . The claims (iii) and (iv) are much deeper. For (iii), see [10] and [12, Thm. IV] or [27, 6.4.7]. Claim (iv) holds for all connected Pro-Lie groups, see [16, Lem. 9.52]. Every locally compact group is a Pro-Lie group by the Approximation Theorem due to Gleason, Montgomery–Zippin, and Yamabe [23, p. 175]. \square

When we are concerned with continuity questions, we view every countable group Γ as a discrete locally compact group. This is justified by the following elementary fact.

Lemma 1.3. *A locally compact countable group is discrete.*

Proof. By the Baire Category Theorem, some and hence every singleton $\{g\} \subseteq G$ is open, see [6, XI.10]. \square

2. THE CASE WHERE G IS LOCALLY COMPACT ABELIAN

For locally compact abelian groups we have the following general result. Note that a finitely generated abelian group is residually finite.

Theorem 2.1. *Suppose that G is a locally compact abelian group, that G/G° is topologically finitely generated, and that Γ is a countable residually finite group. Suppose that*

$$\varphi : G \longrightarrow \Gamma$$

is an abstract surjective group homomorphism. Then φ is continuous and open.

Proof. We proceed in four steps. The steps 1) and 2) below follow loosely the arguments in [24, Thm. 5.1].

0) *We may assume that $G^\circ = \{1\}$ and thus that G is topologically finitely generated.*

Since G° is divisible and since a residually finite group contains no nontrivial divisible elements (see 1.2), φ is constant on G° . Hence φ factors through the continuous open homomorphism $G \longrightarrow G/G^\circ$.

1) *The claim is true if G is compact and if Γ is finite.*

Let $F \subseteq G$ be a finitely generated dense subgroup and let m denote the exponent of Γ . Then $G^m = \{g^m \mid g \in G\} \subseteq G$ is a compact subgroup and the compact abelian group G/G^m has finite exponent. The image of F in G/G^m is finitely generated abelian of finite exponent and therefore finite. On the other hand, F has dense image in G/G^m . Therefore G/G^m is finite and thus G^m is open in G . Now G^m is contained in the kernel of φ , hence $\ker(\varphi)$ is both open and closed. Therefore φ is continuous and open.

2) *The claim is true if G is compact.*

Let \mathcal{N} denote the collection of all normal subgroups of finite index in Γ . For $\Delta \in \mathcal{N}$ let φ_Δ denote the composite $G \longrightarrow \Gamma \longrightarrow \Gamma/\Delta$. By Step 1), $\ker(\varphi_\Delta)$ is closed. Hence

$\ker(\varphi) = \bigcap \{\ker(\varphi_\Delta) \mid \Delta \in \mathcal{N}\}$ is closed. Therefore $G/\ker(\varphi)$ is a countable locally compact group. By Lemma 1.3, $G/\ker(\varphi)$ is discrete and therefore $\ker(\varphi)$ is open and φ is continuous and open.

3) *The claim is true in general.*

We may decompose the locally compact abelian group G topologically as $G = \mathbb{R}^n \times H$, where H has a compact open subgroup K , see [17, Thm. 7.57 (i)]. However, $G^\circ = \{1\}$ by Step 0), whence $G = H$ has a compact open subgroup K . Let $\Gamma_0 = \varphi(K)$. Then $\Gamma_0 \subseteq \Gamma$ is residually finite and countable. The restriction $\varphi : K \rightarrow \Gamma_0$ is continuous and open by Step 2). Since $K \subseteq H$ is open, $\varphi : H \rightarrow \Gamma$ is continuous and open. \square

The following examples show that the assumptions on G and Γ cannot be dropped.

Example 2.2. As an abstract group, the 1-torus $U(1) \cong \mathbb{R}/\mathbb{Z}$ is isomorphic to a product of Prüfer groups and a rational vector space of uncountable dimension, see e.g. [17, A1.43]. In particular, there exist (many) non-continuous surjective homomorphisms $U(1) \rightarrow \mathbb{Q}$. Hence the condition that Γ is residually finite cannot be dropped.

The compact group $G = \prod_{i=0}^{\infty} \mathbb{F}_2$ admits many non-continuous surjective homomorphisms onto \mathbb{F}_2 (where \mathbb{F}_2 denotes the field of two elements). Hence the condition that G/G° is topologically finitely generated can also not be dropped.

On the other hand, it is easy to extend the previous result to Pro-Lie groups.

Corollary 2.3. *Suppose that G is an abelian Pro-Lie group, that G/G° is locally compact and topologically finitely generated, and that Γ is a countable residually finite group. Suppose that*

$$\varphi : G \rightarrow \Gamma$$

is an abstract surjective group homomorphism. Then φ is continuous and open.

Proof. The identity component G° is a connected Pro-Lie group [16, 3.29], and therefore divisible [16, Lem. 9.52]. Hence φ factors through the open homomorphism $G \rightarrow G/G^\circ$. The claim follows now from Theorem 2.1. \square

3. THE CASE WHERE G IS AN ALMOST CONNECTED LIE GROUP

Now we consider an abstract surjective group homomorphism

$$G \rightarrow \Gamma,$$

where G is an almost connected² Lie group and Γ is a finitely generated group. The next result was observed (in different degrees of generality) by Goto [9], Ragozin [26] and George Michael [7]. See also [17, Thm. 9.90] and [28, 94.21]. For the sake of completeness, we include a proof.

²i.e. G/G° is finite

Theorem 3.1. *Let G be a Lie group. If the Lie algebra $\text{Lie}(G)$ is semisimple, then every abstract normal subgroup $N \trianglelefteq G$ is closed.*

Proof. Let $N \trianglelefteq G$ be a normal subgroup. Since $G^\circ \subseteq G$ is closed and open, it suffices to show that $N \cap G^\circ$ is closed in G° in order to show that N is closed. Hence we may assume that G is connected, and we proceed by induction on the dimension of the compact connected semisimple Lie group G . The claim holds for trivial reasons if $\dim(G) = 0$. For $\dim(G) > 0$ we view $N \subseteq G$ as a topological group with respect to the subspace topology. Let $N_1 \subseteq N$ denote the path component of the identity.

If $N_1 = \{1\}$, then the path component of every element $n \in N$ is trivial. Then $\{gng^{-1} \mid g \in G\} = \{n\}$ holds for every $n \in N$. Thus N is contained in the closed discrete group $\text{Cen}(G) \subseteq G$ and therefore N is closed.

If N_1 is nontrivial then there exists, by Yamabe's Theorem (see [9] or [13, Thm. 9.6.1]), a Lie group structure on N_1 such that $N_1 \hookrightarrow G$ is a continuous injection. In other words, $N_1 \hookrightarrow G$ is a nontrivial connected analytic/virtual/integral Lie subgroup of G . Its Lie algebra $\text{Lie}(N_1)$ is then a nontrivial ideal in $\text{Lie}(G)$. Since $\text{Lie}(G)$ is semisimple, the virtual subgroup corresponding to any ideal in $\text{Lie}(G)$ is closed: it is the connected centralizer of the complementary ideal. Therefore $N_1 \subseteq G$ is closed and thus a closed Lie subgroup of G . Now we may apply the induction hypothesis to $N/N_1 \subseteq G/N_1$. The group N is the preimage of N/N_1 under the continuous homomorphism $G \rightarrow G/N_1$ and therefore closed in G . \square

Corollary 3.2. *If G is a connected Lie group whose Lie algebra is simple, then every proper abstract normal subgroup of G is contained in the center of G . In particular, $G/\text{Cen}(G)$ is simple as an abstract group.* \square

Corollary 3.3. *A connected semisimple Lie group has no proper abstract normal subgroups of countable index.* \square

Now we come to the main result of this section. A variation is proved in Proposition 4.3.

Theorem 3.4. *Let G be a Lie group, with G/G° finite. Let Γ be a finitely generated group and let*

$$\varphi : G \longrightarrow \Gamma$$

be an abstract surjective homomorphism. Then Γ is finite and φ is continuous and open.

Proof. The subgroup $\varphi(G^\circ) \subseteq \Gamma$ has finite index and is therefore finitely generated. Hence it suffices to prove that φ is constant if G is connected.

Suppose that G is connected. Let $R \trianglelefteq G$ denote the solvable radical, i.e. the unique maximal connected solvable closed normal subgroup of G , see [13, Prop. 16.2.2.] and [28, Def. 94.18, Prop. 94.19]. Then the Lie algebra $\text{Lie}(G/R)$ is semisimple. We put $N = \ker(\varphi)$. By Theorem 3.1, $NR \subseteq G$ is closed and therefore a Lie subgroup. Since $\Gamma \cong G/N$ is countable, G/NR is a countable and hence discrete Lie group. It follows that $NR \subseteq G$ is open, hence $G = NR$. Now we have $\Gamma \cong NR/N \cong R/R \cap N$. Thus Γ is

solvable. The connected Lie group R is divisibly generated (see 1.2). It follows that the abelianization Γ_{ab} of Γ is both divisible and finitely generated abelian, whence $\Gamma_{ab} = \{1\}$ (again by 1.2). On the other hand, Γ is solvable, hence $\Gamma_{ab} \neq \{1\}$ if $\Gamma \neq \{1\}$. Hence $\Gamma = \{1\}$. \square

Problem 3.5. Is the conclusion of Theorem 3.4 still true if we drop the assumption that G/G° is finite?

4. QUOTIENTS OF LOCALLY COMPACT GROUPS

We now extend the results from the previous sections to locally compact groups. Here the fundamental work of Nikolov–Segal [24], as well as the solution of the 5th Hilbert problem, enter in an essential way. But first we draw a simple consequence from the facts collected in 1.2.

Lemma 4.1. *Let G be a connected locally compact group and let E be a group of finite exponent (e.g. a finite group), or a residually finite group. Then every abstract homomorphism $G \rightarrow E$ is trivial. In particular, a connected locally compact group has no abstract proper subgroups of finite index, and no nontrivial residually finite quotients.*

Proof. By 1.2, the group G is divisibly generated, and E contains no divisible elements. Hence every homomorphism $G \rightarrow E$ is constant. For the last assertion we note that a group G which has a proper finite index subgroup H maps nontrivially into the finite symmetric group $\text{Sym}(G/H)$. Hence the existence of proper finite index subgroups implies the existence of proper finite index normal subgroups. \square

The situation is quite different for non-normal subgroups of countable index: in connected Lie groups, such groups always exist.

Theorem 4.2. *The group $\text{SL}_n\mathbb{C}$ has subgroups of infinite countable index. Therefore every connected Lie group G has subgroups of infinite countable index.*

Proof. This is proved in [32, Thm. 2.1] and in [20, Thm. 1]. The algebraic-geometric reason is the following. Let p be a fixed prime. The group $\text{SL}_n\mathbb{Q}_p$ acts on its Bruhat–Tits building X , which has a countable set of vertices. Thus $\text{SL}_n\mathbb{Q}_p$ has subgroups of countable index, namely the parahoric subgroups.

Now let K denote the algebraic closure of \mathbb{Q}_p . The discrete valuation on \mathbb{Q}_p extends to a nondiscrete valuation on K , and there is a corresponding nondiscrete euclidean building X_K on which SL_nK acts. This building is not simplicial any more, but nevertheless it has a countable invariant subset of ‘vertices’. Thus SL_nK has subgroups of countable index.

Next we note that all algebraically closed fields of characteristic 0 and cardinality 2^{\aleph_0} are isomorphic to \mathbb{C} . Thus $\text{SL}_n\mathbb{C} \cong \text{SL}_nK$ has abstract subgroups of countable index.

Finally, every connected Lie group G admits a nontrivial representation $G \rightarrow \text{SL}_n\mathbb{C}$, for some n . Thus G has nontrivial abstract subgroups of countable index as well, and the index cannot be finite by Lemma 4.1. \square

More generally, one may look for subgroups in a locally compact group G which are not Borel sets, or even not measurable with respect to the Haar measure. Note that the σ -additivity of the Haar measure implies that subgroups of countable index cannot be measurable, so the subgroups appearing in Theorem 4.2 cannot be measurable. The existence of non-measurable subgroups in general compact groups is studied in detail in a recent article by Hernández–Hofmann–Morris [11].

In contrast to Theorem 4.2, we have the following result for *normal* subgroups of countable index. For the next results in this section, ‘countable’ may be replaced by ‘of cardinality strictly less than 2^{\aleph_0} ’ without changing any of the arguments.

Proposition 4.3. *Let G be a Lie group, with G/G° finite. Suppose that $N \trianglelefteq G$ is an abstract normal subgroup of countable index. Then G/N is virtually solvable. If G is connected, then G/N is solvable.*

Proof. Since the image of G° in G/N has finite index, it suffices to show that the image of G° in G/N is solvable. So we may assume that G is connected. As in the proof of Theorem 3.4, let $R \trianglelefteq G$ denote the solvable radical of G . Then NR/R is a normal subgroup of countable index in the semisimple Lie group G/R . By Corollary 3.3 we have $NR = G$. Thus $G/N = NR/R \cong R/N \cap R$ is solvable. \square

In order to extend this result to general locally compact groups, a crucial ingredient is Iwasawa’s splitting theorem. The global version of this result is an extremely useful answer to Hilbert’s 5th problem.

Theorem 4.4 (The Global Splitting Theorem). *Let G be a locally compact group and let $U \subseteq G$ be a neighborhood of the identity. Then there is a compact subgroup $K \subseteq G$ contained in U , a simply connected Lie group L , and an open and continuous homomorphism*

$$\psi : L \times K \longrightarrow G$$

with discrete kernel such that $\psi(1, k) = k$ for all $k \in K$.

Proof. The local version of this result, assuming G to be connected and approximable by Lie groups, is due to Iwasawa [18]. See also Gluškov [8]. The present version (and some more remarks on the history) can be found in Hofmann–Morris [15, Thm. 4.1] and in [14, Thm. 4.4]. \square

The next result by Nikolov–Segal is considerably more difficult to prove than Proposition 4.3. A proof will be given in Section 6 below.

Theorem 4.5 (Nikolov–Segal). *Let G be a compact group, with G/G° finite. Suppose that $N \trianglelefteq G$ is an abstract normal subgroup of countable index. Then G/N is virtually abelian. If G is connected, then G/N is abelian. If G/N is residually finite, then N is closed.*

Our final result about quotients of locally compact groups is as follows.

Theorem 4.6. *Let G be a locally compact group, with G/G° compact. Suppose that Γ is a finitely generated group and that*

$$\varphi : G \longrightarrow \Gamma$$

is an abstract surjective homomorphism. Then Γ is finite. If G/G° is topologically finitely generated, then φ is continuous and open.

Proof. Let $\Gamma_0 = \varphi(G^\circ)$ and $N = \ker(\varphi)$. Since G/G° is compact and since G/NG° is finitely generated, the main result [24, Thm. 5.25] implies that $G/NG^\circ \cong \Gamma/\Gamma_0$ is finite. Thus Γ_0 is also finitely generated. We put $\psi : L \times K \longrightarrow G^\circ$ as in Theorem 4.4 and we note that this homomorphism is necessarily surjective. We consider the composite

$$\begin{array}{ccc} L \times K & \xrightarrow{\psi} & G^\circ \\ & \searrow \varphi & \downarrow \varphi \\ & & \Gamma_0. \end{array}$$

The connected Lie group $L \times \{1\}$ maps onto the finitely generated group $\Gamma_0/\bar{\varphi}(\{1\} \times K)$. By Theorem 3.4, we have $\bar{\varphi}(\{1\} \times K) = \Gamma_0$. In particular, K maps onto the finitely generated group Γ_0 . Again by [24, Thm. 5.25], the quotient Γ_0 is finite. Since G° is divisibly generated (see 1.2), the group Γ_0 is trivial and Γ is finite. Moreover, φ factors through the projection $G \longrightarrow G/G^\circ$. Again by [24, Thm. 5.25], the composite $G \longrightarrow G/G^\circ \longrightarrow \Gamma$ is continuous if G/G° is topologically finitely generated. \square

The previous proof relies heavily on Nikolov–Segal [24, Thm. 5.25], which is stronger than Theorem 4.5 above. This result depends, among other things, on the classification of the finite simple groups, and on their advanced structure theory.

5. PRODUCTS AND ULTRAPRODUCTS OF NONABELIAN SIMPLE GROUPS

In order to prove Theorem 4.5, we will have to consider normal subgroups in infinite products of compact semisimple Lie groups. In this section we take a look at normal subgroups in products of abstract groups. We will see that this leads almost inevitably to questions about ultraproducts.

If G_1, \dots, G_m are nonabelian simple groups, then it is easy to see that the product $G_1 \times \dots \times G_m$ is a perfect group, and that every proper normal subgroup of the product is contained in the kernel of at least one projection map

$$pr_j : G_1 \times \dots \times G_m \longrightarrow G_j.$$

But both questions become considerably more interesting for infinite products of groups.

First we look at products of perfect groups. Given a perfect group H , let $c(H)$ denote the *commutator length* of H , that is, the smallest integer n such that every element $h \in H$ can be written as a product of at most n commutators. If no such integer exists, we put $c(H) = \infty$.

Lemma 5.1. *Let $(G_i)_{i \in I}$ be a family of perfect groups. Put $I_\infty = \{i \in I \mid c(G_i) = \infty\}$. Then the product $\prod_{i \in I} G_i$ is perfect if and only if I_∞ is finite and if the function $i \mapsto c(G_i)$ is bounded on $I_0 = I - I_\infty$.*

Proof. Suppose that I_∞ is finite. A finite product of perfect groups is perfect, hence $L = \prod_{j \in I_\infty} G_j$ is perfect. If the function $k \mapsto c(G_k)$ is bounded on I_0 by a constant $n \in \mathbb{N}$, then every element in $M = \prod_{k \in I_0} G_k$ is a product of at most n commutators, and hence M is perfect as well. Therefore $\prod_{i \in I} G_i = L \times M$ is perfect.

If I_∞ is infinite or if the function $k \mapsto c(G_k)$ is unbounded on I_0 , then we can find an injection $\iota : \mathbb{N} \rightarrow I$ such that for every $n \in \mathbb{N}$ there is an element $g_{\iota(n)} \in G_{\iota(n)}$ which cannot be written as a product of less than n commutators. Put $g_j = 1$ if $j \notin \iota(\mathbb{N})$. Then $(g_i)_{i \in I}$ is an element in $\prod_{i \in I} G_i$ which is not a product of commutators. Therefore $\prod_{i \in I} G_i$ is not perfect. \square

In view of the previous lemma, we should expect to encounter lots of 'strange' normal subgroups in infinite products of nonabelian simple groups. In particular we cannot expect to catch all proper normal subgroups by the projection maps

$$pr_j : \prod_{i \in I} G_i \longrightarrow G_j.$$

But if we replace the finiteness of the index set I by a compactness condition, we gain some control over normal subgroups in the product. This leads directly to ultraproducts, as we will see. We first change our viewpoint slightly.

5.2. Presheaves of groups. Recall that a *presheaf of groups* \mathcal{G} on a topological space X is a contravariant group functor

$$\mathcal{G} : \mathbf{Open}(X) \longrightarrow \mathbf{Group}$$

which assigns to every open set $U \subseteq X$ a group $\mathcal{G}(U)$. For open sets $U \subseteq V \subseteq X$, there are restriction homomorphisms $pr_U^V : \mathcal{G}(V) \rightarrow \mathcal{G}(U)$, subject to the usual compatibility condition $pr_U^V \circ pr_V^W = pr_U^W$, for $U \subseteq V \subseteq W$, and with $pr_U^U = \text{id}_{\mathcal{G}(U)}$. Recall also that the *stalk* at a point $p \in X$ is defined as the direct limit

$$\mathcal{G}(p) = \lim_{\rightarrow} \{\mathcal{G}(U) \mid U \in \mathbf{Open}(X) \text{ and } p \in U\}.$$

For every $p \in U \in \mathbf{Open}(X)$, there is a natural restriction homomorphism

$$pr_p^U : \mathcal{G}(U) \longrightarrow \mathcal{G}(p).$$

Given a continuous map $\varphi : X \rightarrow Y$ between topological spaces, there is the *direct image presheaf* $\varphi_* \mathcal{G}$ on Y which assigns to $W \in \mathbf{Open}(Y)$ the group

$$\varphi_* \mathcal{G}(W) = \mathcal{G}(\varphi^{-1}(W)).$$

We note that $\varphi_* \mathcal{G}(Y) = \mathcal{G}(X)$.

5.3. Products of groups as presheaves. We may view the family of groups $(G_i)_{i \in I}$ as the group-valued presheaf \mathcal{G} on the discrete topological space I which assigns to every nonempty subset $J \subseteq I$ the group $\mathcal{G}_J = \prod_{j \in J} G_j$, and to $\emptyset \subseteq I$ the trivial group. The stalks are then the groups $\mathcal{G}(i) = G_i$. The problem that not all normal subgroups in $\prod_{i \in I} G_i$ can be detected by the projection maps $pr_j : \prod_{i \in I} G_i \rightarrow G_j$ can be rephrased as follows:

There are not enough stalks in the space I in order to detect all normal subgroups in the product $\prod_{i \in I} G_i$.

We remove this deficiency by 'enlarging' I through a map $I \rightarrow X$, where X is a compact space.

Proposition 5.4. *Suppose that $(G_i)_{i \in I}$ is a family of groups and that $N \trianglelefteq \prod_{i \in I} G_i$ is a normal subgroup. Let X be a compact space and let $\iota : I \rightarrow X$ be any map. If N maps under the composite homomorphism*

$$N \hookrightarrow \prod_{i \in I} G_i \xrightarrow{\cong} \iota_* \mathcal{G}(X) \rightarrow \iota_* \mathcal{G}(p)$$

onto $\iota_ \mathcal{G}(p)$, for every $p \in X$, then N contains the derived group of $\prod_{i \in I} G_i$.*

Proof. We identify $\prod_{i \in I} G_i$ with $\iota_* \mathcal{G}(X)$. Assuming that N surjects onto all stalks $\iota_* \mathcal{G}(p)$, and given any two elements $g, h \in \iota_* \mathcal{G}(X)$, we will show that N contains $[g, h]$.

Let $p \in X$. Since N maps under pr_p^X onto $\iota_* \mathcal{G}(p)$, there is an open neighborhood $U_p \subseteq X$ of p and an element $n_p \in N$ such that $pr_{U_p}^X(n_p) = pr_{U_p}^X(g)$. Since X is compact, we can cover X by finitely many such sets U_{p_1}, \dots, U_{p_r} . For $s = 1, \dots, r$, put $n_s = n_{p_s}$, and put

$$I_s = \iota^{-1}(U_{p_s}) \quad \text{and} \quad I'_s = I_s - (I_1 \cup \dots \cup I_{s-1}).$$

Then

$$I = I'_1 \dot{\cup} \dots \dot{\cup} I'_r \quad \text{and} \quad \iota(I_s) \subseteq U_{p_s}.$$

For $s = 1, \dots, r$ we define elements $m_s \in \prod_{i \in I} G_i$ by

$$(m_s)_i = \begin{cases} h_i & \text{if } i \in I'_s \\ 1 & \text{else.} \end{cases}$$

Then

$$([n_s, m_s])_i = \begin{cases} ([g, h])_i & \text{if } i \in I'_s \\ 1 & \text{else.} \end{cases}$$

Therefore $[g, h] = [n_1, m_1] \cdots [n_r, m_r] \in N$. □

In a completely analogous way, we have the following result

Proposition 5.5. *Let X be a compact space, let $(G_i)_{i \in I}$ be a family of groups, and let $\iota : I \rightarrow X$ be a map. Then $\prod_{i \in I} G_i$ is perfect if and only if every stalk $\iota_* \mathcal{G}(p)$ is perfect.*

Proof. First of all we note that it is clear from the construction of $\iota_*\mathcal{G}$ that the restriction homomorphisms pr_V^U are always surjective (such a presheaf is called *flabby*), hence $\mathcal{G}(X)$ maps onto $\mathcal{G}(p)$, for every $p \in X$. The image of a perfect group is perfect, so one implication is clear.

Now let $g \in \prod_{i \in I} G_i$ and assume that every stalk $\iota_*\mathcal{G}(p)$ is perfect. Hence for every $p \in X$ there are elements $a_1, b_1, \dots, a_n, b_n \in \prod_{i \in I} G_i$ such that $pr_p^X([a_1, b_1] \cdots [a_n, b_n]) = pr_p^X(g)$. Again, this relation holds then in some open neighborhood U_p of p . Now X is covered by finitely many such sets $U_1 = U_{p_1}, \dots, U_m = U_{p_m}$, where

$$pr_{U_s}^X(g) = pr_{U_s}^X[a_{s,1}, b_{s,1}] \cdots [a_{s,n}, b_{s,n}].$$

The number n can be chosen uniformly. By a similar argument as in the proof of Proposition 5.4, there is a disjoint decomposition $I = I'_1 \cup \dots \cup I'_m$ with $\iota(I'_s) \subseteq U_s$. We put

$$\widetilde{(a_{s,t})}_i = \begin{cases} (a_{s,t})_i & \text{if } i \in I'_s \\ 1 & \text{else} \end{cases} \quad \text{and} \quad \widetilde{(b_{s,t})}_i = \begin{cases} (b_{s,t})_i & \text{if } i \in I'_s \\ 1 & \text{else} \end{cases}.$$

Then $g = [\widetilde{a_{1,1}}, \widetilde{b_{1,1}}] \cdots [\widetilde{a_{m,n}}, \widetilde{b_{m,n}}]$. □

Now we look at specific choices for X . There are (a least) three obvious candidates.

Example 5.6 (The trivial case). If $X = \{p\}$ is a singleton and ι is the constant map, then $\iota_*\mathcal{G}(p) = \prod_{i \in I} G_i$. Obviously, the Propositions 5.4 and 5.5 tell us nothing new.

Example 5.7 (The cofinite case). If $X = \hat{I} = I \cup \{\infty\}$ is the Alexandrov compactification of I and $\iota : I \rightarrow \hat{I}$ is the inclusion, then $\ker(pr_{\infty}^{\hat{I}})$ is the restricted product

$$\prod_{i \in I}^{res} G_i = \{(g_i)_{i \in I} \mid g_i = 1 \text{ for almost all } i\} \trianglelefteq \prod_{i \in I} G_i.$$

Thus $\prod_{i \in I} G_i$ is perfect if and only if each G_i is perfect, and if in addition $\prod_{i \in I} G_i / \prod_{i \in I}^{res} G_i$ is perfect (compare this with Lemma 5.1).

If $\prod_{i \in I} G_i$ is perfect and if $N \trianglelefteq \prod_{i \in I} G_i$ is a proper normal subgroup with $pr_i^X(N) = G_i$ for all $i \in I$ (an example of such a group is $N = \prod_{i \in I}^{res} G_i$), then $N(\prod_{i \in I}^{res} G_i) \neq \prod_{i \in I} G_i$.

Example 5.8 (The ultraproduct case). The real case of interest is however when $X = \beta I$ is the Čech–Stone compactification of I , with the standard inclusion $\iota : I \rightarrow \beta I$. Recall that βI can be identified with the set of all ultrafilters on I . The points $j \in I$ correspond to the principal ultrafilters via $j \mapsto \{J \subseteq I \mid j \in J\}$, whereas the 'new' points in βI correspond to the free ultrafilters³ on I . For $J \subseteq I$ put $J^* = \{\mu \in \beta I \mid J \in \mu\}$. Then $\{J^* \mid J \subseteq I\}$ is a basis for the topology on βI . Given $\mu \in \beta I$, the set $\{J^* \mid J \in \mu\}$

³also called non-principal ultrafilters

is a neighborhood basis of μ . It follows that $\iota_*\mathcal{G}(\mu) = G_j$ if μ is the principal ultrafilter generated by j . But if μ is a free ultrafilter, then $\iota_*\mathcal{G}(\mu)$ is the *ultraproduct* (see [3, Ch. 5])

$$\iota_*\mathcal{G}(\mu) = \prod_{i \in I} G_i / \mu,$$

where we identify two sequences $(g_i)_{i \in I}, (h_i)_{i \in I}$ if $\{j \mid g_j = h_j\} \in \mu$.⁴ We write

$$pr_\mu = pr_\mu^{\beta I} : \prod_{i \in I} G_i \longrightarrow \prod_{i \in I} G_i / \mu$$

for the corresponding projection map.

By Proposition 5.5, the product $\prod_{i \in I} G_i$ is perfect if and only if each G_i is perfect and if for every free ultrafilter μ on I , the ultraproduct $\prod_{i \in I} G_i / \mu$ is perfect.

If the product $\prod_{i \in I} G_i$ is perfect and if $N \trianglelefteq \prod_{i \in I} G_i$ is a proper normal subgroup, then by Proposition 5.4 either there exists an index $j \in I$ with $pr_j(N) \neq G_j$, or there exists a free ultrafilter μ such that the image of N in $\prod_{i \in I} G_i / \mu$ is different from $\prod_{i \in I} G_i / \mu$.

We record this last result for later use.

Proposition 5.9. *Let $(G_i)_{i \in I}$ be a family of groups. Suppose that the product $\prod_{i \in I} G_i$ is perfect and that $N \trianglelefteq \prod_{i \in I} G_i$ is a proper normal subgroup. Then either there exists an index $j \in I$ with $pr_j(N) \neq G_j$ or there exists a free ultrafilter μ on I such that $pr_\mu(N) \neq \prod_{i \in I} G_i / \mu$. \square*

Products and ultraproducts of finite simple groups and their normal subgroups have been extensively studied. See e.g. [29], [32], [25] and the literature quoted there. The normal subgroup lattice of ultraproducts of compact simple Lie groups is studied in detail in [30].

6. NORMAL GENERATION OF COMPACT SIMPLE LIE GROUPS

We call a Lie group G *almost simple* if it is connected and if its Lie algebra is simple. Let G be a compact almost simple Lie group. Then the center $\text{Cen}(G)$ is finite and $G/\text{Cen}(G)$ is simple as an abstract group by Theorem 3.1. Moreover, the conjugacy class

$$C(a) = \{gag^{-1} \mid g \in G\}$$

of every non-central element $a \in G$ generates G . We will now re-prove this in a completely different way. The new approach has the advantage that it says, in a quantitative way, how fast a conjugacy class generates the whole group.

⁴One may think of the elements of μ as sets of measure 1. Thus two sequences are considered to be equivalent if they differ only on a set of measure 0.

6.1. Some observations. Suppose that $G = \mathrm{SU}(n)$. The conjugacy class of a matrix $a \in \mathrm{SU}(n)$ is determined by its spectrum (with its multiplicities). Given such a spectral datum, the task is therefore to write an arbitrary element $g \in \mathrm{SU}(n)$ as a product $g = a_1 \cdots a_m$ of special unitary matrices a_1, \dots, a_m with a prescribed spectrum.⁵

This viewpoint shows also that there are lower bounds on the number of matrices needed. Consider for example the group $G = \mathrm{SO}(2k+1)$, for $k \geq 1$. Every element in $g \in \mathrm{SO}(2k+1)$ can be written as a product of an even number of reflections. Let $g \in \mathrm{SO}(2k+1)$ be a matrix with exactly one real eigenspace $E \subseteq \mathbb{R}^{2k+1}$, of dimension $\dim(E) = 1$. Then g cannot be written as a product of less than $2k$ reflections. To see this, note that every reflection fixes a hyperplane. The intersection of less than $2k$ hyperplanes in \mathbb{R}^{2k+1} has dimension at least 2, hence a product of less than $2k$ reflections fixes a 2-dimensional subspace pointwise. The reflections themselves are not in $\mathrm{SO}(2k+1)$, but their negatives are. Hence we have here a lower bound on the number of multiplications of a fixed conjugacy class needed, which grows with the rank of the group.

We will prove in Theorem 6.11 that there exists a rational real function f , and on every simply connected almost simple compact Lie group G a nonnegative bounded continuous real class function $\hat{\sigma}$ with the following property. If $\hat{\sigma}(a) > \theta$, then every element in G is a product of less than $f(\theta)$ conjugates of a, a^{-1} . This is a first-order property of compact Lie groups, which extends by Los' Theorem to ultraproducts. Combining Proposition 5.9 with the Structure Theorem for compact connected groups, we obtain then a proof of Theorem 4.5.

In what follows, we put for a nonnegative real number r

$$\lceil r \rceil = \min\{k \in \mathbb{N} \mid k \geq r\}$$

and we note that

$$\lceil r \rceil < r + 1.$$

Our whole approach follows closely the reasoning in Nikolov–Segal [24], pp. 582–595, combined with a tweak introduced in [30]. However we avoid the notion of *asymptotic cones of metric spaces* (the *ultralimits* in *loc.cit.*) altogether. Instead we work bare hand with ultraproducts.

6.2. Some elementary calculations. We first consider the case of the almost simple matrix group

$$G = \mathrm{SU}(2) = \left\{ \begin{pmatrix} c & s \\ -\bar{s} & \bar{c} \end{pmatrix} \in \mathbb{C}^{2 \times 2} \mid c\bar{c} + s\bar{s} = 1 \right\}.$$

For $c = \pm 1$ we obtain the 2×2 identity matrix $\mathbf{1}$ and its negative $-\mathbf{1}$, respectively. Note that $\mathrm{Cen}(\mathrm{SU}(2)) = \{\pm \mathbf{1}\}$. We identify $\mathrm{SU}(2)$ with the round 3-sphere in $\mathbb{C}^2 \cong \mathbb{R}^4$, and

⁵This problem is ultimately related to Horn inequalities, Schubert calculus and buildings, see e.g. [2], [19] and [21].

we endow it with the unique bi-invariant Riemannian metric d of diameter π (so d is the spherical or angular metric). We fix the maximal torus

$$T = \left\{ \begin{pmatrix} c & \\ & \bar{c} \end{pmatrix} \in \mathbb{C}^{2 \times 2} \mid c\bar{c} = 1 \right\} \subseteq \mathrm{SU}(2)$$

and we note that every element in $\mathrm{SU}(2)$ is conjugate to an element in T .

Now we use some elementary spherical geometry. The conjugation action of G on itself is by rotations around the axis passing through $\pm \mathbf{1}$. The conjugacy class $C(a) = \{gag^{-1} \mid g \in G\}$ of an element $a \in G - \{\pm \mathbf{1}\}$ is thus a round 2-sphere of spherical radius θ , where $\frac{\theta}{2} = \min\{d(\mathbf{1}, a), d(-\mathbf{1}, a)\} \in [0, \frac{\pi}{2}]$. Then $C(a)a^{-1}$ is a round sphere of the same shape, but passing through the identity element $\mathbf{1} \in G$. The set of all conjugates of the set $C(a)a^{-1}$, which coincides obviously with $C(a)C(a^{-1})$, is thus the closed metric ball

$$C(a)C(a^{-1}) = B_\theta(\mathbf{1})$$

consisting of all $b \in G$ whose spherical distance from $\mathbf{1}$ is at most θ . Next we note that we have for every $\theta \in [0, \pi]$ that

$$G = \underbrace{B_\theta(\mathbf{1}) \cdots B_\theta(\mathbf{1})}_{k \text{ factors}} \text{ whenever } k\theta \geq \pi.$$

In particular we have shown that every element in $\mathrm{SU}(2)$ is a product not more than $2\lceil \frac{\pi}{\theta} \rceil$ conjugates of a and a^{-1} .

The quantity θ can be defined in a more natural way. We consider the *root* (or character)

$$\delta : T \longrightarrow \mathrm{U}(1), \quad \delta \begin{pmatrix} c & \\ & \bar{c} \end{pmatrix} = c^2.$$

For $c \in \mathrm{U}(1)$ we denote by $\ell(c) \in [0, \pi]$ the spherical (or angular) distance between 1 and c on the unit circle. Thus $\ell(ab) \leq \ell(a) + \ell(b)$ and $\ell(\bar{a}) = \ell(a)$ hold for all $a, b \in \mathrm{U}(1)$. Then we have for all $a \in T$ that the conjugacy class $C(a)$ is a sphere of diameter

$$\theta = \ell(\delta(a))$$

(where a 'sphere of diameter 0' is meant to be a point). Using these elementary observations, we have the following 'bootstrap Lemma'.

Lemma 6.3. *Let $H = H_0 \cdot H_1 \cdots H_s$ be a compact connected Lie group which is a central product of groups $H_i \cong \mathrm{SU}(2)$, for $1 \leq i \leq s$, and a torus H_0 . For each factor $H_i \cong \mathrm{SU}(2)$ with $1 \leq i \leq s$ we fix a torus $T_i \subseteq H_i$ and a root $\delta_i : T_i \longrightarrow \mathrm{U}(1)$ exactly as in 6.2. Suppose that $0 < \theta \leq \pi$ and that $h \in H$ is an element such that $\ell(\delta_i(h)) \geq \theta$ holds for all $1 \leq i \leq s$. Then every element in the subgroup $H_1 \cdot H_2 \cdots H_s \subseteq H$ is a product of at most $2\lceil \frac{\pi}{\theta} \rceil$ conjugates of h and h^{-1} .*

Proof. We write h as a product $h = h_0 h_1 \cdots h_s$, with $h_i \in H_i$. Then $C(h)C(h^{-1})$ is the product of the s commuting sets $(C(h_i)C(h_i^{-1}))$, with $1 \leq i \leq s$. There appears no more factor in H_0 , because H_0 is in the center of H and because h_0 and h_0^{-1} cancel out. The claim follows now from the observations in 6.2, applied to each H_i individually. \square

Now we turn to root systems in compact almost simple Lie groups in general. The books [1], [4], [13] and [17] are excellent references for the facts that we need.

6.4. Roots in a compact Lie group. Suppose that G is a compact simply connected almost simple Lie group. We fix a maximal torus $T \subseteq G$ and its character group $X^*(T) = \text{Hom}(T, \text{U}(1))$. Associated to T we have the root system $\Phi \subseteq X^*(T)$. The special case of $G = \text{SU}(2)$, where $\Phi = \{\pm\delta\}$, was described in 6.2.

Let $\Delta \subseteq \Phi$ be a system of simple roots, $\Delta = \{\delta_1, \dots, \delta_r\}$. Then $r = \dim(T)$ is the *rank* of G . There is a canonical isomorphism $V = X(T)^* \otimes \mathbb{R} \cong \text{Hom}_{\mathbb{R}}(\text{Lie}(T), \mathbb{R})$, and Δ is a basis of this real vector space. There is also a canonical inner product on V , and Φ is a reduced irreducible root system in the vector space V . The reflection group generated by Φ is the Weyl group $W = W(\Phi)$. The Weyl group is canonically isomorphic to N/T , where $N = \text{Nor}_G(T)$ is the normalizer of T in G .

There is also the cocharacter group $X_*(T) = \text{Hom}(\text{U}(1), T)$, and a natural isomorphism $X_*(T) \otimes \mathbb{R} \cong \text{Lie}(T)$. Associated to every root α there is a coroot α^\vee , with $\alpha(\alpha^\vee(c)) = c^2$. The action of $W(\Phi)$ on $\text{Lie}(T)$ coincides with the adjoint action of N/T .

6.5. Some more definitions. For a subset $\Delta' \subseteq \Delta$ we denote by $T_{\Delta'}$ the $\#\Delta'$ -dimensional subtorus spanned by the coroots $\{\delta^\vee \mid \delta \in \Delta'\}$, i.e.

$$T_{\Delta'} = \prod \{\delta^\vee(c) \mid \delta \in \Delta' \text{ and } c \in \text{U}(1)\} \subseteq T.$$

We now define real-valued continuous functions σ and $\hat{\sigma}$ on T by

$$\sigma(h) = \frac{1}{r} \sum_{i=1}^r \ell(\delta_i(h))$$

and

$$\hat{\sigma}(h) = \max\{\sigma(w(h)) \mid w \in W\}.$$

Lemma 6.6. *The function $\sigma : T \rightarrow \mathbb{R}$ has the following properties.*

- (i) $\sigma(h) = 0$ holds if and only if $h \in \text{Cen}(G)$.
- (ii) $\sigma(h) = \sigma(h^{-1})$.
- (iii) $\sigma(hh') \leq \sigma(h) + \sigma(h')$.
- (iv) $\sigma(T) = [0, \pi]$.

The function $\hat{\sigma}$ has the same properties (i)–(iv). In addition, $\hat{\sigma}$ extends uniquely to a continuous class function on G .

Proof. Properties (ii) and (iii) are clear from the definition (both for σ and for $\hat{\sigma}$). It is also clear from the definition that σ and $\hat{\sigma}$ take their values in $[0, \pi]$. Both σ and $\hat{\sigma}$ map the identity element in T to 0. Consider the Lie group homomorphism

$$P : T \rightarrow \text{U}(1) \times \dots \times \text{U}(1), \quad P(h) = (\delta_1(h), \dots, \delta_r(h)).$$

This homomorphism has maximal rank r (this can be seen from the coroots δ_i^\vee —the homomorphism P surjects onto each factor $\text{U}(1)$) and is therefore surjective, because the target group is connected. Now $\sigma(h)$ is the distance between $P(h)$ and the identity

element, with respect to the $\frac{1}{r}$ -scaled ℓ_1 -product metric on the target group. Therefore σ assumes the value π . Since \hat{T} is connected, $\sigma(T) = [0, \pi]$. It follows that $\hat{\sigma}(T) = [0, \pi]$, and hence claim (iv) holds. For claim (i) we note that the kernel of P is precisely the center of G , see [1, Def. 4.38 and Prop 5.3] or [4, Ch. V Prop. 7.16]. Indeed, the adjoint action of T on $\text{Lie}(G) \otimes_{\mathbb{R}} \mathbb{C}$ is precisely given by the characters, and the kernel of this action is the center of G . Finally, the function $\hat{\sigma}$ is by construction invariant under $W = \text{Nor}_G(T)/T$. Therefore it extends uniquely to a continuous class function on G , see [4, Ch. IV Cor. 2.7]. \square

6.7. The root system, continued. We keep the notation and the assumptions from 6.4. Corresponding to every simple root $\delta_i \in \Delta$ there is a connected subgroup H_i of type $\text{Lie}(H_i) \cong \mathfrak{su}(2)$, see [13, Lem. 12.2.15]. If two simple roots δ_i and δ_j are orthogonal, then they are *strongly orthogonal*, i.e. $\delta_i \pm \delta_j \notin \Phi$. It follows from the commutator relations in the complexified Lie algebra $\text{Lie}(G) \otimes_{\mathbb{R}} \mathbb{C}$ that then the groups H_i and H_j commute, $[H_i, H_j] = 1$.

Lemma 6.8. *Since G is simply connected, the groups H_i are isomorphic to $\text{SU}(2)$ and the root δ_i restricts to a root on $T \cap H_i$ as described in 6.2.*

Proof. Let $\Gamma \subseteq \text{Lie}(T)$ denote the subgroup generated by the coroots. Since G is simply connected, this group coincides with the kernel I of the homomorphism $\exp : \text{Lie}(T) \rightarrow T$, see [13, Prop. 12.4.14] or [4, Ch. V Thm. 7.1]. By [13, Prop 12.4.10], we have for every simple root $\delta \in \Delta$ a homomorphism $\rho_\delta : \text{SU}(2) \rightarrow G$ which maps the torus elements $(\begin{smallmatrix} c & \\ & \bar{c} \end{smallmatrix})$ in $\text{SU}(2)$ into T , such that $\delta(\rho_\delta(\begin{smallmatrix} c & \\ & \bar{c} \end{smallmatrix})) = c^2$. The composite $c \mapsto \rho_\delta(\begin{smallmatrix} c & \\ & \bar{c} \end{smallmatrix}))$ coincides with the coroot δ^\vee . Now $\delta^\vee(-1) = \exp(\frac{1}{2}\delta^\vee) \neq 1$, because $\frac{1}{2}\delta^\vee$ is not in $\Gamma = I$. Therefore ρ_δ is injective. \square

Now we get to the following key observation.

Key Lemma 6.9 (Nikolov–Segal). *With the notation and assumptions as in 6.4, let G be a compact simply connected almost simple Lie group of rank r , with maximal torus $T \subseteq G$. Let θ be a positive real number, and let $h \in T$ be an element with $\sigma(h) \geq \theta > 0$. Then there exists a set $\Delta_0 \subseteq \Delta$ of pairwise orthogonal simple roots such that the following conditions are simultaneously satisfied.*

- (a) Δ_0 has at not less than $\frac{r\theta}{4}$ elements.
- (b) Every element $a \in T_{\Delta_0}$ is a product of at most $2\lceil \frac{2\pi}{\theta} \rceil$ conjugates of h and h^{-1} .

Proof. We put $\Delta_1 = \{\delta \in \Delta \mid \ell(\delta(h)) \geq \theta/2\}$ and $t = \#\Delta_1$. Now we count

$$\theta \leq \sigma(h) \leq \frac{1}{r}(t + (r-t)\frac{\theta}{2}) \leq \frac{t}{r} + \frac{\theta}{2},$$

whence $\frac{r\theta}{2} \leq t$.

We claim that Δ_1 contains a subset Δ_0 consisting of pairwise orthogonal roots, with $\frac{t}{2} \leq \#\Delta_0$. We postpone the proof of this claim, which is easy and purely combinatorial, to 6.15. We have thus $\#\Delta_0 \geq \frac{r\theta}{4}$.

Now we apply Lemma 6.3 to the group H generated by T and the H_i with $\delta_i \in \Delta_0$. We conclude that every element in $T_{\Delta_0} \subseteq H$ is a product of at most $2\lceil \frac{2\pi}{\theta} \rceil$ conjugates of h and h^{-1} . \square

The last ingredient that we need is the following representation theoretic fact.

Lemma 6.10. *Let Φ be a reduced irreducible root system of rank r in a real r -dimensional vector space V , let $\Delta \subseteq \Phi$ be a set of simple roots and let $\Delta_0 \subseteq \Delta$ be a nonempty set consisting of s pairwise orthogonal roots. Let W denote the Weyl group of Φ . Then there exist elements $w_1, \dots, w_t \in W$ with the following properties.*

- (1) $w_1(\Delta_0) \cup \dots \cup w_t(\Delta_0)$ generates V .
- (2) $t \leq \frac{2r}{s} + 3$.

Again we postpone the proof to 6.15, and state first the main result.

Theorem 6.11 (Nikolov–Segal). *Let G be a compact simply connected almost simple Lie group of rank r . Suppose that $h \in G$ is an element with $\hat{\sigma}(h) \geq \theta > 0$. Then every element of G can be written as a product of less than $f(\theta) = 2(\frac{8}{\theta} + 3)(\frac{2\pi}{\theta} + 1)$ conjugates of h and h^{-1} .*

Proof. Let $T \subseteq G$ be a maximal torus. Every element in G is conjugate to an element in T , see [1, Thm. 4.21]. Since $\hat{\sigma}$ is a class function, we may assume that $h \in T$. After conjugating h further by an element in the Weyl group $W = \text{Nor}_G(T)/T$, we can assume that $\sigma(h) = \hat{\sigma}(h)$.

We use the notation set up in 6.4. By the Key Lemma 6.9, there is a nonempty set Δ_0 of s pairwise orthogonal simple roots, with $s \geq \frac{r\theta}{4}$. Moreover, we can generate the torus T_{Δ_0} in less than $2(\frac{2\pi}{\theta} + 1)$ steps from conjugates of h and h^{-1} . Let w_1, \dots, w_t be as in Lemma 6.10. Then the t -fold product multiplication map $T_{\Delta_0} \times \dots \times T_{\Delta_0} \rightarrow T$ which maps (a_1, \dots, a_t) to $w_1(a_1)w_2(a_2)\dots w_t(a_t)$ is a surjective homomorphism, because it is a homomorphism of connected Lie groups whose derivative at the identity is surjective. Hence every element in T is a product of not more than $2t(\frac{2\pi}{\theta} + 1)$ conjugates of h and h^{-1} . Moreover, $t \leq \frac{2r}{s} + 3$ and $\frac{1}{s} \leq \frac{4}{r\theta}$, whence $t \leq \frac{8}{\theta} + 3$. Since every element in G is conjugate to an element in T , the claim follows. \square

6.12. A geometric interpretation. The adjoint action of G on itself is a prototype of a *polar action*, as studied in Riemannian geometry. The conjugacy classes in G near the identity element are the flag manifolds for the spherical Tits building Δ associated to the complexification $G_{\mathbb{C}}$ of G . The question that is addressed by Theorem 6.11 can be re-stated as follows. Let $X = G/\text{Cen}(G)$. Endowed with a bi-invariant Riemannian metric, this is a symmetric space of compact type.

Let $\kappa \subseteq X$ be a nontrivial geodesic segment. Show that every element $g \in X$ can be joined to the identity $1 \in X$ by a piecewise geodesic path, whose pieces are conjugates of κ under the isometric action of $G \times G$ on X . Bound the number of pieces needed in terms of κ .

It would be interesting to know if there is a 'Riemannian proof' of Theorem 6.11, starting from this viewpoint of broken geodesics, using Morse Theory and polar foliations.

In any case we now combine Theorem 6.11, Proposition 5.9 and Łos' Theorem 6.13 in order to prove Theorem 4.5. We need some elementary facts from the model theory of ultraproducts. Any old-fashioned logic text book such as [3] and [5] will do as a reference. See also [22, pp. 5–7] for a very brief introduction aimed at metric geometers.

Theorem 6.13 (Łos' Theorem). *Let $(\mathcal{M}_i)_{i \in I}$ be a family of \mathcal{L} -structures, for some first order language \mathcal{L} . Let μ be a free ultrafilter on I and let ${}^*\mathcal{M}$ denote the ultraproduct with respect to μ . Let ψ be a sentence in \mathcal{L} . Then ψ is true in ${}^*\mathcal{M}$ if and only if the set $I_\psi = \{i \in I \mid \psi \text{ is true in } \mathcal{M}_i\}$ is contained in μ .*

The proof of Łos' Theorem is not difficult and well explained e.g. in Bell–Slomson [3, Ch. 5, Thm. 2.1].

Proposition 6.14. *Let $(G_i)_{i \in I}$ be a family of compact simply connected almost simple Lie groups. Let $N \trianglelefteq \prod_{i \in I} G_i$ be an abstract normal subgroup of countable index. Then $N = \prod_{i \in I} G_i$.*

Proof. Assume to the contrary that $N \neq \prod_{i \in I} G_i$ is of countable index. If $pr_j(N) = K_j \neq G_j$ for some index $j \in I$, then $K_j \subseteq G_j$ is contained in the center of G_j by 3.1 and hence G/N maps onto the uncountable group G_j/K_j , which is impossible. Thus $pr_j(N) = G_j$ holds for all $j \in I$. In a compact semisimple Lie group, every element is a commutator by Goto's Theorem, see [17, Cor. 6.56]. Thus $\prod_{i \in I} G_i$ is perfect and we may apply Proposition 5.9. It follows that there exists a free ultrafilter μ on I such that

$$pr_\mu(N) = K_\mu \neq \prod_{i \in I} G_i/\mu.$$

Let $f(\theta) = 2(\frac{8}{\theta} + 3)(\frac{2\pi}{\theta} + 1)$.

We put ${}^*G = \prod_{i \in I} G_i/\mu$ for short. We choose for every $i \in I$ a maximal torus $T_i \subseteq G_i$, a set of simple roots, and we define $\hat{\sigma}_i : G_i \rightarrow [0, \pi]$ as in 6.4. In the ultraproduct we obtain an abelian subgroup ${}^*T = \prod_{i \in I} T_i/\mu \subseteq {}^*G$. The ultraproduct of the $\hat{\sigma}_i$ is denoted by ${}^*\hat{\sigma}$. This function takes its values in the nonstandard reals ${}^*\mathbb{R}$ (the ultraproduct of \mathbb{R} with respect to μ , a real closed non-archimedean ordered field).

Let m be a positive integer. Then by Theorem 6.11, the following first-order sentence is true in each G_i .⁶

If $g \in G$ with $\hat{\sigma}_i(g) \geq \frac{1}{m}$, then every element in G_i is a product of at most $f(\frac{1}{m})$ conjugates of g .

⁶The relevant first-order language contains predicates for elements of groups G, T , of a field R , function symbols for maps $\hat{\sigma}, f$, function symbols $\cdot, +, \times$ for the group multiplication, the addition and multiplication in a field, and a relation symbol $<$ for an ordering on the field. The number m is written out as $1 + 1 + \dots + 1$.

By Łos' Theorem 6.13, the same sentence is true in the ultraproduct *G . Also, every element in *G is conjugate to an element in *T (either by Łos' Theorem, or directly, because this is true in each G_i by [1, Thm. 4.21]). Therefore ${}^*T \cap K_\mu$ consists necessarily of elements whose $\hat{\sigma}$ -value is infinitesimally small. Let $A \subseteq {}^*T$ denote the collection of all elements in *T whose ${}^*\hat{\sigma}$ -value is infinitesimally small. By Łos' Theorem and the properties of $\hat{\sigma}$ stated in Lemma 6.6, $A \subseteq {}^*T$ is a subgroup. Moreover, $|{}^*\hat{\sigma}(a) - {}^*\hat{\sigma}(b)|$ is infinitesimally small if $a, b \in {}^*T$ are elements with $ab^{-1} \in A$. Consider the standard part function $\text{std} : {}^*\mathbb{R}_{\text{fin}} \rightarrow \mathbb{R}$ which assigns to every finite nonstandard real its real part. The map ${}^*\hat{\sigma}$ surjects *T onto the set $Q = \{x \in {}^*\mathbb{R} \mid 0 \leq x \leq \pi\}$ (again by Lemma 6.6 and Łos' Theorem). Then std surjects Q onto $[0, \pi]$ and we have a commuting diagram

$$\begin{array}{ccc} {}^*T & \xrightarrow{{}^*\hat{\sigma}} & Q \\ \downarrow & & \downarrow \text{std} \\ {}^*T/A & \longrightarrow & [0, \pi]. \end{array}$$

Therefore the quotient ${}^*T/A$ is uncountable. Now ${}^*T/{}^*T \cap K_\mu$ injects into ${}^*G/K_\mu$ and surjects onto ${}^*T/A$. It follows that ${}^*G/K_\mu$ is uncountable. Hence G/N is also uncountable, a contradiction. \square

Proof of Theorem 4.5. Let φ denote the projection $G \rightarrow G/N$ and put $\Gamma = G/N$ and $\Gamma_0 = \varphi(G^\circ)$. Then Γ_0 has finite index in Γ .

By the Approximation Theorem for compact connected groups, there exists a compact connected abelian group Z , a family $(G_i)_{i \in I}$ of simply connected compact almost simple Lie groups and a continuous central surjective homomorphism $\psi : Z \times \prod_{i \in I} G_i \rightarrow G^\circ$ with totally disconnected kernel, see [17, Thm. 9.24]. By Theorem 6.11, $\varphi \circ \psi$ annihilates $\prod_{i \in I} G_i$, because Γ_0 is countable. Since $\prod_{i \in I} G_i$ is the derived group of $Z \times \prod_{i \in I} G_i$, the group Γ_0 is abelian, and thus Γ is virtually abelian.

Suppose now that Γ is residually finite. Then Γ_0 is also residually finite. On the other hand, Z is connected and therefore divisible. It follows from 1.2 that φ is constant on Z and hence constant on G° . Thus φ is continuous and open, and Γ is finite. \square

6.15. The remaining combinatorial proofs.

Proof of the claim made in the proof of the Key Lemma 6.9. Let Γ denote the underlying graph of the Dynkin diagram of Φ . The vertices of Γ are the simple roots in Δ . This graph is a tree and admits therefore a coloring, using red and blue, such that no two adjacent nodes have the same color. Simple roots which are not joined by an edge are orthogonal. Thus both the red and the blue roots form sets of pairwise orthogonal roots. In particular we have partitioned Δ_1 into two sets of pairwise orthogonal roots. One of these sets has at least $\frac{\#\Delta_1}{2}$ elements. \square

Proof of Lemma 6.10. Given a set Δ_0 consisting of $s \geq 1$ pairwise orthogonal simple roots, let t be the minimum number of elements $w_1, \dots, w_t \in W$ with the property that $w_1(\Delta_0) \cup \dots \cup w_t(\Delta_0)$ generates V . The action of W on V is irreducible (because Φ is irreducible). Therefore the W -orbit of every root α generates V . Given α , there exist thus r elements $w_1, \dots, w_r \in W$ such that $w_1(\alpha), \dots, w_r(\alpha)$ is a basis for V . This shows that we have always

$$t \leq r,$$

and that $t = r$ if $s = 1$. Now we consider the different types of root systems of rank $r \geq 2$, and we assume that $s > 1$.

For type A_r the Weyl group W is the symmetric group on the set $\{1, \dots, r+1\}$. The reflections determined by the roots are the transpositions (i, j) with $i < j$. The reflections corresponding to the simple roots $\delta_1, \dots, \delta_r$ are the transpositions $(i, i+1)$, for $1 \leq i \leq r$. The reflections corresponding to the roots in Δ_0 are then pairwise commuting transpositions. Given $1 \leq k \leq r - 2s - 1$, there exists therefore a permutation $w \in W$ which conjugates these s commuting transpositions to $(k, k+1), (k+2, k+3), \dots, (k+2s-2, k+2s-1)$. Thus w maps Δ_0 to the set of roots $\varepsilon_1 \delta_k, \varepsilon_2 \delta_{k+2}, \dots, \varepsilon_s \delta_{k+2s-2} \subseteq \Phi$, with $\varepsilon_i = \pm 1$. Since V is spanned by $\delta_1, \dots, \delta_r$, we may simply count how many subdiagrams of type A_{2s} are needed to cover the whole A_r -diagram. We obtain thus an upper bound

$$t \leq 2 \lceil \frac{r}{2s} \rceil < \frac{r}{s} + 2 < \frac{2r}{s} + 3.$$

For the types B_r, C_r, D_r, E_r and $r \geq 5$ we note that there is a root subsystem $\Phi' \subseteq \Phi$ of type A_{r-1} , such that Δ_0 intersects this subsystem in a set of size at least $s - 1 > 0$. The $r - 1$ -dimensional subspace V' generated by Φ' is thus contained in a subspace generated by at most $2 \lceil \frac{r-1}{2(s-1)} \rceil$ W -translates of Δ_0 . Since V' is not invariant under W , we can generate V if we add one more W -translate of Δ_0 , i.e.

$$t \leq 2 \lceil \frac{r-1}{2s-2} \rceil + 1 < \frac{r-1}{s-1} + 3 \leq \frac{2r}{s} + 3.$$

One checks directly that for $2 \leq r \leq 4$ and $2 \leq s < r$ one has $r \leq \frac{2r}{s} + 3$, whence $t \leq \frac{2r}{s} + 3$. \square

Acknowledgement. The author thanks Martin Bridson, Arjeh Cohen, Theo Grundhöfer, Karl Hofmann, Karl-Hermann Neeb, and the anonymous referee for helpful remarks and clarifications.

REFERENCES

- [1] J. F. Adams, *Lectures on Lie groups*, W. A. Benjamin, Inc., New York, 1969. MR0252560 (40 #5780)
- [2] S. Agnihotri and C. Woodward, Eigenvalues of products of unitary matrices and quantum Schubert calculus, *Math. Res. Lett.* **5** (1998), no. 6, 817–836. MR1671192 (2000a:14066)
- [3] J. L. Bell and A. B. Slomson, *Models and ultraproducts: An introduction*, North-Holland, Amsterdam, 1969. MR0269486 (42 #4381)

- [4] T. Bröcker and T. tom Dieck, *Representations of compact Lie groups*, translated from the German manuscript, corrected reprint of the 1985 translation, Graduate Texts in Mathematics, 98, Springer, New York, 1995. MR1410059 (97i:22005)
- [5] C. C. Chang and H. J. Keisler, *Model theory*, third edition, Studies in Logic and the Foundations of Mathematics, 73, North-Holland, Amsterdam, 1990. MR1059055 (91c:03026)
- [6] J. Dugundji, *Topology*, Allyn & Bacon, Boston, MA, 1966. MR0193606 (33 #1824)
- [7] A. A. George Michael, On normal subgroups of semisimple Lie groups, *Results Math.* **58** (2010), no. 1-2, 37–38. MR2672623 (2011i:22005)
- [8] V. M. Gluškov, Structure of locally bicomact groups and Hilbert’s fifth problem, *Uspehi Mat. Nauk (N.S.)* **12** (1957), no. 2 (74), 3–41. MR0101892 (21 #698)
English translation: The structure of locally compact groups and Hilbert’s fifth problem, *Amer. Math. Soc. Transl. (2)* **15** (1960), 55–93. MR0114872 (22 #5690)
- [9] M. Goto, On an arcwise connected subgroup of a Lie group, *Proc. Amer. Math. Soc.* **20** (1969), 157–162. MR0233923 (38 #2244)
- [10] V. S. Guba, A finitely generated complete group, *Izv. Akad. Nauk SSSR Ser. Mat.* **50** (1986), no. 5, 883–924. MR0873654 (88e:20034)
English translation: *Math. USSR-Izv.* **29** (1987), no. 2, 233–277.
- [11] S. Hernández, K. H. Hofmann and S. A. Morris, Nonmeasurable subgroups of compact groups, *J. Group Theory* **19** (2016), no. 1, 179–189. MR3441133
- [12] G. Higman, B. H. Neumann and H. Neumann, Embedding theorems for groups, *J. London Math. Soc.* **24** (1949), 247–254. MR0032641 (11,322d)
- [13] J. Hilgert and K.-H. Neeb, *Structure and geometry of Lie groups*, Springer Monographs in Mathematics, Springer, New York, 2012. MR3025417
- [14] K. H. Hofmann and L. Kramer, Transitive actions of locally compact groups on locally contractible spaces, *J. Reine Angew. Math.* **702** (2015), 227–243. MR3341471
Erratum, *J. Reine Angew. Math.* **702** (2015), 245–246.
- [15] K. H. Hofmann and S. A. Morris, Transitive actions of compact groups and topological dimension, *J. Algebra* **234** (2000), no. 2, 454–479. MR1801101 (2002a:22006)
- [16] K. H. Hofmann and S. A. Morris, *The Lie theory of connected pro-Lie groups*, EMS Tracts in Mathematics, 2, European Mathematical Society (EMS), Zürich, 2007. MR2337107 (2008h:22001)
- [17] K. H. Hofmann and S. A. Morris, *The structure of compact groups*, third edition, revised and augmented., De Gruyter Studies in Mathematics, 25, de Gruyter, Berlin, 2013. MR3114697
- [18] K. Iwasawa, On some types of topological groups, *Ann. of Math. (2)* **50** (1949), 507–558. MR0029911 (10,679a)
- [19] L. C. Jeffrey and A.-L. Mare, Products of conjugacy classes in $SU(2)$, *Canad. Math. Bull.* **48** (2005), no. 1, 90–96. MR2118766 (2005m:20073)
- [20] R. R. Kallman, Every reasonably sized matrix group is a subgroup of S_∞ , *Fund. Math.* **164** (2000), no. 1, 35–40. MR1784652 (2001h:20005)
- [21] M. Kapovich, B. Leeb and J. J. Millson, The generalized triangle inequalities in symmetric spaces and buildings with applications to algebra, *Mem. Amer. Math. Soc.* **192** (2008), no. 896, viii+83 pp. MR2369545 (2009d:22018)
- [22] L. Kramer and R. M. Weiss, Coarse equivalences of Euclidean buildings. With an appendix by Jeroen Schillewaert and Koen Struyve, *Adv. Math.* **253** (2014), 1–49. MR3148544
- [23] D. Montgomery and L. Zippin, *Topological transformation groups*, Interscience Publishers, New York, 1955. MR0073104 (17,383b)
- [24] N. Nikolov and D. Segal, Generators and commutators in finite groups; abstract quotients of compact groups, *Invent. Math.* **190** (2012), no. 3, 513–602. MR2995181

- [25] F. Point, Ultraproducts and Chevalley groups, Arch. Math. Logic **38** (1999), no. 6, 355–372. MR1711404 (2001e:03070)
- [26] D. L. Ragozin, A normal subgroup of a semisimple Lie group is closed, Proc. Amer. Math. Soc. **32** (1972), 632–633. MR0294563 (45 #3633)
- [27] D. J. S. Robinson, *A course in the theory of groups*, second edition, Graduate Texts in Mathematics, 80, Springer, New York, 1996. MR1357169 (96f:20001)
- [28] H. Salzmann et al., *Compact projective planes*, de Gruyter Expositions in Mathematics, 21, de Gruyter, Berlin, 1995. MR1384300 (97b:51009)
- [29] J. Saxl, S. Shelah and S. Thomas, Infinite products of finite simple groups, Trans. Amer. Math. Soc. **348** (1996), no. 11, 4611–4641. MR1376555 (97j:20031)
- [30] A. Stolz and A. Thom, On the lattice of normal subgroups in ultraproducts of compact simple groups, Proc. Lond. Math. Soc. (3) **108** (2014), no. 1, 73–102. MR3162821
- [31] M. Stroppel, *Locally compact groups*, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2006. MR2226087 (2007d:22001)
- [32] S. Thomas, Infinite products of finite simple groups. II, J. Group Theory **2** (1999), no. 4, 401–434. MR1718758 (2000m:20043)

LINUS KRAMER
MATHEMATISCHES INSTITUT, UNIVERSITÄT MÜNSTER
EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY
E-mail address: `linus.kramer@uni-muenster.de`