

IWASAWA THEORY OF HEEGNER CYCLES, I. RANK OVER THE IWASAWA ALGEBRA

MATTEO LONGO AND STEFANO VIGNI

ABSTRACT. Iwasawa theory of Heegner points on abelian varieties of GL_2 type has been studied by, among others, Mazur, Perrin-Riou, Bertolini and Howard. The purpose of this paper, the first in a series of two, is to describe extensions of some of their results in which abelian varieties are replaced by the Galois cohomology of Deligne’s p -adic representation attached to a modular form f of even weight > 2 . In this more general setting, the role of Heegner points is played by higher-dimensional Heegner cycles in the sense of Nekovář. In particular, we prove that the Pontryagin dual of a certain Bloch–Kato Selmer group associated with f has rank 1 over a suitable anticyclotomic Iwasawa algebra.

1. INTRODUCTION

Initiated by Mazur’s paper [20], Iwasawa theory of Heegner points on abelian varieties of GL_2 type (most notably, elliptic curves) has been investigated by, among others, Perrin-Riou ([28]), Bertolini ([1], [2]) and Howard ([12], [13]). A recurrent theme in all these works is the study of pro- p -Selmer groups, where p is a prime number, in terms of Iwasawa modules built out of compatible families of Heegner points over the anticyclotomic \mathbb{Z}_p -extension of an imaginary quadratic field. In particular, several results on the structure of Selmer groups obtained by Kolyvagin by using his theory of Euler systems ([15]) were generalized to an Iwasawa-theoretic setting.

The goal of the present paper, the first in a series of two, is to describe extensions of some of the results of the previously mentioned authors in which abelian varieties are replaced by the Galois cohomology of Deligne’s p -adic representation attached to a modular form f of even weight > 2 . In this context, the role of Heegner points is played by Heegner cycles, which were introduced by Nekovář in [24] in order to extend Kolyvagin’s theory to Chow groups of Kuga–Sato varieties. More precisely, the main result of our article shows that the Pontryagin dual of a certain Bloch–Kato Selmer group associated with f has rank 1 over a suitable anticyclotomic Iwasawa algebra.

Let $N \geq 3$ be an integer, let $k \geq 4$ be an even integer and let f be a normalized newform of weight k and level $\Gamma_0(N)$, whose q -expansion will be denoted by

$$f(q) = \sum_{n \geq 1} a_n q^n.$$

Fix an imaginary quadratic field K of discriminant coprime to Np in which all the prime factors of N split and let p be a prime number not dividing N . For simplicity, we suppose that $\mathcal{O}_K^\times = \{\pm 1\}$, i.e., that $K \neq \mathbb{Q}(\sqrt{-1})$ and $K \neq \mathbb{Q}(\sqrt{-3})$. Fix also embeddings $K \hookrightarrow \mathbb{C}$ and $\mathbb{Q} \hookrightarrow \mathbb{Q}_p$. Write F for the number field generated over \mathbb{Q} by the Fourier coefficients a_n of f and let \mathcal{O}_F be its ring of integers. Let \mathfrak{p} be a prime ideal of \mathcal{O}_F above p and denote by $V_{f,\mathfrak{p}}$ the p -adic representation of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ attached to f and \mathfrak{p} by Deligne ([10]). If $F_{\mathfrak{p}}$

2010 *Mathematics Subject Classification.* 11R23, 11F11.

Key words and phrases. Iwasawa theory, modular forms, Heegner cycles.

The two authors are supported by PRIN 2010–11 “Arithmetic Algebraic Geometry and Number Theory”. The first author is also supported by PRAT 2013 “Arithmetic of Varieties over Number Fields”; the second author is also supported by PRA 2012 “Geometria Algebraica e Geometria Aritmetica”.

is the completion of F at \mathfrak{p} then $V_{f,\mathfrak{p}}$ is an $F_{\mathfrak{p}}$ -vector space of dimension 2 equipped with a continuous action of $\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$. Write $\mathcal{O}_{\mathfrak{p}}$ for the valuation ring of $F_{\mathfrak{p}}$ and set $a'_p := a_p/p^{k/2-1}$.

Throughout this article we shall always work under the following

- Assumption 1.1.** (1) $V_{f,\mathfrak{p}}$ is non-exceptional ([6, Definition 6.1]);
 (2) $p \nmid 6N\phi(N)h_K(k-2)!$ where ϕ is Euler's function and h_K is the class number of K ;
 (3) p does not ramify in F ;
 (4) $a'_p \in \mathcal{O}_{\mathfrak{p}}^\times$;
 (5) $a'_p \not\equiv 2 \pmod{p}$ if p splits in K and $a'_p \not\equiv 1 \pmod{p}$ if p is inert in K .

Additional cohomological conditions that allow us to obtain an analogue of Mazur's control theorem for elliptic curves will be listed and made precise in Assumption 2.4.

In light of results of Serre on eigenvalues of Hecke operators ([30, §7.2]), it seems reasonable to expect that (4) in Assumption 1.1 holds for infinitely many primes p (at least if $F \neq \mathbb{Q}$). In fact, questions of this sort appear to lie in the circle of ideas of the conjectures of Lang and Trotter on the distribution of traces of Frobenius automorphisms acting on elliptic curves ([16]) and of their extensions to higher weight modular forms ([22], [23]).

Let K_∞ be the anticyclotomic \mathbb{Z}_p -extension of K (i.e., the \mathbb{Z}_p -extension of K that is dihedral over \mathbb{Q}), set $G_\infty := \mathrm{Gal}(K_\infty/K)$ and form the Iwasawa algebra $\Lambda := \mathcal{O}_{\mathfrak{p}}[[G_\infty]]$. As in [24], one can define a $\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ -representation A that is a quotient of the $k/2$ -twist of $V_{f,\mathfrak{p}}$; denote by $H_f^1(K_\infty, A)$ the Bloch–Kato Selmer group of A over K_∞ . Finally, let

$$\mathcal{X}_\infty := \mathrm{Hom}_{\mathcal{O}_{\mathfrak{p}}}^{\mathrm{cont}}(H_f^1(K_\infty, A), F_{\mathfrak{p}}/\mathcal{O}_{\mathfrak{p}})$$

be the Pontryagin dual of $H_f^1(K_\infty, A)$, which turns out to be finitely generated over Λ .

Our main result is the following

Theorem 1.2. *The Λ -module \mathcal{X}_∞ has rank 1.*

This is the counterpart of [1, Theorem A]. The key tool in the proof of Theorem 1.2 is the Iwasawa module of Heegner cycles, which is denoted by \mathcal{H}_∞ in the main body of the text. The Λ -module \mathcal{H}_∞ is built out of the systematic supply of Heegner cycles introduced by Nekovář ([24]), which are higher-dimensional analogues of classical Heegner points. Since we are interested in Iwasawa-theoretic results, we work with Heegner cycles of p -power conductor; these cycles live in the Selmer groups $H_f^1(K_m, A)$, where for every $m \geq 1$ the field K_m is the subextension of K_∞/K such that $\mathrm{Gal}(K_m/K) \simeq \mathbb{Z}/p^m\mathbb{Z}$.

As will become apparent later, our strategy follows [1] closely; in fact, we extend to Heegner cycles the Λ -adic Kolyvagin method developed by Bertolini. From a slightly different point of view, analogous results could presumably be obtained by exploiting the theory of Kolyvagin systems due to Mazur and Rubin ([21]). More precisely, we expect that our constructions can easily be adapted to the formalism of [21], as was done by Howard in [12] for Heegner points on elliptic curves and by Fouquet in [11] in the context of big Heegner points, thus leading to one divisibility in the relevant Iwasawa-type Main Conjecture. However, in this paper we preferred to adopt the more direct and explicit approach of Bertolini, and hence to work over the Iwasawa algebra only instead of deducing results for the Iwasawa algebra from specializations to discrete valuation rings as in [21].

We observe that a crucial role in our arguments is played by a result of Howard ([14, Theorem A]) that extends to Heegner cycles and the cohomology of $V_{f,\mathfrak{p}}$ a theorem of Cornut ([9]) asserting the non-triviality of Heegner points on an elliptic curve E/\mathbb{Q} as one ascends K_∞ . We also note that a version of Theorem 1.2 in the context of Hida families of Hilbert modular forms was proved by Nekovář in [25, Theorem 12.9.11, (ii)]. It is worth pointing out that Nekovář's result applies to all but finitely many members of a given family and that his proof, relying on his theory of Selmer complexes and (of course) on Hida-theoretic techniques, is different from ours.

We conclude this introduction by remarking that in the sequel [17] to this paper we shall study the torsion submodule $\mathcal{X}_\infty^{\text{tors}}$ of \mathcal{X}_∞ . In particular, we will generalize a divisibility result due to Howard ([13, Theorem B]) as well as a result of Bertolini concerning the structure of the annihilator of $\mathcal{X}_\infty^{\text{tors}}$ ([1, Theorem B]). Applications to a Main Conjecture *à la* Perrin-Riou will also be given.

For the convenience of the reader, we collect below all the basic notation that will be used throughout this paper.

Notation. • If G is an abelian group and $n \geq 1$ is an integer then G_n denotes the n -torsion subgroup of G .

• For any field K we fix an algebraic closure \bar{K} of K and write $G_K := \text{Gal}(\bar{K}/K)$ for the absolute Galois group of K . In the special case where K is a number field, we shall view K as a subfield of $\bar{\mathbb{Q}}$ (in other words, we choose $\bar{K} = \bar{\mathbb{Q}}$).

• If M is a continuous G_K -module then $H_{\text{cont}}^i(K, M)$ is the i -th continuous cohomology group of G_K with coefficients in M . We will usually write $H^i(K, M)$ as a shorthand for $H_{\text{cont}}^i(K, M)$ and $M(K)$ as a shorthand for $H^0(K, M)$.

• If K/F is an algebraic field extension then we denote by

$$\text{res}_{K/F} : H^i(F, M) \longrightarrow H^i(K, M), \quad \text{cores}_{K/F} : H^i(K, M) \longrightarrow H^i(F, M)$$

the restriction and corestriction maps in cohomology, respectively. We recall that, for K/F finite and Galois, we have the equality

$$(1) \quad \text{res}_{K/F} \circ \text{cores}_{K/F} = \text{tr}_{K/F},$$

where $\text{tr}_{K/F} := \sum_{\sigma \in \text{Gal}(K/F)} \sigma$ is the Galois trace map on $H^i(K, M)$. Further, if v is a place of a number field K then we denote by

$$\text{res}_v : H^i(K, M) \longrightarrow H^i(K_v, M)$$

the restriction (localization) map at v , which is determined by the choice of an embedding of $\bar{\mathbb{Q}}$ into \bar{K}_v .

• If K is a number field and ℓ is a prime number then we denote by $K_\ell := \bigoplus_{\lambda|\ell} K_\lambda$ the direct sum of the completions K_λ of K at all the primes λ above ℓ , and we set $H^i(K_\ell, M) := \bigoplus_{\lambda|\ell} H^i(K_\lambda, M)$.

• If G is a profinite group, p is a prime and R is the ring of integers of a finite extension of \mathbb{Q}_p then $R[[G]]$ is the Iwasawa algebra of G with coefficients in R . For an $R[[G]]$ -module M we write M^G and M_G for the largest R -submodule and quotient of M , respectively, on which G acts trivially.

• If K is a local field then K^{nr} denotes the maximal unramified extension of K inside \bar{K} , so that $I_K := \text{Gal}(\bar{K}/K^{\text{nr}})$ is the inertia group of K . When K is a number field and v is a place of K we also write I_v for I_{K_v} . In particular, for all prime numbers ℓ we let F_ℓ be the arithmetic Frobenius in $\text{Gal}(\mathbb{Q}_\ell^{\text{nr}}/\mathbb{Q}_\ell)$. We also fix field embeddings $\bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_\ell$ and, with an abuse of notation, when dealing with a $G_{\mathbb{Q}}$ -module that is unramified at ℓ we often adopt the same symbol to denote a lift of F_ℓ to $G_{\mathbb{Q}_\ell}$ (and its image in $G_{\mathbb{Q}}$).

• If L/E is a Galois extension of number fields, λ is a prime of E that is unramified in L and λ' is a prime of L above λ then $\text{Frob}_{\lambda'/\lambda} \in \text{Gal}(L/E)$ denotes the Frobenius substitution at λ' ; the conjugacy class of $\text{Frob}_{\lambda'/\lambda}$ in $\text{Gal}(L/E)$ will be denoted by Frob_λ (notation not reflecting dependence on L).

• For an algebraic variety V defined over a field E of characteristic 0, an integer r such that $0 \leq r \leq \dim(V)$ and a prime number p , we let

$$\text{AJ}_{V/E}^{(r)} : \text{CH}^r(V/E) \longrightarrow H^1(E, H_{\text{ét}}^{2r-1}(\bar{V}, \mathbb{Z}_p))$$

denote the corresponding p -adic Abel–Jacobi map, with $\bar{V} := V \otimes_E \bar{E}$. We simply write AJ_E in place of $\text{AJ}_{V/E}^{(r)}$ when V and r are clear from the context.

2. BLOCH–KATO SELMER GROUPS IN \mathbb{Z}_p -EXTENSIONS

Our goal in this section is to introduce the Selmer groups we shall be interested in and state a “control theorem” for them.

2.1. Bloch–Kato Selmer groups. We begin with a general discussion of Selmer groups of p -adic representations.

For a number field E and a p -adic representation V of the Galois group G_E (by which we mean, as usual, a finite-dimensional \mathbb{Q}_p -vector space V equipped with a continuous action of G_E), the *Bloch–Kato Selmer group* of V over E ([7, Sections 3 and 5]) is the group $H_f^1(E, V)$ that makes the sequence

$$0 \longrightarrow H_f^1(E, V) \longrightarrow H^1(E, V) \xrightarrow{\prod_v \partial_v} \prod_v H_s^1(E_v, V)$$

exact. Here the product is taken over all places of E , we set

$$H_s^1(E_v, V) := H^1(E_v, V)/H_f^1(E_v, V),$$

the symbol $H_f^1(E_v, V)$ denotes the local Bloch–Kato condition at v (see [18, §2.4]) and

$$\partial_v : H^1(E, V) \longrightarrow H_s^1(E_v, V)$$

is the composition of the restriction $H^1(E, V) \rightarrow H^1(E_v, V)$ with the canonical projection. If T is a G_E -stable lattice in V then set $A := V/T$ and, for every integer $n \geq 1$, let A_{p^n} denote the p^n -torsion of A . There is a canonical isomorphism $A_{p^n} \simeq T/p^n T$.

The projection $p : V \rightarrow A$ and the inclusion $i : T \hookrightarrow V$ induce maps

$$p : H^1(E, V) \longrightarrow H^1(E, A), \quad i : H^1(E, T) \longrightarrow H^1(E, V);$$

let us define $H_f^1(E, A) := p(H_f^1(E, V))$ and $H_f^1(E, T) := i^{-1}(H_f^1(E, V))$. Furthermore, the inclusion $i_n : A_{p^n} \hookrightarrow A$ and the projection $p_n : T \rightarrow T/p^n T$ induce maps

$$i_n : H^1(E, A_{p^n}) \longrightarrow H^1(E, A), \quad p_n : H^1(E, T) \longrightarrow H^1(E, T/p^n T);$$

we set $H_f^1(E, T/p^n T) := p_n(H_f^1(E, T))$ and $H_f^1(E, A_{p^n}) := i_n^{-1}(H_f^1(E, A))$. It can be checked that the isomorphisms $A_{p^n} \simeq T/p^n T$ induce isomorphisms $H_f^1(E, A_{p^n}) \simeq H_f^1(E, T/p^n T)$ between Selmer groups.

Now let $M \in \{V, T, A, A_{p^n}, T/p^n T\}$. If L/E is a finite extension of number fields then restriction and corestriction induce maps

$$\text{res}_{L/E} : H_f^1(E, M) \longrightarrow H_f^1(L, M), \quad \text{cores}_{L/E} : H_f^1(L, M) \longrightarrow H_f^1(E, M).$$

Finally, if E is a number field and ℓ is a prime number then we set

$$H_f^i(E_\ell, M) := \bigoplus_{\lambda|\ell} H_f^i(E_\lambda, M), \quad H_s^i(E_\ell, M) := \bigoplus_{\lambda|\ell} H_s^i(E_\lambda, M), \quad \partial_\ell := \bigoplus_{\lambda|\ell} \partial_\lambda,$$

the direct sums being taken over the primes λ of E above ℓ .

2.2. The anticyclotomic \mathbb{Z}_p -extension of K . For every integer $m \geq 0$ write H_{p^m} for the ring class field of K of conductor p^m . By assumption, $p \nmid h_K = |\mathrm{Gal}(H_1/K)|$ and, since p is unramified in K , we also have $p \nmid |\mathrm{Gal}(H_p/H_1)|$. Moreover, $\mathrm{Gal}(H_{p^{m+1}}/H_p) \simeq \mathbb{Z}/p^m\mathbb{Z}$ for all $m \geq 1$. It follows that for every $m \geq 1$ there is a splitting

$$\mathrm{Gal}(H_{p^{m+1}}/K) \simeq G_m \times \Delta$$

with $G_m \simeq \mathbb{Z}/p^m\mathbb{Z}$ and $\Delta \simeq \mathrm{Gal}(H_p/K)$ of order prime to p . For every $m \geq 1$ define K_m as the fixed field of G_m ; then K_m is a subfield of $H_{p^{m+1}}$ such that

$$G_m = \mathrm{Gal}(K_m/K) \simeq \mathbb{Z}/p^m\mathbb{Z}.$$

The field $K_\infty := \cup_{m \geq 1} K_m$ is the anticyclotomic \mathbb{Z}_p -extension of K ; equivalently, it is the \mathbb{Z}_p -extension of K that is (generalized) dihedral over \mathbb{Q} . Set

$$G_\infty := \varprojlim_m G_m = \mathrm{Gal}(K_\infty/K) \simeq \mathbb{Z}_p.$$

For every integer $m \geq 0$ define $\Gamma_m := \mathrm{Gal}(K_\infty/K_m)$. Furthermore, for every $m \geq 1$ set $\Lambda_m := \mathcal{O}_p[G_m]$ and define

$$\Lambda := \varprojlim_m \Lambda_m = \mathcal{O}_p[[G_\infty]].$$

Here the inverse limit is taken with respect to the maps induced by the natural projections $G_{m+1} \rightarrow G_m$. For all $m \geq 1$ fix a generator γ_m of G_m in such a way that $\gamma_{m+1}|_{K_m} = \gamma_m$; then $\gamma_\infty := (\gamma_1, \dots, \gamma_m, \dots)$ is a topological generator of G_∞ . It is well known that the map

$$(2) \quad \Lambda \xrightarrow{\simeq} \mathcal{O}_p[[X]], \quad \gamma_\infty \longmapsto 1 + X$$

is an isomorphism of \mathcal{O}_p -algebras (see, e.g., [26, Proposition 5.3.5]).

For an abelian pro- p group M write $M^\vee := \mathrm{Hom}_{\mathbb{Z}_p}^{\mathrm{cont}}(M, \mathbb{Q}_p/\mathbb{Z}_p)$ for its Pontryagin dual, equipped with the compact-open topology (here $\mathrm{Hom}_{\mathbb{Z}_p}^{\mathrm{cont}}$ denotes continuous homomorphisms of \mathbb{Z}_p -modules and $\mathbb{Q}_p/\mathbb{Z}_p$ is discrete). In the rest of the paper it will be convenient to use also the alternative definition $M^\vee := \mathrm{Hom}_{\mathcal{O}_p}^{\mathrm{cont}}(M, F_p/\mathcal{O}_p)$, where $\mathrm{Hom}_{\mathcal{O}_p}^{\mathrm{cont}}$ stands for continuous homomorphisms of \mathcal{O}_p -modules and F_p/\mathcal{O}_p is given the discrete topology. It turns out that the two definitions are equivalent, as there is a non-canonical isomorphism between $\mathrm{Hom}_{\mathcal{O}_p}^{\mathrm{cont}}(M, F_p/\mathcal{O}_p)$ and $\mathrm{Hom}_{\mathbb{Z}_p}^{\mathrm{cont}}(M, \mathbb{Q}_p/\mathbb{Z}_p)$ that depends on the choice of a \mathbb{Z}_p -basis of \mathcal{O}_p . See, e.g., [8, Lemma 2.4] for details.

2.3. Torsion subgroups in compact modules. In this subsection and the next we collect results on Pontryagin duals that will be used in the sequel and for which we were unable to find convenient references in the literature.

Let R be a topological commutative noetherian ring. In the next sections R will always be a commutative noetherian local ring (R, \mathfrak{m}) that is an \mathfrak{m} -adically complete \mathbb{Z}_p -algebra. Moreover, let M denote a compact Hausdorff abelian group, endowed with a structure of a topological R -module. For an ideal $I \subset R$ let $M[I]$ denote the I -torsion submodule of M , that is

$$M[I] := \{m \in M \mid xm = 0 \text{ for all } x \in I\}.$$

Finally, equip the Pontryagin dual M^\vee with the R -module structure given by $(x\varphi)(m) := \varphi(xm)$ for all $\varphi \in M^\vee$, $m \in M$ and $x \in R$.

Lemma 2.1. *There is a canonical isomorphism of R -modules*

$$M^\vee/IM^\vee \simeq M[I]^\vee.$$

Proof. Write $I = (x_1, \dots, x_n)$ and consider the map

$$\xi : M \longrightarrow \prod_{i=1}^n x_i M, \quad m \longmapsto (x_i m)_{i=1, \dots, n},$$

whose kernel is equal to $M[I]$. If $i : M[I] \hookrightarrow M$ denotes inclusion then Pontryagin duality gives an exact sequence of R -modules

$$(3) \quad \left(\prod_{i=1}^n x_i M \right)^\vee \xrightarrow{\xi^\vee} M^\vee \xrightarrow{i^\vee} M[I]^\vee \longrightarrow 0;$$

here the surjectivity of i^\vee is a consequence of $M[I]$ being closed in M , hence compact. On the other hand, sending $(\varphi_1, \dots, \varphi_n)$ to $\sum_i \varphi_i$ gives an isomorphism between $\prod_i (x_i M)^\vee$ and $(\prod_i x_i M)^\vee$, so we can rewrite (3) as

$$(4) \quad \prod_{i=1}^n (x_i M)^\vee \xrightarrow{\xi^\vee} M^\vee \xrightarrow{i^\vee} M[I]^\vee \longrightarrow 0.$$

In light of (4), we want to check that $\text{im}(\xi^\vee) = IM^\vee$. First of all, let $\varphi = \sum_{i=1}^n x_i \varphi_i \in IM^\vee$, with $\varphi_i \in M^\vee$ for all $i = 1, \dots, n$. Then $\varphi = \xi^\vee((\varphi_1|_{x_1 M}, \dots, \varphi_n|_{x_n M}))$, which shows that $\varphi \in \text{im}(\xi^\vee)$. Conversely, let $\varphi \in \text{im}(\xi^\vee)$; by definition, for every $i = 1, \dots, n$ there exists $\varphi_i \in (x_i M)^\vee$ such that $\varphi = \xi^\vee((\varphi_1, \dots, \varphi_n))$. For every i , the R -module $x_i M$ is compact because M is, hence the inclusion $x_i M \hookrightarrow M$ gives a surjection $M^\vee \twoheadrightarrow (x_i M)^\vee$. Now for every $i = 1, \dots, n$ choose a lift $\psi_i \in M^\vee$ of φ_i . It follows that $\varphi = \sum_{i=1}^n x_i \psi_i \in IM^\vee$, and the proof is complete. \square

2.4. Application to Iwasawa algebras. As in §2.2, let γ_∞ be a topological generator of $G_\infty \simeq \mathbb{Z}_p$. Let $I_\infty = (\gamma_\infty - 1)$ be the augmentation ideal of Λ and for every integer $n \geq 0$ consider the ideal $I_n := (\gamma_\infty^{p^n} - 1)$ of Λ ; in particular, $I_0 = I_\infty$.

Now let M be a continuous Λ -module. As before, the dual M^\vee inherits a structure of continuous Λ -module. Since $\gamma_\infty^{p^n}$ is a topological generator of Γ_n , for all $n \geq 0$ there are equalities

$$(5) \quad M[I_n] = M^{\Gamma_n}, \quad M_{\Gamma_n} = M/I_n M.$$

Proposition 2.2. *If M is compact then*

$$(M^\vee)_{\Gamma_n} = M^\vee / I_n M^\vee \simeq (M^{\Gamma_n})^\vee$$

for every $n \geq 0$.

Proof. The equality on the left is just the second equality in (5) applied to M^\vee , while the canonical isomorphism on the right follows upon taking $I = I_n$ in Lemma 2.1 and using the first equality in (5). \square

2.5. The Control Theorem. Fix a normalized newform f of weight k , a prime \mathfrak{p} and an imaginary quadratic field K as in Assumption 1.1.

Write T for the $G_{\mathbb{Q}}$ -representation considered by Nekovář in [24, Proposition 3.1], where it is denoted by $A_{\mathfrak{p}}$. This is a free $\mathcal{O}_{\mathfrak{p}}$ -module of rank 2. The $G_{\mathbb{Q}}$ -representation $V := T \otimes_{\mathcal{O}_{\mathfrak{p}}} F_{\mathfrak{p}}$ is then the $k/2$ -twist of the representation $V_{f, \mathfrak{p}}$. Finally, define the $G_{\mathbb{Q}}$ -representation $A := V/T$. As above, we shall write A_{p^n} for the p^n -torsion submodule of A . Observe that

$$(6) \quad A = \bigcup_{n \geq 1} A_{p^n} = \varinjlim_n A_{p^n}$$

where the direct limit is taken with respect to the natural inclusions $A_{p^n} \hookrightarrow A_{p^{n+1}}$.

Lemma 2.3. $H^0(K_m, A) = 0$ for all integers $m \geq 0$.

Proof. Fix an integer $m \geq 0$. The extension K_m/\mathbb{Q} is solvable, so $H^0(K_m, A_{p^n}) = 0$ for all $n \geq 0$ by [18, Lemma 3.10, (2)]. It follows from (6) that

$$H^0(K_m, A) = H^0\left(K_m, \varinjlim_n A_{p^n}\right) = \varinjlim_n H^0(K_m, A_{p^n}) = 0,$$

as was to be shown. \square

To obtain a version of Mazur's Control Theorem ([19]) in our setting, we make the following

Assumption 2.4. (1) For all integers $m \geq 1$ and all places v of K_m dividing N , the group $H^0(I_{G_{m,v}}, A)$ is p -divisible.

(2) For all m and all places v of K_m above p , the canonical projection

$$\varprojlim_n H_f^1(K_{n,v}, T^*(1)) \longrightarrow H_f^1(K_{m,v}, T^*(1))$$

is surjective, where $T^* := \text{Hom}_{\mathbb{Z}_p}(T, \mathbb{Z}_p)$ is the \mathbb{Z}_p -linear dual of T and the inverse limit is taken with respect to the corestriction maps.

Remark 2.5. Condition (2) in Assumption 2.4 is standard in the case of elliptic curves over \mathbb{Q} (see, e.g., [3, Assumption 2.15] and [4, §1.1]) and is satisfied for all but finitely many primes p . Also, conditions similar to (1) in Assumption 2.4 are discussed in [5, Assumption 2.1] in the case of elliptic curves E . It would be interesting to investigate analogous results for the representation $V_{f,p}$.

Define the discrete Λ -module

$$H_f^1(K_\infty, A) := \varinjlim_m H_f^1(K_m, A),$$

the inductive limit being taken with respect to the restriction maps.

Theorem 2.6 (Control Theorem). *For every integer $m \geq 0$ there is an isomorphism*

$$\text{res}_{K_\infty/K_m} : H_f^1(K_m, A) \xrightarrow{\cong} H_f^1(K_\infty, A)^{\Gamma_m}.$$

Sketch of proof. This is essentially [27, Theorem 2.4]. Fix an integer $m \geq 0$. To begin with, the inflation-restriction exact sequence reads

$$(7) \quad 0 \longrightarrow H^1(\text{Gal}(K_\infty/K_m), H^0(K_\infty, A)) \longrightarrow H^1(K_m, A) \longrightarrow H^1(K_\infty, A)^{\Gamma_m}.$$

On the other hand, $H^0(K_m, A) = 0$ for all m by Lemma 2.3, hence

$$H^0(K_\infty, A) = \varinjlim_m H^0(K_m, A) = 0.$$

Thus sequence (7) gives an injection $H^1(K_m, A) \hookrightarrow H^1(K_\infty, A)^{\Gamma_m}$, which in turn restricts to an injection

$$\text{res}_{K_\infty/K_m} : H_f^1(K_m, A) \hookrightarrow H_f^1(K_\infty, A)^{\Gamma_m}$$

between Selmer groups. Finally, by comparing with the proof of [27, Theorem 2.4] (see, in particular, [27, p. 81]), one can check that Assumption 2.4 forces $\text{res}_{K_\infty/K_m}$ to be surjective as well. \square

Let

$$\mathcal{X}_\infty := \text{Hom}_{\mathcal{O}_p}^{\text{cont}}(H_f^1(K_\infty, A), F_p/\mathcal{O}_p)$$

be the Pontryagin dual of $H_f^1(K_\infty, A)$, equipped with its canonical structure of compact Λ -module. For every integer $m \geq 0$ let

$$\mathcal{X}_m := \text{Hom}_{\mathcal{O}_p}^{\text{cont}}(H_f^1(K_m, A), F_p/\mathcal{O}_p)$$

be the Pontryagin dual of $H_f^1(K_m, A)$. Each \mathcal{X}_m has a natural structure of Λ_m -module and there is a canonical isomorphism of Λ -modules $\mathcal{X}_\infty \simeq \varprojlim_m \mathcal{X}_m$. Note that, since the Galois representation A is unramified outside Np , the \mathcal{O}_p -modules \mathcal{X}_m are finitely generated.

Corollary 2.7. *For every $m \geq 0$ there is a canonical isomorphism $(\mathcal{X}_\infty)_{\Gamma_m} \simeq \mathcal{X}_m$.*

Proof. Fix an integer $m \geq 0$. Thanks to Proposition 2.2, the isomorphism of Theorem 2.6 gives, by duality, a canonical isomorphism

$$(8) \quad \mathcal{X}_\infty / I_m \mathcal{X}_\infty \simeq \mathcal{X}_m.$$

But, again by Proposition 2.2, the quotient $\mathcal{X}_\infty / I_m \mathcal{X}_\infty$ is canonically isomorphic to $(\mathcal{X}_\infty)_{\Gamma_m}$, and we are done. \square

Corollary 2.8. *The Λ -module \mathcal{X}_∞ is finitely generated.*

Proof. By choosing $m = 0$ in (8), we obtain an isomorphism $\mathcal{X}_\infty / I_\infty \mathcal{X}_\infty \simeq \mathcal{X}_0$. Since \mathcal{X}_0 is a finitely generated \mathcal{O}_p -module, the result follows from a topological version of Nakayama's lemma ([8, Corollary 1.5] or [26, Corollary 5.2.18, (ii)]). \square

2.6. Projective Selmer modules. For every integer $m \geq 0$ define the \mathcal{O}_p -module

$$S_m := \varprojlim_n H_f^1(K_m, A_{p^n})$$

where the inverse limit is taken with respect to the multiplication-by- p maps. The Tate module of $H_f^1(K_m, A)$ is the \mathcal{O}_p -module

$$\mathrm{Ta}_p(H_f^1(K_m, A)) := \varprojlim_n H_f^1(K_m, A)_{p^n}$$

where $H_f^1(K_m, A)_{p^n}$ denotes the p^n -torsion submodule of $H_f^1(K_m, A)$ and, again, the inverse limit is taken with respect to the multiplication-by- p maps.

Lemma 2.9. *For all $m, n \geq 0$ there is an isomorphism*

$$H_f^1(K_m, A_{p^n}) \simeq H_f^1(K_m, A)_{p^n}.$$

Proof. Fix integers $m, n \geq 0$. Taking G_{K_m} -cohomology of the short exact sequence

$$0 \longrightarrow A_{p^n} \longrightarrow A \xrightarrow{p^n} A \longrightarrow 0,$$

where p^n denotes the multiplication-by- p^n map, and using Lemma 2.3, we obtain isomorphisms

$$H^1(K_m, A_{p^n}) \simeq H^1(K_m, A)_{p^n}$$

for all $n \geq 0$. These restrict to isomorphisms

$$H_f^1(K_m, A_{p^n}) \simeq H_f^1(K_m, A)_{p^n},$$

as desired. \square

Proposition 2.10. *For every $m \geq 0$ there is a canonical isomorphism*

$$S_m \simeq \mathrm{Ta}_p(H_f^1(K_m, A)).$$

Proof. Fix an integer $m \geq 0$. Passing to inverse limits over n in Lemma 2.9 with respect to the multiplication-by- p maps gives the result. \square

We need one more auxiliary result on Tate modules.

Lemma 2.11. *For every $m \geq 0$ there is a canonical isomorphism*

$$\mathrm{Ta}_p(H_f^1(K_m, A)) \simeq \mathrm{Hom}_{\mathcal{O}_p}(\mathcal{X}_m, \mathcal{O}_p).$$

Proof. Fix an integer $m \geq 0$. First of all, $F_p = \mathcal{O}_p[1/p]$ because p is unramified in F by Assumption 1.1, so there is an identification

$$(9) \quad \mathrm{Ta}_p(H_f^1(K_m, A)) = \mathrm{Hom}_{\mathcal{O}_p}(F_p/\mathcal{O}_p, H_f^1(K_m, A)).$$

On the other hand, taking Pontryagin duals gives a canonical isomorphism

$$(10) \quad \mathrm{Hom}_{\mathcal{O}_p}(F_p/\mathcal{O}_p, H_f^1(K_m, A)) \simeq \mathrm{Hom}_{\mathcal{O}_p}(\mathcal{X}_m, \mathcal{O}_p).$$

The lemma follows by combining (9) and (10). \square

Consider the Λ -modules

$$S_\infty := \varprojlim_m S_m, \quad \varprojlim_m \mathrm{Ta}_p(H_f^1(K_m, A)),$$

the inverse limits being taken with respect to the corestriction maps. There is a sequence of isomorphisms of Λ -modules

$$S_\infty \simeq \varprojlim_m \mathrm{Ta}_p(H_f^1(K_m, A)) \simeq \varprojlim_m \mathrm{Hom}_{\mathcal{O}_p}(\mathcal{X}_m, \mathcal{O}_p) \simeq \varprojlim_m \mathrm{Hom}_{\mathcal{O}_p}((\mathcal{X}_\infty)_{\Gamma_m}, \mathcal{O}_p)$$

where the first isomorphism comes from Proposition 2.10, the second from Lemma 2.11 and the third from Corollary 2.7. Using [28, Lemma 4, (ii)], whose proof can be extended to the case of Iwasawa algebras with coefficients in \mathcal{O}_p , we obtain canonical isomorphisms

$$(11) \quad S_\infty \simeq \varprojlim_m \mathrm{Ta}_p(H_f^1(K_m, A)) \simeq \mathrm{Hom}_\Lambda(\mathcal{X}_\infty, \Lambda)$$

of Λ -modules. It follows from Corollary 2.8 that S_∞ is finitely generated over Λ .

Proposition 2.12. *The Λ -module S_∞ is free of finite rank.*

Proof. By (11), the Λ -module $(S_\infty)_{G_\infty}$ is isomorphic to $\mathrm{Hom}_\Lambda(\mathcal{X}_\infty, \Lambda)_{G_\infty}$, and this injects into the free \mathcal{O}_p -module $\mathrm{Hom}_{\mathcal{O}_p}((\mathcal{X}_\infty)_{G_\infty}, \mathcal{O}_p)$ thanks to a straightforward extension of [28, Lemma 4, (iii)] to our case, where Iwasawa algebras have coefficients in \mathcal{O}_p . Therefore $(S_\infty)_{G_\infty}$ is a free \mathcal{O}_p -module. Since S_∞ is torsion-free over Λ by (11), by [26, Proposition 5.3.19, (ii)] (again, generalized to Iwasawa algebras with coefficients in \mathcal{O}_p) we conclude that S_∞ is free of finite rank over Λ . \square

Definition 2.13. The *pro- p Bloch–Kato Selmer group of f over K_∞* is the Λ -module

$$\hat{H}_f^1(K_\infty, T) := \varprojlim_m H_f^1(K_m, T),$$

where the inverse limit is taken with respect to the corestriction maps.

The following result is a straightforward consequence of Definition 2.13.

Proposition 2.14. *The Λ -modules $\hat{H}_f^1(K_\infty, T)$ and S_∞ are canonically isomorphic.*

Proof. By [31, p. 261, Corollary], there is a canonical isomorphism

$$H^1(K_m, T) \simeq \varprojlim_n H^1(K_m, T/p^n T)$$

where the inverse limit is taken with respect to the maps induced by the canonical projections $T/p^{n+1}T \rightarrow T/p^n T$. One easily shows that this isomorphism restricts to an isomorphism between Selmer groups, so we obtain a sequence of canonical isomorphisms

$$\hat{H}_f^1(K_\infty, T) \simeq \varprojlim_m \varprojlim_n H_f^1(K_m, T/p^n T) \simeq \varprojlim_m \varprojlim_n H_f^1(K_m, A_{p^n}) = S_\infty$$

(here recall that the inverse limit over n of the groups $H_f^1(K_m, A_{p^n})$ is computed with respect to the multiplication-by- p maps, while all the inverse limits over m are taken with respect to the corestriction maps). \square

Combining Propositions 2.12 and 2.14, we obtain

Corollary 2.15. *The Λ -module $\hat{H}_f^1(K_\infty, T)$ is free of finite rank.*

3. IWASAWA MODULES OF HEEGNER CYCLES

3.1. Preliminaries on Heegner cycles. We briefly recall construction and basic properties of Heegner cycles, following [24, Section 5] closely. This will also give us the occasion to fix some notation that will be used in the rest of the paper.

Choose an ideal $\mathcal{N} \subset \mathcal{O}_K$ such that $\mathcal{O}_K/\mathcal{N} \simeq \mathbb{Z}/N\mathbb{Z}$, which exists thanks to the Heegner hypothesis satisfied by K . For every integer $c \geq 1$ prime to N and the discriminant of K , let $\mathcal{O}_c := \mathbb{Z} + c\mathcal{O}_K$ be the order of K of conductor c . Let $X_0(N)$ be the compact modular curve of level $\Gamma_0(N)$; the isogeny $\mathbb{C}/\mathcal{O}_c \rightarrow \mathbb{C}/(\mathcal{O}_c \cap \mathcal{N})^{-1}$ defines a Heegner point $x_c \in X_0(N)$ which, by complex multiplication, is rational over the ring class field H_c of K of conductor c . Furthermore, let $X(N)$ be the compact modular curve of level $\Gamma(N)$.

Pick $\tilde{x}_c \in \kappa^{-1}(x_c)$. The elliptic curve E_c with full level N structure corresponding to \tilde{x}_c has complex multiplication by \mathcal{O}_c . Fix the unique square root $\xi_c = \sqrt{-Dc}$ of the discriminant of \mathcal{O}_c with positive imaginary part under the chosen embedding of K into \mathbb{C} . For all $a \in \mathcal{O}_c$ write $\Gamma_{c,a} \subset E_c \times E_c$ for the graph of a and $i_{\tilde{x}_c} : \pi_{k-2}^{-1}(\tilde{x}_c) \hookrightarrow \tilde{\mathcal{E}}_N^{k-2}$ for the natural inclusion, where $\tilde{\mathcal{E}}_N^{k-2}$ is the Kuga–Sato variety of level N and weight k . One can construct two canonical projectors Π_B and Π_ϵ (see [24, Sections 2 and 3] for definitions) acting on the Chow groups $\mathrm{CH}^{k/2}(\tilde{\mathcal{E}}_N^{k-2}/H_c) \otimes \mathbb{Z}_p$. This gives an element

$$(12) \quad \Pi_B \Pi_\epsilon (i_{\tilde{x}_c})_* \left(\Gamma_{m, \xi_c}^{(k-2)/2} \right) \in \Pi_B \Pi_\epsilon (\mathrm{CH}^{k/2}(\tilde{\mathcal{E}}_N^{k-2}/H_{p^m}) \otimes \mathbb{Z}_p).$$

On the other hand, the Abel–Jacobi map for \mathcal{E}_N^{k-2} , combined with the projectors Π_B and Π_ϵ , gives a map

$$(13) \quad \Pi_B \Pi_\epsilon (\mathrm{CH}^{k/2}(\tilde{\mathcal{E}}_N^{k-2}/H_c) \otimes \mathbb{Z}_p) \longrightarrow H_{\mathrm{cont}}^1(H_c, T)$$

(see [24, Section 4] for details). Then one can define the *Heegner cycle* y_c in $H_{\mathrm{cont}}^1(H_c, T)$ as the image of the cycle (12) via the Abel–Jacobi map (13). This class is independent of the choice of \tilde{x}_c ([24, p. 107]). Finally, since the image of the Abel–Jacobi map is contained in the Bloch–Kato Selmer group, it turns out that

$$y_c \in H_f^1(H_c, T).$$

Remark 3.1. In [24], the Heegner cycle y_c is introduced only for c square-free and coprime to NDp , where D is the discriminant of K . However, one can readily check that the construction of y_c carries over without change to our more general setting (in fact, our interest in Iwasawa-theoretic considerations will lead us to specialize to the case where c is a power of p).

3.2. Hecke action. Corestriction from $H_{p^{m+1}}$ to K_m gives an element

$$\alpha_m := \mathrm{cores}_{H_{p^{m+1}}/K_m}(y_{p^{m+1}}) \in H_f^1(K_m, T).$$

Put $K_0 := K$ and define

$$\alpha_0 := \mathrm{cores}_{H_p/K}(y_p) \in H_f^1(K, T).$$

As in the introduction, set $a'_p := a_p/p^{k/2-1}$. Recall that, by Assumption 1.1, a'_p belongs to \mathcal{O}_p^\times and $a'_p \not\equiv 2 \pmod{p}$ (respectively, $a'_p \not\equiv 1 \pmod{p}$) if p splits in K (respectively, if p is inert in K).

Lemma 3.2. *If $m \geq 2$ then*

$$\mathrm{cores}_{K_{m+1}/K_m}(\alpha_{m+1}) = a'_p \alpha_m - \mathrm{res}_{K_m/K_{m-1}}(\alpha_{m-1}),$$

and

$$\text{cores}_{K_1/K}(\alpha_1) = \begin{cases} (a'_p - a'_p{}^{-1}(p+1))\alpha_0 & \text{if } p \text{ is inert in } K, \\ (a'_p - (a'_p - 2)^{-1}(p-1))\alpha_0 & \text{if } p \text{ splits in } K. \end{cases}$$

Proof. Combine the proof of [24, Proposition 5.4] and [28, Lemma 2, p. 432]. \square

3.3. Projective Heegner modules. In order to introduce Iwasawa modules of Heegner cycles, let us first record the following

Lemma 3.3. *Let $m \geq 1$ be an integer and let G be a non-trivial subgroup of G_m . For all $u \in \Lambda_m$ the element $a_p - u \sum_{\sigma \in G} \sigma$ is invertible in Λ_m .*

Proof. For simplicity, set $\beta := a_p - u \sum_{\sigma \in G} \sigma$. Suppose that $|G| = p^t$ with $1 \leq t \leq m$ and let $b := p^{m-t}$, so that G is generated by γ_m^b . Write $\pi_m : \Lambda \rightarrow \Lambda_m$ for the canonical projection, fix a lift $\tilde{u} \in \Lambda$ of u under π_m and let $G(X) \in \mathcal{O}_p[[X]]$ be the power series corresponding to \tilde{u} under isomorphism (2), which we can assume to be a polynomial. The element

$$\tilde{\beta} := a_p - \tilde{u} \cdot \sum_{i=1}^{p^t} \gamma_\infty^{bi} \in \Lambda$$

is then a lift of β via π_m that is sent by isomorphism (2) to

$$F(X) := a_p - G(X) \cdot \sum_{i=1}^{p^t} (1+X)^{bi} = a_p - p^t \eta + X f(X) \in \mathcal{O}_p[[X]]$$

for certain $\eta \in \mathcal{O}_p$ and $f(X) \in \mathcal{O}_p[X]$. But $a_p - p^t \eta \in \mathcal{O}_p^\times$ because a_p is a p -adic unit by Assumption 1.1, hence $F(X)$ is invertible in $\mathcal{O}_p[[X]]$. It follows that $\tilde{\beta}$ is invertible in Λ , and then $\beta = \pi_m(\tilde{\beta})$ is invertible in Λ_m , as required. \square

For every integer $m \geq 0$ denote by \mathcal{H}_m the Λ_m -submodule of $H_f^1(K_m, T)$ generated by α_m .

Proposition 3.4. *Corestriction induces surjective maps*

$$\text{cores}_{K_{m+1}/K_m} : \mathcal{H}_{m+1} \longrightarrow \mathcal{H}_m$$

for all integers $m \geq 0$.

Proof. As in the proof of [1, Proposition 4], we proceed by induction on m . By Assumption 1.1, the number $a'_p - a'_p{}^{-1}(p+1)$ (respectively, $a'_p - (a'_p - 2)^{-1}(p-1)$) is a p -adic unit when p is inert in K (respectively, p splits in K), and then the second formula in Lemma 3.2 proves the proposition for $m = 0$. Now suppose that the claim is true for $m - 1$. Then

$$(14) \quad \alpha_{m-1} = u \cdot \text{cores}_{K_m/K_{m-1}}(\alpha_m)$$

for some $u \in \Lambda_{m-1}^\times$, and combining this equality with the first formula in Lemma 3.2 gives

$$(15) \quad \text{cores}_{K_{m+1}/K_m}(\alpha_{m+1}) = (a'_p - u \cdot \text{cores}_{K_m/K_{m-1}})(\alpha_m).$$

Since $a'_p - u \cdot \text{cores}_{K_m/K_{m-1}} \in \Lambda_m^\times$ by Lemma 3.3, the claim for m follows from (15). \square

Definition 3.5. The *Iwasawa module of Heegner cycles* is the compact Λ -module

$$\mathcal{H}_\infty := \varprojlim_m \mathcal{H}_m \subset \hat{H}_f^1(K_\infty, T),$$

where the inverse limit is taken with respect to the corestriction maps.

The proof of the following result crucially exploits a theorem of Howard ([14]) that extends to the higher weight setting results of Cornut ([9]) on the generic non-vanishing of Heegner points on elliptic curves over anticyclotomic \mathbb{Z}_p -extensions.

Theorem 3.6. *The Λ -module \mathcal{H}_∞ is free of rank 1.*

Proof. First of all, observe that \mathcal{H}_∞ is a Λ -submodule of $\hat{H}_f^1(K_\infty, T)$, which is free of finite rank over Λ by Corollary 2.15. It follows that \mathcal{H}_∞ , being cyclic, is either trivial or isomorphic to Λ . On the other hand, [14, Theorem A] ensures that there is an integer $m \geq 0$ such that α_m is not \mathcal{O}_p -torsion in $H_f^1(K_m, T)$. Set $x_m := \alpha_m \in \mathcal{H}_m$ and for every $n \geq m + 1$ choose $x_n \in \mathcal{H}_n$ such that

$$\text{cores}_{K_n/K_{n-1}}(x_n) = x_{n-1}.$$

Since x_m is non-torsion, the \mathcal{O}_p -submodule of \mathcal{H}_∞ generated by this compatible sequence is isomorphic to \mathcal{O}_p , so \mathcal{H}_∞ cannot be trivial. \square

3.4. Injective Heegner modules. Now we introduce the Λ -module \mathcal{E}_∞ , which is obtained by taking an inductive limit of Heegner cycles. First note the following easy consequence of Proposition 3.4.

Corollary 3.7. *Restriction induces injective maps*

$$\text{res}_{K_{m+1}/K_m} : \mathcal{H}_m \hookrightarrow \mathcal{H}_{m+1}$$

for all integers $m \geq 0$.

Proof. Since res_{K_{m+1}/K_m} is injective, it suffices to show that its image is contained in \mathcal{H}_{m+1} . By Proposition 3.4, we know that $\alpha_m = u_m \cdot \text{cores}_{K_{m+1}/K_m}(\alpha_{m+1})$ for some $u_m \in \Lambda_m^\times$, and the corollary follows by applying restriction and using (1). \square

For every integer $m \geq 0$ let $\bar{\alpha}_m$ be the image of α_m in $H_f^1(K_m, T/p^m T)$ via the map induced by the projection $T \rightarrow T/p^m T$. Denote by β_m the image of $\bar{\alpha}_m$ under the map induced by the isomorphism $T/p^m T \simeq A_{p^m}$, then define $\mathcal{E}_m := \Lambda_m \beta_m$ as the Λ_m -submodule of $H_f^1(K_m, A_{p^m})$ generated by β_m . By Corollary 3.7, res_{K_{m+1}/K_m} sends the Λ_m -module generated by $\bar{\alpha}_m$ to the Λ_{m+1} -module generated by $\bar{\alpha}_{m+1}$. Therefore composing with the map obtained from the natural inclusion $A_{p^m} \subset A_{p^{m+1}}$ gives canonical maps $\rho_m : \mathcal{E}_m \rightarrow \mathcal{E}_{m+1}$, and we can consider the discrete Λ -module

$$\mathcal{E}_\infty := \varinjlim_m \mathcal{E}_m,$$

the direct limit being taken with respect to the maps ρ_m .

Proposition 3.8. *There is an injection of Λ -modules $\mathcal{E}_\infty \hookrightarrow H_f^1(K_\infty, A)$.*

Proof. By Lemma 2.9, there are isomorphisms

$$H_f^1(K_m, A_{p^m}) \simeq H_f^1(K_m, A)_{p^m}$$

for all integers $m \geq 0$. By definition, every \mathcal{E}_m is a submodule of $H_f^1(K_m, A_{p^m})$, and then the (left) exactness of the direct limit gives injections

$$\begin{aligned} \mathcal{E}_\infty = \varinjlim_m \mathcal{E}_m &\hookrightarrow \varinjlim_m H_f^1(K_m, A_{p^m}) \simeq \varinjlim_m H_f^1(K_m, A)_{p^m} \\ &\hookrightarrow \varinjlim_m H_f^1(K_m, A) = H_f^1(K_\infty, A), \end{aligned}$$

which completes the proof. \square

Taking the Pontryagin dual, we get a compact Λ -module

$$\mathcal{E}_\infty^\vee := \text{Hom}_{\mathcal{O}_p}^{\text{cont}}(\mathcal{E}_\infty, F_p/\mathcal{O}_p) \simeq \varprojlim_m \mathcal{E}_m^\vee$$

where, for each m , the module $\mathcal{E}_m^\vee := \text{Hom}_{\mathcal{O}_p}^{\text{cont}}(\mathcal{E}_m, F_p/\mathcal{O}_p)$ is the Pontryagin dual of \mathcal{E}_m . Proposition 3.8 gives a surjection of Λ -modules

$$(16) \quad \pi : \mathcal{X}_\infty \twoheadrightarrow \mathcal{E}_\infty^\vee.$$

In particular, there is also an injection of Λ -modules

$$(17) \quad \mathrm{Hom}_\Lambda(\mathcal{E}_\infty^\vee, \Lambda) \hookrightarrow \mathrm{Hom}_\Lambda(\mathcal{X}_\infty, \Lambda) \simeq \hat{H}_f^1(K_\infty, T),$$

where the isomorphism on the right is a consequence of (11) and Proposition 2.14. The next result describes the image of this map.

Proposition 3.9. *The image of (17) is equal to \mathcal{H}_∞ .*

Proof. A more or less tautological, albeit somewhat tedious, diagram chasing. We omit the details. \square

It follows from Proposition 3.9 that there is an isomorphism

$$\mathrm{Hom}_\Lambda(\mathcal{E}_\infty^\vee, \Lambda) \simeq \mathcal{H}_\infty$$

of Λ -modules. In particular, Theorem 3.6 implies that \mathcal{E}_∞^\vee has rank 1 over Λ .

4. THE EULER SYSTEM ARGUMENT

The aim of this section is to prove Theorem 1.2; we restate it below.

Theorem 4.1. *The rank of \mathcal{X}_∞ over Λ is 1.*

By Theorem 3.6, the Λ -module \mathcal{H}_∞ of Heegner cycles is free of rank 1. Furthermore, the discussion of §3.4 shows that the Λ -rank of \mathcal{E}_∞^\vee is 1. Recall that, by Corollary 2.8, the Λ -module \mathcal{X}_∞ is finitely generated. The surjection $\pi : \mathcal{X}_\infty \twoheadrightarrow \mathcal{E}_\infty^\vee$ of Λ -modules introduced in (16) is $\mathrm{Gal}(K/\mathbb{Q})$ -equivariant. Since \mathcal{E}_∞^\vee has rank 1 over Λ , proving Theorem 4.1 is equivalent to showing that the Λ -module $\ker(\pi)$ is torsion. Let τ be the generator of $\mathrm{Gal}(K/\mathbb{Q})$. In order to prove Theorem 4.1 it suffices to show that every $y \in \ker(\pi)$ lying in an eigenspace for τ (i.e., such that $\tau(y) = \pm y$) is Λ -torsion.

As in §2.2, let γ_∞ be a topological generator of G_∞ . We need two lemmas.

Lemma 4.2. *There is a generator ω of the ideal $(\gamma_\infty - 1)\Lambda$ such that $\omega^\tau = -\omega$.*

Proof. Define $\omega := \gamma_\infty - \gamma_\infty^{-1} \in \Lambda$. Since $\gamma_\infty^\tau = \gamma_\infty^{-1}$, one has $\omega^\tau = -\omega$. On the other hand, one can write $\omega = \gamma_\infty^{-1}(\gamma_\infty + 1)(\gamma_\infty - 1)$. The isomorphism (2) sends $\gamma_\infty + 1$ to $2 + X$, which is invertible in $\mathcal{O}_p[[X]]$ because $p \neq 2$, and it follows that ω is a generator of $(\gamma_\infty - 1)\Lambda$. \square

Lemma 4.3. *Let $\epsilon \in \{\pm\}$ and suppose that every $y \in \ker(\pi)^\epsilon$ is Λ -torsion. Then every $y \in \ker(\pi)$ lying in an eigenspace for τ is Λ -torsion.*

Proof. By Lemma 4.2, there is a generator ω of $(\gamma_\infty - 1)\Lambda$ such that $\omega^\tau = -\omega$. Now pick $y \in \ker(\pi)^{-\epsilon}$. Then $\omega y \in \ker(\pi)$ and $\tau(\omega y) = \epsilon \omega y$, so $\omega y \in \ker(\pi)^\epsilon$. It follows that ωy is Λ -torsion, hence y is Λ -torsion as well. \square

Choose an element $x \in \mathcal{X}_\infty$ such that $\tau(x) = \epsilon x$ for some $\epsilon \in \{\pm\}$ and $\pi(x) \neq 0$. Thanks to Lemma 4.3, in order to prove Theorem 4.1 it is enough to show that every $y \in \ker(\pi)^{-\epsilon}$ is Λ -torsion. To do this, we will adapt the Λ -adic Euler system argument of [1], as explained in the next subsections.

4.1. Kolyvagin primes. Denote by

$$\rho_m : G_\mathbb{Q} \longrightarrow \mathrm{Aut}(A_{p^m}) \simeq \mathrm{GL}_2(\mathcal{O}_p/p^m\mathcal{O}_p)$$

the Galois representation on A_{p^m} and let $K(A_{p^m})$ be the composite of K and the field cut out by ρ_m ; in other words, $K(A_{p^m})$ is the composite of K and $\bar{\mathbb{Q}}^{\ker(\rho_m)}$. In particular, $K(A_{p^m})$ is Galois over \mathbb{Q} .

Definition 4.4. A prime number ℓ is a *Kolyvagin prime* relative to p^m if $\ell \nmid N D p$ and $\mathrm{Frob}_\ell = [\tau]$ in $\mathrm{Gal}(K(A_{p^m})/\mathbb{Q})$.

In particular, Kolyvagin primes are inert in K . Define

$$R_m := \mathcal{O}_{\mathfrak{p}}/p^m \mathcal{O}_{\mathfrak{p}}[G_m].$$

Likewise, set $D_m := \text{Gal}(K_m/\mathbb{Q})$, let \tilde{R}_m be the quotient of Λ_m (and of $\mathcal{O}_{\mathfrak{p}}[D_m]$) given by

$$\tilde{R}_m := \mathcal{O}_{\mathfrak{p}}/p^m \mathcal{O}_{\mathfrak{p}}[D_m]$$

and, for each choice of sign \pm , let $R_m^{(\pm)}$ be the \tilde{R}_m -module R_m with τ acting on group-like elements by $\gamma^\tau := \pm\gamma^{-1}$. In particular, $R_m^{(+)}$ corresponds to the linear extension of the natural action of τ on G_m .

Let ℓ be a Kolyvagin prime relative to p^m and let λ be the unique prime of K above ℓ ; let K_λ be the completion of K at λ . Evaluation at Frobenius gives a $\text{Gal}(K/\mathbb{Q})$ -equivariant isomorphism

$$H_f^1(K_\lambda, A_{p^m}) \simeq A_{p^m}$$

for every $m \geq 1$ (see, e.g., [6, Lemma 6.8]). Since λ splits completely in K_m , for each choice of sign \pm we obtain an isomorphism of \tilde{R}_m -modules

$$(18) \quad H_f^1(K_{m,\ell}, A_{p^m})^{(\pm)} \simeq R_m^{(\pm)}.$$

If ℓ is a Kolyvagin prime relative to p^m and $M \in \{V, T, A, A_{p^m}, T/p^m T\}$ then we let

$$\text{res}_\ell = \text{res}_{m,\ell} : H^1(K_m, M) \longrightarrow H^1(K_{m,\ell}, M)$$

be the direct sum $\bigoplus_{\lambda|\ell} \text{res}_\lambda$ of the local restrictions res_λ where the sum ranges over all the primes λ of K_m above ℓ . We will also use the same symbol for the restriction of res_ℓ to subgroups of $H^1(K_m, M)$.

4.2. Action of complex conjugation. In this subsection we study the action of $\text{Gal}(K/\mathbb{Q})$ on Selmer groups. These results will be used in §4.5 to show the existence of suitable families of Kolyvagin primes.

Define $\tilde{\Lambda} := \mathcal{O}_{\mathfrak{p}}[[D_\infty]]$ to be the Iwasawa algebra of $D_\infty := \text{Gal}(K_\infty/\mathbb{Q})$ with coefficients in $\mathcal{O}_{\mathfrak{p}}$. As before, for each sign \pm write $\tilde{\Lambda}^{(\pm)}$ for the ring $\tilde{\Lambda}$ viewed as a module over $\tilde{\Lambda}$ via the action of τ given by $\gamma^\tau = \pm\gamma^{-1}$ for all $\gamma \in G_\infty$. In particular, $\tilde{\Lambda}^{(+)}$ corresponds to the linear extension of the natural action of τ on G_∞ .

The canonical action of τ on \mathcal{X}_∞ makes it into a $\tilde{\Lambda}$ -module. Recall the element $x \in \mathcal{X}_\infty$ chosen at the beginning of this section such that $\pi(x) \neq 0$, where π is the map in (16), and $\tau(x) = \epsilon x$ for some $\epsilon \in \{\pm 1\}$. Now pick an element $y \in \ker(\pi)^{-\epsilon}$ and consider the surjection of $\tilde{\Lambda}$ -modules

$$\tilde{\Lambda}^{(\epsilon)} \oplus \tilde{\Lambda}^{(-\epsilon)} \longrightarrow \tilde{\Lambda}x \oplus \tilde{\Lambda}y \subset \mathcal{X}_\infty \oplus \mathcal{X}_\infty, \quad (\xi, \eta) \longmapsto (\xi x, \eta y).$$

Since \mathcal{E}_∞^\vee is torsion-free, $\ker(\pi) \cap \tilde{\Lambda}x = \{0\}$, hence $\tilde{\Lambda}x \cap \tilde{\Lambda}y = \{0\}$. Therefore the canonical map of $\tilde{\Lambda}$ -modules $\tilde{\Lambda}x \oplus \tilde{\Lambda}y \rightarrow \mathcal{X}_\infty$ given by the sum is injective. Composing the last two maps, we get a map of $\tilde{\Lambda}$ -modules

$$(19) \quad \vartheta : \tilde{\Lambda}^{(\epsilon)} \oplus \tilde{\Lambda}^{(-\epsilon)} \longrightarrow \mathcal{X}_\infty.$$

that sends (α, β) to $\alpha x + \beta y$.

Lemma 4.5. *For every $m \geq 1$ there is a canonical isomorphism*

$$\mathcal{X}_m/p^m \mathcal{X}_m \simeq H_f^1(K_m, A_{p^m})^\vee.$$

Proof. By definition, \mathcal{X}_m is the Pontryagin dual of $H_f^1(K_m, A)$. Applying Lemma 2.1 with $M = H_f^1(K_m, A)$ and $I = (p^m)$, we see that $\mathcal{X}_m/p^m \mathcal{X}_m$ is isomorphic to the Pontryagin dual of $H_f^1(K_m, A)_{p^m}$. But $H_f^1(K_m, A)_{p^m}$ is isomorphic to $H_f^1(K_m, A_{p^m})$ by Lemma 2.9, and the claim follows. \square

For every integer $m \geq 1$ consider the surjection

$$(20) \quad p_m : \mathcal{X}_\infty \twoheadrightarrow (\mathcal{X}_\infty)_{\Gamma_m} \simeq \mathcal{X}_m \twoheadrightarrow \mathcal{X}_m/p^m \mathcal{X}_m \simeq H_f^1(K_m, A_{p^m})^\vee,$$

where the first isomorphism comes from Corollary 2.7 and the second from Lemma 4.5. Let us define the following \tilde{R}_m -submodules of $H_f^1(K_m, A_{p^m})^\vee$:

$$\begin{aligned} Z_m &:= ((p_m \circ \vartheta)(\Lambda^{(\epsilon)} \oplus \{0\})) \cap ((p_m \circ \vartheta)(\{0\} \oplus \Lambda^{(-\epsilon)})), \\ W_m^{(\epsilon)} &:= ((p_m \circ \vartheta)(\Lambda^{(\epsilon)} \oplus \{0\}))/Z_m, \\ W_m^{(-\epsilon)} &:= ((p_m \circ \vartheta)(\{0\} \oplus \Lambda^{(-\epsilon)}))/Z_m. \end{aligned}$$

Now set

$$\Sigma_m := \left(H_f^1(K_m, A_{p^m})^\vee / Z_m \right)^\vee,$$

so that there is a canonical identification $\Sigma_m^\vee = H_f^1(K_m, A_{p^m})^\vee / Z_m$. Since the submodule Z_m is closed in $H_f^1(K_m, A_{p^m})^\vee$, Pontryagin duality applied to $H_f^1(K_m, A_{p^m})$ yields a natural injection $\Sigma_m \hookrightarrow H_f^1(K_m, A_{p^m})$ of \tilde{R}_m -modules, which we shall often view as an inclusion. Therefore we obtain a chain of maps of \tilde{R}_m -modules

$$(21) \quad \Lambda^{(\epsilon)} \oplus \Lambda^{(-\epsilon)} \twoheadrightarrow W_m^{(\epsilon)} \oplus W_m^{(-\epsilon)} \hookrightarrow \Sigma_m^\vee,$$

where the surjection is induced by $p_m \circ \vartheta$ and the injection is given by the sum. By construction, the composition in (21) factors through the surjection $\Lambda^{(\epsilon)} \oplus \Lambda^{(-\epsilon)} \twoheadrightarrow R_m^{(\epsilon)} \oplus R_m^{(-\epsilon)}$. Let

$$\vartheta_m : R_m^{(\epsilon)} \oplus R_m^{(-\epsilon)} \twoheadrightarrow W_m^{(\epsilon)} \oplus W_m^{(-\epsilon)} \hookrightarrow \Sigma_m^\vee$$

be the resulting map of \tilde{R}_m -modules; if \bar{x} and \bar{y} denote the images of x and y in Σ_m^\vee then $\vartheta_m((\alpha, \beta)) = \alpha\bar{x} + \beta\bar{y}$. Since there is a non-canonical isomorphism of \mathcal{O}_p -modules $R_m^{(\pm)} \simeq (\mathcal{O}_p/p^m \mathcal{O}_p)^{p^m}$, it turns out that $R_m^{(\pm)}$ is (non-canonically) isomorphic to $(R_m^{(\pm)})^\vee$. Fix isomorphisms

$$(22) \quad i_m^{(\pm)} : (R_m^{(\pm)})^\vee \xrightarrow{\simeq} R_m^{(\pm)}$$

of R_m -modules and set $i_m := i_m^{(\epsilon)} \oplus i_m^{(-\epsilon)}$. Composing the Pontryagin dual ϑ_m^\vee of ϑ_m with i_m , we get a map $i_m \circ \vartheta_m^\vee$ of \tilde{R}_m -modules that we still denote by

$$(23) \quad \vartheta_m^\vee : \Sigma_m \twoheadrightarrow R_m^{(\epsilon)} \oplus R_m^{(-\epsilon)}.$$

If $\bar{\Sigma}_m := \Sigma_m / \ker(\vartheta_m^\vee)$ then there is an injection $\bar{\vartheta}_m^\vee : \bar{\Sigma}_m \hookrightarrow R_m^{(\epsilon)} \oplus R_m^{(-\epsilon)}$ of \tilde{R}_m -modules. Define $\bar{\Sigma}_m^{(\epsilon)} := (\bar{\vartheta}_m^\vee)^{-1}(R_m^{(\epsilon)} \oplus \{0\})$ and $\bar{\Sigma}_m^{(-\epsilon)} := (\bar{\vartheta}_m^\vee)^{-1}(\{0\} \oplus R_m^{(-\epsilon)})$. It follows that there is a splitting

$$(24) \quad \bar{\Sigma}_m = \bar{\Sigma}_m^{(\epsilon)} \oplus \bar{\Sigma}_m^{(-\epsilon)}$$

of \tilde{R}_m -modules. Taking G_m -invariants, we obtain an injection

$$(\bar{\Sigma}_m^{(\pm)})^{G_m} \hookrightarrow (R_m^{(\pm)})^{G_m} \simeq \mathcal{O}_p/p^m \mathcal{O}_p$$

of $\mathcal{O}_p/p^m \mathcal{O}_p$ -modules, and the structure theorem for finitely generated \mathcal{O}_p -modules implies that $(\bar{\Sigma}_m^{(\pm)})^{G_m} \simeq \mathcal{O}_p/p^{m^{(\pm)}} \mathcal{O}_p$ for suitable integers $m^{(\pm)} \leq m$.

4.3. Compatibility of the maps. In order to ensure compatibility of the various maps appearing in the previous subsection as m varies, in the sequel it will be useful to make a convenient choice of the isomorphism $i_m^{(\epsilon)}$ introduced in (22).

Let

$$\pi_m : H_f^1(K_m, A_{p^m})^\vee \longrightarrow \mathcal{E}_m^\vee$$

be the dual of the inclusion $\mathcal{E}_m \subset H_f^1(K_m, A_{p^m})$; since $H_f^1(K_m, A_{p^m})$ is discrete, the map π_m is surjective. Since $y \in \ker(\pi)$, we have $\pi_m(Z_m) = 0$, so there is a surjection $\bar{\pi}_m : \Sigma_m^\vee \twoheadrightarrow \mathcal{E}_m^\vee$ showing that \mathcal{E}_m is actually a submodule of Σ_m . Since $(\bar{\pi}_m \circ \vartheta_m)(\{0\} \oplus R_m^{(-\epsilon)}) = \{0\}$, again because $y \in \ker(\pi)$, the dual of $\bar{\pi}_m \circ \vartheta_m$ factors through a map $\tilde{\psi}_m : \mathcal{E}_m \rightarrow (R_m^{(\epsilon)})^\vee$. Using (14), one easily checks the relation

$$u_m \beta_m = p \cdot \text{cores}_{K_{m+1}/K_m}(\beta_{m+1})$$

for some $u_m \in R_m^\times$ (recall that we denote by β_m the image of the Heegner cycle α_m in \mathcal{E}_m). Using this relation, one can prove the existence of isomorphisms $i_m^{(\epsilon)} : (R_m^{(\epsilon)})^\vee \xrightarrow{\sim} R_m^{(\epsilon)}$ such that if ψ_m is the composition

$$\psi_m : \mathcal{E}_m \xrightarrow{\tilde{\psi}_m} (R_m^{(\epsilon)})^\vee \xrightarrow{i_m^{(\epsilon)}} R_m^{(\epsilon)}$$

then the cyclic R_m -modules $\psi_m(\mathcal{E}_m)$ are generated by elements $\theta_m \in R_m$ satisfying $\theta_\infty := (\theta_m)_m \in \Lambda$. From now on, fix $i_m^{(\epsilon)}$ as before, so that $\theta_\infty \in \Lambda$. In the following, we will identify $R_m^{(\epsilon)}$ and its Pontryagin dual by means of this map without making it explicit. We will also implicitly identify $R_m^{(-\epsilon)}$ with its Pontryagin dual, but we will not need to specify a convenient isomorphism in this case.

4.4. Galois extensions. In this subsection we introduce several Galois extensions attached to the modules defined in §4.2. We start with the following general discussion.

For any $\mathcal{O}_p/p^m\mathcal{O}_p$ -submodule $S \subset H_f^1(K_m, A_{p^m})$, we define the extension M_S of $K_m(A_{p^m})$ cut out by S as follows. Set

$$\mathcal{G}_m := \text{Gal}(K_m(A_{p^m})/K_m).$$

With a slight abuse, we shall often view \mathcal{G}_m as a subgroup of $\text{GL}_2(\mathcal{O}_p/p^m\mathcal{O}_p)$, according to convenience. By [6, Lemma 6.2], the image of ρ_m in $\text{GL}_2(\mathcal{O}_p/p^m\mathcal{O}_p)$ contains a subgroup that is conjugate to $\text{GL}_2(\mathbb{Z}/p^m\mathbb{Z})$. Therefore

$$(25) \quad \mathcal{G}_m \text{ contains a subgroup that is conjugate to } \text{GL}_2(\mathbb{Z}/p^m\mathbb{Z}) \text{ in } \text{GL}_2(\mathcal{O}_p/p^m\mathcal{O}_p).$$

We need a variant of [6, Proposition 6.3, (2)], which we prove in the next

Lemma 4.6. $H^1(\text{Gal}(K_m(A_{p^m})/K_m), A_{p^m}) = 0$.

Proof. Restriction of automorphisms injects $\text{Gal}(K_m(A_{p^m})/K_m)$ into $\text{Gal}(K(A_{p^m})/K)$; this induces a natural identification

$$(26) \quad \text{Gal}(K_m(A_{p^m})/K_m) = \text{Gal}(K(A_{p^m})/K(A_{p^m}) \cap K_m).$$

From the inflation-restriction exact sequence we can extract the exact sequence

$$(27) \quad \begin{aligned} H^1(\text{Gal}(K(A_{p^m})/K), A_{p^m}) &\longrightarrow H^1(\text{Gal}(K(A_{p^m})/K(A_{p^m}) \cap K_m), A_{p^m})^{\text{Gal}(K(A_{p^m}) \cap K_m/K)} \\ &\longrightarrow H^2(\text{Gal}(K(A_{p^m}) \cap K_m/K), A_{p^m}(K(A_{p^m}) \cap K_m)), \end{aligned}$$

where the first map is restriction and the second is transgression. By [6, Proposition 6.3, (2)], $H^1(\text{Gal}(K(A_{p^m})/K), A_{p^m}) = 0$. On the other hand, the extension K_m/\mathbb{Q} is solvable, hence $A_{p^m}(K(A_{p^m}) \cap K_m) = 0$ by [18, Lemma 3.10, (2)]. As a consequence, the middle term in (27)

is trivial. But $H^1(\text{Gal}(K(A_{p^m})/K(A_{p^m}) \cap K_m), A_{p^m})$ and $\text{Gal}(K(A_{p^m}) \cap K_m/K)$ are both p -groups, hence [29, Lemma 3] implies that

$$H^1(\text{Gal}(K(A_{p^m})/K(A_{p^m}) \cap K_m), A_{p^m}) = 0.$$

Now the lemma follows from (26). \square

Thanks to the vanishing result of Lemma 4.6, restriction gives an injection

$$H_f^1(K_m, A_{p^m}) \hookrightarrow H_f^1(K_m(A_{p^m}), A_{p^m})^{\mathcal{G}_m}.$$

Define $G_{K_m(A_{p^m})}^{\text{ab}} := \text{Gal}(K_m(A_{p^m})^{\text{ab}}/K_m(A_{p^m}))$ where $K_m(A_{p^m})^{\text{ab}}$ is the maximal abelian extension of $K_m(A_{p^m})$. It follows that there is an identification

$$H^1(K_m(A_{p^m}), A_{p^m})^{\mathcal{G}_m} = \text{Hom}_{\mathcal{G}_m}(G_{K_m(A_{p^m})}^{\text{ab}}, A_{p^m})$$

of $\mathcal{O}_p/p^m\mathcal{O}_p$ -modules, where $\text{Hom}_{\mathcal{G}_m}(\bullet, \star)$ stands for the group of \mathcal{G}_m -homomorphisms. Thus we obtain an injection of $\mathcal{O}_p/p^m\mathcal{O}_p$ -modules

$$(28) \quad S \hookrightarrow \text{Hom}_{\mathcal{G}_m}(G_{K_m(A_{p^m})}^{\text{ab}}, A_{p^m}), \quad s \mapsto \varphi_s,$$

and for every $s \in S$ we let M_s be the subfield of $K_m(A_{p^m})^{\text{ab}}$ fixed by $\ker(\varphi_s)$. In other words, M_s is the smallest abelian extension of $K_m(A_{p^m})$ such that the restriction of φ_s to $\text{Gal}(K_m(A_{p^m})^{\text{ab}}/M_s)$ is trivial. The maps φ_s induce injections

$$\varphi_s : \text{Gal}(M_s/K_m(A_{p^m})) \hookrightarrow A_{p^m}$$

of \mathcal{G}_m -modules. Let $M_S \subset K_m(A_{p^m})^{\text{ab}}$ denote the composite of all the fields M_s for $s \in S$.

Now we prove that the map

$$(29) \quad \text{Gal}(M_S/K_m(A_{p^m})) \longrightarrow \text{Hom}(S, A_{p^m}), \quad g \mapsto (s \mapsto \varphi_s(g|_{M_s}))$$

is a \mathcal{G}_m -isomorphism; here $\text{Hom}(\bullet, \star)$ is a shorthand for $\text{Hom}_{\mathcal{O}_p/p^m\mathcal{O}_p}(\bullet, \star)$. Furthermore, we show that the map (28) induces an isomorphism

$$(30) \quad S \xrightarrow{\cong} \text{Hom}_{\mathcal{G}_m}(\text{Gal}(M_S/K_m(A_{p^m})), A_{p^m})$$

of $\mathcal{O}_p/p^m\mathcal{O}_p$ -modules. First of all, note that (28) gives an injection

$$(31) \quad S \hookrightarrow \text{Hom}_{\mathcal{G}_m}(\text{Gal}(M_S/K_m(A_{p^m})), A_{p^m}), \quad s \mapsto (g \mapsto \varphi_s(g|_{M_s}))$$

of $\mathcal{O}_p/p^m\mathcal{O}_p$ -modules.

Since S is a finite \mathcal{O}_p -module of exponent p^m , there is an isomorphism

$$\xi : S \xrightarrow{\cong} \prod_{i=1}^t \mathcal{O}_p/p^{m_i}\mathcal{O}_p$$

of $\mathcal{O}_p/p^m\mathcal{O}_p$ -modules for suitable integers $m_i \leq m$. For every $i \in \{1, \dots, t\}$ set $S_i := \xi^{-1}(\mathcal{O}_p/p^{m_i}\mathcal{O}_p)$ and choose a generator s_i of the free $\mathcal{O}_p/p^{m_i}\mathcal{O}_p$ -module S_i . The map φ_{s_i} is killed by p^{m_i} , so it induces an injection

$$\varphi_{s_i} : \text{Gal}(M_{s_i}/K_m(A_{p^m})) \hookrightarrow A_{p^{m_i}}.$$

On the other hand, for every i there is a \mathcal{G}_m -equivariant injection

$$\text{Gal}(M_{s_i}/K_m(A_{p^m})) \hookrightarrow \text{Hom}(S_i, A_{p^m}), \quad g \mapsto (s_i \mapsto \varphi_{s_i}(g)).$$

But $\text{Hom}(S_{s_i}, A_{p^m}) \simeq A_{p^{m_i}}$, hence there is a \mathcal{G}_m -equivariant injection

$$(32) \quad \text{Gal}(M_S/K_m(A_{p^m})) \hookrightarrow \text{Hom}(S, A_{p^m}) \simeq \prod_{i=1}^t A_{p^{m_i}}.$$

Using the fact that the Galois representation ρ_1 on A_p is irreducible ([18, Lemma 3.8]), it can be shown that the image of (32) is of the form $\prod_{i=1}^{t'} A_{p^{m'_i}}$ for integers $t' \leq t$ and $m'_i \leq m_i$, hence there is a \mathcal{G}_m -equivariant isomorphism $\text{Gal}(M_S/K_m(A_{p^m})) \simeq \prod_{i=1}^{t'} A_{p^{m'_i}}$.

Proposition 4.7. $\text{Hom}_{\mathcal{G}_m}(A_{p^{m'_i}}, A_{p^m}) \simeq \mathcal{O}_p/p^{m'_i}\mathcal{O}_p$.

Proof. First of all, there is a canonical identification

$$\text{Hom}_{\mathcal{G}_m}(A_{p^{m'_i}}, A_{p^m}) = \text{Hom}_{\mathcal{G}_m}(A_{p^{m'_i}}, A_{p^{m'_i}}).$$

To ease the notation, set $n := m'_i$. Since A_{p^n} is free of rank 2 over $\mathcal{O}_p/p^n\mathcal{O}_p$, there is an isomorphism $\text{Hom}_{\mathcal{O}_p/p^n\mathcal{O}_p}(A_{p^n}, A_{p^n}) \simeq \text{M}_2(\mathcal{O}_p/p^n\mathcal{O}_p)$; to obtain it, we fix a basis of A_{p^n} over $\mathcal{O}_p/p^n\mathcal{O}_p$. Observe that the group $\mathcal{G}_m \subset \text{GL}_2(\mathcal{O}_p/p^m\mathcal{O}_p)$ acts on A_{p^n} via the reduction map $\text{GL}_2(\mathcal{O}_p/p^m\mathcal{O}_p) \rightarrow \text{GL}_2(\mathcal{O}_p/p^n\mathcal{O}_p)$. Let $\mathcal{G}_m^{(n)}$ be the image of \mathcal{G}_m in $\text{GL}_2(\mathcal{O}_p/p^n\mathcal{O}_p)$, so that

$$(33) \quad \text{Hom}_{\mathcal{G}_m}(A_{p^n}, A_{p^m}) = \text{Hom}_{\mathcal{G}_m^{(n)}}(A_{p^n}, A_{p^n}).$$

By (25), \mathcal{G}_m contains a subgroup that is conjugate to $\text{GL}_2(\mathbb{Z}/p^m\mathbb{Z})$, hence $\mathcal{G}_m^{(n)}$ contains a subgroup that is conjugate to $\text{GL}_2(\mathbb{Z}/p^n\mathbb{Z})$. Write $g \text{GL}_2(\mathbb{Z}/p^n\mathbb{Z}) g^{-1}$ with $g \in \text{GL}_2(\mathcal{O}_p/p^n\mathcal{O}_p)$ for this subgroup. In particular, we want to find the matrices $A \in \text{M}_2(\mathcal{O}_p/p^n\mathcal{O}_p)$ such that

$$(gXg^{-1})A(gX^{-1}g^{-1}) = A$$

for all $X \in \text{GL}_2(\mathbb{Z}/p^n\mathbb{Z})$. Upon setting $B := g^{-1}Ag$, this is equivalent to describing the matrices $B \in \text{M}_2(\mathcal{O}_p/p^n\mathcal{O}_p)$ such that

$$XBX^{-1} = B$$

for all $X \in \text{GL}_2(\mathbb{Z}/p^n\mathbb{Z})$. Clearly, these are exactly the diagonal matrices, so we conclude that $\text{Hom}_{\mathcal{G}_m^{(n)}}(A_{p^n}, A_{p^m}) \simeq \mathcal{O}_p/p^n\mathcal{O}_p$. The proposition follows from (33). \square

It follows from Proposition 4.7 that there is an isomorphism of $\mathcal{O}_p/p^m\mathcal{O}_p$ -modules

$$\text{Hom}_{\mathcal{G}_m}(\text{Gal}(M_S/K_m(A_{p^m})), A_{p^m}) \simeq \prod_{i=1}^{t'} \mathcal{O}_p/p^{m'_i}\mathcal{O}_p.$$

By (31), S injects into $\text{Hom}_{\mathcal{G}_m}(\text{Gal}(M_S/K_m(A_{p^m})), A_{p^m})$, and then comparing cardinalities gives $t = t'$ and $m_i = m'_i$ for all i , thus proving (29) and (30) simultaneously. From this we also deduce the following

Proposition 4.8. *Given subgroups $S' \subset S \subset H_f^1(K_m, A_{p^m})$, there is a canonical isomorphism of groups*

$$\text{Gal}(M_S/M_{S'}) \simeq \text{Hom}(S/S', A_{p^m}).$$

Conversely, for every subgroup \bar{S} of S/S' there is a subextension $M_{\bar{S}}/M_{S'}$ of $M_S/M_{S'}$ such that

$$\text{Gal}(M_{\bar{S}}/M_{S'}) \simeq \text{Hom}(\bar{S}, A_{p^m}).$$

Proof. The short exact sequence $0 \rightarrow S' \rightarrow S \rightarrow S/S' \rightarrow 0$ gives a short exact sequence

$$0 \longrightarrow \text{Hom}(S/S', A_{p^m}) \longrightarrow \text{Hom}(S, A_{p^m}) \longrightarrow \text{Hom}(S', A_{p^m}) \longrightarrow 0,$$

where the surjectivity on the right comes from isomorphism (29) and the surjectivity of the canonical map from $\text{Gal}(M_S/K_m(A_{p^m}))$ to $\text{Gal}(M_{S'}/K_m(A_{p^m}))$. The first claim follows from the fact that, again by (29), the kernel of the third arrow is naturally isomorphic to $\text{Gal}(M_S/M_{S'})$.

As for the second assertion, suppose that \bar{S} is a subgroup of S/S' . Then, as before, one can define a subextension $M_{\bar{S}}/M_{S'}$ of $M_S/M_{S'}$; namely, an element $s \in \bar{S}$ cuts out an extension

$M_S/M_{S'}$ and we let $M_{\bar{S}}$ be the composite of all the fields M_S . Replacing S with \bar{S} , $K_m(A_{p^m})$ with $M_{S'}$ and \mathcal{G}_m with $\text{Gal}(M_{S'}/K_m)$ in the discussion above, one checks that there are isomorphisms of groups

$$\bar{S} \simeq \text{Hom}_{\text{Gal}(M_{S'}/K_m)}(\text{Gal}(M_{\bar{S}}/M_{S'}), A_{p^m}), \quad \text{Gal}(M_{\bar{S}}/M_{S'}) \simeq \text{Hom}(\bar{S}, A_{p^m}),$$

as required. \square

With notation as in Proposition 4.8, we say that $M_S/M_{S'}$ is the extension associated with the quotient S/S' . Now we apply these constructions to the setting of §4.2.

Lemma 4.9. *There is an injection*

$$H_f^1(K_m, A_{p^m}) \hookrightarrow H_f^1(K_{m+1}, A_{p^{m+1}})$$

induced by the restriction map and the inclusion $A_{p^m} \hookrightarrow A_{p^{m+1}}$.

Proof. The extension K_{m+1}/\mathbb{Q} is solvable, hence $A_{p^m}(K_{m+1}) = 0$ by [18, Lemma 3.10, (2)]. It follows that restriction induces an injection

$$(34) \quad H^1(K_m, A_{p^m}) \hookrightarrow H^1(K_{m+1}, A_{p^m}).$$

Now consider the short exact sequence of $G_{K_{m+1}}$ -modules

$$(35) \quad 0 \longrightarrow A_{p^m} \longrightarrow A_{p^{m+1}} \longrightarrow A_{p^{m+1}}/A_{p^m} \longrightarrow 0.$$

Multiplication by p^m yields a Galois-equivariant isomorphism $A_{p^{m+1}}/A_{p^m} \simeq A_p$, therefore $(A_{p^{m+1}}/A_{p^m})(K_{m+1}) = 0$ again by [18, Lemma 3.10, (2)]. In light of this vanishing, passing to the long exact $G_{K_{m+1}}$ -cohomology sequence associated with (35) gives an injection

$$(36) \quad H^1(K_{m+1}, A_{p^m}) \hookrightarrow H^1(K_{m+1}, A_{p^{m+1}}).$$

Combining (34) and (36), we get an injection

$$H_f^1(K_m, A_{p^m}) \hookrightarrow H_f^1(K_{m+1}, A_{p^{m+1}})$$

that restricts to the desired injection between Selmer groups. \square

Thanks to Lemma 4.9, in the sequel we shall often view $H_f^1(K_m, A_{p^m})$ as a subgroup of $H_f^1(K_{m+1}, A_{p^{m+1}})$. Let M_m denote the field cut out by the whole group $S = H_f^1(K_m, A_{p^m})$; then $M_m \subset M_{m+1}$.

Proposition 4.10. *There is a canonical surjection*

$$\text{Gal}(M_{m+1}/K_{m+1}(A_{p^{m+1}})) \twoheadrightarrow \text{Gal}(M_m/K_m(A_{p^m})).$$

Proof. View $H_f^1(K_m, A_{p^m})$ as a subgroup of $H_f^1(K_{m+1}, A_{p^{m+1}})$ and consider the extension H of $K_{m+1}(A_{p^{m+1}})$ cut out by it inside M_{m+1} . Then

$$\text{Gal}(H/K_{m+1}(A_{p^{m+1}})) \simeq \text{Hom}(H_f^1(K_m, A_{p^m}), A_{p^{m+1}}) = \text{Hom}(H_f^1(K_m, A_{p^m}), A_{p^m}),$$

where the equality is a consequence of $H_f^1(K_m, A_{p^m})$ being p^m -torsion. On the other hand, by definition of M_m there is an isomorphism $\text{Gal}(M_m/K_m(A_{p^m})) \simeq \text{Hom}(H_f^1(K_m, A_{p^m}), A_{p^m})$. Comparing, we find an isomorphism

$$\text{Gal}(H/K_{m+1}(A_{p^{m+1}})) \simeq \text{Gal}(M_m/K_m(A_{p^m}))$$

that can be combined with the canonical surjection

$$\text{Gal}(M_{m+1}/K_{m+1}(A_{p^{m+1}})) \twoheadrightarrow \text{Gal}(H/K_{m+1}(A_{p^{m+1}}))$$

to complete the proof. \square

Graphically, there is a diagram of field extensions

$$\begin{array}{ccc}
 & M_{m+1} & \\
 & \downarrow & \searrow \\
 & H & \\
 & \downarrow & \\
 K_{m+1}(A_{p^{m+1}}) & & M_m \\
 & \searrow & \downarrow \\
 & & K_m(A_{p^m})
 \end{array}$$

with $\text{Gal}(H/K_{m+1}(A_{p^{m+1}})) \simeq \text{Gal}(M_m/K_m(A_{p^m}))$. Let us define

$$M_\infty := \varprojlim_m M_m, \quad K_\infty(A) := \varprojlim_m K_m(A_{p^m}),$$

so that

$$\text{Gal}(M_\infty/K_\infty(A)) := \varprojlim_m \text{Gal}(M_m/K_m(A_{p^m})),$$

the inverse limit being taken with respect to the maps of Proposition 4.10. By (29), for every $m \geq 0$ there is an isomorphism $\text{Gal}(M_m/K_m(A_{p^m})) \simeq \text{Hom}(H_f^1(K_m, A_{p^m}), A_{p^m})$ of $\mathcal{O}_p/p^m \mathcal{O}_p[\mathcal{G}_m]$ -modules, hence there is an isomorphism of $\mathcal{O}_p[[\mathcal{G}_\infty]]$ -modules

$$\text{Gal}(M_\infty/K_\infty(A)) \simeq \text{Hom}(H_f^1(K_\infty, A), A)$$

where $\mathcal{O}_p[[\mathcal{G}_\infty]] := \varprojlim_m \mathcal{O}_p[\mathcal{G}_m]$ is defined with respect to the natural maps $\mathcal{G}_{m+1} \rightarrow \mathcal{G}_m$.

Now recall the map ϑ_m^\vee of (23) and let $L_m \subset M_m$ be the extension of $K_m(A_{p^m})$ cut out by $\ker(\vartheta_m^\vee)$. Then there are canonical \mathcal{G}_m -isomorphisms

$$\text{Gal}(L_m/K_m(A_{p^m})) \simeq \text{Hom}(\ker(\vartheta_m^\vee), A_{p^m})$$

and

$$(37) \quad \text{Gal}(M_m/L_m) \simeq \text{Hom}(\overline{\Sigma}_m, A_{p^m}) \simeq \text{Hom}(\overline{\Sigma}_m^{(\epsilon)}, A_{p^m}) \oplus \text{Hom}(\overline{\Sigma}_m^{(-\epsilon)}, A_{p^m});$$

here (37) is a consequence of Proposition 4.8. Let $L_m^{(\pm)}/L_m$ be the subextension of M_m/L_m corresponding to $\overline{\Sigma}_m^{(\pm)}$ (cf. Proposition 4.8); then $L_m^{(\epsilon)} \cap L_m^{(-\epsilon)} = L_m$ and $M_m = L_m^{(\epsilon)} \cdot L_m^{(-\epsilon)}$. Finally, let $\tilde{L}_m^{(\pm)}$ be the extension of $L_m^{(\pm)}$ corresponding to $(\overline{\Sigma}_m^{(\pm)})^{G_m}$. We have $\tilde{L}_m^{(+)} \cap \tilde{L}_m^{(-)} = L_m$ and

$$\text{Gal}(\tilde{L}_m^{(\pm)}/L_m) \simeq \text{Hom}\left(\left(\overline{\Sigma}_m^{(\pm)}\right)^{G_m}, A_{p^m}\right).$$

Moreover, if $\tilde{L}_m := \tilde{L}_m^{(+)} \cdot \tilde{L}_m^{(-)}$ then

$$\begin{aligned}
 \text{Gal}(\tilde{L}_m/L_m) &\simeq \text{Hom}\left(\left(\overline{\Sigma}_m^{(+)}\right)^{G_m}, A_{p^m}\right) \oplus \text{Hom}\left(\left(\overline{\Sigma}_m^{(-)}\right)^{G_m}, A_{p^m}\right) \\
 &\simeq \text{Hom}\left(\left(\overline{\Sigma}_m\right)^{G_m}, A_{p^m}\right),
 \end{aligned}$$

where the second isomorphism follows by taking G_m -invariants in (24). Since, by Lemma 4.9, $H_f^1(K_m, A_{p^m})$ injects via restriction into $H_f^1(K_{m+1}, A_{p^{m+1}})$, restriction induces an injection $(\overline{\Sigma}_m)^{G_m} \hookrightarrow (\overline{\Sigma}_{m+1})^{G_{m+1}}$. It follows that for every $m \geq 0$ there is a canonical projection

$$(38) \quad \text{Gal}(\tilde{L}_{m+1}/L_{m+1}) \twoheadrightarrow \text{Gal}(\tilde{L}_m/L_m).$$

To introduce the last field extensions that we need, we dualize the exact sequence

$$R_m^{(\epsilon)} \oplus R_m^{(-\epsilon)} \xrightarrow{\vartheta_m} \Sigma_m^\vee \longrightarrow \Sigma_m^\vee/\text{im}(\vartheta_m) \longrightarrow 0$$

and get an isomorphism $\ker(\vartheta_m^\vee) \simeq (\Sigma_m^\vee / \text{im}(\vartheta_m))^\vee$. Furthermore, dualizing the short exact sequence

$$0 \longrightarrow \text{im}(\vartheta_m) \longrightarrow \Sigma_m^\vee \longrightarrow \ker(\vartheta_m^\vee)^\vee \longrightarrow 0$$

gives a short exact sequence

$$(39) \quad 0 \longrightarrow \ker(\vartheta_m^\vee) \longrightarrow \Sigma_m \xrightarrow{\vartheta_m^\vee} \text{im}(\vartheta_m)^\vee \longrightarrow 0.$$

Finally, with maps ϑ and p_m defined as in (19) and (20), let U_m be the \tilde{R}_m -submodule of $H_f^1(K_m, A_{p^m})$ such that there is an identification

$$(40) \quad \text{im}(p_m \circ \vartheta) = \left(H_f^1(K_m, A_{p^m}) / U_m \right)^\vee.$$

Namely, set for simplicity $\mathcal{I}_m := \text{im}(p_m \circ \vartheta)$ and consider the short exact sequence

$$0 \longrightarrow \mathcal{I}_m \longrightarrow H_f^1(K_m, A_{p^m})^\vee \longrightarrow H_f^1(K_m, A_{p^m})^\vee / \mathcal{I}_m \longrightarrow 0.$$

Since \mathcal{I}_m is compact, hence closed in $H_f^1(K_m, A_{p^m})^\vee$, dualizing the sequence above gives a short exact sequence

$$(41) \quad 0 \longrightarrow \left(H_f^1(K_m, A_{p^m})^\vee / \mathcal{I}_m \right)^\vee \longrightarrow H_f^1(K_m, A_{p^m}) \longrightarrow \mathcal{I}_m^\vee \longrightarrow 0.$$

Define $U_m := \left(H_f^1(K_m, A_{p^m})^\vee / \mathcal{I}_m \right)^\vee$ and regard U_m as an \tilde{R}_m -submodule of $H_f^1(K_m, A_{p^m})$ via (41). Then there is a natural identification

$$(42) \quad H_f^1(K_m, A_{p^m}) / U_m = \mathcal{I}_m^\vee,$$

and dualizing (42) gives (40).

Now write \tilde{M}_m for the field cut out by U_m . As $p_m \circ \theta$ factors through $R_m^{(\epsilon)} \oplus R_m^{(-\epsilon)}$, there is a commutative diagram

$$\begin{array}{ccc} \Lambda^{(\epsilon)} \oplus \Lambda^{(-\epsilon)} & \xrightarrow{\vartheta} & \mathcal{X}_\infty \xrightarrow{p_m} H_f^1(K_m, A_{p^m})^\vee \\ \downarrow & & \downarrow \\ R_m^{(\epsilon)} \oplus R_m^{(-\epsilon)} & \xrightarrow{\vartheta_m} & \Sigma_m^\vee \end{array}$$

which induces a surjection $\mathcal{I}_m \twoheadrightarrow \text{im}(\vartheta_m)$ and then, by duality, an injection $\text{im}(\vartheta_m)^\vee \hookrightarrow \mathcal{I}_m^\vee$. From this we obtain a commutative diagram with exact rows

$$(43) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \ker(\vartheta_m^\vee) & \longrightarrow & \Sigma_m & \longrightarrow & \text{im}(\vartheta_m)^\vee \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & U_m & \longrightarrow & H_f^1(K_m, A_{p^m}) & \longrightarrow & \mathcal{I}_m^\vee \longrightarrow 0 \end{array}$$

whose upper row is (39). Denote by L_m^* the field corresponding to Σ_m , so that $\tilde{L}_m \subset L_m^*$ by Proposition 4.8. Observe that \tilde{M}_m and \tilde{L}_m are linearly disjoint over L_m . To check this, note that $\tilde{M}_m \cap \tilde{L}_m \subset \tilde{M}_m \cap L_m^*$ and that the second intersection corresponds to the subgroup $U_m \cap \Sigma_m$ inside $H_f^1(K_m, A_{p^m})$. But diagram (43) shows that $\ker(\vartheta_m^\vee) = U_m \cap \Sigma_m$, hence $\tilde{M}_m \cap L_m^* = L_m$; we conclude that $\tilde{M}_m \cap \tilde{L}_m = L_m$. It follows that

$$\text{Gal}(\tilde{M}_m \cdot \tilde{L}_m / \tilde{M}_m) \simeq \text{Gal}(\tilde{L}_m / \tilde{M}_m \cap \tilde{L}_m) \simeq \text{Gal}(\tilde{L}_m / L_m),$$

and then the inclusion $\tilde{M}_m \cdot \tilde{L}_m \subset M_m$ induces a surjection

$$(44) \quad \text{Gal}(M_m / \tilde{M}_m) \twoheadrightarrow \text{Gal}(\tilde{L}_m / L_m).$$

It follows that for every m there is a commutative square of surjective maps

$$(45) \quad \begin{array}{ccc} \mathrm{Gal}(M_{m+1}/\tilde{M}_{m+1}) & \twoheadrightarrow & \mathrm{Gal}(\tilde{L}_{m+1}/L_{m+1}) \\ \downarrow & & \downarrow \\ \mathrm{Gal}(M_m/\tilde{M}_m) & \twoheadrightarrow & \mathrm{Gal}(\tilde{L}_m/L_m) \end{array}$$

where the horizontal arrows are given by (44) and the right vertical arrow is given by (38). The only things that remain to be checked are the surjectivity of the left vertical map and the commutativity of (45). First of all, we want to show that $\tilde{M}_{m+1} \cap M_m = \tilde{M}_m$; this intersection corresponds to the subgroup $U_{m+1} \cap H_f^1(K_m, A_{p^m})$ inside $H_f^1(K_{m+1}, A_{p^{m+1}})$, with U_{m+1} defined as in (42). Set $\mathcal{I}_{m+1} := \mathrm{im}(p_{m+1} \circ \vartheta)$; the injection between Selmer groups of Lemma 4.9 gives a surjection $\mathcal{I}_{m+1} \twoheadrightarrow \mathcal{I}_m$ and then, by duality, an injection $\mathcal{I}_m^\vee \hookrightarrow \mathcal{I}_{m+1}^\vee$. Therefore there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & U_m & \longrightarrow & H_f^1(K_m, A_{p^m}) & \longrightarrow & \mathcal{I}_m^\vee \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & U_{m+1} & \longrightarrow & H_f^1(K_{m+1}, A_{p^{m+1}}) & \longrightarrow & \mathcal{I}_{m+1}^\vee \longrightarrow 0 \end{array}$$

where the injectivity of the map $U_m \rightarrow U_{m+1}$ is forced by the injectivity of the map between Selmer groups. We conclude that the subgroup $U_{m+1} \cap H_f^1(K_m, A_{p^m})$, which corresponds to \tilde{M}_m , is equal to the image of U_m inside $H_f^1(K_{m+1}, A_{p^{m+1}})$, and then $\tilde{M}_{m+1} \cap M_m = \tilde{M}_m$, as claimed above. It follows that

$$\mathrm{Gal}(\tilde{M}_{m+1} \cdot M_m / \tilde{M}_{m+1}) \simeq \mathrm{Gal}(M_m / \tilde{M}_{m+1} \cap M_m) \simeq \mathrm{Gal}(M_m / \tilde{M}_m),$$

and then the inclusion $\tilde{M}_{m+1} \cdot M_m \subset M_{m+1}$ yields the desired surjection

$$\mathrm{Gal}(M_{m+1} / \tilde{M}_{m+1}) \twoheadrightarrow \mathrm{Gal}(M_m / \tilde{M}_m).$$

This explains the existence and the surjectivity of the maps in diagram (45). Finally, one checks that the commutativity of (45) is an immediate consequence of the constructions.

4.5. Families of Kolyvagin primes. The purpose of this subsection is to show that one can manufacture a compatible sequence $(\ell_m)_{m \geq 1}$ of Kolyvagin primes, each ℓ_m being coprime with p^m ; here the word ‘‘compatibility’’ refers to the canonical maps of Galois theory. More precisely, our goal is to prove the following

Proposition 4.11. *There is a sequence $\ell_\infty = (\ell_m)_{m \geq 1}$ of Kolyvagin primes satisfying the following conditions:*

(1) $\mathrm{Frob}_{\ell_m} = [\tau g_m]$ in $\mathrm{Gal}(M_m/\mathbb{Q})$ with $g_m \in \mathrm{Gal}(M_m/K_m(A_{p^m}))$ such that

$$(g_m)_m \in \mathrm{Gal}(M_\infty/K_\infty(A));$$

(2) restriction induces an injective group homomorphism

$$\overline{\mathrm{res}}_{\ell_m} : \overline{\Sigma}_m \hookrightarrow H_f^1(K_{m, \ell_m}, A_{p^m});$$

(3) $(\ell_m + 1 \pm a_{\ell_m})/p^m$ are invertible in $\mathcal{O}_{\mathfrak{p}}/p^m \mathcal{O}_{\mathfrak{p}}$.

Proof. For each sign \pm choose $h_m^{(\pm)} \in \mathrm{Gal}(\tilde{L}_m^{(\pm)}/L_m)$ such that the period of $(h_m^{(\pm)})^\tau h_m^{(\pm)}$ is $p^{m(\pm)}$. To justify the existence of an element with this property, observe that if $h_m^{(\pm)}$ corresponds to the homomorphism $\phi : (\overline{\Sigma}_m^{(\pm)})^{G_m} \rightarrow A_{p^m}$ then $(h_m^{(\pm)})^\tau h_m^{(\pm)}$ corresponds to $x \mapsto \pm \tau \phi(x) + \phi(x)$. In light of this, to show the existence of such an $h_m^{(\pm)}$ it suffices to choose a ϕ that takes a generator of $(\overline{\Sigma}_m^{(\pm)})^{G_m}$ to an element of order $p^{m(\pm)}$ in $A_{p^m}^{(\pm)}$. Define

$h_m := (h_m^{(+)}, h_m^{(-)}) \in \text{Gal}(\tilde{L}_m/L_m)$ and choose the sequence $(h_m)_{m \geq 1}$ so that the image of h_{m+1} via surjection (38) is h_m . Using diagram (45), select also a compatible sequence of elements $g_m \in \text{Gal}(M_m/K_m(A_{p^m}))$ such that the image of g_m in $\text{Gal}(\tilde{L}_m/L_m)$ is h_m . For every integer $m \geq 1$ choose a prime number ℓ_m such that

$$(46) \quad \text{Frob}_{\ell_m} = [\tau g_m] \quad \text{in } \text{Gal}(M_m/\mathbb{Q}).$$

Clearly, ℓ_m is a Kolyvagin prime and the required compatibility conditions are fulfilled by construction, so (1) is satisfied. To check (2), we must show that the restriction is injective. For this, fix a prime \mathfrak{l}_m of M_m above ℓ_m satisfying $\text{Frob}_{\mathfrak{l}_m/\ell_m} = \tau g_m$. Then the restriction of $\text{Frob}_{\mathfrak{l}_m/\ell_m}$ to $\text{Gal}(\tilde{L}_m/L_m)$ corresponds to an injective homomorphism

$$\phi_{\mathfrak{l}_m/\ell_m} : (\overline{\Sigma}_m)^{G_m} \hookrightarrow A_{p^m}$$

consisting in the evaluation at $\text{Frob}_{\mathfrak{l}_m/\ell_m}$; namely, one has

$$\phi_{\mathfrak{l}_m/\ell_m}(s) = s(\text{Frob}_{\mathfrak{l}_m/\ell_m})$$

for all $s \in (\overline{\Sigma}_m)^{G_m}$. The choice of \mathfrak{l}_m determines a prime $\tilde{\lambda}_m$ of K_m above ℓ_m , and the completion of K_m at $\tilde{\lambda}_m$ is canonically isomorphic to the completion K_{λ_m} of K at the unique prime λ_m of K above ℓ_m . It follows that the canonical restriction map

$$(47) \quad (\overline{\Sigma}_m)^{G_m} \hookrightarrow H_f^1(K_{\lambda_m}, A_{p^m})$$

is injective, since the same is true of the composition of this map with the isomorphism

$$H_f^1(K_{\lambda_m}, A_{p^m}) \simeq A_{p^m}$$

that is given precisely by evaluation at Frobenius (recall (18)). Suppose now that $s \in \overline{\Sigma}_m$ is non-zero and $\overline{\text{res}}_{\ell_m}(s) = 0$. In particular, the submodule $(R_m s)^{G_m}$ of $(\overline{\Sigma}_m)^{G_m}$ is sent to zero, via (47), in the direct summand $H_f^1(K_{\lambda_m}, A_{p^m})$ of $H_f^1(K_{m, \ell_m}, A_{p^m})$ corresponding to $\tilde{\lambda}_m$. Up to multiplying s by a suitable power of p , we may assume that s is p -torsion. Now $R_m s$ is a non-trivial $\mathcal{O}_p/p\mathcal{O}_p$ -vector space on which the p -group G_m acts. By [29, Proposition 26], the submodule $(R_m s)^{G_m}$ is non-trivial, and this contradicts the injectivity of (47). Summing up, we have proved that all choices of a sequence $\ell_\infty = (\ell_m)_{m \geq 1}$ satisfying (46) enjoy properties (1) and (2) in the statement of the proposition.

The finer choice of a sequence ℓ_∞ satisfying (3) as well can be made by arguing as in the proof of [24, Proposition 12.2, (3)]; see the proof of [18, Proposition 3.26] for details. \square

4.6. Local duality. The aim of this subsection is to bound the Λ -rank of $\Lambda x \oplus \Lambda y$ by a Λ -module $V(\ell_\infty)$ that surjects onto $\Lambda x \oplus \Lambda y$; the module $V(\ell_\infty)$ depends on the choice of a compatible family $\ell_\infty = (\ell_m)_{m \geq 1}$ of Kolyvagin primes as in §4.5.

By [7, Proposition 3.8], if E is a number field and v is a prime of E then there is a perfect pairing

$$\langle \cdot, \cdot \rangle_{E,v} : H^1(E_v, T/p^m T) \times H^1(E_v, A_{p^m}) \longrightarrow \mathbb{Q}_p/\mathbb{Z}_p$$

under which $H_f^1(E_v, T/p^m T)$ and $H_f^1(E_v, A_{p^m})$ are exact annihilators of each other. From this one deduces the existence (see, e.g., [18, eq. (34)]) of a τ -antiequivariant isomorphism

$$\delta_{E,v} : H_s^1(E_v, T/p^m T) \xrightarrow{\simeq} H_f^1(E_v, A_{p^m})^\vee.$$

Now let ℓ be a Kolyvagin prime and write $\delta_{m,\lambda}$ as a shorthand for $\delta_{K_m,\lambda}$. Taking the direct sum of the maps $\delta_{m,\lambda}$ over all the primes $\lambda | \ell$, we get a τ -antiequivariant isomorphism

$$(48) \quad \delta_{m,\ell} : H_s^1(K_{m,\ell}, T/p^m T) \xrightarrow{\simeq} H_f^1(K_{m,\ell}, A_{p^m})^\vee.$$

Composing $\delta_{m,\ell}$ with the dual of the restriction $\text{res}_{m,\ell} : H_f^1(K_m, A_{p^m}) \rightarrow H_f^1(K_{m,\ell}, A_{p^m})$, we get a map

$$H_s^1(K_{m,\ell}, T/p^m T) \xrightarrow{\delta_{m,\ell}} H_f^1(K_{m,\ell}, A_{p^m})^\vee \xrightarrow{\text{res}_{m,\ell}^\vee} H_f^1(K_m, A_{p^m})^\vee$$

whose image we denote by $V_m(\ell)$. When m is understood, we shall simply write δ_ℓ for $\delta_{m,\ell}$ and $V(\ell)$ for $V_m(\ell)$.

Proposition 4.12. *Let ℓ_∞ be the sequence of Kolyvagin primes of Proposition 4.11.*

- (1) *For every $m \geq 1$ there is a canonical surjection $V(\ell_m) \twoheadrightarrow W_m^{(\epsilon)} \oplus W_m^{(-\epsilon)}$.*
- (2) *For every $m \geq 1$ there is a canonical surjection $V(\ell_{m+1}) \twoheadrightarrow V(\ell_m)$.*

Proof. Fix an integer $m \geq 1$. Composing the isomorphism δ_{m,ℓ_m} in (48) with the dual of the map $\overline{\text{res}}_{\ell_m}$ introduced in part (2) of Proposition 4.11, we get a surjection

$$H_s^1(K_{m,\ell_m}, T/p^m T) \xrightarrow{\delta_{\ell_m}} H_f^1(K_{m,\ell}, A_{p^m})^\vee \xrightarrow{\overline{\text{res}}_{\ell_m}^\vee} \overline{\Sigma}_m^\vee$$

that, by definition, factors through $V(\ell_m) = V_m(\ell_m)$. Now $\overline{\Sigma}_m^\vee \simeq \text{im}(\vartheta_m)$, which is isomorphic to $W_m^{(\epsilon)} \oplus W_m^{(-\epsilon)}$. Therefore we get an \tilde{R}_m -equivariant surjection

$$V(\ell_m) \twoheadrightarrow W_m^{(\epsilon)} \oplus W_m^{(-\epsilon)},$$

which proves part (1).

As for part (2), let us define the map $V_{m+1}(\ell_{m+1}) \rightarrow V_m(\ell_m)$. There is a commutative diagram

$$\begin{array}{ccccc} H_s^1(K_{m+1}, T/p^{m+1}T) & \twoheadrightarrow & V_{m+1}(\ell_{m+1}) & \hookrightarrow & H_f^1(K_{m+1}, A_{p^{m+1}})^\vee \\ \downarrow \text{cores} & & & & \downarrow \text{res}^\vee \\ H_s^1(K_m, T/p^m T) & \twoheadrightarrow & V_m(\ell_m) & \hookrightarrow & H_f^1(K_m, A_{p^m})^\vee \end{array}$$

in which we have written simply “res” for the map $H_f^1(K_m, A_{p^m}) \hookrightarrow H_f^1(K_{m+1}, A_{p^{m+1}})$ of Lemma 4.9 and “cores” for the analogous map $H_s^1(K_{m+1}, T/p^{m+1}T) \rightarrow H_s^1(K_m, T/p^m T)$ induced by corestriction. This shows that the dual of the restriction map gives a map $V_{m+1}(\ell_{m+1}) \rightarrow V_m(\ell_m)$. Furthermore, by definition, $\text{im}(\vartheta_m)$ is a quotient of $\text{im}(\vartheta_{m+1})$, hence the map $\overline{\Sigma}_{m+1}^\vee \rightarrow \overline{\Sigma}_m^\vee$ is surjective. Now there is yet another commutative diagram

$$\begin{array}{ccccc} R_{m+1}^{(+)} \oplus R_{m+1}^{(-)} \simeq H_s^1(K_{m+1,\ell_{m+1}}, T/p^{m+1}T) & \twoheadrightarrow & V_{m+1}(\ell_{m+1}) & \twoheadrightarrow & \overline{\Sigma}_{m+1}^\vee \\ \downarrow & & \downarrow & & \downarrow \\ R_m^{(+)} \oplus R_m^{(-)} \simeq H_s^1(K_{m,\ell_m}, T/p^m T) & \twoheadrightarrow & V_m(\ell_m) & \twoheadrightarrow & \overline{\Sigma}_m^\vee \end{array}$$

where the vertical maps are (co)restrictions or their duals and the leftmost vertical arrow is surjective because the same is true of the map $R_{m+1} \rightarrow R_m$ (here we are implicitly using (18) and the identification provided by local duality). The desired surjectivity follows. \square

In the rest of the paper, let $\ell_\infty = (\ell_m)_{m \geq 1}$ be the sequence of Kolyvagin primes constructed in Proposition 4.11. It follows from part (2) of Proposition 4.12 that we can define the Λ -module

$$V(\ell_\infty) := \varprojlim_m V(\ell_m).$$

Proposition 4.13. *There is a surjection*

$$V(\ell_\infty) \twoheadrightarrow \Lambda x \oplus \Lambda y.$$

Proof. The inverse limit of the maps in part (1) of Proposition 4.12 gives a surjection

$$(49) \quad V(\ell_\infty) \twoheadrightarrow \varprojlim_m (W_m^{(\epsilon)} \oplus W_m^{(-\epsilon)}),$$

where we use the fact that the projective system satisfies the Mittag–Leffler condition, as all the modules involved are finite. On the other hand, there is a short exact sequence

$$0 \longrightarrow Z_m \longrightarrow \text{im}(p_m \circ \vartheta) \longrightarrow W_m^{(\epsilon)} \oplus W_m^{(-\epsilon)} \longrightarrow 0,$$

and passing to inverse limits shows that there is a short exact sequence

$$0 \longrightarrow \varprojlim_m Z_m \longrightarrow \Lambda x \oplus \Lambda y \longrightarrow \varprojlim_m (W_m^{(\epsilon)} \oplus W_m^{(-\epsilon)}) \longrightarrow 0.$$

Since $\Lambda x \cap \Lambda y = \{0\}$ and $\varprojlim_m Z_m \subset \Lambda x \cap \Lambda y$, we have $\varprojlim_m Z_m = 0$, therefore $\Lambda x \oplus \Lambda y$ is isomorphic to $\varprojlim_m (W_m^{(\epsilon)} \oplus W_m^{(-\epsilon)})$. Combining this with (49) gives the result. \square

4.7. Kolyvagin classes. We briefly review the construction of Kolyvagin classes attached to Heegner cycles.

Let ℓ be a Kolyvagin prime relative to p^m ; in particular, $p^m \mid \ell + 1$ and $p^m \mid a_\ell$. Assume also that $p^{m+1} \nmid \ell + 1 \pm a_\ell$. Let H_ℓ be the ring class field of K of conductor ℓ . The fields K_m and H_ℓ are linearly disjoint over the Hilbert class field H_1 of K , and $\text{Gal}(H_\ell/H_1)$ is cyclic of order $\ell + 1$. Let $H_{m,\ell}^{(p)}$ be the maximal subextension of the composite $K_m H_\ell$ having p -power degree over K_m and let

$$\alpha(\ell) := \text{cores}_{H_{\ell p^{m+1}}/H_{m,\ell}^{(p)}}(y_{\ell p^{m+1}}) \in H_f^1(H_{m,\ell}^{(p)}, T)$$

be the corestriction from $H_{\ell p^{m+1}}$ to $H_{m,\ell}^{(p)}$ of the Heegner cycle $y_{\ell p^{m+1}} \in H_f^1(H_{\ell p^{m+1}}, T)$.

Set $\mathfrak{G}_\ell := \text{Gal}(H_{m,\ell}^{(p)}/K_m)$. By class field theory, if $n_\ell := \text{ord}_p(\ell + 1)$ then $\mathfrak{G}_\ell \simeq \mathbb{Z}/p^{n_\ell}\mathbb{Z}$; in particular, $m \mid n_\ell$. Fix a generator σ_ℓ of \mathfrak{G}_ℓ and consider the Kolyvagin operator

$$\mathbf{D}_\ell := \sum_{i=1}^{p^{n_\ell}-1} i \sigma_\ell^i \in \mathbb{Z}/p^m \mathbb{Z}[\mathfrak{G}_\ell].$$

One has $(\sigma_\ell - 1)\mathbf{D}_\ell = -\text{tr}_{H_{m,\ell}^{(p)}/K_m}$, and therefore

$$\mathbf{D}_\ell(\alpha(\ell)) \in H_f^1(H_{m,\ell}^{(p)}, T/p^m T)^{\mathfrak{G}_\ell}.$$

Moreover, restriction induces an isomorphism

$$(50) \quad \text{res}_{H_{m,\ell}^{(p)}/K_m} : H_f^1(K_m, T/p^m T) \xrightarrow{\simeq} H_f^1(H_{m,\ell}^{(p)}, T/p^m T)^{\mathfrak{G}_\ell}.$$

We can give the following

Definition 4.14. The *Kolyvagin class* $d(\ell) \in H^1(K_m, T/p^m T)$ attached to ℓ is the class corresponding to $\mathbf{D}_\ell(\alpha(\ell))$ under isomorphism (50).

If ℓ is a Kolyvagin prime relative to p^m then there exists ([24, p.116]) a τ -antiequivariant isomorphism of $\mathcal{O}_p/p^m \mathcal{O}_p$ -modules

$$\phi_{m,\ell} : H_s^1(K_{m,\ell}, T/p^m T) \xrightarrow{\simeq} H_f^1(K_{m,\ell}, T/p^m T).$$

Finally, recall the map ∂_ℓ that was defined at the end of §2.1, which consists of the sum of the projections to the singular local cohomology groups for all primes of K_m dividing ℓ .

Proposition 4.15. *The class $d(\ell)$ enjoys the following properties:*

- (1) if v is a prime of K_m not dividing ℓ then $\text{res}_v(d(\ell))$ belongs to $H_f^1(K_m, T/p^m T)$;
- (2) $\phi_{m,\ell}(\partial_\ell(d(\ell))) = u_\ell \cdot \text{res}_\ell(\alpha_m)$ with $u_\ell \in \mathcal{O}_p^\times$.

Proof. This follows from [24, Proposition 10.2] for primes $v \nmid p$, while for primes $v \mid p$ it is a consequence of the de Rham conjecture for open varieties proved by Faltings. For details, see [18, Proposition 3.17 and Lemma 3.19]. \square

4.8. Global duality. In this subsection we use Kolyvagin classes, combined with global reciprocity laws, to bound the rank of the Λ -module $V(\ell_\infty)$, which was introduced in §4.6 and surjects onto $\Lambda x \oplus \Lambda y$.

Recall that if E is a number field, $s \in H^1(E, A_{p^m})$ and $t \in H^1(E, T/p^m T)$ then

$$(51) \quad \sum_v \langle \text{res}_v(s), \text{res}_v(t) \rangle_{E,v} = 0$$

where v ranges over all finite places of E and $\langle \cdot, \cdot \rangle_{E,v}$ is the local Tate pairing at v .

Proposition 4.16. *The Λ -rank of $V(\ell_\infty)$ is at most 1.*

Proof. For every $m \geq 1$ consider the Kolyvagin class $d(\ell_m) \in H^1(K_m, T/p^m T)$. Since there is an isomorphism of \tilde{R}_m -modules $\text{res}_{\ell_m}(\mathcal{E}_m) \simeq R_m^{(\epsilon)} \theta_m$, where $\theta_m \in R_m$ is defined as in §4.3, part (2) of Proposition 4.15 implies that

$$\text{res}_{\ell_m}(R_m d(\ell_m)) \simeq R_m^{(-\epsilon)} \theta_m$$

as \tilde{R}_m -modules. If we let $\xi_m^{(\pm)}$ be generators of $H^1(K_{m,\ell_m}, T/p^m T)^{(\pm)}$ as $R_m^{(\pm)}$ -module then

$$(52) \quad \text{res}_{\ell_m}(R_m d(\ell_m)) \cap \left(H^1(K_{m,\ell_m}, T/p^m T)^{(\epsilon)} \oplus \{0\} \right) = \{0\}.$$

Using [1, §1.2, Lemma 7], write $\text{res}_{\ell_m}(d(\ell_m)) = (\rho_m \theta_m \xi_m^{(\epsilon)}, \nu_m \theta_m \xi_m^{(-\epsilon)})$ for suitable $\rho_m \in R_m$ and $\nu_m \in R_m^\times$. Define $W_m := R_m(\rho_m \xi_m^{(\epsilon)}, \nu_m \xi_m^{(-\epsilon)})$. Then (52) gives a decomposition

$$H^1(K_{m,\ell_m}, T/p^m T) \simeq H^1(K_{m,\ell_m}, T/p^m T)^{(\epsilon)} \oplus W_m,$$

from which we deduce that

$$\theta_m H^1(K_{m,\ell_m}, T/p^m T) \simeq \theta_m H^1(K_{m,\ell_m}, T/p^m T)^{(\epsilon)} \oplus \text{res}_{\ell_m}(R_m d(\ell_m)).$$

By part (1) of Proposition 4.15, the class $d(\ell_m)$ is trivial at all the primes not dividing ℓ_m , and then (51) implies that $(\delta_{\ell_m} \circ \text{res}_{\ell_m})(d(\ell_m)) = 0$. Summing up, we get

$$\theta_m V(\ell_m) \simeq \delta_{\ell_m}(\theta_m H^1(K_{m,\ell_m}, T/p^m T)) \simeq \delta_{\ell_m}(\theta_m H^1(K_{m,\ell_m}, T/p^m T)^{(\epsilon)}).$$

It follows that $\theta_m V(\ell_m)$ is a cyclic R_m -module for all m . Since $\theta_\infty \in \Lambda$ and $\Lambda = \varprojlim_m R_m$, we conclude that $\theta_\infty V(\ell_\infty)$ is a cyclic Λ -module, and then $V(\ell_\infty)$ is cyclic over Λ as well. \square

4.9. Completion of the proof of the main result. Recall from the beginning of Section 4 that our goal is to show that the element $y \in \ker(\pi)$ is Λ -torsion. But this is immediate: since Λx is free of rank 1, combining Propositions 4.13 and 4.16 shows that Λy is Λ -torsion, which concludes the proof.

Finally, we record the following consequence of Theorem 3.6.

Corollary 4.17. *The Λ -module $V(\ell_\infty)$ has rank 1.*

REFERENCES

1. M. Bertolini, *Selmer groups and Heegner points in anticyclotomic \mathbf{Z}_p -extensions*, Compos. Math. **99** (1995), no. 2, 153–182.
2. ———, *Iwasawa theory for elliptic curves over imaginary quadratic fields*, J. Théor. Nombres Bordeaux **13** (2001), no. 1, 1–25.
3. M. Bertolini and H. Darmon, *Derived heights and generalized Mazur–Tate regulators*, Duke Math. J. **76** (1994), no. 1, 75–111.
4. ———, *Derived p -adic heights*, Amer. J. Math. **117** (1995), no. 6, 1517–1554.

5. ———, *Iwasawa's Main Conjecture for elliptic curves over anticyclotomic \mathbb{Z}_p -extensions*, Ann. of Math. (2) **162** (2005), no. 1, 1–64.
6. A. Besser, *On the finiteness of III for motives associated to modular forms*, Doc. Math. **2** (1997), 31–46.
7. S. Bloch and K. Kato, *L-functions and Tamagawa numbers of motives*, The Grothendieck Festschrift, Vol. I, Progr. Math., vol. 86, Birkhäuser Boston, Boston, MA, 1990, pp. 333–400.
8. A. Brumer, *Pseudocompact algebras, profinite groups and class formations*, J. Algebra **4** (1966), no. 3, 442–470.
9. C. Cornut, *Mazur's conjecture on higher Heegner points*, Invent. Math. **148** (2002), no. 3, 495–523.
10. P. Deligne, *Formes modulaires et représentations l -adiques*, Séminaire Bourbaki. Vol. 1968/69: Exposés 347–363, Lecture Notes in Math., vol. 175, Springer, Berlin, 1971, pp. 139–172.
11. O. Fouquet, *Dihedral Iwasawa theory of nearly ordinary quaternionic automorphic forms*, Compos. Math. **149** (2013), no. 3, 356–416.
12. B. Howard, *The Heegner point Kolyvagin system*, Compos. Math. **140** (2004), no. 6, 1439–1472.
13. ———, *Iwasawa theory of Heegner points on abelian varieties of GL_2 type*, Duke Math. J. **124** (2004), no. 1, 1–45.
14. ———, *Special cohomology classes for modular Galois representations*, J. Number Theory **117** (2006), no. 2, 406–438.
15. V. A. Kolyvagin, *Euler systems*, The Grothendieck Festschrift, Vol. II, Progr. Math., vol. 87, Birkhäuser Boston, Boston, MA, 1990, pp. 435–483.
16. S. Lang and H. Trotter, *Frobenius distributions in GL_2 -extensions*, Lecture Notes in Mathematics, Vol. 504, Springer-Verlag, Berlin-New York, 1976.
17. M. Longo and S. Vigni, *Iwasawa theory of Heegner cycles, II. Annihilators and Main Conjecture*, in preparation.
18. ———, *A refined Beilinson–Bloch conjecture for motives of modular forms*, submitted (2013).
19. B. Mazur, *Rational points of abelian varieties with values in towers of number fields*, Invent. Math. **18** (1972), no. 3–4, 183–266.
20. ———, *Modular curves and arithmetic*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983), PWN, Warsaw, 1984, pp. 185–211.
21. B. Mazur and K. Rubin, *Kolyvagin systems*, Mem. Amer. Math. Soc. **168** (2004), no. 799, viii+96.
22. M. R. Murty and V. K. Murty, *A variant of the Lang–Trotter conjecture*, Number theory, analysis and geometry, Springer, New York, 2012, pp. 461–474.
23. M. R. Murty, V. K. Murty, and N. Saradha, *Modular forms and the Chebotarev density theorem*, Amer. J. Math. **110** (1988), no. 2, 253–281.
24. J. Nekovář, *Kolyvagin's method for Chow groups of Kuga–Sato varieties*, Invent. Math. **107** (1992), no. 1, 99–125.
25. ———, *Selmer complexes*, Astérisque (2006), no. 310, viii+559.
26. J. Neukirch, A. Schmidt, and K. Wingberg, *Cohomology of number fields*, Grundlehren der Mathematischen Wissenschaften, vol. 323, Springer-Verlag, Berlin, 2000.
27. T. Ochiai, *Control theorem for Bloch–Kato's Selmer groups of p -adic representations*, J. Number Theory **82** (2000), no. 1, 69–90.
28. B. Perrin-Riou, *Fonctions L p -adiques, théorie d'Iwasawa et points de Heegner*, Bull. Soc. Math. France **115** (1987), no. 4, 399–456.
29. J.-P. Serre, *Linear representations of finite groups*, Graduate Texts in Mathematics, vol. 42, Springer-Verlag, New York, 1977.
30. ———, *Quelques applications du théorème de densité de Chebotarev*, Inst. Hautes Études Sci. Publ. Math. **54** (1981), 323–401.
31. J. Tate, *Relations between K_2 and Galois cohomology*, Invent. Math. **36** (1976), no. 1, 257–274.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PADOVA, VIA TRIESTE 63, 35121 PADOVA, ITALY
E-mail address: `mlongo@math.unipd.it`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, VIA DODECANESO 35, 16146 GENOVA, ITALY
E-mail address: `vigni@dima.unige.it`