

EXTREMES OF A CLASS OF NONHOMOGENEOUS GAUSSIAN RANDOM FIELDS

BY KRZYSZTOF DEBICKI^{1,2,*},
ENKELEJD HASHORVA^{1,†} AND LANPENG JI^{1,†}

University of Wrocław and University of Lausanne†*

This contribution establishes exact tail asymptotics of $\sup_{(s,t) \in \mathbf{E}} X(s,t)$ for a large class of nonhomogeneous Gaussian random fields X on a bounded convex set $\mathbf{E} \subset \mathbb{R}^2$, with variance function that attains its maximum on a segment on \mathbf{E} . These findings extend the classical results for homogeneous Gaussian random fields and Gaussian random fields with unique maximum point of the variance. Applications of our result include the derivation of the exact tail asymptotics of the Shepp statistics for stationary Gaussian processes, Brownian bridge and fractional Brownian motion as well as the exact tail asymptotic expansion for the maximum loss and span of stationary Gaussian processes.

1. Introduction. Consider the fractional Brownian motion (fBm) incremental random field

$$X_\alpha(s,t) = B_\alpha(s+t) - B_\alpha(s), \quad (s,t) \in [0,\infty)^2,$$

where $\{B_\alpha(t), t \in \mathbb{R}\}$ is a standard fBm with Hurst index $\alpha/2 \in (0,1]$ which is a centered self-similar Gaussian process with stationary increments and covariance function

$$\text{Cov}(B_\alpha(t), B_\alpha(s)) = \frac{1}{2}(|t|^\alpha + |s|^\alpha - |t-s|^\alpha), \quad s, t \in \mathbb{R}.$$

For the case $\alpha = 1$, both $X_\alpha(s,t)$ and its standardized version $X_\alpha^*(s,t) = X_\alpha(s,t)/t^{\alpha/2}$ appear naturally as limit models; see, for example, [8]. In the

Received August 2013; revised October 2014.

¹Supported by SNSF Grant 200021-140633/1.

²Supported by NCN Grant No. 2013/09/B/ST1/01778 (2014-2016).

AMS 2000 subject classifications. Primary 60G15; secondary 60G70.

Key words and phrases. Extremes, nonhomogeneous Gaussian random fields, Shepp statistics, fractional Brownian motion, maximum loss, span of Gaussian processes, Pickands constant, Piterbarg constant, generalized Pickands–Piterbarg constant.

This is an electronic reprint of the original article published by the [Institute of Mathematical Statistics](#) in *The Annals of Probability*, 2016, Vol. 44, No. 2, 984–1012. This reprint differs from the original in pagination and typographic detail.

literature,

$$Y_\alpha(t) = \sup_{s \in [0, S]} X_\alpha(s, t)$$

is referred to as the Shepp statistics of fBm, whereas $Y_\alpha^*(t) = \sup_{s \in [0, S]} X_\alpha^*(s, t)$ as the standardized Shepp statistics. Distributional results for Y_1^* are derived in [28]; see also [27] and Theorem 3.2 in [8]. Other important results for the Shepp statistics of Brownian motion and related quantities are presented in [11, 14, 29]. The first known result for the extremes of the Shepp statistics of Brownian motion goes back to [32], which is complemented in [17] for the case of fBm with $\alpha \in (0, 1)$. In view of the aforementioned papers for any $\alpha \in (0, 1]$,

$$(1) \quad \mathbb{P}\left(\sup_{(s,t) \in [0,1]^2} X_\alpha(s, t) > u\right) = C_\alpha u^{4/\alpha-2} \Psi(u)(1 + o(1))$$

holds as $u \rightarrow \infty$ with C_α a positive constant and $\Psi(\cdot)$ the survival function of an $N(0, 1)$ random variable. There is no result for the case $\alpha \in (1, 2)$ in the literature; we shall cover this gap in Proposition 3.5.

Results for the tail asymptotics of supremum of the standardized Shepp statistics can be derived using the findings of [7] and [20]; see also [18, 19]. However, this is not the case for the tail asymptotics of the supremum of the Shepp statistics Y_α ; no theoretical results in the literature can be applied for this case. This is due to the fact that on $[0, 1]^2$ the variance of X_α attains its maximum at an infinite number of points, that is, its maximal value is attained for any $s \in [0, 1]$ and $t = 1$.

In the asymptotic theory of Gaussian random fields, if the random field has a nonconstant variance function, which attains its maximum at a unique (or finite) number of points, then under the so-called Piterbarg conditions, the exact tail asymptotics of supremum of Gaussian random fields with certain (E, α) structures for the variance and the correlation functions are derived by relying on the Double-Sum method; see, for example, the standard monograph [24].

The principle aim of this contribution is to extend Piterbarg's asymptotic theory for Gaussian random fields to the case where the maximum of the variance function on a bounded convex set \mathbf{E} is attained on finite number of disjoint segments on \mathbf{E} . In particular, we assume that $\{X(s, t), (s, t) \in \mathbf{E}\}$, $\mathbf{E} = [0, S] \times [0, T]$, $S, T > 0$, is a centered Gaussian random field with variance function $\sigma^2(s, t) = \text{Var}(X(s, t))$ that satisfies the following assumption.

ASSUMPTION A1. There exists some positive function $\sigma(t)$ which attains its unique maximum on $[0, T]$ at T , and further

$$(2) \quad \begin{aligned} \sigma(s, t) &= \sigma(t) \quad \forall (s, t) \in \mathbf{E}, \\ \sigma(t) &= 1 - b(T - t)^\beta(1 + o(1)), \quad t \uparrow T \end{aligned}$$

hold for some $\beta, b > 0$.

We shall impose the following assumption on the correlation function $r(s, t, s', t') = \mathbb{E}(\overline{X}(s, t)\overline{X}(s', t'))$ where $\overline{X}(s, t) = X(s, t)/\sigma(s, t)$:

ASSUMPTION A2. There exist constants $a_1 > 0$, $a_2 > 0, a_3 \neq 0$ and $\alpha_1, \alpha_2 \in (0, 2]$ such that

$$(3) \quad \begin{aligned} r(s, t, s', t') \\ = 1 - (|a_1(s - s')|^{\alpha_1} + |a_2(t - t') + a_3(s - s')|^{\alpha_2})(1 + o(1)) \end{aligned}$$

holds uniformly with respect to $s, s' \in [0, S]$, as $|s - s'| \rightarrow 0, t, t' \uparrow T$, and further, there exists some constant $\delta_0 \in (0, T)$ such that

$$(4) \quad r(s, t, s', t') < 1$$

holds for any $s, s' \in [0, S]$ satisfying $s \neq s'$, and $t, t' \in [\delta_0, T]$.

Note that in A2 we assume that $a_3 \neq 0$, which includes a large class of correlation functions with (E, α) structure dealt with in [24]; the classical case $a_3 = 0$ is discussed in Remark 2.3.

Our main result, presented in Theorem 2.2 (and stated in higher generality in Remarks 2.4), derives the exact tail asymptotic behavior of supremum of nonhomogeneous Gaussian random fields X satisfying A1 and A2 and a Hölder condition formulated below in Assumption A3. As an illustration to the derived theory, we analyze exact asymptotics of the tail distribution of extremes of Shepp statistics, the maximum loss and the span for a large class of Gaussian processes.

Organization of the paper: Our principal findings are presented in Section 2 followed by two sections dedicated to applications and examples. All the proofs are relegated to Section 5 and the Appendix.

2. Main results. In this section, we are concerned about the asymptotics of

$$\mathbb{P}\left(\sup_{(s,t) \in \mathbf{E}} X(s, t) > u\right), \quad u \rightarrow \infty$$

discussing first the case that $\mathbf{E} = [0, S] \times [0, T]$.

The Pickands and Piterbarg lemmas (cf. [24]) are fundamental in the analysis of the tail asymptotic behavior of supremum of nonsmooth centered Gaussian processes and Gaussian random fields. Restricting ourselves to the case that $\{X(t), t \geq 0\}$ is a centered stationary Gaussian process with a.s. continuous sample paths and correlation function $r(t)$, such that

$r(t) = 1 - t^\alpha(1 + o(1))$ as $t \rightarrow 0$, with $\alpha \in (0, 2]$, and $r(t) < 1$ for all $t > 0$, in view of the seminal papers by J. Pickands III (see [21, 22]), for any $T \in (0, \infty)$

$$(5) \quad \mathbb{P}\left(\sup_{t \in [0, T]} X(t) > u\right) = \mathcal{H}_\alpha T u^{2/\alpha} \Psi(u)(1 + o(1)), \quad u \rightarrow \infty.$$

Here, \mathcal{H}_α is the *Pickands constant* defined by

$$\mathcal{H}_\alpha = \lim_{T \rightarrow \infty} \frac{1}{T} \mathcal{H}_\alpha[0, T] \in (0, \infty)$$

with

$$\mathcal{H}_\alpha[0, T] = \mathbb{E}\left(\exp\left(\sup_{t \in [0, T]} (\sqrt{2}B_\alpha(t) - t^\alpha)\right)\right).$$

The derivation of (5) is based on *Pickand's lemma* which states that

$$(6) \quad \mathbb{P}\left(\sup_{t \in [0, u^{-2/\alpha}T]} X(t) > u\right) = \mathcal{H}_\alpha[0, T] \Psi(u)(1 + o(1)), \quad u \rightarrow \infty.$$

In [23], Piterbarg rigorously proved Pickand's theorem and further derived a crucial extension of (6) which we shall refer to as the *Piterbarg lemma*; it states that

$$(7) \quad \mathbb{P}\left(\sup_{t \in [0, u^{-2/\alpha}T]} \frac{X(t)}{1 + bt^\alpha} > u\right) = \mathcal{P}_\alpha^b[0, T] \Psi(u)(1 + o(1)), \quad u \rightarrow \infty$$

holds for any $b > 0$ with

$$\mathcal{P}_\alpha^b[0, T] = \mathbb{E}\left(\exp\left(\sup_{t \in [0, T]} (\sqrt{2}B_\alpha(t) - (1 + b)t^\alpha)\right)\right) \in (0, \infty).$$

The positive constant (referred to as the *Piterbarg constant*) given by

$$\mathcal{P}_\alpha^b = \lim_{T \rightarrow \infty} \mathcal{P}_\alpha^b[0, T] \in (0, \infty)$$

appears naturally when dealing with the extremes of nonstationary Gaussian processes or Gaussian random fields; see, for example, [24] and our main result below. It is known that $\mathcal{H}_1 = 1$, $\mathcal{H}_2 = 1/\sqrt{\pi}$, and

$$(8) \quad \mathcal{P}_1^b = 1 + \frac{1}{b}, \quad \mathcal{P}_2^b = \frac{1}{2} \left(1 + \sqrt{1 + \frac{1}{b}}\right), \quad b > 0$$

see, for example, [2, 10, 12, 13, 15, 16].

We note in passing that for stationary Gaussian processes [3] and [5] presented new elegant proofs of (5) without using the Pickands lemma. The following extension of the Pickands and Piterbarg lemmas plays an important role in our analysis. Hereafter, we denote by \tilde{B}_α and B_α two independent fBm's defined on \mathbb{R} with Hurst index $\alpha/2 \in (0, 1]$. Recall that $\Psi(\cdot)$ denotes the survival function of an $N(0, 1)$ random variable; we write below $\Gamma(\cdot)$ for the Euler Gamma function.

LEMMA 2.1. *Let $\{\eta(s, t), (s, t) \in [0, \infty)^2\}$ be a centered homogeneous Gaussian random field with covariance function*

$$r_\eta(s, t) = \exp(-|a_1 s|^{\alpha_1} - |a_2 t - a_3 s|^{\alpha_2}), \quad (s, t) \in [0, \infty)^2,$$

where constants $\alpha_i \in (0, 2], i = 1, 2, a_1 > 0, a_2 > 0, a_3 \in \mathbb{R}$. Let further b, S, T be three positive constants. If $\beta \geq \alpha_2 \geq \alpha_1$, then for any positive measurable function $g(u), u > 0$ satisfying $\lim_{u \rightarrow \infty} g(u)/u = 1$

$$(9) \quad \begin{aligned} & \mathbb{P} \left(\sup_{(s,t) \in [0, Su^{-2/\alpha_1}] \times [0, Tu^{-2/\alpha_2}]} \frac{\eta(s, t)}{1 + bt^\beta} > g(u) \right) \\ &= \mathcal{H}_Y^b[S, T] \Psi(g(u)) (1 + o(1)), \quad u \rightarrow \infty, \end{aligned}$$

where

$$(10) \quad \begin{aligned} \mathcal{H}_Y^d[S, T] &= \mathbb{E} \left(\exp \left(\sup_{(s,t) \in [0, S] \times [0, T]} (\sqrt{2}Y(s, t) - \sigma_Y^2(s, t) - d(t)) \right) \right) \\ &\in (0, \infty) \end{aligned}$$

with $\sigma_Y^2(s, t) = \text{Var}(Y(s, t))$ and

$$(11) \quad \begin{aligned} Y(s, t) &= \begin{cases} Y_1(s, t) := \tilde{B}_{\alpha_1}(a_1 s) + B_{\alpha_2}(a_2 t - a_3 s), & \alpha_1 = \alpha_2, \\ Y_2(s, t) := \tilde{B}_{\alpha_1}(a_1 s) + B_{\alpha_2}(a_2 t), & \alpha_1 < \alpha_2, \end{cases} \\ d(t) &= \begin{cases} 0, & \beta > \alpha_2, \\ bt^\beta, & \beta = \alpha_2, \end{cases} \quad (s, t) \in [0, \infty)^2. \end{aligned}$$

Using the definition of Y_1 and Y_2 appearing in (11) we shall determine, for given a_i 's, α_i 's and b, β as above, the following constants (referred to as *generalized Pickands–Piterbarg constants*):

$$\mathcal{M}_{Y, \beta}^b = \lim_{T \rightarrow \infty} \lim_{S \rightarrow \infty} \frac{1}{S} \mathcal{H}_Y^b[S, T] \in (0, \infty)$$

and

$$\begin{aligned} \widetilde{\mathcal{M}}_{Y, \beta}^b &= \lim_{T \rightarrow \infty} \lim_{S \rightarrow \infty} \frac{1}{S} \mathbb{E} \left(\exp \left(\sup_{(s,t) \in [0, S] \times [-T, T]} (\sqrt{2}Y(s, t) - \sigma_Y^2(s, t) - bt^\beta) \right) \right) \\ &\in (0, \infty). \end{aligned}$$

Here, $\mathcal{M}_{Y, \beta}^b$ and $\widetilde{\mathcal{M}}_{Y, \beta}^b$ are defined only for $\beta = \alpha_2$. Note that we suppress a_i 's and α_i 's in the definition of $\mathcal{M}_{Y, \beta}^b$ and $\widetilde{\mathcal{M}}_{Y, \beta}^b$ since they appear directly in the definition of Y .

Additional to [A1](#) and [A2](#) we shall impose the following Hölder condition, which in the literature is called *regularity*; see [\[24\]](#).

ASSUMPTION A3. There exist positive constants $\rho_1, \rho_2, \gamma, \mathcal{Q}$ such that

$$\mathbb{E}((X(s, t) - X(s', t'))^2) \leq \mathcal{Q}(|t - t'|^\gamma + |s - s'|^\gamma)$$

holds for all $t, t' \in [\rho_1, T], s, s' \in [0, S]$ satisfying $|s - s'| < \rho_2$.

We present next our main result.

THEOREM 2.2. *Let $\{X(s, t), (s, t) \in \mathbf{E}\}$, $\mathbf{E} = [0, S] \times [0, T]$ be a centered Gaussian random field with a.s. continuous sample paths. Suppose that Assumptions A1–A3 are satisfied with the parameters mentioned therein. Then, as $u \rightarrow \infty$,*

(i) *if $\beta > \max(\alpha_1, \alpha_2)$*

$$(12) \quad \mathbb{P}\left(\sup_{(s,t) \in \mathbf{E}} X(s, t) > u\right) = S\Gamma(1/\beta + 1) \prod_{k=1}^2 (a_k \mathcal{H}_{\alpha_k}) b^{-1/\beta} \\ \times u^{2/\alpha_2 + 2/\alpha_1 - 2/\beta} \Psi(u)(1 + o(1));$$

(ii) *if $\beta = \alpha_2 = \alpha_1$*

$$(13) \quad \mathbb{P}\left(\sup_{(s,t) \in \mathbf{E}} X(s, t) > u\right) = S\mathcal{M}_{Y_1, \alpha_1}^b u^{2/\alpha_1} \Psi(u)(1 + o(1));$$

(iii) *if $\beta = \alpha_2 > \alpha_1$*

$$(14) \quad \mathbb{P}\left(\sup_{(s,t) \in \mathbf{E}} X(s, t) > u\right) = Sa_1 a_2 \mathcal{P}_{\alpha_2}^{ba_2^{-\alpha_2}} \mathcal{H}_{\alpha_1} u^{2/\alpha_1} \Psi(u)(1 + o(1));$$

(iv) *if $\beta < \alpha_2 = \alpha_1$*

$$(15) \quad \mathbb{P}\left(\sup_{(s,t) \in \mathbf{E}} X(s, t) > u\right) = S(a_1^{\alpha_1} + |a_3|^{\alpha_1})^{1/\alpha_1} \mathcal{H}_{\alpha_1} u^{2/\alpha_1} \Psi(u)(1 + o(1));$$

(v) *if $\beta < \alpha_2$ and $\alpha_1 < \alpha_2$*

$$(16) \quad \mathbb{P}\left(\sup_{(s,t) \in \mathbf{E}} X(s, t) > u\right) = Sa_1 \mathcal{H}_{\alpha_1} u^{2/\alpha_1} \Psi(u)(1 + o(1));$$

(vi) *if $\beta = \alpha_1 > \alpha_2$*

$$(17) \quad \mathbb{P}\left(\sup_{(s,t) \in \mathbf{E}} X(s, t) > u\right) = Sa_1 \mathcal{P}_{\alpha_1}^{b(|a_3|/(a_1 a_2))^{\alpha_1}} \mathcal{H}_{\alpha_2} u^{2/\alpha_2} \Psi(u)(1 + o(1));$$

(vii) *if $\beta < \alpha_1$ and $\alpha_2 < \alpha_1$*

$$\mathbb{P}\left(\sup_{(s,t) \in \mathbf{E}} X(s, t) > u\right) = S|a_3| \mathcal{H}_{\alpha_2} u^{2/\alpha_2} \Psi(u)(1 + o(1)).$$

REMARK 2.3. If $a_3 = 0$, then there are only three scenarios to be considered. In particular if $\beta > \alpha_2$, then (12) holds. If $\beta = \alpha_2$, then (14) holds, whereas if $\beta < \alpha_2$, then (16) is valid.

REMARK 2.4. (a) Let \mathbf{E} be any bounded convex subset of \mathbb{R}^2 . Assume that on \mathbf{E} the maximum of the standard deviation $\sigma(s, t)$ is attained only on a segment \mathbf{L} which is inside of \mathbf{E} , parallel to s -axis and of length ℓ . Then the claims of Theorem 2.2 are still valid, by replacing S with ℓ in cases (i)–(vii), $\Gamma(\cdot)$ with $2\Gamma(\cdot)$ in case (i), $\mathcal{M}_{Y_1, \alpha_1}^b$ with $\widetilde{\mathcal{M}}_{Y_1, \alpha_1}^b$ in cases (ii), $\mathcal{P}_{\alpha_2}^{ba_2^{-\alpha_2}}$ with $\widetilde{\mathcal{P}}_{\alpha_2}^{ba_2^{-\alpha_2}}$ in case (iii), and $\mathcal{P}_{\alpha_1}^{b(|a_3|/(a_1 a_2))^{\alpha_1}}$ with $\widetilde{\mathcal{P}}_{\alpha_1}^{b(|a_3|/(a_1 a_2))^{\alpha_1}}$ in case (vi), respectively. Here, $\widetilde{\mathcal{P}}_{\alpha}^b$, with $b > 0$ and $\alpha \in (0, 2]$ is the Piterbarg constant defined on the real line, that is,

$$\widetilde{\mathcal{P}}_{\alpha}^b = \lim_{T \rightarrow \infty} \mathbb{E} \left(\exp \left(\sup_{t \in [-T, T]} (\sqrt{2} B_{\alpha}(t) - (1+b)t^{\alpha}) \right) \right) \in (0, \infty).$$

(b) Assume that on \mathbf{E} the maximum of the standard deviation $\sigma(s, t)$ is attained only on n segments $\{\mathbf{L}_i\}_{i=1}^n$ which are inside or on the boundary of \mathbf{E} , and parallel to s -axis. By the convexity of \mathbf{E} , we can always find n nonadjacent convex sets $\{\mathbf{E}_i\}_{i=1}^n$ such that $\mathbf{L}_i \subset \mathbf{E}_i \subset \mathbf{E}$, $i = 1, \dots, n$. If further for any $i \neq j$

$$(18) \quad \sup_{(s, t) \in \mathbf{E}_i, (s', t') \in \mathbf{E}_j} r(s, t, s', t') < 1$$

holds, then

$$(19) \quad \mathbb{P} \left(\sup_{(s, t) \in \mathbf{E}} X(s, t) > u \right) = \sum_{i=1}^n \mathbb{P} \left(\sup_{(s, t) \in \mathbf{E}_i} X(s, t) > u \right) (1 + o(1))$$

as $u \rightarrow \infty$. Additionally, suppose that on each $\{\mathbf{E}_i\}_{i=1}^n$ the Assumptions A1–A3 are satisfied. Then an explicit expression for (19) can be established by applying the results in Theorem 2.2 and Remark 2.4(a) above.

(c) Similar results can also be obtained when the segments $\{\mathbf{L}_i\}_{i=1}^n$, where the maximum of $\sigma(s, t)$ is attained, are nonparallel and disjoint. Specifically, we see from Remark 2.4(b) that it is sufficient to consider the asymptotics of

$$\mathbb{P} \left(\sup_{(s, t) \in \mathbf{E}_i} X(s, t) > u \right), \quad u \rightarrow \infty, i = 1, \dots, n,$$

respectively. Let $(s, t)^{\top}$ be the transpose of (s, t) . Then, for any $i = 1, \dots, n$, there is a nondegenerate lower triangular (rotation) matrix $A_i \in \mathbb{R}^{2 \times 2}$ such that the maximum of the variance of $X((A_i(s, t)^{\top})^{\top})$ on $A_i^{-1} \mathbf{E}_i = \{(\tilde{s}, \tilde{t}) : (\tilde{s}, \tilde{t})^{\top} = A_i^{-1}(s, t)^{\top}, (s, t) \in \mathbf{E}_i\}$ is attained on a line parallel to s -axis or

t -axis. Consequently, similar results as in Theorem 2.2 can be obtained if certain Assumptions as A1–A3 are satisfied by each $\{X((A_i(s, t)^\top)^\top), (s, t) \in A_i^{-1}\mathbf{E}_i\}$.

We conclude this section with an example, which illustrates the existence of all the cases discussed in Theorem 2.2.

EXAMPLE 2.5. Consider a Gaussian random field defined as

$$Z(s, t) = \frac{1}{\sqrt{2}}(Y(s+t) - X(s))(1 - b(T-t)^\beta), \quad (s, t) \in [0, S] \times [0, T],$$

where b, β are two positive constants, and X, Y are two independent centered stationary Gaussian processes with covariance functions r_X, r_Y satisfying as $t \rightarrow 0$

$$r_X(t) = 1 - a_1 t^{\alpha_1}(1 + o(1)), \quad r_Y(t) = 1 - a_2 t^{\alpha_2}(1 + o(1))$$

for some constants $a_i > 0, \alpha_i \in (0, 2], i = 1, 2$. Further, assume that

$$r_X(s) < 1, \forall s \in (0, S] \quad r_Y(t) < 1 \quad \forall t \in (0, S+T].$$

It follows that the assumptions of Theorem 2.2 are satisfied by $\{Z(s, t), (s, t) \in [0, S] \times [0, T]\}$.

3. Extremes of Shepp statistics. For a given centered Gaussian process $\{X(t), t \geq 0\}$, we shall define the incremental random field Z by

$$(20) \quad Z(s, t) = X(s+t) - X(s), \quad (s, t) \in [0, S] \times [0, T].$$

The asymptotic analysis of the supremum of the Shepp statistics

$$Y(t) = \sup_{s \in [0, S]} Z(s, t), \quad t \in [0, T]$$

boils down to the study of the tail asymptotics of the double-supremum $\sup_{(s, t) \in [0, S] \times [0, T]} Z(s, t)$. In this section, we shall consider several important examples which can be analysed utilising the theory developed in Section 2.

3.1. *Stationary Gaussian processes.* Consider the Gaussian random field Z as in (20) where X is a centered stationary Gaussian process with covariance function r_X satisfying the following conditions:

- S1: $r_X(t)$ attains its minimum on $[0, T]$ at the unique point $t = T$;
- S2: there exist positive constants α_1, a_1, a_2 and $\alpha_2 \in (0, 2)$ such that

$$r_X(t) = r_X(T) + a_1(T-t)^{\alpha_1}(1 + o(1)), \quad t \rightarrow T,$$

$$r_X(t) = 1 - a_2 t^{\alpha_2}(1 + o(1)), \quad t \rightarrow 0;$$

- S3: $r_X(s) < 1$ for any $s \in (0, S+T]$.

PROPOSITION 3.1. *Let $\{Z(s, t), (s, t) \in [0, S] \times [0, T]\}$ be an incremental random field given as in (20) with r_X satisfying S1–S3. Suppose that r_X is twice continuously differentiable on $[\mu, T]$ for some $\mu \in (0, T)$, $|r_X''(T)| \in (0, \infty)$, and let $b_i = a_i/\rho_T^2, i = 1, 2$ with $\rho_T = \sqrt{2(1 - r_X(T))}$. Then, as $u \rightarrow \infty$,*

(i) *if $\alpha_1 > \alpha_2$*

$$\begin{aligned} & \mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [0,T]} Z(s, t) > u\right) \\ &= S\Gamma(1/\alpha_1 + 1)\mathcal{H}_{\alpha_2}^2 b_2^{2/\alpha_2} b_1^{-1/\alpha_1} \left(\frac{u}{\rho_T}\right)^{4/\alpha_2 - 2/\alpha_1} \Psi\left(\frac{u}{\rho_T}\right) (1 + o(1)); \end{aligned}$$

(ii) *if $\alpha_1 = \alpha_2$*

$$\mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [0,T]} Z(s, t) > u\right) = S\mathcal{M}_{Y, \alpha_1}^{b_1} \left(\frac{u}{\rho_T}\right)^{2/\alpha_2} \Psi\left(\frac{u}{\rho_T}\right) (1 + o(1)),$$

where

$$Y(s, t) := \tilde{B}_{\alpha_2}(b_2^{1/\alpha_2} s) + B_{\alpha_2}(b_2^{1/\alpha_2} t - b_2^{1/\alpha_2} s), \quad (s, t) \in [0, \infty)^2;$$

(iii) *if $\alpha_1 < \alpha_2$*

$$\mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [0,T]} Z(s, t) > u\right) = S(2b_2)^{1/\alpha_2} \mathcal{H}_{\alpha_2} \left(\frac{u}{\rho_T}\right)^{2/\alpha_2} \Psi\left(\frac{u}{\rho_T}\right) (1 + o(1)).$$

We present two important examples that illustrate Proposition 3.1.

EXAMPLE 3.2 (Slepian process). Consider X to be the Slepian process, that is,

$$X(t) = B_1(t+1) - B_1(t), \quad t \in [0, \infty),$$

with B_1 the standard Brownian motion. It follows that the assumptions of Proposition 3.1 are satisfied, hence as $u \rightarrow \infty$

$$\mathbb{P}\left(\sup_{(s,t) \in [0,1] \times [0,1/2]} Z(s, t) > u\right) = \mathcal{M}_{Y,1}^1 u^2 \Psi(u) (1 + o(1))$$

holds with $Y(s, t) := \tilde{B}_1(s) + B_1(t - s), (s, t) \in (0, \infty)^2$.

EXAMPLE 3.3 (Ornstein–Uhlenbeck process). Consider a centered stationary Gaussian process X with covariance function $r(t) = e^{-t}, t \geq 0$. Then following Proposition 3.1,

$$\mathbb{P}\left(\sup_{(s,t) \in [0,1]^2} Z(s, t) > u\right) = \mathcal{M}_{Y,1}^{b_1} b_1 u^2 \Psi(\sqrt{b_1} u) (1 + o(1)), \quad u \rightarrow \infty,$$

with $b_1 = e^{-1}/(2(1 - e^{-1}))$, $b_2 = 1/(2(1 - e^{-1}))$ and $Y(s, t) := \tilde{B}_1(b_2 s) + B_1(b_2 t - b_2 s), (s, t) \in (0, \infty)^2$.

3.2. *Brownian bridge.* In this section, we analyze

$$(21) \quad Z(s, t) = X(s+t) - X(s), \quad s, s+t \in [0, 1],$$

where $X(s) := B_1(s) - sB_1(1)$, $s \in [0, 1]$ is a Brownian bridge (recall B_1 is a standard Brownian motion). Clearly, X is nonstationary and, therefore, we cannot apply Proposition 3.1 for this case.

PROPOSITION 3.4. *If $\{Z(s, t), (s, t) \in [0, 1/2]^2\}$ is given by (21), then*

$$(22) \quad \mathbb{P}\left(\sup_{(s,t) \in [0, 1/2]^2} Z(s, t) > u\right) = 2^{5/2} \sqrt{\pi} u^3 \Psi(2u)(1 + o(1)), \quad u \rightarrow \infty.$$

3.3. *Fractional Brownian motion.* Consider the fBm incremental random field

$$(23) \quad Z(s, t) = B_\alpha(s+t) - B_\alpha(s), \quad (s, t) \in [0, S] \times [0, 1],$$

where B_α is the fBm with Hurst index $\alpha/2 \in (0, 1)$.

The following proposition extends the main result of [17] to the whole range of $\alpha \in (0, 2)$.

PROPOSITION 3.5. *Let $\{Z(s, t), (s, t) \in [0, S] \times [0, 1]\}$ be given as in (23). We have, as $u \rightarrow \infty$,*

(i) *if $\alpha \in (0, 1)$*

$$(24) \quad \mathbb{P}\left(\sup_{(s,t) \in [0, S] \times [0, 1]} Z(s, t) > u\right) = S 2^{1-2/\alpha} \alpha^{-1} \mathcal{H}_\alpha^2 u^{4/\alpha-2} \Psi(u)(1 + o(1));$$

(ii) *if $\alpha = 1$*

$$(25) \quad \mathbb{P}\left(\sup_{(s,t) \in [0, S] \times [0, 1]} Z(s, t) > u\right) = S \mathcal{M}_{Y,1}^{1/2} u^2 \Psi(u)(1 + o(1)),$$

with

$$Y(s, t) := \tilde{B}_1(2^{-1}s) + B_1(2^{-1}(t-s)), \quad (s, t) \in [0, \infty)^2;$$

(iii) *if $\alpha \in (1, 2)$*

$$(26) \quad \mathbb{P}\left(\sup_{(s,t) \in [0, S] \times [0, 1]} Z(s, t) > u\right) = S \mathcal{H}_\alpha u^{2/\alpha} \Psi(u)(1 + o(1)).$$

4. Extremes of maximum loss and span of Gaussian processes. Let $\{\xi(t), t \in [0, 1]\}$ be a Gaussian process with a.s. continuous sample paths. The maximum loss of the process ξ is given by

$$\chi_1(\xi) = \max_{0 \leq s \leq t \leq 1} (\xi(s) - \xi(t)),$$

and its span is defined as

$$\chi_2(\xi) = \max_{t \in [0, 1]} \xi(t) - \min_{t \in [0, 1]} \xi(t).$$

The notion of the maximum loss of certain Gaussian processes (e.g., Brownian motion and fBm, etc.) plays an important role in finance and insurance modelling; see, for example, [30], [31] and references therein.

In this section, as an application of Theorem 2.2 and Remark 2.4, we derive exact tail asymptotics of the maximum loss for both stationary Gaussian process (in Proposition 4.1) and for Brownian bridge (in Proposition 4.2). The exact tail asymptotics of the span $\chi_2(\xi)$ when ξ is a centered stationary Gaussian process with covariance function that satisfies certain regular conditions is obtained in [26]. The same result should be retrieved, using first a time scaling and then resorting to Remark 2.4. This observation is confirmed in Proposition 4.1 below.

Hereafter, assume that $\{\xi(t), t \in [0, 1]\}$ is a centered stationary Gaussian process with covariance function $r_\xi(s)$ satisfying the following conditions:

- S1': $r_\xi(t)$ attains its minimum on $[0, 1]$ at unique point $t_m \in (0, 1)$;
- S2': there exist positive constants a_1, a_2, α_1 and $\alpha_2 \in (0, 2)$ such that

$$r_\xi(t) = r_\xi(t_m) + a_1|t - t_m|^{\alpha_1}(1 + o(1)), \quad t \rightarrow t_m$$

and

$$r_\xi(t) = 1 - a_2 t^{\alpha_2}(1 + o(1)), \quad t \rightarrow 0;$$

- S3': $r_\xi(t) < 1$ for any $t \in (0, 1]$.

PROPOSITION 4.1. *Let $\{\xi(t), t \in [0, 1]\}$ be a centered stationary Gaussian process with covariance function $r_\xi(t)$ satisfying S1'–S3'. If $r_\xi(t)$ is twice continuously differentiable on interval $[t_m - \mu, t_m + \mu]$ for some positive small constant μ , then, as $u \rightarrow \infty$,*

$$\begin{aligned} \mathbb{P}(\chi_2(\xi) > u) &= 2\mathbb{P}(\chi_1(\xi) > u) \\ (27) \quad &= 2^{2-4/\alpha_2+2/\alpha_1} (1 - r_\xi(t_m)) \mathcal{H}_{\alpha_2}^2 a_2^{2/\alpha_2} (1 - r_\xi(t_m))^{2-4/\alpha_2+2/\alpha_1} \\ &\quad \times u^{4/\alpha_2-2/\alpha_1} \Psi\left(\frac{u}{\sqrt{2(1 - r_\xi(t_m))}}\right) (1 + o(1)). \end{aligned}$$

PROPOSITION 4.2. *If $\{X(t), t \in [0, 1]\}$ is the Brownian bridge given in (21), then, as $u \rightarrow \infty$,*

$$(28) \quad \mathbb{P}(\chi_2(X) > u) = 2\mathbb{P}(\chi_1(X) > u) = 2^{9/2}\sqrt{\pi}u^3\Psi(2u)(1 + o(1)).$$

REMARKS 4.3. (a) The claim in (27) is consistent with Theorem 2.1 in [26].

(b) Let B_α be a standard fBm and consider its maximum loss $\chi_1(B_\alpha)$ and span $\chi_2(B_\alpha)$. The variance function of the random field $X_1(s, t) := B_\alpha(t) - B_\alpha(s)$ is given by

$$\sigma_{X_1}^2(s, t) = |t - s|^\alpha, \quad (s, t) \in [0, 1]^2$$

and attains its maximum only at points $(0, 1)$ and $(1, 0)$. Therefore, Theorem 8.2 in [24] yields that, as $u \rightarrow \infty$,

(i) if $\alpha \in (0, 1)$

$$\mathbb{P}(\chi_2(B_\alpha) > u) = 2\mathbb{P}(\chi_1(B_\alpha) > u) = 2^{3-2/\alpha}\alpha^{-2}\mathcal{H}_\alpha^2 u^{4/\alpha-4}\Psi(u)(1 + o(1));$$

(ii) if $\alpha = 1$

$$\mathbb{P}(\chi_2(B_\alpha) > u) = 2\mathbb{P}(\chi_1(B_\alpha) > u) = 8\Psi(u)(1 + o(1));$$

(iii) if $\alpha \in (1, 2)$

$$\mathbb{P}(\chi_2(B_\alpha) > u) = 2\mathbb{P}(\chi_1(B_\alpha) > u) = 2\Psi(u)(1 + o(1)).$$

5. Proofs.

PROOF OF LEMMA 2.1. The claim follows by a direct application of Lemma A.1 given in the Appendix. \square

PROOF OF THEOREM 2.2. As it will be seen at the end of the proof, by symmetry, cases (vi) and (vii) follow from the claims of cases (iii) and (v), respectively. Thus, we shall first focus on the proof of cases (i)–(v). In view of Assumption A1 there exist some $\theta \in (0, 1)$ and $\rho_0 \geq \rho_1$ (ρ_1 is as in A3) such that

$$\sup_{(s,t) \in [0,S] \times [0,\rho_0]} \sigma(s, t) < \theta.$$

For $\delta(u) = (\ln u/u)^{2/\beta}$, $u > 0$, we may write

$$\begin{aligned} & \mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [T-\delta(u), T]} X(s, t) > u\right) \\ & \leq \mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [0, T]} X(s, t) > u\right) \\ & \leq \mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [T-\delta(u), T]} X(s, t) > u\right) + \pi_1(u) + \pi_2(u), \end{aligned}$$

where

$$\begin{aligned}\pi_1(u) &:= \mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [0,\rho_0]} X(s,t) > u\right), \\ \pi_2(u) &:= \mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [\rho_0, T-\delta(u)]} X(s,t) > u\right).\end{aligned}$$

We shall mainly focus on the analysis of

$$(29) \quad \pi(u) := \mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [T-\delta(u), T]} X(s,t) > u\right), \quad u \rightarrow \infty$$

and show that for $i = 1, 2$

$$(30) \quad \pi_i(u) = o(\pi(u)), \quad u \rightarrow \infty,$$

which then implies

$$\mathbb{P}\left(\sup_{(s,t) \in [0,S] \times [0,T]} X(s,t) > u\right) = \pi(u)(1 + o(1)), \quad u \rightarrow \infty.$$

The asymptotics of (29) will be investigated for the cases (i)–(v) separately by using a case-specific approach.

Case (i) $\beta > \max(\alpha_1, \alpha_2)$: For space saving, we consider only the case that $\alpha_1 = \alpha_2 =: \alpha$; the other cases can be shown with similar arguments. Following the idea of [25] choose first a constant $\alpha_0 \in (\alpha, \beta)$, and denote

$$\Delta_{ij} = \Delta_i \times \Delta_j, \quad \Delta_{ij}^T = \Delta_i \times (T - \Delta_j)$$

with

$$\Delta_i = [iu^{-2/\alpha_0}, (i+1)u^{-2/\alpha_0}], \quad i = 0, 1, \dots$$

Set further

$$\tilde{N}_1(u) = \lfloor Su^{2/\alpha_0} \rfloor + 1, \quad \tilde{N}_2(u) = \lfloor (\ln u)^{2/\beta} u^{2/\alpha_0 - 2/\beta} \rfloor + 1,$$

where $\lfloor \cdot \rfloor$ stands for the ceiling function. By Bonferroni's inequality, we have that

$$(31) \quad \begin{aligned} & \sum_{i=0}^{\tilde{N}_1(u)} \sum_{j=0}^{\tilde{N}_2(u)} \mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}^T} X(s,t) > u\right) \\ & \geq \pi(u) \\ & \geq \sum_{i=0}^{\tilde{N}_1(u)-1} \sum_{j=0}^{\tilde{N}_2(u)-1} \mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}^T} X(s,t) > u\right) - \Sigma_1(u), \end{aligned}$$

with

$$\Sigma_1(u) = \sum_{\substack{0 \leq i, i' \leq \tilde{N}_1(u)-1 \\ (i, j) \neq (i', j')}} \sum_{\substack{0 \leq j, j' \leq \tilde{N}_2(u)-1 \\ (i, j) \neq (i', j')}} \mathbb{P} \left(\sup_{(s, t) \in \Delta_{ij}^T} X(s, t) > u, \right. \\ \left. \sup_{(s, t) \in \Delta_{i'j'}^T} X(s, t) > u \right).$$

For any $\varepsilon \in (0, 1)$ and all u large [set $b_{\pm\varepsilon} := b(1 \pm \varepsilon)$]

$$\mathbb{P} \left(\sup_{(s, t) \in \Delta_{ij}^T} X(s, t) > u \right) \leq \mathbb{P} \left(\sup_{(s, t) \in \Delta_{ij}} \frac{X(s, T-t)}{\sigma(s, T-t)} > u_{j-} \right), \\ \mathbb{P} \left(\sup_{(s, t) \in \Delta_{ij}^T} X(s, t) > u \right) \geq \mathbb{P} \left(\sup_{(s, t) \in \Delta_{ij}} \frac{X(s, T-t)}{\sigma(s, T-t)} > u_{j+} \right),$$

with

$$u_{j-} = u(1 + b_{-\varepsilon}(ju^{-2/\alpha_0})^\beta), \quad u_{j+} = u(1 + b_{+\varepsilon}((j+1)u^{-2/\alpha_0})^\beta).$$

Let $\{\eta_{\pm\varepsilon}(s, t), (s, t) \in [0, \infty)^2\}$ with ε as above be centered stationary Gaussian random fields with covariance functions

$$r_{\eta_{\pm\varepsilon}}(s, t) = \exp(-(1 \pm \varepsilon)^\alpha (|a_1 s|^\alpha + |a_2 t + a_3 s|^\alpha)), \quad (s, t) \in [0, \infty)^2,$$

respectively. By Slepian's lemma (see, e.g., [6] or [4]) for all u large

$$\mathbb{P} \left(\sup_{(s, t) \in \Delta_{ij}} \frac{X(s, T-t)}{\sigma(s, T-t)} > u_{j-} \right) \leq \mathbb{P} \left(\sup_{(s, t) \in \Delta_{ij}} \eta_{+\varepsilon}(s, T-t) > u_{j-} \right).$$

In view of Theorem 7.2 in [24], as $u \rightarrow \infty$,

$$\begin{aligned} \pi(u) &\leq \sum_{i=0}^{\tilde{N}_1(u)} \sum_{j=0}^{\tilde{N}_2(u)} \mathbb{P} \left(\sup_{(s, t) \in \Delta_{ij}} \eta_{+\varepsilon}(s, T-t) > u_{j-} \right) \\ &= (1 + \varepsilon)^2 a_1 a_2 \mathcal{H}_\alpha^2 u^{-4/\alpha_0} \sum_{i=0}^{\tilde{N}_1(u)} \sum_{j=0}^{\tilde{N}_2(u)} u_{j-}^{4/\alpha} \Psi(u_{j-}) (1 + o(1)) \\ (32) \quad &= (1 + \varepsilon)^2 a_1 a_2 \mathcal{H}_\alpha^2 S u^{-2/\alpha_0 + 4/\alpha} \Psi(u) \sum_{j=0}^{\tilde{N}_2(u)} \exp(-b_{-\varepsilon}(ju^{2/\beta - 2/\alpha_0})^\beta) \\ &\quad \times (1 + o(1)) \\ &= (1 + \varepsilon)^2 a_1 a_2 \mathcal{H}_\alpha^2 S u^{4/\alpha - 2/\beta} \Psi(u) \int_0^\infty \exp(-b_{-\varepsilon} x^\beta) dx (1 + o(1)). \end{aligned}$$

Similarly, we obtain

$$\begin{aligned}
& \sum_{i=0}^{\tilde{N}_1(u)-1} \sum_{j=0}^{\tilde{N}_2(u)-1} \mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}^T} X(s,t) > u\right) \\
(33) \quad & \geq \sum_{i=0}^{\tilde{N}_1(u)-1} \sum_{j=0}^{\tilde{N}_2(u)-1} \mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}} \eta_{-\varepsilon}(s, T-t) > u_{j+}\right) \\
& \geq (1-\varepsilon)^2 a_1 a_2 \mathcal{H}_\alpha^2 S u^{4/\alpha-2/\beta} \Psi(u) \int_0^\infty \exp(-b_{+\varepsilon} x^\beta) dx (1+o(1)).
\end{aligned}$$

Next, we deal with the double sum part $\Sigma_1(u)$. Denote the distance of two nonempty sets $A, B \subset \mathbb{R}^n$ by

$$\rho(A, B) = \inf_{x \in A, y \in B} \|x - y\|,$$

with $\|\cdot\|$ the Euclidean distance. We see from (3) that there exists a positive constant ρ_3 such that

$$\begin{aligned}
(34) \quad & \frac{3}{2} (|a_1(s-s')|^\alpha + |a_2(t-t') + a_3(s-s')|^\alpha) \\
& \geq 1 - r(s, t, s', t') \\
& \geq \frac{1}{2} (|a_1(s-s')|^\alpha + |a_2(t-t') + a_3(s-s')|^\alpha)
\end{aligned}$$

for $|s-s'| \leq 2\rho_3$, $|T-t| \leq 2\rho_3$ and $|T-t'| \leq 2\rho_3$. It follows further from (4) that there exists some $\theta_0 \in (0, 1)$ such that

$$\sup_{\substack{0 \leq i, i' \leq \tilde{N}_1(u)-1, 0 \leq j, j' \leq \tilde{N}_2(u)-1 \\ \rho(\Delta_i, \Delta_{i'}) > \rho_3}} \sup_{\substack{(s,t) \in \Delta_{ij}^T \\ (s',t') \in \Delta_{i'j'}^T}} r(s, t, s', t') < \theta_0.$$

Next, we divide the double sum part $\Sigma_1(u)$ as follows:

$$\Sigma_1(u) = \Sigma_{1,1}(u) + \Sigma_{1,2}(u) + \Sigma_{1,3}(u), \quad u \geq 0,$$

where $\Sigma_{1,1}(u)$ is the sum taken on $\rho(\Delta_i, \Delta_{i'}) > \rho_3$, $\Sigma_{1,2}(u)$ is the sum taken on $\rho(\Delta_{ij}^T, \Delta_{i'j'}^T) = 0$ and $\Sigma_{1,3}(u)$ is the sum taken on $u^{-2/\alpha_0} \leq \rho(\Delta_{ij}^T, \Delta_{i'j'}^T)$ and $\rho(\Delta_i, \Delta_{i'}) \leq \rho_3$. We first give the estimation of $\Sigma_{1,1}(u)$. For $\xi(s, t, s', t') := \overline{X}(s, t) + \overline{X}(s', t')$ we have

$$(35) \quad \mathbb{E}(\xi^2(s, t, s', t')) = 4 - 2(1 - r(s, t, s', t'))$$

implying

$$\sup_{\substack{0 \leq i, i' \leq \tilde{N}_1(u)-1, 0 \leq j, j' \leq \tilde{N}_2(u)-1 \\ \rho(\Delta_i, \Delta_{i'}) > \rho_3}} \sup_{\substack{(s,t) \in \Delta_{ij}^T \\ (s',t') \in \Delta_{i'j'}^T}} \mathbb{E}(\xi^2(s, t, s', t')) \leq 4 - 2(1 - \theta_0) < 4.$$

Further, we have

$$\begin{aligned}
& \mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}^T} X(s,t) > u, \sup_{(s,t) \in \Delta_{i'j'}^T} X(s,t) > u\right) \\
& \leq \mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}^T} \bar{X}(s,t) > u, \sup_{(s,t) \in \Delta_{i'j'}^T} \bar{X}(s,t) > u\right) \\
& \leq \mathbb{P}\left(\sup_{\substack{(s,t) \in \Delta_{ij}^T \\ (s',t') \in \Delta_{i'j'}^T}} \xi(s,t,s',t') > 2u\right).
\end{aligned}$$

By Borell–TIS inequality (see [1] or [24]), for u sufficiently large

$$\mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}^T} X(s,t) > u, \sup_{(s,t) \in \Delta_{i'j'}^T} X(s,t) > u\right) \leq \exp\left(-\frac{(u-a)^2}{2-(1-\theta_0)}\right),$$

where $a = \mathbb{E}(\sup_{(s,t),(s',t') \in [0,S] \times [0,T]} \xi(s,t,s',t')) < \infty$. Thus

$$(36) \quad \limsup_{u \rightarrow \infty} \frac{\Sigma_{1,1}(u)}{u^{4/\alpha-2/\beta}\Psi(u)} = 0.$$

The summand of $\Sigma_{1,2}(u)$ is equal to

$$\begin{aligned}
& \mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}^T} X(s,t) > u\right) + \mathbb{P}\left(\sup_{(s,t) \in \Delta_{i'j'}^T} X(s,t) > u\right) \\
& \quad - \mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}^T \cup \Delta_{i'j'}^T} X(s,t) > u\right).
\end{aligned}$$

Since $\rho(\Delta_{ij}^T, \Delta_{i'j'}^T) = 0$, we have for $(s,t) \in \Delta_{ij}^T \cup \Delta_{i'j'}^T$ and sufficiently large u

$$\begin{aligned}
u(1 + b_{-\varepsilon}((j-1)_+ u^{-2/\alpha_0})^\beta) &=: \tilde{u}_{j-} \leq \frac{u}{\sigma(s,t)} \leq \tilde{u}_{j+} \\
&:= u(1 + b_{+\varepsilon}((j+2)u^{-2/\alpha_0})^\beta).
\end{aligned}$$

Using again Theorem 7.2 in [24] for the last term, we have

$$\mathbb{P}\left(\sup_{(s,t) \in \Delta_{ij}^T \cup \Delta_{i'j'}^T} X(s,t) > u\right) \geq 2(1-\varepsilon)^2 a_1 a_2 \mathcal{H}_\alpha^2 u^{-4/\alpha_0} \tilde{u}_{j+}^{4/\alpha} \Psi(\tilde{u}_{j+})(1+o(1))$$

as $u \rightarrow \infty$. Consequently, noting that for any Δ_{ij}^T there are at most 8 sets of the form $\Delta_{i'j'}^T$ in $[0, S] \times [T - \delta(u), T]$ adjacent with it, we conclude that

$$\begin{aligned}
\Sigma_{1,2}(u) &\leq 8 \sum_{i=0}^{\tilde{N}_1(u)} \sum_{j=0}^{\tilde{N}_2(u)} (2(1+\varepsilon)^2 a_1 a_2 \mathcal{H}_\alpha^2 u^{-4/\alpha_0} \tilde{u}_{j-}^{4/\alpha} \Psi(\tilde{u}_{j-}) \\
&\quad - 2(1-\varepsilon)^2 a_1 a_2 \mathcal{H}_\alpha^2 u^{-4/\alpha_0} \tilde{u}_{j+}^{4/\alpha} \Psi(\tilde{u}_{j+}))(1+o(1))
\end{aligned}$$

and thus similar arguments as in (32) yield

$$(37) \quad \limsup_{\varepsilon \rightarrow 0} \limsup_{u \rightarrow \infty} \frac{\Sigma_{1,2}(u)}{u^{4/\alpha-2/\beta} \Psi(u)} = 0.$$

Finally, we estimate $\Sigma_{1,3}(u)$. Since $u^{-2/\alpha_0} \leq \rho(\Delta_{ij}^T, \Delta_{i'j'}^T)$ and $\rho(\Delta_i, \Delta_{i'}) \leq \rho_3$, it follows in view of (34) that

$$\inf_{\substack{0 \leq i, i' \leq \tilde{N}_1(u)-1, \\ \rho(\Delta_i, \Delta_{i'}) \leq \rho_3}} \inf_{\substack{0 \leq j, j' \leq \tilde{N}_2(u)-1 \\ u^{-2/\alpha_0} \leq \rho(\Delta_{ij}^T, \Delta_{i'j'}^T)}} (1 - r(s, t, s', t')) \geq \frac{1}{2} \nu u^{-2\alpha/\alpha_0}$$

for some positive constant ν , and thus

$$\sup_{\substack{0 \leq i, i' \leq \tilde{N}_1(u)-1, \\ \rho(\Delta_i, \Delta_{i'}) \leq \rho_3}} \sup_{\substack{0 \leq j, j' \leq \tilde{N}_2(u)-1 \\ u^{-2/\alpha_0} \leq \rho(\Delta_{ij}^T, \Delta_{i'j'}^T)}} \mathbb{E}(\xi^2(s, t, s', t')) \leq 4 - \nu u^{-2\alpha/\alpha_0}.$$

Consequently, using the Piterbarg inequality (cf. Theorem 8.1 in [24] or Theorem 8.1 in [25]) for the summand of $\Sigma_{1,3}(u)$ we obtain

$$\begin{aligned} & \mathbb{P} \left(\sup_{(s,t) \in \Delta_{ij}^T} X(s, t) > u, \sup_{(s,t) \in \Delta_{i'j'}^T} X(s, t) > u \right) \\ & \leq \mathbb{P} \left(\sup_{\substack{(s,t) \in \Delta_{ij}^T \\ (s',t') \in \Delta_{i'j'}^T}} \xi(s, t, s', t') > 2u \right) \\ & = o \left(\exp \left(-\frac{1}{16} \nu u^{-2((\alpha_0-\alpha)/\alpha_0)} \right) \right) u^{4/\alpha-2/\beta} \Psi(u), \end{aligned}$$

which implies that

$$(38) \quad \begin{aligned} & \limsup_{u \rightarrow \infty} \frac{\Sigma_{1,3}(u)}{u^{4/\alpha-2/\beta} \Psi(u)} \\ & \leq \limsup_{u \rightarrow \infty} \sum_{\substack{0 \leq i, i' \leq \tilde{N}_1(u)-1 \\ (i,j) \neq (i',j')}} \sum_{\substack{0 \leq j, j' \leq \tilde{N}_2(u)-1}} o \left(\exp \left(-\frac{1}{16} \nu u^{-2((\alpha_0-\alpha)/\alpha_0)} \right) \right) \\ & = 0. \end{aligned}$$

Hence, in view of (31)–(33), (36)–(38) and by letting $\varepsilon \rightarrow 0$ we conclude that

$$\pi(u) = a_1 a_2 \mathcal{H}_\alpha^2 S u^{4/\alpha-2/\beta} \Psi(u) \int_0^\infty \exp(-bx^\beta) dx (1 + o(1)), \quad u \rightarrow \infty.$$

Case (ii) $\beta = \alpha_1 = \alpha_2$: In order to simplify notation, we set $\alpha := \alpha_1 = \alpha_2$. Let S_1, T_1 be two positive constants and define

$$\begin{aligned}\widehat{\Delta}_i &= [iS_1u^{-2/\alpha}, (i+1)S_1u^{-2/\alpha}], & i = 0, \dots, N_1(u), \\ \widetilde{\Delta}_i &= [iT_1u^{-2/\alpha}, (i+1)T_1u^{-2/\alpha}], & i = 0, \dots, N_2(u), \\ \overline{\Delta}_{ij} &= \widehat{\Delta}_i \times \widetilde{\Delta}_j, & \overline{\Delta}_{ij}^T = \widehat{\Delta}_i \times (T - \widetilde{\Delta}_j),\end{aligned}$$

where

$$N_1(u) = \left\lfloor \frac{S}{S_1} u^{2/\alpha} \right\rfloor + 1, \quad N_2(u) = \left\lfloor \frac{(\ln u)^{2/\beta}}{T_1} \right\rfloor + 1.$$

Again, Bonferroni's inequality implies

$$\begin{aligned}(39) \quad \Sigma_2(u) + \sum_{i=0}^{N_1(u)} \mathbb{P}\left(\sup_{(s,t) \in \overline{\Delta}_{i0}^T} X(s,t) > u \right) \\ \geq \pi(u) \\ \geq \sum_{i=0}^{N_1(u)-1} \mathbb{P}\left(\sup_{(s,t) \in \overline{\Delta}_{i0}^T} X(s,t) > u \right) - \Sigma_3(u),\end{aligned}$$

where

$$\begin{aligned}\Sigma_2(u) &= \sum_{i=0}^{N_1(u)} \sum_{j=1}^{N_2(u)} \mathbb{P}\left(\sup_{(s,t) \in \overline{\Delta}_{ij}^T} X(s,t) > u \right), \\ \Sigma_3(u) &= \sum_{0 \leq i < i' \leq N_1(u)-1} \sum \mathbb{P}\left(\sup_{(s,t) \in \overline{\Delta}_{i0}^T} X(s,t) > u, \sup_{(s,t) \in \overline{\Delta}_{i'0}^T} X(s,t) > u \right).\end{aligned}$$

Since our approach is of asymptotic nature, for any fixed $0 \leq i \leq N_1(u)$, the local structures of the variance and correlation of the Gaussian random field X on $\overline{\Delta}_{i0}^T$ are the only necessary properties influencing the asymptotics. Therefore,

$$\begin{aligned}\mathbb{P}\left(\sup_{(s,t) \in \overline{\Delta}_{i0}^T} X(s,t) > u \right) \\ = \mathbb{P}\left(\sup_{(s,t) \in \overline{\Delta}_{i0}^T} \frac{\eta(s,t)}{1 + bt^\beta} > u \right) (1 + o(1))\end{aligned}$$

as $u \rightarrow \infty$, where $\{\eta(s,t), (s,t) \in [0, S] \times [0, T]\}$ is the same as in Lemma 2.1. Hence, Lemma 2.1 implies

$$(40) \quad \sum_{i=0}^{N_1(u)} \mathbb{P}\left(\sup_{(s,t) \in \overline{\Delta}_{i0}^T} X(s,t) > u \right) = \frac{S}{S_1} u^{2/\alpha} \mathcal{H}_{Y_1}^b[S_1, T_1] \Psi(u) (1 + o(1))$$

as $u \rightarrow \infty$. Similarly,

$$(41) \quad \begin{aligned} & \sum_{i=0}^{N_1(u)-1} \mathbb{P} \left(\sup_{(s,t) \in \overline{\Delta}_{i0}^T} X(s,t) > u \right) \\ &= \frac{S}{S_1} u^{2/\alpha} \mathcal{H}_{Y_1}^b [S_1, T_1] \Psi(u) (1 + o(1)) \end{aligned}$$

as $u \rightarrow \infty$. Note that, for any $c, d \in \mathbb{R}$

$$\begin{aligned} |c+d|^p &\leq |c|^p + |d|^p, & \text{if } p \in (0, 1], \\ |c+d|^p &\leq 2^{p-1}(|c|^p + |d|^p) & \text{if } p \in (1, \infty). \end{aligned}$$

In view of Slepian's lemma,

$$\begin{aligned} & \mathbb{P} \left(\sup_{(s,t) \in \overline{\Delta}_{ij}^T} X(s,t) > u \right) \\ & \leq \mathbb{P} \left(\sup_{(s,t) \in \overline{\Delta}_{ij}^T} \eta(s,t) > u(1 + b(jT_1 u^{-2/\alpha})^\beta) \right) (1 + o(1)) \\ & \leq \mathbb{P} \left(\sup_{(s,t) \in \overline{\Delta}_{ij}^T} \tilde{\eta}(s,t) > u(1 + b(jT_1 u^{-2/\alpha})^\beta) \right) (1 + o(1)) \end{aligned}$$

as $u \rightarrow \infty$, where $\{\tilde{\eta}(s,t), (s,t) \in [0, S] \times [0, T]\}$ is a centered homogeneous Gaussian random field with covariance function

$$r_{\tilde{\eta}}(s,t) = \exp(-|\tilde{a}_1 s|^\alpha - |\tilde{a}_2 t|^\alpha), \quad (s,t) \in [0, S] \times [0, T],$$

with $\tilde{a}_1 = (a_1^\alpha + 2|a_3|^\alpha)^{1/\alpha}$ and $\tilde{a}_2 = 2^{1/\alpha} a_2$. It follows further, using Lemma 2.1 that

$$\begin{aligned} & \mathbb{P} \left(\sup_{(s,t) \in \overline{\Delta}_{ij}^T} X(s,t) > u \right) \\ & \leq \mathbb{P} \left(\sup_{(s,t) \in \overline{\Delta}_{ij}^T} \tilde{\eta}(s,t) > u(1 + b(jT_1 u^{-2/\alpha})^\beta) \right) (1 + o(1)) \\ & = \mathcal{H}_{Y_2}^0 [S_1, T_1] \frac{1}{\sqrt{2\pi}u} \exp \left(-\frac{u^2(1 + 2b(jT_1 u^{-2/\alpha})^\beta)}{2} \right) (1 + o(1)) \\ & = \mathcal{H}_{Y_2}^0 [S_1, T_1] \exp(-b(jT_1)^\beta) \Psi(u) (1 + o(1)) \end{aligned}$$

as $u \rightarrow \infty$, where $\mathcal{H}_{Y_2}^0 [S_1, T_1]$ is defined in a similar way as $\mathcal{H}_{Y_2}^0 [S_1, T_1]$ with $a_i, i = 1, 2$ replaced by $\tilde{a}_i, i = 1, 2$. Consequently, as $u \rightarrow \infty$,

$$(42) \quad \Sigma_2(u) \leq \sum_{j=1}^{\infty} \frac{S}{S_1} u^{2/\alpha} \mathcal{H}_{Y_2}^0 [S_1, T_1] \exp(-b(jT_1)^\beta) \Psi(u) (1 + o(1)).$$

From (4), there exists some $\theta_1 \in (0, 1)$ such that

$$\sup_{\substack{1 \leq i < i' \leq N_1(u) \\ \rho(\widehat{\Delta}_i, \widehat{\Delta}_{i'}) > \rho_3}} \sup_{\substack{s \in \widehat{\Delta}_i, s' \in \widehat{\Delta}_{i'} \\ t, t' \in [0, T]}} r(s, t, s', t') < \theta_1,$$

where ρ_3 is the same as in (34). Below we shall re-write $\Sigma_3(u)$ as

$$\Sigma_3(u) = \Sigma_{3,1}(u) + \Sigma_{3,2}(u) + \Sigma_{3,3}(u), \quad u \geq 0,$$

where $\Sigma_{3,1}(u)$ is the sum taken on $\rho(\widehat{\Delta}_i, \widehat{\Delta}_{i'}) > \rho_3$, $\Sigma_{3,2}(u)$ is the sum taken on $i' = i + 1$, and $\Sigma_{3,3}(u)$ is the sum taken on $i' > i + 1$ and $\rho(\widehat{\Delta}_i, \widehat{\Delta}_{i'}) \leq \rho_3$. First, note that the estimation of $\Sigma_{3,1}(u)$ can be derived similarly to that of $\Sigma_{1,1}(u)$ in case (a), and thus for u sufficiently large

$$(43) \quad \Sigma_{3,1}(u) \leq \frac{S^2}{S_1^2} u^{4/\alpha} \exp\left(-\frac{(u-a)^2}{2-(1-\theta_1)}\right),$$

where a is the same as in (36). Next, we consider $\Sigma_{3,3}(u)$. In view of (34) and (35), it follows that for $s \in \widehat{\Delta}_i, s' \in \widehat{\Delta}_{i'}, t, t' \in T - \widehat{\Delta}_0$ and u large enough

$$(44) \quad 2 \leq \mathbb{E}(\xi^2(s, t, s', t')) \leq 4 - |a_1(i' - i)S_1|^\alpha u^{-2}.$$

Further set $\bar{\xi}(s, t, s', t') = \xi(s, t, s', t') / \sqrt{\text{Var}(\xi(s, t, s', t'))}$. Following similar argument as in the proof of Lemma 6.3 in [24], we obtain that

$$\begin{aligned} & \mathbb{E}(\bar{\xi}(s, t, s', t') - \bar{\xi}(v, w, v', w'))^2 \\ & \leq 4(\mathbb{E}(\bar{X}(s, t) - \bar{X}(v, w))^2 + \mathbb{E}(\bar{X}(s', t') - \bar{X}(v', w'))^2). \end{aligned}$$

Moreover, from (34) we see that, for u sufficiently large

$$\mathbb{E}(\bar{X}(s, t) - \bar{X}(v, w))^2 \leq 3(|\tilde{a}_1(s-v)|^\alpha + |\tilde{a}_2(t-w)|^\alpha)$$

implying thus

$$(45) \quad \begin{aligned} & \mathbb{E}(\bar{\xi}(s, t, s', t') - \bar{\xi}(v, w, v', w'))^2 \\ & \leq 2(1 - r_\zeta(s-v, t-w, s'-v', t'-w')), \end{aligned}$$

where

$$r_\zeta(s, t, s', t') = \exp(-7(|\tilde{a}_1 s|^\alpha + |\tilde{a}_2 t|^\alpha + |\tilde{a}_1 s'|^\alpha + |\tilde{a}_2 t'|^\alpha))$$

is the covariance function of the homogeneous Gaussian random field $\{\zeta(s, t, s', t'), (s, t, s', t') \in (0, \infty)^4\}$. Consequently, (44), (45) and Slepian's lemma imply

$$\begin{aligned} & \mathbb{P}\left(\sup_{(s,t) \in \overline{\Delta}_{i0}^T} X(s, t) > u, \sup_{(s,t) \in \overline{\Delta}_{i'0}^T} X(s, t) > u\right) \\ & \leq \mathbb{P}\left(\sup_{\substack{(s,t) \in \overline{\Delta}_{i0}^T \\ (s',t') \in \overline{\Delta}_{i'0}^T}} \zeta(s, t, s', t') > \frac{2u}{\sqrt{4 - |a_1(i' - i)S_1|^\alpha u^{-2}}}\right). \end{aligned}$$

We obtain further from a similar lemma as Lemma 2.1 (cf. Lemma 6.1 in [24]) that

$$\begin{aligned} & \mathbb{P}\left(\sup_{\substack{(s,t) \in \bar{\Delta}_{i0}^T \\ (s',t') \in \bar{\Delta}_{i'0}^T}} \zeta(s,t,s',t') > \frac{2u}{\sqrt{4 - |a_1(i' - i)S_1|^{\alpha u^{-2}}}}\right) \\ &= (\tilde{\mathcal{H}}_{\tilde{Y}_2}^0[S_1, T_1])^2 \frac{1}{\sqrt{2\pi}u} \exp\left(-\frac{4u^2}{2(4 - |a_1(i' - i)S_1|^{\alpha u^{-2}})}\right) (1 + o(1)), \end{aligned}$$

where $\tilde{\mathcal{H}}_{\tilde{Y}_2}^0[S_1, T_1]$ is defined in a similar way as $\mathcal{H}_{Y_2}^0[S_1, T_1]$ with a_1, a_2 replaced by $7^{1/\alpha}\tilde{a}_1, 7^{1/\alpha}\tilde{a}_2$, respectively. Consequently, for all large u ,

$$(46) \quad \Sigma_{3,3}(u) \leq \frac{S}{S_1} \sum_{j \geq 1} (\tilde{\mathcal{H}}_{\tilde{Y}_2}^0[S_1, T_1])^2 \exp\left(-\frac{1}{8}|a_1 j S_1|^\alpha\right) u^{2/\alpha} \Psi(u) (1 + o(1)).$$

Next, we consider $\Sigma_{3,2}(u)$. For any u positive,

$$\begin{aligned} & \mathbb{P}\left(\sup_{(s,t) \in \bar{\Delta}_{i0}^T} X(s,t) > u, \quad \sup_{(s,t) \in \bar{\Delta}_{(i+1)0}^T} X(s,t) > u\right) \\ & \leq \mathbb{P}\left(\sup_{(s,t) \in \bar{\Delta}_{i0}^T} X(s,t) > u, \right. \\ & \quad \left. \sup_{(s,t) \in [(i+1)S_1 u^{-2/\alpha}, (i+1)S_1 u^{-2/\alpha} + \sqrt{S_1} u^{-2/\alpha}] \times (T - \tilde{\Delta}_0)} X(s,t) > u\right) \\ & + \mathbb{P}\left(\sup_{(s,t) \in \bar{\Delta}_{i0}^T} X(s,t) > u, \right. \\ & \quad \left. \sup_{(s,t) \in [(i+1)S_1 u^{-2/\alpha} + \sqrt{S_1} u^{-2/\alpha}, (i+2)S_1 u^{-2/\alpha}] \times (T - \tilde{\Delta}_0)} X(s,t) > u\right) \end{aligned}$$

and further

$$\begin{aligned} & \mathbb{P}\left(\sup_{(s,t) \in \bar{\Delta}_{i0}^T} X(s,t) > u, \quad \sup_{(s,t) \in \bar{\Delta}_{(i+1)0}^T} X(s,t) > u\right) \\ & \leq \mathcal{H}_{\tilde{Y}_2}^0[\sqrt{S_1}, T_1] \Psi(u) (1 + o(1)) \\ & \quad + (\tilde{\mathcal{H}}_{\tilde{Y}_2}^0[\sqrt{S_1}, T_1])^2 \exp\left(-\frac{1}{8}|a_1 \sqrt{S_1}|^\alpha\right) \Psi(u) (1 + o(1)). \end{aligned}$$

Therefore, for all large u

$$\Sigma_{3,2}(u) \leq \frac{S}{S_1} \left(\mathcal{H}_{\tilde{Y}_2}^0[\sqrt{S_1}, T_1] \right)$$

$$(47) \quad + (\tilde{\mathcal{H}}_{Y_2}^0[\sqrt{S_1}, T_1])^2 \exp\left(-\frac{1}{8}|a_1\sqrt{S_1}|^\alpha\right) u^{2/\alpha} \Psi(u)(1+o(1)).$$

Consequently, from (39)–(43) and (46)–(47), we conclude that for any $S_i, T_i, i = 1, 2$

$$\begin{aligned} & S_1^{-1} \mathcal{H}_{Y_1}^b[S_1, T_1] + \sum_{j=1}^{\infty} S_1^{-1} \mathcal{H}_{Y_2}^0[S_1, T_1] \exp(-b(jT_1)^\beta) \\ & \geq \limsup_{u \rightarrow \infty} \frac{\pi(u)}{Su^{\alpha/2} \Psi(u)} \geq \liminf_{u \rightarrow \infty} \frac{\pi(u)}{Su^{\alpha/2} \Psi(u)} \\ & \geq S_2^{-1} \mathcal{H}_{Y_1}^b[S_2, T_2] - S_2^{-1} (\tilde{\mathcal{H}}_{Y_2}^0[S_2, T_2])^2 \sum_{j \geq 1} \exp\left(-\frac{1}{8}|a_1 j S_2|^\alpha\right) \\ & \quad - S_2^{-1} \left(\mathcal{H}_{Y_2}^0[\sqrt{S_2}, T_2] + (\tilde{\mathcal{H}}_{Y_2}^0[\sqrt{S_2}, T_2])^2 \exp\left(-\frac{1}{8}|a_1 \sqrt{S_2}|^\alpha\right) \right). \end{aligned}$$

Therefore, by similar arguments as in the proof of Theorem D.2 in [24], we conclude that

$$0 < \mathcal{M}_{Y_1, \alpha_1}^b \leq \limsup_{u \rightarrow \infty} \frac{\pi(u)}{Su^{\alpha/2} \Psi(u)} \leq \liminf_{u \rightarrow \infty} \frac{\pi(u)}{Su^{\alpha/2} \Psi(u)} \leq \mathcal{M}_{Y_1, \alpha_1}^b < \infty$$

establishing the claim.

Case (iii) $\beta = \alpha_2 > \alpha_1$: Note that $\mathcal{M}_{Y_2, \beta}^b$ can be given in terms of Piterbarg and Pickands constants as

$$\mathcal{M}_{Y_2, \beta}^b = \lim_{T \rightarrow \infty} \lim_{S \rightarrow \infty} \frac{1}{S} \mathcal{H}_{Y_2}^b[S, T] = a_1 a_2 \mathcal{P}_{\alpha_2}^{ba_2^{-\alpha_2}} \mathcal{H}_{\alpha_1}.$$

The proof for this case can be established using step-by-step the same arguments as in case (ii).

Case (iv) $\beta < \alpha_2 = \alpha_1$: In order to make use of the notation introduced in case (ii) we set $\alpha := \alpha_1 = \alpha_2$. First, note that $\delta(u) < T_1 u^{-2/\alpha}$, which implies

$$\begin{aligned} \pi(u) & \leq \mathbb{P}\left(\sup_{(s,t) \in [0, S] \times (T - \tilde{\Delta}_0)} X(s, t) > u\right) \\ & \leq \sum_{i=0}^{N_1(u)} \mathbb{P}\left(\sup_{(s,t) \in \tilde{\Delta}_{i0}^T} X(s, t) > u\right) \\ & \leq \frac{S}{S_1} u^{2/\alpha} \mathcal{H}_{Y_1}^0[S_1, T_1] \Psi(u)(1+o(1)) \end{aligned}$$

as $u \rightarrow \infty$. Further, by Assumptions A1 and A2 we have that $\mathbb{E}((X(s, T))^2) = 1, \forall s \in [0, S]$ and

$$r(s, T, s', T) = 1 - (a_1^\alpha + |a_3|^\alpha) |s - s'|^\alpha (1 + o(1))$$

holds uniformly with respect to $s, s' \in [0, S]$, as $|s - s'| \rightarrow 0$. This means that $\{X(s, T), s \in [0, S]\}$ is a locally stationary Gaussian process. Therefore, in view of Theorem 7.1 in [24],

$$\begin{aligned} \pi(u) &\geq \mathbb{P}\left(\sup_{s \in [0, S]} X(s, T) > u\right) \\ &= S(a_1^\alpha + |a_3|^\alpha)^{1/\alpha} \mathcal{H}_\alpha u^{2/\alpha} \Psi(u)(1 + o(1)), \quad u \rightarrow \infty. \end{aligned}$$

Letting $T_1 \rightarrow 0, S_1 \rightarrow \infty$, we conclude that

$$0 < \lim_{u \rightarrow \infty} \frac{\pi(u)}{S u^{2/\alpha} \Psi(u)} = (a_1^\alpha + |a_3|^\alpha)^{1/\alpha} \mathcal{H}_\alpha < \infty.$$

Case (v) $\beta < \alpha_2$ and $\alpha_1 < \alpha_2$: The claim follows with identical arguments as in the proof of case (iv).

In order to complete the proof of cases (i)–(v) we only need to show (30), for which it is sufficient to give the following upper bounds for $\pi_1(u)$ and $\pi_2(u)$. By Borell–TIS inequality, for u large enough

$$(48) \quad \pi_1(u) \leq \exp\left(-\frac{(u - \mathbb{E}(\sup_{(s,t) \in [0, S] \times [0, \rho_0]} X(s, t)))^2}{2\theta^2}\right).$$

Further, by Assumption A3 applying the Piterbarg inequality we obtain, as $u \rightarrow \infty$

$$(49) \quad \begin{aligned} \pi_2(u) &\leq \mathcal{Q} u^{4/\gamma-1} \exp\left(-\frac{u^2}{2\sigma^2(T - \delta(u))}\right) \\ &= \mathcal{Q} u^{4/\gamma-1} \exp\left(-\frac{u^2}{2}\right) \exp(-b(\ln u)^2)(1 + o(1)), \end{aligned}$$

where \mathcal{Q} is some positive constant not depending on u . Therefore, the proof of cases (i)–(v) is complete.

Next, we consider cases (vi)–(vii). We introduce a time scaling of the Gaussian random field $\{X(s, t), (s, t) \in \mathbf{E}\}$ by matrix $B = \begin{pmatrix} a_3 & a_2 \\ 0 & a_2 \end{pmatrix}$, that is, let $Z(s, t) := X((s - t)/a_3, t/a_2)$. By this time scaling, we have

$$(50) \quad \mathbb{P}\left(\sup_{(s,t) \in \mathbf{E}} X(s, t) > u\right) = \mathbb{P}\left(\sup_{(s,t) \in \mathbf{K}} Z(s, t) > u\right),$$

where \mathbf{K} is a region on \mathbb{R}^2 with vertices at points $(0, 0)$, $(a_2 T, a_2 T)$, $(a_3 S, 0)$ and $(a_3 S + a_2 T, a_2 T)$. The Gaussian random field $\{Z(s, t), (s, t) \in \mathbf{K}\}$ has the following properties:

(P1) The standard deviation function $\sigma_Z(s, t)$ of $\{Z(s, t), (s, t) \in \mathbf{K}\}$ satisfies

$$\sigma_Z(s, t) = 1 - \frac{b}{a_2^\beta} (a_2 T - t)^\beta (1 + o(1)), \quad t \uparrow a_2 T.$$

(P2) The correlation function $r_Z(s, t, s', t')$ of $\{Z(s, t), (s, t) \in \mathbf{K}\}$ satisfies

$$r_Z(s, t, s', t') = 1 - \left(|s - s'|^{\alpha_2} + \left| \frac{a_1}{a_3}(t - t') - \frac{a_1}{a_3}(s - s') \right|^{\alpha_1} \right) (1 + o(1))$$

for any $(s, t), (s', t') \in \mathbf{K}$ such that $|s - s'| \rightarrow 0$ and $t, t' \uparrow a_2 T$, and further there exists some $\delta_0 \in (0, T)$ such that

$$r(s, t, s', t') < 1$$

holds for any $(s, t), (s', t') \in \mathbf{K}_0$ satisfying $s \neq s'$. Here, \mathbf{K}_0 is a region on \mathbb{R}^2 with vertices at points $(a_2 \delta_0, a_2 \delta_0)$, $(a_2 T, a_2 T)$, $(a_3 S + a_2 \delta_0, a_2 \delta_0)$ and $(a_3 S + a_2 T, a_2 T)$.

(P3) There exist positive constants $\mathcal{Q}, \gamma, \rho_1$ and ρ_2 such that

$$\mathbb{E}((Z(s, t) - Z(s', t'))^2) \leq \mathcal{Q}(|s - s'|^\gamma + |t - t'|^\gamma)$$

holds for any $(s, t), (s', t') \in \mathbf{K}$ satisfying $a_2 T - t < \rho_1$, $a_2 T - t' < \rho_1$ and $|s - s'| < \rho_2$.

Note that in the above proof the most important structural property of the set \mathbf{E} is that the segment $\mathbf{L} = \{(s, t) \in \mathbf{E} : t = T\}$ is on the boundary of \mathbf{E} , which is also the case for $\{Z(s, t), (s, t) \in \mathbf{K}\}$. Therefore, in view of the above properties of $\{Z(s, t), (s, t) \in \mathbf{K}\}$, the claims of the cases (vi) and (vii) follow by an application of the claims of cases (iii) and (v). The proof is complete. \square

PROOF OF PROPOSITION 3.1. The variance function of Z is given by

$$\sigma_Z^2(s, t) = 2(1 - r_X(t))$$

and attains its maximum on $[0, S] \times \{T\}$. Therefore, it is sufficient to consider the asymptotics of

$$\Pi(u) := \mathbb{P}\left(\sup_{(s, t) \in [0, S] \times [0, T]} Z^*(s, t) > \tilde{u}\right), \quad u \rightarrow \infty,$$

with

$$\tilde{u} := \frac{u}{\rho_T} \quad \text{and} \quad Z^*(s, t) := \frac{Z(s, t)}{\rho_T},$$

where $\rho_T = \sqrt{2(1 - r_X(T))} > 0$. The asymptotics of $\Pi(u)$ follows from Theorem 2.2 by checking the Assumptions A1–A3. The standard deviation function of Z^* satisfies

$$\begin{aligned} \sigma_{Z^*}(s, t) &= \frac{\sqrt{2(1 - r_X(t))}}{\rho_T} \\ &= 1 - \frac{a_1}{2(1 - r_X(T))} (T - t)^{\alpha_1} (1 + o(1)), \quad t \rightarrow T, \end{aligned}$$

whereas for its correlation function we have

$$(51) \quad R_{Z^*}(s, t, s', t') = \frac{r_X(|s+t-s'-t'|) - r_X(|s-s'-t'|)}{2\sqrt{(1-r_X(t))(1-r_X(t'))}} + \frac{-r_X(|s+t-s'|) + r_X(|s-s'|)}{2\sqrt{(1-r_X(t))(1-r_X(t'))}}.$$

Since $r_X(t)$ is twice continuously differentiable in $[\mu, T]$ and $|r_X''(T)| \in (0, \infty)$ for some constant \mathcal{Q}_1 , we have

$$\begin{aligned} & |r_X(t') - r_X(|s-s'-t'|) + r_X(t) - r_X(|s+t-s'|)| \\ & \leq \mathcal{Q}_1(|t-t'+s-s'|^2 + |s-s'|^2)(1+o(1)) \end{aligned}$$

as $t, t' \rightarrow T, |s-s'| \rightarrow 0$. Consequently, $\alpha_2 \in (0, 2)$ implies

$$(52) \quad \begin{aligned} & R_{Z^*}(s, t, s', t') \\ & = 1 - \frac{a_2}{\rho_T^2}(|t-t'+s-s'|^{\alpha_2} + |s-s'|^{\alpha_2})(1+o(1)) \end{aligned}$$

as $t, t' \rightarrow T, |s-s'| \rightarrow 0$. Next, for any fixed $\varepsilon_0 > 0$, we have from S3 that there exists some θ_0 such that

$$r_X(|s-s'|) \leq \theta_0 < 1$$

for any $s, s' \in [0, S]$ satisfying $|s-s'| > \varepsilon_0$. Further, from S2 we obtain that there exists some positive constant δ_0 such that

$$2\sqrt{(1-r_X(t))(1-r_X(t'))} \geq \rho_T^2 - \frac{1-\theta_0}{2} > 0$$

for any $t, t' \in [\delta_0, T]$. Hence,

$$(53) \quad R_{Z^*}(s, t, s', t') \leq \frac{1+\theta_0-2r_X(T)}{\rho_T^2 - (1-\theta_0)/2} < 1$$

for any $t, t' \in [\delta_0, T], s, s' \in [0, S]$ satisfying $|s-s'| > \varepsilon_0$, and thus both A1 and A2 are satisfied. It follows that

$$\begin{aligned} \mathbb{E}(Z^*(s, t) - Z^*(s', t'))^2 & \leq 2\mathbb{E}(\bar{Z}(s, t) - \bar{Z}(s', t'))^2 \\ & \quad + \frac{2}{\rho_T^2}(\sigma_Z(s, t) - \sigma_Z(s', t'))^2. \end{aligned}$$

Therefore, the differentiability of $r_X(t)$, assumption S2 and (52) imply that there exist some positive constants $\rho_1, \rho_2, \mathcal{Q}_3, \mathcal{Q}_4$ such that

$$\begin{aligned} & \mathbb{E}(Z^*(s, t) - Z^*(s', t'))^2 \\ & \leq \mathcal{Q}_3(|t-t'+s-s'|^{\alpha_2} + |s-s'|^{\alpha_2} + |t-t'|^{2\min(\alpha_1, 1)}) \\ & \leq \mathcal{Q}_4(|t-t'|^{\min(2\alpha_1, \alpha_2)} + |s-s'|^{\min(2\alpha_1, \alpha_2)}) \end{aligned}$$

for all $s, s' \in [0, S], t, t' \in [\rho_1, T]$ satisfying $|s - s'| < \rho_2$, hence the proof is complete. \square

PROOFS OF PROPOSITIONS 3.4 AND 3.5. Note first that the standard deviation of the incremental random field Z of the Brownian bridge satisfies

$$(54) \quad \sigma_Z(s, t) = (t(1-t))^{1/2} = \frac{1}{2} - (t - \frac{1}{2})^2(1 + o(1)), \quad t \rightarrow \frac{1}{2}.$$

Furthermore, for its correlation function we have

$$(55) \quad r_Z(s, s', t, t') = 1 - 2(|t - t' + s - s'| + |s - s'|)(1 + o(1))$$

as $t, t' \rightarrow 1/2, |s - s'| \rightarrow 0$.

For the fBm incremental random field Z , we have for its standard deviation

$$\sigma_Z(s, t) = t^{\alpha/2} = 1 - \frac{\alpha}{2}(1-t)(1 + o(1)), \quad t \rightarrow 1.$$

As shown in [25], the correlation function r_Z of Z satisfies

$$r_Z(s, s', t, t') = 1 - \frac{1}{2}(|t - t' + s - s'|^\alpha + |s - s'|^\alpha)(1 + o(1))$$

as $t, t' \rightarrow 1, |s - s'| \rightarrow 0$. Hence, for both cases A1–A3 are fulfilled, and thus the claims follow by a direct application of Theorem 2.2. \square

PROOFS OF PROPOSITIONS 4.1 AND 4.2. By a linear time change using the matrix $A \in \mathbb{R}^{2 \times 2}$ given by

$$A = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$$

we have for any $u > 0$

$$\mathbb{P}(\chi_2(\xi) > u) = \mathbb{P}\left(\sup_{(s,t) \in A[0,1]^2} (\xi(t+s) - \xi(s)) > u\right).$$

Here, the set $A[0, 1]^2 = \{(\tilde{s}, \tilde{t}) : (\tilde{s}, \tilde{t})^\top = A(s, t)^\top, (s, t) \in [0, 1]^2\}$ is bounded and convex. The variance function of the random field $\{\xi(t+s) - \xi(s), (s, t) \in A[0, 1]^2\}$ is $2(1 - r_\xi(|t|))$ which attains its unique maximum on the set $A[0, 1]^2$ on two lines $\mathbf{L}_1 = \{(s, t) \in A[0, 1]^2 : t = t_m\}$ and $\mathbf{L}_2 = \{(s, t) \in A[0, 1]^2 : t = -t_m\}$. Note that the differentiability of $r_\xi(t)$ implies $\alpha_1 \geq 2 > \alpha_2$. Therefore, the claim in (27) follows from Remark 2.4(b); the conditions therein can be established directly as in the proof of Proposition 3.1 except (18) for $i = 1, j = 2$, which can also be confirmed by a similar argument as in (53). Further, since

$$\mathbb{P}(\chi_2(X) > u) = \mathbb{P}\left(\sup_{(s,t) \in A[0,1]^2} (X(t+s) - X(s)) > u\right)$$

in view of (54) and (55) we conclude that the claim in (28) follows immediately from Remark 2.4(b), and thus the proof is complete. \square

APPENDIX

Let \mathbf{D} be a compact set in \mathbb{R}^2 such that $(0, 0) \in \mathbf{D}$, and let $\{\xi_u(s, t), (s, t) \in \mathbf{D}\}$, $u > 0$ be a family of centered Gaussian random fields with a.s. continuous sample paths. The next lemma is proved based on the classical approach rooted in the ideas of [21, 22] (see also [9]), Lemma 1; in particular, it implies the claim of Lemma 2.1.

LEMMA A.1. *Let $d(\cdot)$ be a nonnegative continuous function on $[0, \infty)$ and let $g(u), u > 0$ be a positive function satisfying $\lim_{u \rightarrow \infty} g(u)/u = 1$. Assume that the variance function $\sigma_{\xi_u}^2$ of ξ_u satisfies the following conditions:*

$$\sigma_{\xi_u}(0, 0) = 1 \quad \text{for all large } u, \quad \lim_{u \rightarrow \infty} \sup_{(s, t) \in \mathbf{D}} |u^2(1 - \sigma_{\xi_u}(s, t)) - d(t)| = 0,$$

and there exist some positive constants G, ν, u_0 such that, for all $u > u_0$

$$u^2 \text{Var}(\xi_u(s, t) - \xi_u(s', t')) \leq G(|s - s'|^\nu + |t - t'|^\nu)$$

holds uniformly with respect to $(s, t), (s', t') \in \mathbf{D}$. If further there exists a centered Gaussian random field $\{Y(s, t), (s, t) \in (0, \infty)^2\}$ with a.s. continuous sample paths and $Y(0, 0) = 0$ such that

$$\lim_{u \rightarrow \infty} u^2 \text{Var}(\xi_u(s, t) - \xi_u(s', t')) = 2 \text{Var}(Y(s, t) - Y(s', t'))$$

holds for all $(s, t), (s', t') \in \mathbf{D}$, then

$$(56) \quad \mathbb{P}\left(\sup_{(s, t) \in \mathbf{D}} \xi_u(s, t) > g(u)\right) = \mathcal{H}_Y^d[\mathbf{D}] \Psi(g(u))(1 + o(1))$$

as $u \rightarrow \infty$, where

$$\mathcal{H}_Y^d[\mathbf{D}] = \mathbb{E}\left(\exp\left(\sup_{(s, t) \in \mathbf{D}} (\sqrt{2}Y(s, t) - \sigma_Y^2(s, t) - d(t))\right)\right).$$

PROOF. For large u , we have

$$(57) \quad \begin{aligned} & \mathbb{P}\left(\sup_{(s, t) \in \mathbf{D}} \xi_u(s, t) > g(u)\right) \\ &= \frac{1}{\sqrt{2\pi}g(u)} \exp\left(-\frac{(g(u))^2}{2}\right) \int_{-\infty}^{\infty} e^{w-w^2/(2(g(u))^2)} \\ & \quad \times \mathbb{P}\left(\sup_{(s, t) \in \mathbf{D}} \xi_u(s, t) > g(u) \mid \xi_u(0, 0) = g(u) - \frac{w}{g(u)}\right) dw. \end{aligned}$$

Let

$$R_{\xi_u}(s, t, s', t') = \mathbb{E}(\xi_u(s, t)\xi_u(s', t')), \quad (s, t), (s', t') \in \mathbf{D}$$

be the covariance function of ξ_u . The conditional random field

$$\left\{ \xi_u(s, t) \mid \xi_u(0, 0) = g(u) - \frac{w}{g(u)}, (s, t) \in \mathbf{D} \right\}$$

has the same distribution as

$$\left\{ \xi_u(s, t) - R_{\xi_u}(s, t, 0, 0)\xi_u(0, 0) + R_{\xi_u}(s, t, 0, 0)\left(g(u) - \frac{w}{g(u)}\right), (s, t) \in \mathbf{D} \right\}.$$

Thus, the integrand in (57) can be rewritten as

$$\begin{aligned} & \mathbb{P}\left(\sup_{(s, t) \in \mathbf{D}} \left(\xi_u(s, t) - R_{\xi_u}(s, t, 0, 0)\xi_u(0, 0) \right. \right. \\ & \quad \left. \left. + R_{\xi_u}(s, t, 0, 0)\left(g(u) - \frac{w}{g(u)}\right) \right) > g(u) \right) \\ &= \mathbb{P}\left(\sup_{(s, t) \in \mathbf{D}} (\chi_u(s, t) - (g(u))^2(1 - R_{\xi_u}(s, t, 0, 0)) \right. \\ & \quad \left. + w(1 - R_{\xi_u}(s, t, 0, 0))) > w \right), \end{aligned}$$

where

$$\chi_u(s, t) = g(u)(\xi_u(s, t) - R_{\xi_u}(s, t, 0, 0)\xi_u(0, 0)).$$

Next, the following convergence

$$(g(u))^2(1 - R_{\xi_u}(s, t, 0, 0)) - w(1 - R_{\xi_u}(s, t, 0, 0)) \rightarrow \sigma_Y^2(s, t) + d(t)$$

holds as $u \rightarrow \infty$, for any $w \in \mathbb{R}$, uniformly with respect to $(s, t) \in \mathbf{D}$. Moreover,

$$\begin{aligned} & \mathbb{E}((\chi_u(s, t) - \chi_u(s', t'))^2) \\ &= (g(u))^2(\mathbb{E}((\xi_u(s, t) - \xi_u(s', t'))^2) - (R_{\xi_u}(s, t, 0, 0) - R_{\xi_u}(s', t', 0, 0))^2) \\ &\rightarrow 2\text{Var}(Y(s, t) - Y(s', t')), \quad u \rightarrow \infty \end{aligned}$$

holds for any $(s, t), (s', t') \in \mathbf{D}$. Hence, the claim follows by using the same arguments as in the proof of Lemma 6.1 in [24] or those in the proof of Lemma 1 in [9]. \square

Acknowledgements. We are thankful to the Editor, the Associate Editor and the referees for their comments and suggestions.

REFERENCES

- [1] ADLER, R. J. and TAYLOR, J. E. (2007). *Random Fields and Geometry*. Springer, New York. [MR2319516](#)
- [2] ALBIN, J. M. P. (1990). On extremal theory for stationary processes. *Ann. Probab.* **18** 92–128. [MR1043939](#)
- [3] ALBIN, J. M. P. and CHOI, H. (2010). A new proof of an old result by Pickands. *Electron. Commun. Probab.* **15** 339–345. [MR2685014](#)
- [4] AZAÏS, J.-M. and WSCHEBOR, M. (2009). *Level Sets and Extrema of Random Processes and Fields*. Wiley, Hoboken, NJ. [MR2478201](#)
- [5] BERMAN, S. M. (1982). Sojourns and extremes of stationary processes. *Ann. Probab.* **10** 1–46. [MR0637375](#)
- [6] BERMAN, S. M. (1992). *Sojourns and Extremes of Stochastic Processes*. Wadsworth & Brooks/Cole Advanced Books & Software, Pacific Grove, CA. [MR1126464](#)
- [7] CHAN, H. P. and LAI, T. L. (2006). Maxima of asymptotically Gaussian random fields and moderate deviation approximations to boundary crossing probabilities of sums of random variables with multidimensional indices. *Ann. Probab.* **34** 80–121. [MR2206343](#)
- [8] CRESSIE, N. (1980). The asymptotic distribution of the scan statistic under uniformity. *Ann. Probab.* **8** 828–840. [MR0577319](#)
- [9] DĘBICKI, K. and KOSIŃSKI, K. M. (2014). On the infimum attained by the reflected fractional Brownian motion. *Extremes* **17** 431–446. [MR3252820](#)
- [10] DĘBICKI, K. and MANDJES, M. (2003). Exact overflow asymptotics for queues with many Gaussian inputs. *J. Appl. Probab.* **40** 704–720. [MR1993262](#)
- [11] DEHEUVELS, P. and DEVROYE, L. (1987). Limit laws of Erdős–Rényi–Shepp type. *Ann. Probab.* **15** 1363–1386. [MR0905337](#)
- [12] DIEKER, A. B. (2005). Extremes of Gaussian processes over an infinite horizon. *Stochastic Process. Appl.* **115** 207–248. [MR2111193](#)
- [13] DIEKER, A. B. and YAKIR, B. (2014). On asymptotic constants in the theory of extremes for Gaussian processes. *Bernoulli* **20** 1600–1619. [MR3217455](#)
- [14] DÜMBGEN, L. and SPOKOINY, V. G. (2001). Multiscale testing of qualitative hypotheses. *Ann. Statist.* **29** 124–152. [MR1833961](#)
- [15] HARPER, A. J. (2013). Bounds on the suprema of Gaussian processes, and omega results for the sum of a random multiplicative function. *Ann. Appl. Probab.* **23** 584–616. [MR3059269](#)
- [16] HASHORVA, E., JI, L. and PITERBARG, V. I. (2013). On the supremum of γ -reflected processes with fractional Brownian motion as input. *Stochastic Process. Appl.* **123** 4111–4127. [MR3091101](#)
- [17] HASHORVA, E. and TAN, Z. (2013). Large deviations of Shepp statistics for fractional Brownian motion. *Statist. Probab. Lett.* **83** 2242–2247. [MR3093809](#)
- [18] KABLUCHKO, Z. (2011). Extremes of the standardized Gaussian noise. *Stochastic Process. Appl.* **121** 515–533. [MR2763094](#)
- [19] KABLUCHKO, Z. and WANG, Y. (2014). Limiting distribution for the maximal standardized increment of a random walk. *Stochastic Process. Appl.* **124** 2824–2867. [MR3217426](#)
- [20] MIKHALEVA, T. L. and PITERBARG, V. I. (1996). On the distribution of the maximum of a Gaussian field with constant variance on a smooth manifold. *Teor. Veroyatn. Primen.* **41** 438–451. [MR1445766](#)
- [21] PICKANDS, J. III (1969). Asymptotic properties of the maximum in a stationary Gaussian process. *Trans. Amer. Math. Soc.* **145** 75–86. [MR0250368](#)

- [22] PICKANDS, J. III (1969). Upcrossing probabilities for stationary Gaussian processes. *Trans. Amer. Math. Soc.* **145** 51–73. [MR0250367](#)
- [23] PITERBARG, V. I. (1972). On the paper by J. Pickands “Upcrossing probabilities for stationary Gaussian processes”. *Vestnik Moskov. Univ. Ser. I Mat. Meh.* **27** 25–30. [MR0334321](#)
- [24] PITERBARG, V. I. (1996). *Asymptotic Methods in the Theory of Gaussian Processes and Fields. Translations of Mathematical Monographs* **148**. Amer. Math. Soc., Providence, RI. [MR1361884](#)
- [25] PITERBARG, V. I. (2001). Large deviations of a storage process with fractional Brownian motion as input. *Extremes* **4** 147–164. [MR1893870](#)
- [26] PITERBARG, V. I. and PRISJAŽNJUK, V. P. (1981). Exact asymptotic behavior of the probability of large span of a Gaussian stationary process. *Teor. Veroyatn. Primen.* **26** 480–495. [MR0627856](#)
- [27] SHEPP, L. A. (1966). Radon–Nikodým derivatives of Gaussian measures. *Ann. Math. Statist.* **37** 321–354. [MR0190999](#)
- [28] SHEPP, L. A. (1971). First passage time for a particular Gaussian process. *Ann. Math. Statist.* **42** 946–951. [MR0278375](#)
- [29] SIEGMUND, D. and VENKATRAMAN, E. S. (1995). Using the generalized likelihood ratio statistic for sequential detection of a change-point. *Ann. Statist.* **23** 255–271. [MR1331667](#)
- [30] VARDAR-ACAR, C. and CAKAR, H. (2013). Bounds on the expected value of maximum loss of fractional Brownian motion. Preprint.
- [31] VARDAR-ACAR, C., ZIRBEL, C. L. and SZÉKELY, G. J. (2013). On the correlation of the supremum and the infimum and of maximum gain and maximum loss of Brownian motion with drift. *J. Comput. Appl. Math.* **248** 61–75. [MR3029217](#)
- [32] ZHOLUD, D. (2008). Extremes of Shepp statistics for the Wiener process. *Extremes* **11** 339–351. [MR2453344](#)

K. DĘBICKI
MATHEMATICAL INSTITUTE
UNIVERSITY OF WROCLAW
PL. GRUNWALDZKI 2/4, 50-384 WROCLAW
POLAND
E-MAIL: Krzysztof.Debicki@math.uni.wroc.pl

E. HASHORVA
L. JI
DEPARTMENT OF ACTUARIAL SCIENCE
UNIVERSITY OF LAUSANNE
UNIL-DORIGNY, 1015 LAUSANNE
SWITZERLAND
E-MAIL: Enkelejd.Hashorva@unil.ch
Lanpeng.Ji@unil.ch