

# LOWER BOUNDS FOR THE CONSTANTS OF THE HARDY-LITTLEWOOD INEQUALITIES

GUSTAVO ARAÚJO AND DANIEL PELLEGRINO

ABSTRACT. Given an integer  $m \geq 2$ , the Hardy–Littlewood inequality (for real scalars) says that for all  $2m \leq p \leq \infty$ , there exists a constant  $C_{m,p}^{\mathbb{R}} \geq 1$  such that, for all continuous  $m$ -linear forms  $A : \ell_p^N \times \cdots \times \ell_p^N \rightarrow \mathbb{R}$  and all positive integers  $N$ ,

$$\left( \sum_{j_1, \dots, j_m=1}^N |A(e_{j_1}, \dots, e_{j_m})|^{\frac{2mp}{mp+p-2m}} \right)^{\frac{mp+p-2m}{2mp}} \leq C_{m,p}^{\mathbb{R}} \|A\|.$$

The limiting case  $p = \infty$  is the well-known Bohnenblust–Hille inequality; the behavior of the constants  $C_{m,p}^{\mathbb{R}}$  is an open problem. In this note we provide nontrivial lower bounds for these constants.

## 1. INTRODUCTION

Let  $\mathbb{K}$  denote the field of real or complex scalars. The multilinear Bohnenblust–Hille inequality asserts that for all positive integers  $m \geq 2$  there exists a constant  $C_m^{\mathbb{K}} \geq 1$  such that

$$\left( \sum_{j_1, \dots, j_m=1}^N |A(e_{j_1}, \dots, e_{j_m})|^{\frac{2m}{m+1}} \right)^{\frac{m+1}{2m}} \leq C_m^{\mathbb{K}} \|A\|$$

for all continuous  $m$ -linear forms  $A : \ell_{\infty}^N \times \cdots \times \ell_{\infty}^N \rightarrow \mathbb{K}$  and all positive integers  $N$ . The precise growth of the constants  $C_m^{\mathbb{K}}$  is important for applications (see [6]) and remains a big open problem. Only very recently, in [2, 8] it was shown that the constants have a subpolynomial growth. For real scalars, in 2014 (see [4]) it was shown that the optimal constant for  $m = 2$  is  $\sqrt{2}$  and in general  $C_m^{\mathbb{R}} \geq 2^{1-\frac{1}{m}}$ . In the case of complex scalars it is still an open problem whether the optimal constants are strictly greater than 1; in the polynomial case, in 2013 D. Núñez-Alarcón proved that the complex constants are strictly greater than 1 (see [7]).

Even basic questions related to the constants  $C_m^{\mathbb{K}}$  remain unsolved. For instance:

- Is the sequence of optimal constants  $(C_m^{\mathbb{K}})_{m=1}^{\infty}$  increasing?
- Is the sequence of optimal constants  $(C_m^{\mathbb{K}})_{m=1}^{\infty}$  bounded?
- Is  $C_m^{\mathbb{C}} = 1$ ?

The Hardy–Littlewood inequalities are a generalization of the Bohnenblust–Hille inequality to  $\ell_p$  spaces. The bilinear case was proved by Hardy and Littlewood in 1934 (see [5]) and in 1981 it was extended to multilinear operators by Praciano–Pereira (see [9]).

**Theorem 1.1** (Hardy and Littlewood inequalities ([5, 9])). *Let  $m \geq 2$  be a positive integer and  $p \geq 2m$ . Then there exists a constant  $C_{m,p}^{\mathbb{K}} \geq 1$  such that, for every continuous  $m$ -linear form  $A : \ell_p^N \times \cdots \times \ell_p^N \rightarrow \mathbb{K}$ ,*

$$\left( \sum_{j_1, \dots, j_m=1}^N |A(e_{j_1}, \dots, e_{j_m})|^{\frac{2mp}{mp+p-2m}} \right)^{\frac{mp+p-2m}{2mp}} \leq C_{m,p}^{\mathbb{K}} \|A\|$$

for all positive integers  $N$ .

*Key words and phrases.* Absolutely summing operators.

The authors are supported by CNPq Grant 313797/2013-7 - PVE - Linha 2.

2010 Mathematics Subject Classification. Primary 46G25. Secondary 11J13, 30C10.

It was recently shown that  $(C_{m,p}^{\mathbb{R}})_{m=1}^{\infty}$  is sublinear for  $p \geq m^2$ . More precisely, it was shown that (see [1, page 1887])

$$\begin{aligned} C_{m,p}^{\mathbb{C}} &\leq \left(\frac{2}{\sqrt{\pi}}\right)^{\frac{2m(m-1)}{p}} \left(\prod_{j=2}^m \Gamma\left(2 - \frac{1}{j}\right)^{\frac{j}{2-2j}}\right)^{\frac{p-2m}{p}}, \\ C_{m,p}^{\mathbb{R}} &\leq (\sqrt{2})^{\frac{2m(m-1)}{p}} \left(2^{\frac{446381}{55440} - \frac{m}{2}} \prod_{j=14}^m \left(\frac{\Gamma\left(\frac{3}{2} - \frac{1}{j}\right)}{\sqrt{\pi}}\right)^{\frac{j}{2-2j}}\right)^{\frac{p-2m}{p}}, \quad \text{for } m \geq 14, \\ C_{m,p}^{\mathbb{R}} &\leq (\sqrt{2})^{\frac{2m(m-1)}{p}} \left(\prod_{j=2}^m 2^{\frac{1}{2j-2}}\right)^{\frac{p-2m}{p}}, \quad \text{for } 2 \leq m \leq 13. \end{aligned}$$

The precise estimates of the constants of the Hardy–Littlewood inequalities are unknown and even its asymptotic growth is a mystery (as it happens with the Bohnenblust–Hille inequality). In this note we provide nontrivial lower bounds for these inequalities. Following the lines of [4], it is possible to prove that

$$(1) \quad C_{m,p}^{\mathbb{R}} \geq 2^{\frac{mp+2m-2m^2-p}{mp}} > 1$$

when  $p > 2m$ , but note that when  $p = 2m$  we have  $2^{\frac{mp+2m-2m^2-p}{mp}} = 1$  and thus we do not have nontrivial information. In this paper we also treat the extreme case  $p = 2m$ .

**Theorem 1.2.** *The optimal constants of the Hardy–Littlewood inequalities satisfy*

$$C_{m,p}^{\mathbb{R}} > 1$$

## 2. THE PROOF OF THEOREM 1.2

All that is left to prove is the case  $p = 2m$ . We divide the proof in five steps.

**Step 1.** Induction.

This first step follows the lines of [4]. For  $2m \leq p \leq \infty$ , consider

$$\begin{aligned} T_{2,p} &: \ell_p^2 \times \ell_p^2 \rightarrow \mathbb{R} \\ (x^{(1)}, x^{(2)}) &\mapsto x_1^{(1)} x_1^{(2)} + x_1^{(1)} x_2^{(2)} + x_2^{(1)} x_1^{(2)} - x_2^{(1)} x_2^{(2)} \end{aligned}$$

and

$$\begin{aligned} T_{m,p} &: \ell_p^{2^{m-1}} \times \cdots \times \ell_p^{2^{m-1}} \rightarrow \mathbb{R} \\ (x^{(1)}, \dots, x^{(m)}) &\mapsto \left(x_1^{(m)} + x_2^{(m)}\right) T_{m-1,p}(x^{(1)}, \dots, x^{(m)}) \\ &\quad + \left(x_1^{(m)} - x_2^{(m)}\right) T_{m-1,p}\left(B^{2^{m-1}}(x^{(1)}), B^{2^{m-2}}(x^{(2)}), \dots, B^2(x^{(m-1)})\right), \end{aligned}$$

where  $x^{(k)} = \left(x_j^{(k)}\right)_{j=1}^{2^{m-1}} \in \ell_p^{2^{m-1}}$ ,  $1 \leq k \leq m$ , and  $B$  is the backward shift operator in  $\ell_p^{2^{m-1}}$ . Observe that

$$\begin{aligned} \left|T_{m,p}(x^{(1)}, \dots, x^{(m)})\right| &\leq \left|x_1^{(m)} + x_2^{(m)}\right| \left|T_{m-1,p}(x^{(1)}, \dots, x^{(m)})\right| \\ &\quad + \left|x_1^{(m)} - x_2^{(m)}\right| \left|T_{m-1,p}\left(B^{2^{m-1}}(x^{(1)}), B^{2^{m-2}}(x^{(2)}), \dots, B^2(x^{(m-1)})\right)\right| \\ &\leq \|T_{m-1,p}\| \left(\left|x_1^{(m)} + x_2^{(m)}\right| + \left|x_1^{(m)} - x_2^{(m)}\right|\right) \\ &= \|T_{m-1,p}\| 2 \max\left\{\left|x_1^{(m)}\right|, \left|x_2^{(m)}\right|\right\} \\ &\leq 2 \|T_{m-1,p}\| \left\|x^{(m)}\right\|_p. \end{aligned}$$

Therefore,

$$(2) \quad \|T_{m,p}\| \leq 2^{m-2} \|T_{2,p}\|.$$

**Step 2.** Estimating  $\|T_{2,4}\|$ .

Note that

$$\|T_{2,4}\| = \sup \left\{ \left\| T_{2,4}^{(x^{(1)})} \right\| : \|x^{(1)}\|_4 = 1 \right\},$$

where  $T_{2,4}^{(x^{(1)})} : \ell_4^2 \rightarrow \mathbb{R}$  is given by  $x^{(2)} \mapsto T_{2,4}(x^{(1)}, x^{(2)})$ . Thus we have the operator

$$T_{2,4}^{(x^{(1)})}(x^{(2)}) = (x_1^{(1)} + x_2^{(1)})x_1^{(2)} + (x_1^{(1)} - x_2^{(1)})x_2^{(2)}.$$

Since  $(\ell_4)^* = \ell_{\frac{4}{3}}$ , we obtain  $\|T_{2,4}^{(x^{(1)})}\| = \left\| (x_1^{(1)} + x_2^{(1)}, x_1^{(1)} - x_2^{(1)}, 0, 0, \dots) \right\|_{\frac{4}{3}}$ . Therefore

$$\|T_{2,4}\| = \sup \left\{ \left( \left| x_1^{(1)} + x_2^{(1)} \right|^{\frac{4}{3}} + \left| x_1^{(1)} - x_2^{(1)} \right|^{\frac{4}{3}} \right)^{\frac{3}{4}} : |x_1^{(1)}|^4 + |x_2^{(1)}|^4 = 1 \right\}.$$

We can verify that it is enough to maximize the above expression when  $x_1^{(1)}, x_2^{(1)} \geq 0$ . Then

$$\begin{aligned} \|T_{2,4}\| &= \sup \left\{ \left( \left( x + (1-x^4)^{\frac{1}{4}} \right)^{\frac{4}{3}} + \left| x - (1-x^4)^{\frac{1}{4}} \right|^{\frac{4}{3}} \right)^{\frac{3}{4}} : x \in [0, 1] \right\} \\ &= \max \left\{ \sup \left\{ f(x) : x \in [0, 2^{-\frac{1}{4}}] \right\}, \sup \left\{ g(x) : x \in [2^{-\frac{1}{4}}, 1] \right\} \right\} \end{aligned}$$

where

$$f(x) := \left( \left( x + (1-x^4)^{\frac{1}{4}} \right)^{\frac{4}{3}} + \left( (1-x^4)^{\frac{1}{4}} - x \right)^{\frac{4}{3}} \right)^{\frac{3}{4}}$$

and

$$g(x) := \left( \left( x + (1-x^4)^{\frac{1}{4}} \right)^{\frac{4}{3}} + \left( x - (1-x^4)^{\frac{1}{4}} \right)^{\frac{4}{3}} \right)^{\frac{3}{4}}.$$

Examining the maps  $f$  and  $g$  we easily conclude that

$$\|T_{2,4}\| < 1.74.$$

In fact, the precise value seems to be graphically  $\sqrt{3}$ .




FIGURE 1. Graphs of the functions  $f$  and  $g$ , respectively.

**Step 3.** Estimating  $\|T_{2,p}\|$  for  $p \geq 4$ .

For  $1 \leq p \leq +\infty$  and  $A : \ell_p^N \times \ell_p^N \rightarrow \mathbb{K}$ , consider  $\tilde{A} : \ell_p^N \rightarrow (\ell_p^N)^*$  given by  $\tilde{A}(x) = A(x, \cdot) \in (\ell_p^N)^*$ . It is easy to see that  $\|A\| = \|\tilde{A}\|$ .

From [4] we know that

$$\|T_{2,\infty}\| = 2.$$

Let us suppose, for a moment, that we have, in this case, the Riesz-Thorin Theorem for real scalars with constant 1, as in the case of complex scalars. By considering

$$\theta = \frac{p-4}{p},$$

we would conclude from [3, Theorem 1.1.1] that

$$\left\| \widetilde{T_{2,p}} \right\| \leq \left\| \widetilde{T_{2,4}} \right\|^{1-\theta} \left\| \widetilde{T_{2,\infty}} \right\|^\theta,$$

i.e.,

$$(3) \quad \|T_{2,p}\| \leq \|T_{2,4}\|^{1-\theta} \|T_{2,\infty}\|^\theta < (1.74)^{\frac{4}{p}} 2^{\frac{p-4}{p}}.$$

**Step 4.** Estimating the constants.

From (2) and (3) we would conclude that  $\|T_{m,p}\| < 2^{m-2} (1.74)^{\frac{4}{p}} 2^{\frac{p-4}{p}}$ . On the other hand, from Theorem 1.1 we have

$$(4^{m-1})^{\frac{mp+p-2m}{2mp}} = \left( \sum_{j_1, \dots, j_m=1}^{2^{m-1}} |T_{m,p}(e_{j_1}, \dots, e_{j_m})|^{\frac{2mp}{mp+p-2m}} \right)^{\frac{mp+p-2m}{2mp}} < C_{m,p}^{\mathbb{R}} 2^{m-2} (1.74)^{\frac{4}{p}} 2^{\frac{p-4}{p}}.$$

and thus

$$C_{m,p}^{\mathbb{R}} > \frac{(4^{m-1})^{\frac{mp+p-2m}{2mp}}}{2^{m-2} (1.74)^{\frac{4}{p}} 2^{\frac{p-4}{p}}} = 2^{\frac{mp+(6-4 \log_2(1.74))m-2m^2-p}{mp}}.$$

**Step 5.** Verifying that  $2^{\frac{mp+(6-4 \log_2(1.74))m-2m^2-p}{mp}} > 1$ .

Indeed,

$$2^{\frac{mp+(6-4 \log_2(1.74))m-2m^2-p}{mp}} = 2^{\frac{mp+6m-2m^2-p}{mp}} = \frac{2^{\frac{mp+6m-2m^2-p}{mp}}}{(1.74)^{\frac{4}{p}}} > 2^{\frac{mp+2m-2m^2-p}{mp}} \geq 1.$$

However, if  $p = 2m$  it remains to consider the case in which the Riesz-Thorin Theorem holds with a constant bigger than 1; for this particular case, we may repeat the Step 2 for other values of  $p$ . We just need to observe that  $\|T_{2,p}\| < 2$  for  $4 \leq p < \infty$ . In fact, in this case,

$$C_{m,p}^{\mathbb{R}} \geq \frac{(4^{m-1})^{\frac{mp+p-2m}{2mp}}}{2^{m-2} \|T_{m,p}\|} > 2^{\frac{mp+2m-2m^2-p}{mp}} \geq 1.$$

**Acknowledgement:** The authors thank the referees for important suggestions that improved the final version of this paper.

## REFERENCES

- [1] G. Araújo, D. Pellegrino, and D.D.P. Silva, On the upper bounds for the constants of the Hardy–Littlewood inequality, *J. Funct. Anal.* **267** (2014), 1878–1888.
- [2] F. Bayart, D. Pellegrino, and J.B. Seoane-Sepúlveda, The Bohr radius of the  $n$ -dimensional polydisk is equivalent to  $\sqrt{(\log n)/n}$ , *Advances in Math.* **264** (2014) 726–746.
- [3] J. Bergh and J. Löfström, Interpolation spaces: an introduction, *Grundlehren der Mathematischen Wissenschaften* **223**, Berlin-New York: Springer-Verlag, ISBN 3-540-07875-4.
- [4] D. Diniz, G. A. Muñoz-Fernández, D. Pellegrino, and J. B. Seoane-Sepúlveda, Lower Bounds for the constants in the Bohnenblust–Hille inequality: the case of real scalars, *Proc. Amer. Math. Soc.* **142** (2014), n. 2, 575–580.
- [5] G. Hardy and J. E. Littlewood, Bilinear forms bounded in space  $[p, q]$ , *Quart. J. Math.* **5** (1934).
- [6] A. Montanaro, Some applications of hypercontractive inequalities in quantum information theory, *J. Math. Phys.* **53** (2012).
- [7] D. Núñez-Alarcón, A note on the polynomial Bohnenblust–Hille inequality, *J. Math. Anal. Appl.*, **407** (2013) 179–181.
- [8] D. Núñez-Alarcón, D. Pellegrino, D. M. Serrano-Rodríguez, and J. B. Seoane-Sepúlveda, There exist multilinear Bohnenblust–Hille constants  $(C_n)_{n=1}^\infty$  with  $\lim_{n \rightarrow \infty} (C_{n+1} - C_n) = 0$ , *J. Funct. Anal.*, **264** (2013) 429–463.
- [9] T. Praciano-Pereira, On bounded multilinear forms on a class of  $\ell_p$  spaces, *J. Math. Anal. Appl.* **81** (1981), n. 2, 561–568.

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, 58.051-900 - JOÃO PESSOA, BRAZIL.  
E-mail address: gdasaraujo@gmail.com

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, 58.051-900 - JOÃO PESSOA, BRAZIL.  
E-mail address: pellegrino@pq.cnpq.br and dpellegrino@gmail.com