arXiv:1405.3016v1 [math.AP] 13 May 2014

NON-LOCAL OPERATORS, NON-ARCHIMEDEAN PARABOLIC-TYPE EQUATIONS WITH VARIABLE COEFFICIENTS AND MARKOV PROCESSES

L. F. CHACÓN-CORTES AND W. A. ZÚÑIGA-GALINDO

ABSTRACT. In this article, we introduce a new class of parabolic-type pseudo differential equations with variable coefficients over the *p*-adics. We establish the existence and uniqueness of solutions for the Cauchy problem associated with these equations. The fundamental solutions of these equations are connected with Markov processes. Some of these equations are related to new models of complex systems.

1. INTRODUCTION

Stochastic processes on *p*-adic spaces, or more generally on ultrametric spaces, have been studied extensively due to its connections with models of complex systems, see e.g. [2]-[3], [4]-[6], [8], [9], [11], [13], [14], [15], [16], [22], and the references therein. In [4]-[6], Avetisov et al. introduced a new class of models for complex systems based on p-adic analysis, these models can be applied, for instance, to the study the relaxation of biological complex systems. From a mathematical point view, in these models the time-evolution of a complex system is described by a *p*-adic master equation (a parabolic-type pseudodifferential equation) which controls the time-evolution of a transition function of a random walk on an ultrametric space, and the random walk describes the dynamics of the system in the space of configurational states which is approximated by an ultrametric space (\mathbb{Q}_n) . The simplest type of master equation is the one-dimensional *p*-adic heat equation. This equation was introduced in the book of Vladimirov, Volovich and Zelenov [22, Section XVI]. In [13, Chapters 4, 5] Kochubei presented a general theory for onedimensional parabolic-type pseudodifferential equations with variable coefficients, whose fundamental solutions are transition density functions for Markov processes in the p-adic line, see also [17], [18], [21]. In [23], the second author introduced padic analogs for the *n*-dimensional elliptic operators and studied the corresponding heat equations and the associated Markov processes, see also [7], [21].

In [8], the authors introduced a new type of non-local operators which are naturally connected with parabolic-type pseudodifferential equations. Building up on [8] and [13]-[12], in this article, we introduce a new class of parabolic-type pseudodifferential equations with variable coefficients, which contains the one-dimensional p-adic heat equation of [22], the equations studied by Kochubei in [13], and the equations studied by Rodríguez-Vega in [17]. Our theory is not applicable to the

²⁰⁰⁰ Mathematics Subject Classification. Primary 35K90, 60J25; Secondary 26E30.

Key words and phrases. Parabolic-type equations, diffusion, dynamics of disordered systems, Markov processes, p-adic fields, non-Archimedean analysis.

The second author was partially supported by Conacyt (Mexico), Grant # 127794.

equations studied in [23], [7]. We establish the existence and uniqueness of solutions for the Cauchy problem for such equations, see Theorems 4.1, 5.5, 6.3. We show that the fundamental solutions of these equations are transition density functions of Markov processes, see Theorem 7.4. Finally, we study the well-possednes of the Cauchy problem, see Theorem 8.1.

2. Preliminaries

In this section we fix the notation and collect some basic results on *p*-adic analysis that we will use through the article. For a detailed exposition the reader may consult [1], [19], [22].

2.1. The field of *p*-adic numbers. Along this article *p* will denote a prime number. The field of *p*-adic numbers \mathbb{Q}_p is defined as the completion of the field of rational numbers \mathbb{Q} with respect to the *p*-adic norm $|\cdot|_p$, which is defined as

$$|x|_p = \begin{cases} 0 & \text{if } x = 0\\ p^{-\gamma} & \text{if } x = p^{\gamma} \frac{a}{b} \end{cases}$$

where a and b are integers coprime with p. The integer $\gamma := ord(x)$, with $ord(0) := +\infty$, is called the p-adic order of x. We extend the p-adic norm to \mathbb{Q}_p^n by taking

$$||x||_p := \max_{1 \le i \le n} |x_i|_p, \quad \text{for } x = (x_1, \dots, x_n) \in \mathbb{Q}_p^n.$$

We define $ord(x) = \min_{1 \le i \le n} \{ ord(x_i) \}$, then $||x||_p = p^{-ord(x)}$. The set $(\mathbb{Q}_p^n, || \cdot ||_p)$ is a complete ultrametric space. As a topological space \mathbb{Q}_p is homeomorphic to a Cantor-like subset of the real line.

Any *p*-adic number $x \neq 0$ has a unique expansion $x = p^{ord(x)} \sum_{j=0}^{\infty} x_j p^j$, where $x_j \in \{0, 1, 2, \ldots, p-1\}$ and $x_0 \neq 0$. By using this expansion, we define the fractional part of $x \in \mathbb{Q}_p$, denoted $\{x\}_p$, as the rational number

$$\{x\}_p = \begin{cases} 0 & \text{if } x = 0 \text{ or } ord(x) \ge 0\\ p^{\operatorname{ord}(x)} \sum_{j=0}^{-ord(x)-1} x_j p^j & \text{if } ord(x) < 0. \end{cases}$$

For $\gamma \in \mathbb{Z}$, denote by $B_{\gamma}^{n}(a) = \{x \in \mathbb{Q}_{p}^{n} : ||x - a||_{p} \leq p^{\gamma}\}$ the ball of radius p^{γ} with center at $a = (a_{1}, \ldots, a_{n}) \in \mathbb{Q}_{p}^{n}$, and take $B_{\gamma}^{n}(0) := B_{\gamma}^{n}$. Notice that $B_{\gamma}^{n}(a) = B_{\gamma}(a_{1}) \times \cdots \times B_{\gamma}(a_{n})$, where $B_{\gamma}(a_{i}) := \{x \in \mathbb{Q}_{p} : |x - a_{i}|_{p} \leq p^{\gamma}\}$ is the one-dimensional ball of radius p^{γ} with center at $a_{i} \in \mathbb{Q}_{p}$. The ball B_{0}^{n} equals the product of n copies of $B_{0} := \mathbb{Z}_{p}$, the ring of p-adic integers. We denote by $\Omega(||x||_{p})$ the characteristic function of B_{0}^{n} . For more general sets, say Borel sets, we use $1_{A}(x)$ to denote the characteristic function of A.

2.2. The Bruhat-Schwartz space. A complex-valued function φ defined on \mathbb{Q}_p^n is called locally constant if for any $x \in \mathbb{Q}_p^n$ there exists an integer $l = l(x) \in \mathbb{Z}$ such that

(2.1)
$$\varphi(x+x') = \varphi(x) \text{ for } x' \in B_l^n.$$

The set of all locally constant functions φ , for which the integer l(x) is independent of x, form \mathbb{C} -vector space denoted by $\widetilde{\mathcal{E}}(\mathbb{Q}_p^n) := \widetilde{\mathcal{E}}$. Given $\varphi \in \widetilde{\mathcal{E}}$, we call the largest possible $l = l(\varphi)$, the parameter of local constancy of φ .

A function $\varphi : \mathbb{Q}_p^n \to \mathbb{C}$ is called a *Bruhat-Schwartz function (or a test function)* if it is locally constant with compact support. The \mathbb{C} -vector space of Bruhat-Schwartz functions is denoted by $S(\mathbb{Q}_p^n) := S$. Notice that $S \subset \widetilde{\mathcal{E}}$.

Let $S'(\mathbb{Q}_p^n) := S'$ denote the set of all functionals (distributions) on $S(\mathbb{Q}_p^n)$. All functionals on $S(\mathbb{Q}_p^n)$ are continuous.

Set $\chi_p(y) = \exp(2\pi i \{y\}_p)$ for $y \in \mathbb{Q}_p$. The map $\chi_p(\cdot)$ is an additive character on \mathbb{Q}_p , i.e. a continuous map from \mathbb{Q}_p into the unit circle satisfying $\chi_p(y_0 + y_1) = \chi_p(y_0)\chi_p(y_1), y_0, y_1 \in \mathbb{Q}_p$.

Given $\xi = (\xi_1, \ldots, \xi_n)$ and $x = (x_1, \ldots, x_n) \in \mathbb{Q}_p^n$, we set $\xi \cdot x := \sum_{j=1}^n \xi_j x_j$. The Fourier transform of $\varphi \in S(\mathbb{Q}_p^n)$ is defined as

$$(\mathcal{F}\varphi)(\xi) = \int_{\mathbb{Q}_p^n} \Psi(-\xi \cdot x)\varphi(x)d^n x \quad \text{for } \xi \in \mathbb{Q}_p^n,$$

where $d^n x$ is the Haar measure on \mathbb{Q}_p^n normalized by the condition $vol(B_0^n) = 1$. The Fourier transform is a linear isomorphism from $S(\mathbb{Q}_p^n)$ onto itself satisfying $(\mathcal{F}(\mathcal{F}\varphi))(\xi) = \varphi(-\xi)$. We will also use the notation $\mathcal{F}_{x\to\xi}\varphi$ and $\widehat{\varphi}$ for the Fourier transform of φ .

2.2.1. Fourier transform. The Fourier transform $\mathcal{F}[T]$ of a distribution $T \in S'(\mathbb{Q}_p^n)$ is defined by

$$(\mathcal{F}[T], \varphi) = (T, \mathcal{F}[\varphi]) \text{ for all } \varphi \in S(\mathbb{Q}_p^n)$$

The Fourier transform $f \to \mathcal{F}[T]$ is a linear isomorphism from $S'(\mathbb{Q}_p^n)$ onto $S'(\mathbb{Q}_p^n)$. Furthermore, $T = \mathcal{F}[\mathcal{F}[T](-\xi)]$.

3. A class of non-local operators

Denote by \mathfrak{M}_{λ} , with $\lambda \geq 0$, the \mathbb{C} -vector space of all the functions $\varphi \in \widetilde{\mathcal{E}}$ satisfying $|\varphi(x)| \leq C(1 + ||x||_p^{\lambda})$. If the function φ depends also on a parameter t, we shall say that φ belongs to \mathfrak{M}_{λ} uniformly with respect to t, if its constant C and its parameter of local constancy do not depend on t. Notice that, if $0 \leq \lambda_1 \leq \lambda_2$, then $\mathfrak{M}_0 \subseteq \mathfrak{M}_{\lambda_1} \subseteq \mathfrak{M}_{\lambda_2}$, and that $S(\mathbb{Q}_p^n) \subseteq \mathfrak{M}_0$.

Take $\mathbb{R}_+ := \{x \in \mathbb{R}; x \ge 0\}$, and fix a function

$$w_{\alpha}: \mathbb{Q}_p^n \to \mathbb{R}_+$$

having the following properties:

(i) $w_{\alpha}(y)$ is a radial (i.e. $w_{\alpha}(y) = w_{\alpha}(\|y\|_{p})$) and continuous function; (ii) $w_{\alpha}(y) = 0$ if and only if y = 0;

(iii) there exist constants $C_0, C_1 > 0$, and $\alpha > n$ such that

$$C_0 \|y\|_p^{\alpha} \le w_{\alpha}(\|y\|_p) \le C_1 \|y\|_p^{\alpha} \text{ for any } y \in \mathbb{Q}_p^n.$$

Set

$$A_{w_{\alpha}}\left(\xi\right) := \int_{\mathbb{Q}_{p}^{n}} \frac{1 - \Psi\left(-y \cdot \xi\right)}{w_{\alpha}\left(\left\|y\right\|_{p}\right)} d^{n}y.$$

In [8], we establish that function $A_{w_{\alpha}}$ is radial, positive, continuous, $A_{w_{\alpha}}(0) = 0$, and $A_{w_{\alpha}}(\xi) = A_{w_{\alpha}}\left(\|\xi\|_{p}\right) = A_{w_{\alpha}}\left(p^{-ord(\xi)}\right)$ is a decreasing function of $ord(\xi)$, cf. [8, Lemma 3.2]. In addition, we introduce the following operator:

(3.1)
$$(\boldsymbol{W}_{\alpha}\varphi)(x) = \int_{\mathbb{Q}_{p}^{n}} \frac{\varphi(x-y) - \varphi(x)}{w_{\alpha}\left(\|y\|_{p}\right)} d^{n}y, \, \varphi \in S(\mathbb{Q}_{p}^{n})$$

Lemma 3.1. If $\alpha - n > \lambda$, then \mathbf{W}_{α} can be extended to \mathfrak{M}_{λ} and formula (3.1) holds. Furthermore, $\mathbf{W}_{\alpha} : \mathfrak{M}_{\lambda} \to \mathfrak{M}_{\lambda}$.

Proof. Notice that if $\varphi \in \mathfrak{M}_{\lambda}$, there exists a constant $l = l(\varphi) \in \mathbb{Z}$, such that

(3.2)
$$(\boldsymbol{W}_{\alpha}\varphi)(x) = \int_{\|\boldsymbol{y}\|_{p} \ge p^{l}} \frac{\varphi(x-\boldsymbol{y}) - \varphi(x)}{w_{\alpha}\left(\|\boldsymbol{y}\|_{p}\right)} d^{n}\boldsymbol{y}$$

We now show that $|(\boldsymbol{W}_{\alpha}\varphi)(x)| \leq A(1+||x||_{p}^{\lambda})$. By using that $\varphi \in \mathfrak{M}_{\lambda}$, and $\alpha > n$,

$$|(\boldsymbol{W}_{\alpha}\varphi)(x)| \leq C \int_{\|y\|_{p} \geq p^{l}} \frac{(1+\|x-y\|_{p}^{\lambda})}{\|y\|_{p}^{\alpha}} d^{n}y + C'(1+\|x\|_{p}^{\lambda}).$$

Hence, it is sufficient to show that the above integral can be bounded by $A(1+||x||_p^{\lambda})$, for some positive constant A. If $||x||_p > ||y||_p$,

$$\int_{\|y\|_{p} \ge p^{l}} \frac{(1 + \|x - y\|_{p}^{\lambda})}{\|y\|_{p}^{\alpha}} d^{n}y \le (1 + \|x\|_{p}^{\lambda}) \int_{\|y\|_{p} \ge p^{l}} \frac{1}{\|y\|_{p}^{\alpha}} d^{n}y$$
$$= B(1 + \|x\|_{p}^{\lambda}),$$

where B is a positive constant. If $||x||_p < ||y||_p$, by using $\alpha - n > \lambda$,

$$\int_{\|y\|_{p} \ge p^{l}} \frac{(1 + \|x - y\|_{p}^{\lambda})}{\|y\|_{p}^{\alpha}} d^{n}y \le \int_{\|y\|_{p} \ge p^{l}} \frac{(1 + \|y\|_{p}^{\lambda})}{\|y\|_{p}^{\alpha}} d^{n}y < \infty.$$

If $||x||_p = ||y||_p \ge p^l$, we take $x = p^L u$, $y = p^L v$, with $||v||_p = ||u||_p = 1$, $L \in \mathbb{Z}$, then

$$\int_{\|y\|_{p}=\|x\|_{p}} \frac{(1+\|x-y\|_{p}^{\lambda})}{\|y\|_{p}^{\alpha}} d^{n}y = p^{-L(n-\alpha)} \int_{\|v\|_{p}=1} (1+p^{-L\lambda} \|u-v\|_{p}^{\lambda}) d^{n}v$$
$$\leq A\left(\|x\|_{p}^{-(\alpha-n)} + \|x\|_{p}^{-(\alpha-n-\lambda)}\right) \leq A'(p,l,\alpha,n,\lambda),$$

where A, A' are positive constants.

Finally, by (3.2) $\boldsymbol{W}_{\alpha}\varphi$ is locally constant.

4. PARABOLIC-TYPE EQUATIONS WITH CONSTANT COEFFICIENTS

Consider the following Cauchy problem:

(4.1)
$$\begin{cases} \frac{\partial u}{\partial t}(x,t) - \kappa \cdot (\boldsymbol{W}_{\alpha}u)(x,t) = f(x,t), & x \in \mathbb{Q}_{p}^{n}, t \in (0,T] \\ u(x,0) = \varphi(x), \end{cases}$$

where, $\alpha > n$, κ , T are positive constants, $\varphi \in Dom(\mathbf{W}_{\alpha}) := \mathfrak{M}_{\lambda}$, with $\alpha - n > \lambda$, f is continuous in (x, t) and belongs to \mathfrak{M}_{λ} uniformly with respect to t, and $u : \mathbb{Q}_{p}^{n} \times [0, T] \to \mathbb{C}$ is an unknown function.

We say that u(x,t) is a solution of (4.1), if u(x,t) is continuous in $(x,t), u(\cdot,t) \in Dom(W_{\alpha})$ for $t \in [0,T], u(x,\cdot)$ is continuously differentiable for $t \in (0,T], u(x,t) \in \mathfrak{M}_{\lambda}$ uniformly in t, and u satisfies (4.1) for all t > 0.

Cauchy problem (4.1) was studied in [8] using semigroup theory. In this article, we study this problem in the space \mathfrak{M}_{λ} , which is not contained in L^{ρ} for any $\rho \in [1, \infty]$, and thus we cannot use semigroup theory, see e.g. [8, Theorem 6.5]. We define

(4.2)
$$Z(x,t;w_{\alpha},\kappa) := Z(x,t) = \int_{\mathbb{Q}_p^n} e^{-\kappa t A_{w_{\alpha}}(\|\xi\|_p)} \Psi(x \cdot \xi) d^n \xi,$$

for t > 0 and $x \in \mathbb{Q}_p^n$. Notice that, $Z(x,t) = \mathcal{F}_{\xi \to x}^{-1}[e^{-\kappa t A_{w_\alpha}(\|\xi\|_p)}] \in L^1 \cap L^2$ for t > 0, since $C' \|\xi\|_p^{\alpha-n} \leq A_{w_\alpha}(\|\xi\|_p) \leq C'' \|\xi\|_p^{\alpha-n}$, cf. [8, Lemma 3.4]. Furthermore, $Z(x,t) \geq 0$, for t > 0, $x \in \mathbb{Q}_p^n$, cf. [8, Theorem 4.3 (i)]. These functions are called *heat kernels*. When considering Z(x,t) as a function of x for t fixed we will write $Z_t(x)$.

We set

$$u_1(x,t) := \int_{\mathbb{Q}_p^n} Z(x-y,t)\varphi(y)d^n y,$$
$$u_2(x,t) := \int_0^t \int_{\mathbb{Q}_p^n} Z(x-y,t-\theta)f(y,\theta)d^n yd\theta$$

for $\varphi, f \in \mathfrak{M}_{\lambda}$ with $\alpha - n > \lambda$, for $0 \leq t \leq T$, and $x \in \mathbb{Q}_p^n$. The main result of this section is the following:

Theorem 4.1. The function

$$u(x,t) = u_1(x,t) + u_2(x,t)$$

is a solution of Cauchy Problem (4.1).

The proof requires several steps.

4.1. Claim $u(x,t) \in \mathfrak{M}_{\lambda}$. In order to prove this claim, we need some preliminary results.

Remark 4.2. The function $Z_t(x)$ is radial since it is the inverse Fourier transform of the radial function $e^{-\kappa t A_{w_\alpha}(\|\xi\|_p)}$. Then $Z_t(x)$ is locally constant in $\mathbb{Q}_p^n \setminus \{0\}$. Furthermore, $Z_t(x+y) = Z_t(x)$ if $\|y\|_p < \|x\|_p$ for any $y \in \mathbb{Q}_p^n$ and $x \in \mathbb{Q}_p^n \setminus \{0\}$, and t > 0.

Lemma 4.3. There exist positive constants C_1, C_2 such that Z(x,t) satisfies the following conditions:

$$\begin{array}{l} (i) \ Z(x,t) \leq C_{1}t^{-\frac{n}{\alpha-n}}, \ for \ t > 0 \ and \ x \in \mathbb{Q}_{p}^{n}; \\ (ii) \ Z(x,t) \leq C_{2}t \ \|x\|_{p}^{-\alpha}, \ for \ t > 0 \ and \ x \in \mathbb{Q}_{p}^{n} \setminus \{0\}; \\ (iii) \ Z(x,t) \leq \max\{2^{\alpha}C_{1}, 2^{\alpha}C_{2}\}t \left(\|x\|_{p} + t^{\frac{1}{\alpha-n}}\right)^{-\alpha}, \ for \ t > 0 \ and \ x \in \mathbb{Q}_{p}^{n}; \\ (iv) \ \int_{\mathbb{Q}_{p}^{n}} Z(x,t)d^{n}x = 1, \ for \ t > 0. \end{array}$$

Proof. (i) By (4.2) and Lemma 3.4 in [8],

$$Z(x,t) \leq \int_{\mathbb{Q}_p^n} e^{-\kappa t A_{w_\alpha}(\|\xi\|_p)} d^n \xi \leq \int_{\mathbb{Q}_p^n} e^{-C_0 t \|\xi\|_p^{\alpha-n}} d^n \xi.$$

Let *m* be an integer such that $p^{m-1} \leq t^{\frac{1}{\alpha-n}} \leq p^m$, then

$$Z(x,t) \leq \int_{\mathbb{Q}_p^n} e^{-C_0 t \left\| p^{-(m-1)} \xi \right\|_p^{\alpha-n}} d^n \xi,$$

now, by changing variables as $z = p^{-(m-1)}\xi$, we have

$$Z(x,t) \le p^{-(m-1)n} \int_{\mathbb{Q}_p^n} e^{-C_0 t \|z\|_p^{\alpha-n}} d^n z \le C_1 t^{-\frac{n}{\alpha-n}}.$$

(ii) It follows from [8, Lemma 4.1]. (iii) The results is obtained from the two following inequalities. If $||x||_p \ge t^{\frac{1}{\alpha-n}}$, then $||x||_p \ge \frac{||x||_p}{2} + \frac{t^{\frac{1}{\alpha-n}}}{2}$ and $||x||_p^{-\alpha} \le 2^{\alpha} \left(||x||_p + t^{\frac{1}{\alpha-n}} \right)^{-\alpha}$, multiplying by $C_2 t$ and using (ii),

$$Z(x,t) \le 2^{\alpha} C_2 t \left(\left\| x \right\|_p + t^{\frac{1}{\alpha - n}} \right)^{-\alpha}$$

If $\|x\|_p \leq t^{\frac{1}{\alpha-n}}$, then $\frac{\|x\|_p}{2} + \frac{t^{\frac{1}{\alpha-n}}}{2} \leq t^{\frac{1}{\alpha-n}}$ and $\left(\|x\|_p + t^{\frac{1}{\alpha-n}}\right)^{-\alpha} \geq 2^{-\alpha}t^{\frac{-\alpha}{\alpha-n}} = 2^{-\alpha}t^{-1-\frac{n}{\alpha-n}}$, multiplying by C_1 and using (i),

$$Z(x,t) \le 2^{\alpha} C_1 t \left(\left\| x \right\|_p + t^{\frac{1}{\alpha - n}} \right)^{-\alpha}$$

(iv) By (iii), $Z_t(x) \in L^1(\mathbb{Q}_p^n)$ for t > 0. Now, the announced identity follows by applying the Fourier inversion formula.

Proposition 4.4 ([18], Proposition 2). If b > 0, $0 \le \lambda < \alpha$, and $x \in \mathbb{Q}_p^n$, then

$$\int_{\mathbb{Q}_{p}^{n}} \left(b + \|x - \xi\|_{p} \right)^{-\alpha - n} \|\xi\|_{p}^{\lambda} d^{n}\xi \leq Cb^{-\alpha} \left(1 + \|x\|_{p}^{\lambda} \right),$$

where the constant C does not depend on b or x.

Lemma 4.5. The functions u_1, u_2 belong to \mathfrak{M}_{λ} uniformly in t, for $\lambda + n < \alpha$.

Proof. By Lemma 4.3 (iii), and Proposition 4.4,

$$\begin{aligned} |u_1(x,t)| &\leq \int_{\mathbb{Q}_p^n} Z(x-y,t) \left| \varphi(y) \right| d^n y \leq C \int_{\mathbb{Q}_p^n} t \left(t^{\frac{1}{\alpha-n}} + \left\| x - y \right\|_p \right)^{-\alpha} \left(1 + \left\| y \right\|_p^{\lambda} \right) d^n y \\ &\leq C' \left(1 + \left\| x \right\|_p^{\lambda} \right). \end{aligned}$$

On the other hand, since

$$u_1(x,t) = \int_{\mathbb{Q}_p^n} Z(w,t)\varphi(x-w)d^n w,$$

 u_1 is locally constant and $l(u_1) = l(\varphi)$ uniformly in t. The proof for u_2 is similar.

Remark 4.6. Notice that $u_1, u_2, W_{\gamma}u_1, W_{\gamma}u_2 \in \mathfrak{M}_{\lambda}$, for any γ satisfying $\lambda + n < \gamma \leq \alpha$.

4.2. Claim u(x,t) satisfies the initial condition. This claim follows from Lemma 4.5 by using the following result.

Lemma 4.7. If $\varphi \in \mathfrak{M}_{\lambda}$, with $\alpha > \lambda + n$, then

$$\lim_{t\to 0^+} \int_{\mathbb{Q}_p^n} Z(x-\xi,t)\varphi(\xi) d^n\xi = \varphi(x).$$

Proof. By Lemma 4.3 (iv),

(4.3)
$$\int_{\mathbb{Q}_p^n} Z(x-\xi,t)\varphi(\xi)d^n\xi = \int_{\mathbb{Q}_p^n} Z(x-\xi,t)\left[\varphi(\xi)-\varphi(x)\right]d^n\xi + \varphi(x).$$

Now, by Lemma 4.3 (iii) and the local constancy of φ ,

$$\begin{split} \int_{\mathbb{Q}_{p}^{n}} & \int_{\mathbb{Q}_{p}^{n}} Z(x-\xi,t) \left[\varphi(\xi) - \varphi(x)\right] d^{n}\xi \leq Ct \int_{\|x-\xi\|_{p} \geq p^{l}} (t^{\frac{1}{\alpha-n}} + \|x-\xi\|_{p})^{-\alpha} \left|\varphi(\xi) - \varphi(x)\right| d^{n}\xi \\ & \leq Ct \int_{\|z\|_{p} \geq p^{l}} (t^{\frac{1}{\alpha-n}} + \|z\|_{p})^{-\alpha} \left|\varphi(x-z) - \varphi(x)\right| d^{n}z \\ & \leq Ct \int_{\|z\|_{p} \geq p^{l}} \|z\|_{p}^{-\alpha} (1 + \|x-z\|^{\lambda}) d^{n}z + C't \left|\varphi(x)\right| \leq th \left(x\right). \end{split}$$

The formula is obtained by taking limit when $t \to 0^+$ in (4.3).

4.3. Claim u(x,t) is a solution of Cauchy problem (4.1). The proof of this claim is a consequence of Corollary 4.10, Lemmas 4.11 and 4.12. Several preliminary results are required.

Lemma 4.8. There exist positive constants C_3, C_4 such that Z(x, t) satisfies the following conditions:

$$\begin{aligned} (i) \quad &\frac{\partial Z(x,t)}{\partial t} = -\kappa \int_{\mathbb{Q}_p^n} A_{w_\alpha}(\|\xi\|_p) e^{-\kappa t A_{w_\alpha}(\|\xi\|_p)} \Psi(x \cdot \xi) d^n \xi, \text{ for } t > 0 \text{ and } x \in \mathbb{Q}_p^n; \\ (ii) \quad & \left| \frac{\partial Z(x,t)}{\partial t} \right| \le C_3 t^{-\frac{\alpha}{\alpha-n}}, \text{ for } t > 0 \text{ and } x \in \mathbb{Q}_p^n; \\ (iii) \quad & \left| \frac{\partial Z(x,t)}{\partial t} \right| \le C_4 t \, \|x\|_p^{n-2\alpha}, \text{ for } t > 0 \text{ and } x \in \mathbb{Q}_p^n \setminus \{0\}; \\ (iv) \quad & \left| \frac{\partial Z(x,t)}{\partial t} \right| \le 2^{\alpha} C_3 \left(\|x\|_p + t^{\frac{1}{\alpha-n}} \right)^{-\alpha}, \text{ for } t > 0 \text{ and } x \in \mathbb{Q}_p^n \setminus \{0\}. \end{aligned}$$

Proof. (i) The formula is obtained by the Lebesgue Dominated Convergence Theorem, and the fact that $-\kappa A_{w_{\alpha}}(\|\xi\|_p)e^{-\kappa\tau A_{w_{\alpha}}(\|\xi\|_p)}\Psi(x\cdot\xi) \in L^1(\mathbb{Q}_p^n)$, for $\tau > 0$ fixed, cf. [8, Lemma 3.4]. (ii) By using (i) and Lemma 3.4 in [8],

$$\left|\frac{\partial Z(x,t)}{\partial t}\right| \leq \int_{\mathbb{Q}_p^n} C_1 \left\|\xi\right\|_p^{\alpha-n} e^{-\kappa C_2 t} \|\xi\|_p^{\alpha-n} d^n \xi.$$

We now pick an integer m such that $p^{m-1} \leq t^{\frac{1}{\alpha-n}} \leq p^m$, and proceed as in the proof of Lemma 4.3 (i), to obtain

$$\left|\frac{\partial Z(x,t)}{\partial t}\right| \le C_1 p^{-(m-1)n-(m-1)(\alpha-n)} \int_{\mathbb{Q}_p^n} \|z\|_p^{\alpha-n} e^{-\kappa C_2 t} \|z\|_p^{\alpha-n} d^n z \le C_3 t^{-\frac{\alpha}{\alpha-n}}.$$

(iii) Set $||x||_p = p^{\beta}$. Now, since $A_{w_{\alpha}}(||\xi||_p)e^{-\kappa t A_{w_{\alpha}}(||\xi||_p)} \in L^1 \cap L^2$ for t > 0, then $\frac{\partial Z(x,t)}{\partial t} \in L^1 \cap L^2$ for t > 0, and by applying the formula for the Fourier Transform of a radial function, we get

$$\frac{\partial Z(x,t)}{\partial t} = \|x\|_p^{-n}$$

$$\times \left((1-p^{-n}) \sum_{j=0}^{\infty} A_{w_\alpha}(p^{-\beta-j}) e^{-\kappa t A_{w_\alpha}(p^{-\beta-j})} p^{-nj} - A_{w_\alpha}(p^{-\beta+1}) e^{-\kappa t A_{w_\alpha}(p^{-\beta+1})} \right).$$

Now, by using that $A_{w_{\alpha}}(\xi)$ is a decreasing function of $ord(\xi)$,

$$\left| \frac{\partial Z(x,t)}{\partial t} \right| \le \left\| x \right\|_p^{-n} A_{w_\alpha}(p^{-\beta+1}) \left| (1-p^{-n}) \sum_{j=0}^{\infty} p^{-nj} - e^{-\kappa t A_{w_\alpha}(p^{-\beta+1})} \right|$$
$$\le \left\| x \right\|_p^{-n} A_{w_\alpha}(p^{-\beta+1}) \left(1 - e^{-\kappa t A_{w_\alpha}(p^{-\beta+1})} \right)$$

By using Mean Value Theorem and Lemma 3.4 in [8], we have

$$\left|\frac{\partial Z(x,t)}{\partial t}\right| \le C_4 \|x\|_p^{n-2\alpha} t.$$
(iv) If $\|x\|_p \le t^{\frac{1}{\alpha-n}}$, then $\frac{\|x\|_p}{2} + \frac{t^{\frac{1}{\alpha-n}}}{2} \le t^{\frac{1}{\alpha-n}}$ and $t^{\frac{-\alpha}{\alpha-n}} \le 2^{\alpha} \left(\|x\|_p + t^{\frac{1}{\alpha-n}}\right)^{-\alpha}$, multiplying by C_3 and using (ii), we have

$$\left|\frac{\partial Z(x,t)}{\partial t}\right| \le 2^{\alpha} C_3 \left(\left\|x\right\|_p + t^{\frac{1}{\alpha-n}}\right)^{-\alpha}.$$

Now, if $||x||_n \ge t^{\frac{1}{\alpha-n}}$, by using (iii),

(iv) If $||x||_p \leq$

(4.4)
$$\left|\frac{\partial Z(x,t)}{\partial t}\right| \le C_3 \|x\|_p^{-\alpha},$$

and since $\|x\|_p \ge t^{\frac{1}{\alpha-n}}$, then $\|x\|_p \ge \left(\frac{\|x\|_p}{2} + \frac{t^{\frac{1}{\alpha-n}}}{2}\right)$ and $2^{\alpha} \left(\|x\|_p + t^{\frac{1}{\alpha-n}}\right)^{-\alpha} \ge t^{\alpha}$ $||x||_p^{-\alpha}$, multiplying by C_3 and using (4.4), we have

$$\left|\frac{\partial Z(x,t)}{\partial t}\right| \le 2^{\alpha} C_3 \left(\left\|x\right\|_p + t^{\frac{1}{\alpha-n}}\right)^{-\alpha}.$$

Lemma 4.9. $(\mathbf{W}_{\gamma}Z_t)(x)$, with $\gamma \leq \alpha$, satisfies the following conditions:

 $(i) (\mathbf{W}_{\gamma} Z_t) (x) = -\int_{\mathbb{Q}_p^n} A_{w_{\gamma}} (\|\xi\|_p) e^{-\kappa t A_{w_{\alpha}}(\|\xi\|_p)} \Psi(x \cdot \xi) d^n \xi, \text{ for } t > 0 \text{ and } x \in \mathbb{Q}_p^n;$ (ii) $|(\mathbf{W}_{\gamma}Z_t)(x)| \leq 2^{\gamma}C\left(||x||_p + t^{\frac{1}{\alpha-n}}\right)^{-\gamma}$, for t > 0 and $x \in \mathbb{Q}_p^n$ and some positive constant C; (iii) $\int_{\mathbb{Q}_p^n} (\mathbf{W}_{\gamma} Z_t) (x) d^n x = 0.$

Proof. (i) Define

(4.5)
$$Z_t^{(M)}(x) = \int_{\|\eta\|_p \le p^M} \Psi(x \cdot \eta) e^{-\kappa t A_{w\alpha}(\|\eta\|_p)} d^n \eta, \text{ for } M \in \mathbb{N}.$$

This function is locally constant on \mathbb{Q}_p^n . Indeed, if $\|\xi\|_p \leq p^{-M}$, then $Z_t^{(M)}(x+\xi) = Z_t^{(M)}(x)$. Furthermore, $Z_t^{(M)}(x)$ is bounded, and thus $Z_t^{(M)}(x) \in \mathfrak{M}_0 \subset Dom(\mathbf{W}_{\gamma})$. We now use formula (3.1) and Fubini's Theorem to compute $(\mathbf{W}_{\gamma}Z_t^{(M)})(x)$ as follows:

$$\begin{aligned} (\mathbf{W}_{\gamma}Z_{t}^{(M)})(x) &= \int_{\mathbb{Q}_{p}^{n}} \frac{Z_{t}^{(M)}(x-\xi) - Z_{t}^{(M)}(x)}{w_{\gamma}(\|\xi\|_{p})} d^{n}\xi \\ &= \int_{\|\xi\|_{p} > p^{-M}} \int_{\|\eta\|_{p} \le p^{M}} e^{-\kappa t A_{w_{\alpha}}(\|\eta\|_{p})} \Psi(x\cdot\eta) \frac{(\Psi(\xi\cdot\eta)-1)}{w_{\gamma}(\|\xi\|_{p})} d^{n}\eta d^{n}\xi \\ &= \int_{\|\eta\|_{p} \le p^{M}} e^{-\kappa t A_{w_{\alpha}}(\|\eta\|_{p})} \Psi(x\cdot\eta) \int_{\|\xi\|_{p} > p^{-M}} \frac{(\Psi(\xi\cdot\eta)-1)}{w_{\gamma}(\|\xi\|_{p})} d^{n}\xi d^{n}\eta \\ &= -\int_{\|\eta\|_{p} \le p^{M}} e^{-\kappa t A_{w_{\alpha}}(\|\eta\|_{p})} \Psi(x\cdot\eta) A_{w_{\gamma}}(\|\eta\|_{p}) d^{n}\eta. \end{aligned}$$

By using that $e^{-\kappa t A_{w_{\alpha}}(\|\xi\|_p)} A_{w_{\gamma}}(\|\xi\|_p) \in L^1(\mathbb{Q}_p^n)$ for t > 0, cf. [8, Lemma 3.4] and the Dominated Convergence Theorem, we obtain

(4.6)
$$\lim_{M \to \infty} (\mathbf{W}_{\gamma} Z_t^{(M)})(x) = -\int_{\mathbb{Q}_p^n} A_{w_{\gamma}}(\|\eta\|_p) e^{-\kappa t A_{w_{\alpha}}(\|\eta\|_p)} \Psi(x \cdot \eta) d^n \eta.$$

On the other hand, by fixing $x \neq 0$ and for t > 0, $Z_t(x - \xi) - Z_t(x)$ is locally constant, cf. Remark 4.2, and bounded, cf. Lemma 4.3 (iii), then $(\mathbf{W}_{\gamma}Z_t)(x)$ is well-defined, and since $Z_t^{(M)}(x)$ is radial,

$$(\mathbf{W}_{\gamma} Z_t^{(M)})(x) = \int_{\|\xi\|_p > \|x\|_p} \frac{Z_t^{(M)}(x-\xi) - Z_t^{(M)}(x)}{w_{\gamma}(\|\xi\|_p)} d^n \xi,$$

and by Dominated Convergence Theorem, $\lim_{M\to\infty} (\mathbf{W}_{\gamma} Z_t^{(M)})(x) = (\mathbf{W}_{\gamma} Z_t)(x)$. Therefore by (4.6), we have

$$(\mathbf{W}_{\gamma}Z_t)(x) = -\int_{\mathbb{Q}_p^n} A_{w_{\gamma}}(\|\eta\|_p) e^{-\kappa t A_{w_{\alpha}}(\|\eta\|_p)} \Psi(x \cdot \eta) d^n \eta.$$

Finally, we note the right-hand side in the above formula is continuous at x = 0.

(ii) By (i) and Lemma 3.4 in [8],

$$|(\mathbf{W}_{\gamma}Z_t)(x)| \le C_0 \int_{\mathbb{Q}_p^n} \|\xi\|_p^{\gamma-n} e^{-\kappa C_1 t} \|\xi\|_p^{\alpha-n} d^n \xi.$$

We now pick an integer m such that $p^{m-1} \leq t^{\frac{1}{\alpha-n}} \leq p^m$, and proceed as in the proof of Lemma 4.3 (i), to obtain

(4.7)
$$|(\mathbf{W}_{\gamma}Z_t)(x)| \le Ct^{-\frac{\gamma}{\alpha-n}}.$$

Now, if $||x||_p \leq t^{\frac{1}{\alpha-n}}$, then $\frac{||x||_p}{2} + \frac{t^{\frac{1}{\alpha-n}}}{2} \leq t^{\frac{1}{\alpha-n}}$ and $t^{\frac{-\gamma}{\alpha-n}} \leq 2^{\gamma} \left(||x||_p + t^{\frac{1}{\alpha-n}} \right)^{-\gamma}$, multiplying by C and by using (4.7), we have

$$\left| (\mathbf{W}_{\gamma} Z_t)(x) \right| \le 2^{\gamma} C \left(\left\| x \right\|_p + t^{\frac{1}{\alpha - n}} \right)^{-\gamma}$$

On the other hand, let $||x||_p = p^{\beta}$, since $A_{w_{\gamma}}(||\xi||_p)e^{-\kappa t A_{w_{\alpha}}(||\xi||_p)} \in L^1 \cap L^2$ for t > 0, then $(\mathbf{W}_{\gamma}Z_t)(x) \in L^1 \cap L^2$ for t > 0, by proceeding as in the proof of Lemma 4.8 (iii), we obtain

$$\left| (\mathbf{W}_{\gamma} Z_t)(x) \right| \le Ct \left\| x \right\|_p^{n-\alpha-\gamma}.$$

Now, if $||x||_p \ge t^{\frac{1}{\alpha-n}}$, then

(4.8)
$$|(\mathbf{W}_{\gamma}Z_t)(x)| \leq C ||x||_p^{-\gamma}.$$

If $\|x\|_p \ge t^{\frac{1}{\alpha-n}}$, then $\|x\|_p \ge \left(\frac{\|x\|_p}{2} + \frac{t^{\frac{1}{\alpha-n}}}{2}\right)$ and $2^{\gamma} \left(\|x\|_p + t^{\frac{1}{\alpha-n}}\right)^{-\gamma} \ge \|x\|_p^{-\gamma}$, multiplying by C and using (4.8), we have

$$\left| (\mathbf{W}_{\gamma} Z_t)(x) \right| \le 2^{\gamma} C \left(\left\| x \right\|_p + t^{\frac{1}{\alpha - n}} \right)^{-\gamma}$$

(iii) It follows from (i) by the inversion formula for the Fourier transform. $\hfill \Box$

Corollary 4.10. $\frac{\partial Z(x,t)}{\partial t} = \kappa \cdot (\mathbf{W}_{\alpha} Z_t)(x)$ for t > 0 and $x \in \mathbb{Q}_p^n$. *Proof.* The formula follows from Lemma 4.8 (i) and Lemma 4.9 (i).

Proposition 4.11. Assume that $\varphi \in \mathfrak{M}_{\lambda}$, then the following assertions hold: (i) $\frac{\partial u_1}{\partial t}(x,t) = \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,t)}{\partial t} \varphi(y) d^n y$, for t > 0 and $x \in \mathbb{Q}_p^n \setminus \{0\}$; (ii) $(\mathbf{W}_{\gamma} u_1)(x,t) = \int_{\mathbb{Q}_p^n} (\mathbf{W}_{\gamma} Z_t)(x-y) \varphi(y) d^n y$, for $n + \lambda < \gamma \leq \alpha$, t > 0 and $x \in \mathbb{Q}_p^n \setminus \{0\}$.

Proof. (i) By using the Mean Value Theorem, $\frac{\partial u_1}{\partial t}(x,t)$ equals

$$\lim_{h\to 0} \int_{\mathbb{Q}_p^n} \left[\frac{Z(x-y,t+h) - Z(x-y,t)}{h} \right] \varphi(y) d^n y = \lim_{h\to 0} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial Z(x-y,\tau)}{\partial t} \varphi(y) d^n y = \lim_{h\to 0} \frac{\partial$$

where τ is between t and t + h. Now, the result follows by applying the Dominated Converge Theorem and Lemma 4.8 (iv).

(ii) By Remark 4.6, if $n + \lambda < \gamma$, then $u_1 \in Dom(\mathbf{W}_{\gamma})$ for t > 0. Then for any $L \in \mathbb{N}$, the following integral exists:

$$\int_{\|y\|_{p} > p^{-L}} \frac{u_{1}(x-y,t) - u_{1}(x,t)}{w_{\gamma}(\|y\|_{p})} d^{n}y$$
$$= \int_{\|y\|_{p} > p^{-L}} \frac{1}{w_{\gamma}(\|y\|_{p})} \int_{\mathbb{Q}_{p}^{n}} (Z_{t}(x-y-\xi) - Z_{t}(x-\xi)) \varphi(\xi) d^{n}\xi d^{n}y,$$

now, by using Fubini's Theorem, cf. Lemma 4.3 (iii),

$$\int_{\mathbb{Q}_p^n} \varphi(\xi) \int_{\|y\|_p > p^{-L}} \frac{(Z_t(x-\xi-y) - Z_t(x-\xi))}{w_\gamma(\|y\|_p)} d^n y d^n \xi.$$

We now fix a positive integer M, such that $\|y\|_p < p^{-L} < p^{-M} < \|x - \xi\|_p$, and use Remark 4.2,

$$(\mathbf{W}_{\gamma}u_{1})(x,t) = \lim_{L \to \infty} \int_{\|y\|_{p} > p^{-L}} \frac{u_{1}(x-y,t) - u_{1}(x,t)}{w_{\gamma}(\|y\|_{p})} d^{n}y$$

$$= \int_{\|x-\xi\|_{p} > p^{-M}} \varphi(\xi) (\mathbf{W}_{\gamma}Z_{t})(x-\xi) d^{n}\xi$$

$$+ \lim_{L \to \infty} \int_{\|x-\xi\|_{p} \le p^{-M}} \varphi(\xi) \int_{\|y\|_{p} > p^{-L}} \frac{(Z_{t}(x-\xi-y) - Z_{t}(x-\xi))}{w_{\gamma}(\|y\|_{p})} d^{n}y d^{n}\xi$$

$$= \int_{\|x-\xi\|_{p} > p^{-M}} \varphi(\xi) (\mathbf{W}_{\gamma}Z_{t})(x-\xi) d^{n}\xi + \int_{\|x-\xi\|_{p} \le p^{-M}} \varphi(\xi) (\mathbf{W}_{\gamma}Z_{t})(x-\xi) d^{n}\xi.$$

 Set

$$u_2(x,t,\tau) := \int_{\tau}^t \int_{\mathbb{Q}_p^n} Z(x-y,t-\theta) f(y,\theta) d^n y d\theta,$$

for $f \in \mathfrak{M}_{\lambda}$ with $\alpha - n > \lambda$, for $0 \le \tau \le t \le T$, and $x \in \mathbb{Q}_p^n$. By reasoning as in the proof of Lemma 4.5, we have $u_2(x, t, \tau) \in \mathfrak{M}_{\lambda}$ uniformly in t and τ .

Proposition 4.12. Assume that $f \in \mathfrak{M}_{\lambda}$, with $\alpha - n > \lambda$, then the following assertions hold:

$$(i) \frac{\partial u_2}{\partial t}(x,t,\tau) = f(x,t) + \int_{\tau}^t \left(\int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,t-\theta)}{\partial t} \left[f(y,\theta) - f(x,\theta) \right] d^n y \right) d\theta, \text{ for } t > 0$$

and $x \in \mathbb{Q}_p^n;$

(ii) $(\mathbf{W}_{\gamma}u_2)(x,t,\tau) = \int_{\tau}^{t} \int_{\mathbb{Q}_p^n} (\mathbf{W}_{\gamma}Z)(x-y,t-\theta)f(y,\theta)d^nyd\theta$, for $n+\lambda < \gamma \leq \alpha$, t > 0 and $x \in \mathbb{Q}_p^n$.

Proof. Set

$$u_{2,h}(x,t,\tau) := \int_{\tau}^{t-h} \int_{\mathbb{Q}_p^n} Z(x-y,t-\theta) f(y,\theta) d^n y d\theta, \ 0 < h < t-\tau.$$

By using a standard reasoning, one shows that

$$\begin{aligned} \frac{\partial u_{2,h}}{\partial t}(x,t,\tau) &= \\ & \int_{\tau}^{t-h} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,t-\theta)}{\partial t} f(y,\theta) d^n y d\theta + \int_{\mathbb{Q}_p^n} Z(x-y,h) f(y,t-h) d^n y d\theta. \end{aligned}$$

This formula can be rewritten as

$$\begin{split} \frac{\partial u_{2,h}}{\partial t}(x,t,\tau) &= \int_{\tau}^{t-h} \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,t-\theta)}{\partial t} \left[f(y,\theta) - f(x,\theta) \right] d^n y d\theta \\ &+ \int_{\tau}^{t-h} f(x,\theta) \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,t-\theta)}{\partial t} d^n y d\theta + \int_{\mathbb{Q}_p^n} Z(x-y,h) \left[f(y,t-h) - f(y,t) \right] d^n y \\ &+ \int_{\mathbb{Q}_p^n} Z(x-y,h) f(y,t) d^n y. \end{split}$$

The first integral contains no singularity at $t = \theta$ due to Lemma 4.8 (iv) and the local constancy of f. By Lemma 4.3 (iv), the second integral is equal to zero. The third integral can be written as a sum of the integrals over $\left\{y \in \mathbb{Q}_p^n \mid ||x - y||_p \ge p^M\right\}$ and the complement of this set, one of these integrals is estimated on the basis of the uniform continuity of f, while the other contains no singularity, see Lemma 4.8 (iv). Finally, the fourth integral tends to f(x,t) as $h \to 0^+$, cf. Lemma 4.7.

(ii) By Lemma 4.5, $\mathbf{W}_{\gamma}u_{2,h}$ is well-defined for any γ satisfying $n + \lambda < \gamma \leq \alpha$. Then, for any $L \in \mathbb{N}$, the following integral exists:

(4.9)
$$\int_{\|\xi\|_{p} \ge p^{-L}} \frac{[u_{2,h}(x-\xi,t,\tau) - u_{2,h}(x,t,\tau)]}{w_{\gamma}(\|\xi\|_{p})} d^{n}\xi$$
$$= \int_{\tau}^{t-h} \int_{\mathbb{Q}_{p}^{n}\|\xi\|_{p} \ge p^{-L}} \frac{[Z(x-\xi-y,t-\theta) - Z(x-y,t-\theta)]}{w_{\gamma}(\|\xi\|_{p})} f(y,\theta) d^{n}\xi d^{n}y d\theta.$$

On the other hand, by Fubini's Theorem,

$$\int_{\|\xi\|_p \ge p^{-L}} \frac{\left[Z(x-\xi-y,t-\theta) - Z(x-y,t-\theta)\right]}{w_{\gamma}(\|\xi\|_p)} d^n \xi$$
$$= \int_{\mathbb{Q}_p^n} \Psi((x-y) \cdot \eta) e^{-\kappa(t-\theta)A_{w_{\gamma}}(\|\eta\|_p)} P_k(\eta) d^n \eta,$$

where

$$P_{k}(\eta) = \int_{\|\xi\|_{p} \ge p^{-L}} \frac{[\Psi(-\xi \cdot \eta) - 1]}{w_{\gamma}(\|\xi\|_{p})} d^{n}\xi.$$

A simple calculation shows that

 $\|\xi\|$

$$|P_k(\eta)| \le C' \, \|\eta\|_p^{\gamma-n} \, ,$$

and then

$$\int_{\|_p \ge p^{-L}} \frac{\left[Z(x-\xi-y,t-\theta) - Z(x-y,t-\theta)\right]}{w_{\gamma}(\|\xi\|_p)} d^n \xi \le C,$$

where the constant does not depend on $x, t \ge h + \tau, L$.

Now, by expressing the right integral of (4.9) as

$$\int_{\tau}^{t-h} \int_{\|x-\xi\|_{p} > p^{-M}} \int_{\|\xi\|_{p} \ge p^{-L}} \frac{\left[Z(x-\xi-y,t-\theta) - Z(x-y,t-\theta)\right]}{w_{\gamma}(\|\xi\|_{p})} f(y,\theta) d^{n}\xi d^{n}y d\theta$$

$$+ \int_{\tau}^{t-h} \int_{\|x-\xi\|_{p} \le p^{-M}} \int_{\|\xi\|_{p} \ge p^{-L}} \frac{\left[Z(x-\xi-y,t-\theta) - Z(x-y,t-\theta)\right]}{w_{\gamma}(\|\xi\|_{p})} f(y,\theta) d^{n}\xi d^{n}y d\theta$$

where M is a positive integer, such that $\|\xi\|_p < p^{-L} < p^{-M} < \|x - \xi\|_p$, and using the same reasoning as in the final part of the proof of Proposition 4.11 (ii), we obtain

(4.10)
$$(\boldsymbol{W}_{\gamma}\boldsymbol{u}_{2,h})(\boldsymbol{x},t) = \int_{\tau}^{t-h} \int_{\mathbb{Q}_p^n} (\boldsymbol{W}_{\gamma}\boldsymbol{Z})(\boldsymbol{x}-\boldsymbol{\xi},t-\theta)f(\boldsymbol{y},\theta)d^n\boldsymbol{\xi}d\theta.$$

Now, by Lemma 4.9 (ii), the fact that $f \in \mathfrak{M}_{\lambda}$, Proposition 4.4, and the Dominated Convergence Theorem, we can take limit as $h \to 0^+$, which completes the proof when $\gamma < \alpha$. If $\gamma = \alpha$, formula (4.10) remains valid. By using Lemma 4.9 (iii), formula (4.10) can be rewritten as

$$(\boldsymbol{W}_{\gamma}\boldsymbol{u}_{2,h})(\boldsymbol{x},t) = \int_{\tau}^{t-h} \int_{\mathbb{Q}_p^n} (\boldsymbol{W}_{\gamma}\boldsymbol{Z})(\boldsymbol{x}-\boldsymbol{\xi},t-\theta) \left[f(\boldsymbol{y},\theta) - f(\boldsymbol{x},\theta)\right] d^n \boldsymbol{\xi} d\theta.$$

Now, by using the local constancy of f, we can justify the passage to the limit as $h \to 0^+$, which completes the proof.

Remark 4.13. By Lemma 4.3 (iv) and Lemma 4.8 (i), $\int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,t-\theta)}{\partial t} d^n y = 0$, then

$$\frac{\partial u_2}{\partial t}(x,t,\tau) = f(x,t) + \int_{\tau}^t \left(\int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y,t-\theta)}{\partial t} f(y,\theta) d^n y \right) d\theta,$$

for t > 0 and $x \in \mathbb{Q}_p^n$.

5. PARABOLIC-TYPE EQUATIONS WITH VARIABLE COEFFICIENTS

First, we fix the notation that will be used through this section. We fix N + 1 positive real numbers satisfying $n < \alpha_1 < \alpha_2 < \cdots < \alpha_N < \alpha$. We fix N + 2 functions $a_k(x,t), k = 0, \ldots, N$ and b(x,t) from $\mathbb{Q}_p^n \times [0,T]$ to \mathbb{R} , here T is a positive constant. We assume that: (i) b(x,t) and $a_k(x,t)$, for $k = 0, \ldots, N$, belong (with respect to x) to \mathfrak{M}_0 uniformly with respect to $t \in [0,T]$; (ii) $a_0(x,t)$ satisfies the Hölder condition in t with exponent $v \in (0,1)$ uniformly in x. We also assume the uniform parabolicity condition $a_0(x,t) \ge \mu > 0$ and that $\alpha_{N+1} := n + (\alpha - n)(1 - v) > \alpha_N$. Notice that $\alpha_{N+1} < \alpha$.

 $v > \alpha_N$. Notice that $\alpha_{N+1} < \alpha$. Set $\widetilde{W} := \sum_{k=1}^N a_k(x,t) \mathbf{W}_{\alpha_k} - b(x,t) \mathbf{I}$ with domain \mathfrak{M}_{λ} , and $0 \le \lambda + n < \alpha_1$. Notice that $\widetilde{W} : \mathfrak{M}_{\lambda} \to \mathfrak{M}_{\lambda}$. In this section we construct a solution for the following initial value problem:

(5.1)
$$\begin{cases} \frac{\partial u}{\partial t}(x,t) - a_0(x,t)(\mathbf{W}_{\alpha}u)(x,t) - (\mathbf{W}_{\alpha}u)(x,t) = f(x,t) \\ u(x,0) = \varphi(x), \end{cases}$$

where $x \in \mathbb{Q}_p^n$, $t \in (0, T]$, $\varphi(x) \in \mathfrak{M}_{\lambda}$, $f(x, t) \in \mathfrak{M}_{\lambda}$ uniformly with respect to t, with $0 \leq \lambda < \alpha_1 - n$, and f(x, t) is continuous in (x, t) (if $a_1(x, t) = \cdots = a_N(x, t) \equiv 0$ then we shall assume that $0 \leq \lambda < \alpha - n$).

5.1. **Parametrized Cauchy problem.** We first study the following Cauchy problem:

(5.2)
$$\begin{cases} \frac{\partial u}{\partial t}(x,t) - a_0(y,\theta)(\mathbf{W}_{\alpha}u)(x,t) = 0, \ x \in \mathbb{Q}_p^n, t \in (0,T] \\ u(x,0) = \varphi(x), \end{cases}$$

where $y \in \mathbb{Q}_p^n$, $\theta > 0$ are parameters. By taking $\kappa = a_0(y, \theta) \ge \mu > 0$ and applying the results of Section 4, Cauchy problem (5.2) has a fundamental solution given by

$$Z(x,t;y,\theta,w_{\alpha},\kappa) := Z(x,t;y,\theta) = \int_{\mathbb{Q}_p^n} \Psi(x\cdot\xi) e^{-a_0(y,\theta)tA_{w_{\alpha}}(\|\xi\|_p)} d^n\xi,$$

for t > 0 and $x \in \mathbb{Q}_p^n$.

Remark 5.1. All statements from the Lemmas 4.3, 4.8, 4.9 hold for $Z(x, t; y, \theta)$ and the involved constants do not depend of y and θ . Thus, we have the following estimates:

(5.3)
$$Z(x,t;y,\theta) \le C_1 t \left(\|x\|_p + t^{\frac{1}{\alpha-n}} \right)^{-\alpha}, \text{ for } t > 0;$$

(5.4)
$$\left|\frac{\partial Z(x,t;y,\theta)}{\partial t}\right| \le C_2 \left(\|x\|_p + t^{\frac{1}{\alpha-n}}\right)^{-\alpha}, \text{ for } t > 0;$$

(5.5)
$$|(\boldsymbol{W}_{\gamma}Z)(x,t;y,\theta)| \le C_3 \left(||x||_p + t^{\frac{1}{\alpha-n}} \right)^{-\gamma}, \text{ for } t > 0 \text{ and } \gamma \le \alpha.$$

And the identities:

/- -->

(5.6)
$$\int_{\mathbb{Q}_p^n} Z(x,t;y,\theta) d^n x = 1, \text{ for } t > 0;$$

~

$$\frac{\partial Z(x,t;y,\theta)}{\partial t} = -a_0\left(y,\theta\right) \int_{\mathbb{Q}_p^n} A_{w_\alpha}(\|\xi\|_p) e^{-a_0\left(y,\theta\right)tA_{w_\alpha}\left(\|\xi\|_p\right)} \Psi(x\cdot\xi) d^n\xi, \text{ for } t > 0;$$

(5.8)
$$(\mathbf{W}_{\gamma}Z)(x,t;y,\theta) = -\int_{\mathbb{Q}_p^n} A_{w_{\gamma}}(\|\xi\|_p) e^{-a_0(y,\theta)tA_{w_{\alpha}}(\|\xi\|_p)} \Psi(x\cdot\xi) d^n\xi,$$

for t > 0 and $\gamma \leq \alpha$;

(5.9)
$$\int_{\mathbb{Q}_p^n} (\boldsymbol{W}_{\gamma} Z_t) (x, t; y, \theta) d^n x = 0.$$

Lemma 5.2. There exists a positive constant C, such that

(5.10)
$$\left| \int_{\mathbb{Q}_p^n} \frac{\partial Z(x-y;t,y,\theta)}{\partial t} d^n y \right| \le C.$$

Proof. See proof Lemma 4.5 in [13].

5.2. Heat potentials. We define the parameterized heat potentials as follows:

$$u(x,t,\tau) := \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} Z(x-y,t-\theta;y,\theta) f(y,\theta) d^{n}y d\theta,$$

where $f \in \mathfrak{M}_{\lambda}$, $0 \leq \lambda < \alpha - n$, f continuous in (y, θ) . By using the same argument given to prove Lemma 4.5, one proves that $u \in \mathfrak{M}_{\lambda}$ uniformly in t and τ .

We now calculate the derivative with respect to t and the action of the operator \mathbf{W}_{γ} on $u(x, t, \tau)$ for $n + \lambda < \gamma \leq \alpha$.

Proposition 5.3. Assume that $f \in \mathfrak{M}_{\lambda}$, $0 \leq \lambda < \alpha - n$, f continuous in (y, θ) . Then the following assertions hold:

$$(i) \ \frac{\partial u(x,t,\tau)}{\partial t} = f(x,t) + \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} \frac{\partial Z(x-y,t-\theta;y,\theta)}{\partial t} f(y,\theta) d^{n} y d\theta;$$

(ii) $(\mathbf{W}_{\gamma}u)(x,t,\tau) = \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} (\mathbf{W}_{\gamma}Z)(x-y,t-\theta;y,\theta) f(y,\theta) d^{n} y d\theta, \ \gamma \leq \alpha.$

Proof. It is a simple variation of the proof given for Proposition 4.12.

The following technical result will be used later on.

Lemma 5.4 ([13], Lemma 4.6). Let

$$J(x,\xi,t,\tau) = \int_{\tau}^{t} (t-\theta)^{-\rho/\beta} (\theta-\tau)^{-\sigma/\beta} \\ \times \left(\int_{\mathbb{Q}_{p}^{n}} \left((t-\theta)^{1/\beta} + \|x-\eta\|_{p} \right)^{-n-b_{1}} \left((\theta-\tau)^{1/\beta} + \|\eta-\xi\|_{p} \right)^{-n-b_{2}} d^{n}\eta \right) d\theta,$$

where $x, \xi \in \mathbb{Q}_p^n, 0 \le \tau < t, b_1, b_2 > 0, \rho + b_1 < \beta, \sigma + b_2 < \beta$. Then

$$J(x,\xi,t,\tau) \leq C\left\{B\left(1-\frac{\rho}{\beta},1-\frac{\sigma+b_2}{\beta}\right)\left((t-\tau)^{1/\beta}+\|x-\xi\|_p\right)^{-n-b_1}(t-\tau)^{-\frac{(\rho+\sigma+b_2-\beta)}{\beta}} + B\left(1-\frac{\rho+b_1}{\beta},1-\frac{\sigma}{\beta}\right)\left((t-\tau)^{1/\beta}+\|x-\xi\|_p\right)^{-n-b_2}(t-\tau)^{-\frac{(\rho+\sigma+b_1-\beta)}{\beta}}\right\},$$

where C is a positive constant depends only on b_{1,b_2} and $B(\cdot, \cdot)$ denotes the Archimedean Beta function.

The proof is a simple variation of that given by Kochubei for Lemma 4.6 in [13].

5.3. Construction of a solution.

Theorem 5.5. The Cauchy problem (5.1), has a solution, which can be represented in the form

(5.11)
$$u(x,t) = \int_{0}^{t} \int_{\mathbb{Q}_{p}^{n}} \Lambda(x,t,\xi,\tau) f(\xi,\tau) d^{n}\xi d\tau + \int_{\mathbb{Q}_{p}^{n}} \Lambda(x,t,\xi,0)\varphi(\xi) d^{n}\xi,$$

where the fundamental solution $\Lambda(x,t,\xi,\tau)$, $x, \xi \in \mathbb{Q}_p^n$, $0 \le \tau < t \le T$, has the form

(5.12)
$$\Lambda(x,t,\xi,\tau) = Z(x-\xi,t-\tau;\xi,\tau) + \mathcal{W}(x,t,\xi,\tau),$$

with

(5.13)
$$|\mathcal{W}(x,t,\xi,\tau)| \leq C \left\{ (t-\tau)^{2-\frac{\lambda}{\alpha-n}} \left[(t-\tau)^{\frac{1}{\alpha-n}} + ||x-\xi||_p \right]^{-\alpha} + (t-\tau) \sum_{k=1}^{N+1} \left[(t-\tau)^{\frac{1}{\alpha-n}} + ||x-\xi||_p \right]^{-\alpha_k} \right\}.$$

Furthermore $Z(x, t; y, \theta)$ satisfies the estimates (5.3), (5.4), (5.5), (5.10).

Proof. We use the usual scheme of Levi's method. Thus, we look for a fundamental solution of (5.1) having form (5.12), with

$$\mathcal{W}(x,t,\xi,\tau) = \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} Z(x-\eta,t-\theta;\eta,\theta) \Phi(\eta,\theta,\xi,\tau) d^{n}\eta d\theta.$$

Then, we require that

$$\frac{\partial \Lambda}{\partial t}(x,t,\xi,\tau) - a_0(x,t)(\mathbf{W}_{\alpha}\Lambda)(x,t,\xi,\tau) - \sum_{k=1}^N a_k(x,t)(\mathbf{W}_{\alpha_k}\Lambda)(x,t,\xi,\tau) + b(x,t)\Lambda(x,t,\xi,\tau) = 0,$$
(5.14)

$$\begin{split} \frac{\partial Z}{\partial t}(x-\xi,t-\tau,\xi,\tau) &+ \Phi(x,t,\xi,\tau) + \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} \frac{\partial Z(x-\eta,t-\theta;\eta,\theta)}{\partial t} \Phi(\eta,\theta,\xi,\tau) d^{n} \eta d\theta \\ &- a_{0}(x,t) \big\{ (\mathbf{W}_{\alpha}Z)(x-\xi,t-\tau;\xi,\tau) + \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} (\mathbf{W}_{\alpha}Z)(x-\eta,t-\theta;\eta,\theta) \times \\ &\Phi(\eta,\theta,\xi,\tau) d^{n} \eta d\theta \big\} - \sum_{k=1}^{N} a_{k}(x,t) \big\{ (\mathbf{W}_{\alpha_{k}}Z)(x-\xi,t-\tau;\xi,\tau) \\ &+ \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} (\mathbf{W}_{\alpha_{k}}Z)(x-\eta,t-\theta;\eta,\theta) \Phi(\eta,\theta,\xi,\tau) d^{n} \eta d\theta \big\} \\ &+ b(x,t) \big\{ Z(x-\xi,t-\tau;\xi,\tau) + \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} Z(x-\eta,t-\theta;\eta,\theta) \Phi(\eta,\theta,\xi,\tau) d^{n} \eta d\theta \big\} = 0. \end{split}$$

By taking

$$R(x,t,\xi,\tau) := (a_0(x,t) - a_0(\xi,\tau))(\mathbf{W}_{\alpha}Z)(x-\xi,t-\tau;\xi,\tau) + \sum_{k=1}^N a_k(x,t)(\mathbf{W}_{\alpha_k}Z)(x-\xi,t-\tau;\xi,\tau) - b(x,t)Z(x-\xi,t-\tau;\xi,\tau),$$

one gets that $\Phi(x, t, \xi, \tau)$ satisfies the integral equation

(5.15)
$$\Phi(x,t,\xi,\tau) = R \ (x,t,\xi,\tau) + \int_{\tau}^{t} \int_{\mathbb{Q}_p^n} R(x,t,\eta,\theta) \Phi(\eta,\theta,\xi,\tau) d^n \eta d\theta.$$

Now, by using (5.5) and (5.3), we obtain

$$|R(x,t,\xi,\tau)| \leq C_0 \left(|a_0(x,t) - a_0(\xi,\tau)| \left((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p \right)^{-\alpha} + \sum_{k=1}^N \left((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p \right)^{-\alpha_k} + \left((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p \right)^{-\alpha} (t-\tau) \right).$$

$$(6) \qquad \qquad + \left((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p \right)^{-\alpha} (t-\tau) \right).$$

(5.16)

Claim A.

 $|a_0(x,t) - a_0(\xi,\tau)| \left((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p \right)^{-\alpha} \le C_1' \left((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p \right)^{-\alpha_{N+1}},$ where $\alpha_{N+1} = n + (\alpha-n) (1-v) > \alpha_N.$

Indeed, by the Hölder condition

$$|a_0(x,t) - a_0(\xi,\tau)| \le C_1(t-\tau)^v + |a_0(x,\tau) - a_0(\xi,\tau)|.$$

Let $l(a_0)$ be the parameter of local constancy of a_0 . Thus, if $||x - \xi||_p \le p^{l(a_0)}$, then $|a_0(x,t) - a_0(\xi,\tau)| \le C_1(t-\tau)^v$. In the case $||x - \xi||_p \le p^{l(a_0)}$, the inequality follows from the fact that $(t-\tau)^v((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p)^{-\alpha+\alpha_{N+1}}$ is bounded, which in turn follows from $\lim_{t\to\tau}(t-\tau)^{v+\frac{-\alpha+\alpha_{N+1}}{\alpha-n}} = 1$. In the case $\|x-\xi\|_p > p^{l(a_0)}$, taking $|a_0(x,t) - a_0(\xi,\tau)| \le C_0$, the inequality follows from

$$\left((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p\right)^{-\alpha+\alpha_{N+1}} \le \|x-\xi\|_p^{-\alpha+\alpha_{N+1}} \le p^{(-\alpha+\alpha_{N+1})l(a_0)}.$$

Claim B.

$$(t-\tau)\left((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p\right)^{-\alpha} \le C_2\left((t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p\right)^{-\alpha_{N+1}}$$

This assertion is a consequence of the fact that $\lim_{t\to\tau} (t-\tau)^{1+\frac{-\alpha+\alpha_{N+1}}{\alpha-n}} = 0$. Now from (5.16), and Claims A-B, we have

(5.17)
$$|R(x,t,\xi,\tau)| \le C \sum_{k=1}^{N+1} \left[(t-\tau)^{\frac{1}{\alpha-n}} + ||x-\xi||_p \right]^{-\alpha_k}$$

We solve integral equation (5.15) by the method of successive approximations:

(5.18)
$$\Phi(x,t,\xi,\tau) = \sum_{m=1}^{\infty} R_m(x,t,\eta,\theta),$$

where $R_1 \equiv R$ and

$$R_{m+1}(x,t,\xi,\tau) = \int_{\tau}^{t} \int_{\mathbb{Q}_p^n} R(x,t,\eta,\theta) R_m(\eta,\theta,\xi,\tau) d^n \eta d\theta, \text{ for } m \ge 1.$$

Claim C.

$$|R_{m+1}(x,t,\xi,\tau)| \le C(2N+2)^m (t-\tau)^{mv} \frac{(\Gamma(v))^{m+1}}{\Gamma((m+1)v)} \times \sum_{j=1}^{N+1} \left[(t-\tau)^{\frac{1}{\alpha-n}} + \|x-\xi\|_p \right]^{-\alpha_j},$$

for $m \geq 0$, where $\Gamma(\cdot)$ denotes the Archimedean Gamma function.

The proof of this assertion will be given later.

It follows from Claim A, by the Stirling formula, that series (5.18) is convergent and that

(5.19)
$$|\Phi(x,t,\xi,\tau)| \le C_0 \sum_{k=1}^{N+1} \left[(t-\tau)^{\frac{1}{\alpha-n}} + ||x-\xi||_p \right]^{-\alpha_k}.$$

Now (5.13) follows from (5.19) and Lemma 5.4.

Denote by $u_1(x,t)$ and $u_2(x,t)$ the first and second terms in the right hand side of (5.11). Substituting (5.12) into (5.11), we find that

$$u_1(x,t) = \int_0^t \int_{\mathbb{Q}_p^n} Z(x-\xi,t-\tau;\xi,\tau) f(\xi,\tau) d^n \xi d\tau + \int_0^t \int_{\mathbb{Q}_p^n} Z(x-\eta,t-\theta;\eta,\theta) F(\eta,\theta) d^n \eta d\theta,$$

and

$$u_2(x,t) = \int_{\mathbb{Q}_p^n} Z(x-\xi,t;\xi,0)\varphi(\xi)d^n\xi + \int_0^t \int_{\mathbb{Q}_p^n} Z(x-\eta,t-\theta;\eta,\theta)G(\eta,\theta)d^n\eta d\theta,$$

where

(5.20)
$$F(\eta,\theta) = \int_{0}^{\theta} \int_{\mathbb{Q}_p^n} \Phi(\eta,\theta,\xi,\tau) f(\xi,\tau) d^n \xi d\tau$$

0

(5.21)
$$G(\eta,\theta) = \int_{\mathbb{Q}_p^n} \Phi(\eta,\theta,\xi,0)\varphi(\xi)d^n\xi.$$

Now, by Proposition 4.4 and (5.19), it follows that

$$|F(\eta,\theta)| \le C_0(1+\|\eta\|_p^{\lambda}), \quad |G(\eta,\theta)| \le C_1(1+\|\eta\|_p^{\lambda}),$$

for all $\eta \in \mathbb{Q}_p^n$ and $\theta \in (0,T]$.

Claim D. The functions F and G belong to $\tilde{\mathcal{E}}$, and their parameters of local constancy do not depend on θ .

We first note that by (5.20)-(5.21), it is sufficient to show that $\Phi(\cdot, \theta, \star, \tau)$ is a locally constant function on $(\mathbb{Q}_p^{\times})^n \times \mathbb{Q}_p^n$ and that its parameter of local constancy do not depend on θ and τ . Now, by the recursive definition of the function Φ we see that if L is the parameter local constancy for all the functions $a_k(\cdot, t)$, $b(\cdot, t)$, $(\mathbf{W}_{\alpha_k}Z)(\cdot, t - \tau; \star, \tau)$ and $Z(\cdot, t - \tau; \star, \tau)$ on $(\mathbb{Q}_p^{\times})^n \times \mathbb{Q}_p^n$, and if $\|\delta\|_p \leq p^{-L}$, we have

$$R(x+\delta, t, \xi+\delta, \tau) = R(x, t, \xi, \tau).$$

Furthermore, we successively obtain

$$R_{m+1}(x+\delta,t,\xi+\delta,\tau) = \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} R(x+\delta,t,\eta,\theta) R_{m}(\eta,\theta,\xi+\delta,\tau) d^{n} \eta d\theta$$
$$= \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} R(x+\delta,t,\zeta+\delta,\theta) R_{m}(\zeta+\delta,\theta,\xi+\delta,\tau) d^{n} \zeta d\theta$$
$$= R_{m+1}(x,t,\xi,\tau),$$

so that $\Phi(x+\delta,t,\xi+\delta,\tau) = \Phi(x,t,\xi,\tau)$, and hence

$$\begin{split} F(\eta+\delta,\theta) &= \int_{0}^{\theta} \int_{\mathbb{Q}_{p}^{n}} \Phi(\eta+\delta,\theta,\xi,\tau) f(\xi,\tau) d^{n}\xi d\tau \\ &= \int_{0}^{\theta} \int_{\mathbb{Q}_{p}^{n}} \Phi(\eta+\delta,\theta,\xi+\delta,\tau) f(\xi+\delta,\tau) d^{n}\xi d\tau = F(\eta,\theta). \end{split}$$

Similarly, $G(\eta + \delta, \theta) = G(\eta, \theta)$ when $|\delta|_p \leq p^{-L}$. Thus $u_1(x, t), u_2(x, t) \in \mathfrak{M}_{\lambda}$ uniformly in t. Thus the potentials in the expressions for $u_1(x, t), u_2(x, t)$ satisfy the conditions to use the differentiation formulas given in Proposition 5.3. By using these formulas along with Proposition 5.3, (5.7)-(5.8) and (5.15), ones verifies after simple transformations that u(x, t) is a solution of Cauchy problem (5.1).

Let us show that $u(x,t) \to \varphi(x)$ as $t \to 0^+$. Due to (5.12) and (5.13), it is sufficient to verify that

$$v(x,t) := \int_{\mathbb{Q}_p^n} Z(x-\xi,t;\xi,0)\varphi(\xi)d^n\xi \to \varphi(x) \text{ as } t \to 0^+.$$

By virtue of formula (5.6), we have

$$\begin{aligned} v(x,t) &= \int_{\mathbb{Q}_p^n} \left[Z(x-\xi,t;\xi,0) - Z(x-\xi,t;x,0) \right] \varphi(\xi) d^n \xi \\ &+ \int_{\mathbb{Q}_p^n} Z(x-\xi,t;x,0) \left[\varphi(\xi) - \varphi(x) \right] d^n \xi + \varphi(x). \end{aligned}$$

Now, since $Z(x - \xi, t; \cdot, 0)$ and $\varphi(\cdot)$ are locally constant functions, it follows that in both integrals the integration is actually performed over the set

$$\left\{\xi \in \mathbb{Q}_p^n : \left\|\xi - x\right\|_p \ge p^{-L}\right\}.$$

By applying (5.3) on this set, we see that both integrals tend to zero as $t \to 0^+$. **Proof of Claim C.** We use induction on m. The case m = 0 is (5.17). We assume the case m, then

$$\begin{aligned} |R_{m+1}(x,t,\xi,\tau)| &\leq \int_{\tau}^{t} \int_{\mathbb{Q}_{p}^{n}} |R(x,t,\eta,\theta)| \left| R_{m}(\eta,\theta,\xi,\tau) \right| d^{n}\eta d\theta \\ &\leq C_{0}(2N+2)^{m-1} \frac{(\Gamma(v))^{m}}{\Gamma(mv)} \sum_{j,k=1}^{N+1} \int_{\tau}^{t} (\theta-\tau)^{(m-1)v} \times \\ &\int_{\mathbb{Q}_{p}^{n}} \left[(\theta-\tau)^{\frac{1}{\alpha-n}} + \|\eta-\xi\|_{p} \right]^{-\alpha_{j}} \left[(t-\theta)^{\frac{1}{\alpha-n}} + \|x-\eta\|_{p} \right]^{-\alpha_{k}} d^{n}\eta d\theta. \end{aligned}$$

Now by Lemma 5.4, with $-\sigma = (m-1)(\alpha - n)v$, $\rho = 0$, $-n - b_2 = -\alpha_j$, $-n - b_1 = -\alpha_k$, $\beta = \alpha - n$, we have

$$\begin{aligned} &|R_{m+1}(x,t,\xi,\tau)| \\ &\leq C_0 \left\{ (2N+2)^{m-1} \frac{(\Gamma(v))^m}{\Gamma(mv)} \sum_{j,k=1}^{N+1} B\left(1, \frac{\alpha+(m-1)(\alpha-n)v-\alpha_j}{\alpha-n}\right) \times \\ &\left((t-\tau)^{1/(\alpha-n)} + \|x-\xi\|_p\right)^{-\alpha_k} (t-\tau)^{\frac{(m-1)(\alpha-n)v-\alpha_j+\alpha)}{\alpha-n}} \\ &+ B\left(\frac{\alpha-\alpha_k}{\alpha-n}, \frac{\alpha-n+(m-1)(\alpha-n)v}{\alpha-n}\right) \left((t-\tau)^{1/(\alpha-n)} + \|x-\xi\|_p\right)^{-\alpha_j} \times \\ &\left(t-\tau\right)^{\frac{(m-1)(\alpha-n)v-\alpha_k+\alpha)}{\alpha-n}} \right\}. \end{aligned}$$

By using $B(z_1 + \epsilon, z_2 + \delta) \leq B(z_1, z_2)$, for $\epsilon, \delta \geq 0$,

$$B\left(1, \frac{\alpha + (m-1)(\alpha - n)v - \alpha_j}{\alpha - n}\right) \le B(v, mv),$$

$$B\left(\frac{\alpha - \alpha_k}{\alpha - n}, \frac{\alpha - n + (m-1)(\alpha - n)v}{\alpha - n}\right) \le B(v, mv),$$

and

$$(t-\tau)^{\frac{(m-1)(\alpha-n)v-\alpha_r+\alpha)}{\alpha-n}} = (t-\tau)^{mv-\frac{(\alpha-n)v+\alpha_r-\alpha}{\alpha-n}} \le C(t-\tau)^{mv},$$

for $1 \leq r \leq N+1$, we get

$$R_{m+1}(x,t,\xi,\tau) \leq C(2N+2)^m \frac{(\Gamma(v))^m}{\Gamma((m+1)v)} (t-\tau)^{mv} \\ \times \sum_{k=1}^{N+1} \left((t-\tau)^{1/(\alpha-n)} + \|x-\xi\|_p \right)^{-\alpha_k}.$$

6. UNIQUENESS OF THE SOLUTION

We recall that $\widetilde{\mathcal{E}}$ is the \mathbb{C} -vector space of all functions $\varphi : \mathbb{Q}_p^n \to \mathbb{C}$, such that there exist a ball B_l^n , with l depending only on φ , and $\varphi(x+x') = \varphi(x)$ for any $x' \in B_l^n$. Notice that $\mathfrak{M}_{\lambda} \subset \widetilde{\mathcal{E}}$ for any λ . We identify any element of $\widetilde{\mathcal{E}}$ with a distribution on \mathbb{Q}_p^n . We now recall the following fact: $T \in S'$ with $supp(T) \subset B_N^n$ if only if $\widehat{T} \in \widetilde{\mathcal{E}}$ and its parameter of local constancy is greater than -N, cf. [22, pg 109] or [19, Proposition 3.17].

Lemma 6.1. $\mathbf{W}_{\alpha}: \widetilde{\mathcal{E}} \to \widetilde{\mathcal{E}}$ is a well-defined linear operator. Furthermore,

$$(\mathbf{W}_{\alpha}\varphi)(x) = -\mathcal{F}_{\xi \to x}^{-1} \left(A_{w_{\alpha}}(\|\xi\|_{p})\mathcal{F}_{x \to \xi}\varphi \right).$$

Proof. Let l be a parameter of locally constancy of φ , then

$$\begin{aligned} (\mathbf{W}_{\alpha}\varphi)(x) &= \int\limits_{\|y\|_{p} \ge p^{l}} \frac{\varphi(x-y) - \varphi(x)}{w_{\alpha}(\|y\|_{p})} d^{n}y \\ &= \frac{1_{\mathbb{Q}_{p}^{n} \smallsetminus B_{l}^{n}}(x)}{w_{\alpha}\left(\|x\|_{p}\right)} * \varphi\left(x\right) - \varphi\left(x\right) \left(\int\limits_{\|y\|_{p} \ge p^{l}} \frac{d^{n}y}{w_{\alpha}(\|y\|_{p})}\right). \end{aligned}$$

Then by taking the Fourier transform in S':

$$\mathcal{F}(\mathbf{W}_{\alpha}\varphi)(\xi) = \left(\int_{\mathbb{Q}_{p}^{n}} 1_{\mathbb{Q}_{p}^{n} \smallsetminus B_{l}^{n}}\left(x\right) \frac{\left(\Psi(x \cdot \xi) - 1\right)}{w_{\alpha}(\|x\|_{p})} d^{n}x\right) (\mathcal{F}\varphi)(\xi),$$

and since $\mathcal{F}\varphi \in S'$ with $supp(\mathcal{F}\varphi) \subset B^n_{-l}$,

$$\mathcal{F}(\mathbf{W}_{\alpha}\varphi)(\xi) = \left(\int_{\mathbb{Q}_{p}^{n}} \frac{(\Psi(x\cdot\xi)-1)}{w_{\alpha}(\|x\|_{p})} d^{n}x\right) \mathcal{F}\varphi(\xi).$$

Therefore,

$$(\mathbf{W}_{\alpha}\varphi)(x) = -\mathcal{F}_{\xi \to x}^{-1} \left(A_{w_{\alpha}}(\|\xi\|_{p}) \mathcal{F}_{x \to \xi}\varphi \right) \in \widetilde{\mathcal{E}}.$$

Take γ be a real number such that $\lambda < \gamma < \alpha_1 - n < \ldots < \alpha_N - n < \alpha - n$, and fix a integer L, and set $\psi(x) := p^{Ln} \Omega(p^L ||x||_p) * ||x||_p^{\gamma}$, then

$$\psi(x) = \begin{cases} \|x\|_{p}^{\gamma} & \text{if } \|x\|_{p} > p^{-L} \\ \\ C & \text{if } \|x\|_{p} \le p^{-L} \end{cases}$$

and thus $\psi \in \widetilde{\mathcal{E}}$.

Lemma 6.2. With above notation, there exist positive constants C_1 and C_2 such that (i) $|(\mathbf{W}_{\alpha}\psi)(x)| \leq C_1 ||x||_p^{\alpha-\gamma+n}$ and (ii) $|(\mathbf{W}_{\alpha_k}\psi)(x)| \leq C_2 ||x||_p^{\alpha_k-\gamma+n}$, for $k=1,\ldots,N.$

Proof. By Lemma 6.1,

$$(\mathbf{W}_{\alpha}\psi)(x) = -\mathcal{F}_{\xi \to x}^{-1} \left(A_{w_{\alpha}}(\|\xi\|_{p}) \Omega(p^{-L} \|\xi\|_{p}) \frac{\|\xi\|_{p}^{-\gamma-n}}{\Gamma_{n}(n+\gamma)} \right) \text{ in } S',$$

where $\Gamma_n(n+\gamma) = \frac{1-p^{\gamma}}{1-p^{-\gamma-n}}$. Now, since $A_{w_{\alpha}}(\|\xi\|_p)\Omega(p^{-L}\|\xi\|_p)\frac{\|\xi\|_p^{-\gamma-n}}{\Gamma_n(n+\gamma)}$ is radial and locally integrable, by applying the formula for Fourier transform of radial function,

$$\begin{aligned} (\mathbf{W}_{\alpha}\varphi)(x) &= \\ &\frac{-\|x\|_{p}^{-n}}{\Gamma_{n}(n+\gamma)} [(1-p^{-n}) \|x\|_{p}^{\gamma+n} \sum_{j=0}^{\infty} A_{w_{\alpha}}(\|x\|_{p}^{-1} p^{-j}) \Omega(\|x\|_{p}^{-1-j} p^{-j}) p^{j(\gamma+n)-jn} \\ &- A_{w_{\alpha}}(\|x\|_{p} p^{-j}) \Omega(\|x\|_{p}^{-1} p^{-L+1}) \|x\|_{p}^{\gamma+n}], \end{aligned}$$

as a distribution on $\mathbb{Q}_p^n \setminus \{0\}$, now by using Lemma 3.4 in [8],

$$|(\mathbf{W}_{\alpha}\varphi)(x)| \leq C' \left[(1-p^{-n}) \sum_{j=0}^{\infty} p^{-j(\alpha-n)+j\gamma} - p^{(-L+1)(\alpha-n)} \right] ||x||_{p}^{-\alpha+n+\gamma}.$$
proof of (ii) is similar.

The proof of (ii) is similar.

Theorem 6.3. Assume that the coefficients $a_k(x,t)$, $k = 0, 1, \dots, N$ are nonnegative bounded continuous functions, b(x,t) is a bounded continuous function, $0 \leq \lambda < \alpha_1 - n$ (if $a_1(x, t) = \cdots = a_k(x, t) \equiv 0$, we shall suppose that $0 \leq \lambda < \alpha - n$) and u(x,t) is a solution of Cauchy problem (5.1) with $f(x,t) = \varphi(x) \equiv 0$ that belongs to class \mathfrak{M}_{λ} . Then $u(x,t) \equiv 0$.

Proof. We may assume that $b(x,t) \geq 0$, otherwise we take $u(x,t)e^{\lambda t}$ with $\lambda > 0$ b(x,t). We prove that $u(x,t) \ge 0$. By contradiction, suppose that u(x',t') < 0, for some $x' \in \mathbb{Q}_p^n$ and $t' \in (0,T]$. By Lemma 6.2, it follows that $(\mathbf{W}_{\alpha}\psi)(x)$ and $(\mathbf{W}_{\alpha_k}\psi)(x) \to 0$ as $\|x\|_p \to \infty$, and thus

$$M := \sup_{\substack{0 \le t \le T, \\ x \in \mathbb{Q}_p^n}} \left\{ a_0(x,t) \left| (\mathbf{W}_{\alpha}\psi)(x) \right| + \sum_{k=1}^N a_k(x,t) \left| (\mathbf{W}_{\alpha_k}\psi)(x) \right| \right\} < \infty.$$

We pick $\rho > 0$ such that $u(x', t') + T\rho < 0$, and then $\sigma > 0$ such that

(6.1)
$$u(x',t') + T\rho + \sigma\psi(x') < 0$$

$$(6.2) \qquad \qquad \rho - \sigma M < 0.$$

We now consider the function

$$v(x,t) := u(x,t) + t\rho + \sigma\psi(x).$$

From (6.1), it follows that v(x', t') < 0, so that

$$\inf_{\substack{0 \le t \le T, \\ x \in \mathbb{Q}_p^n}} v\left(x, t\right) < 0$$

On the other hand, since $u(x,t) \in \mathfrak{M}_{\lambda}$, $\lim_{\|x\|_{p}\to\infty} \frac{u(x,t)}{\psi(x)} = 0$ and thus $\lim_{\|x\|_{p}\to\infty} v(x,t) > 0$ for any t > 0. This implies that there exist $x_{0} \in \mathbb{Q}_{p}^{n}$ and $t_{0} \in (0,T]$, such that

$$\inf_{\substack{0 \leq t \leq T, \\ x \in \mathbb{Q}_p^n}} v\left(x, t\right) = \min_{\substack{0 \leq t \leq T, \\ x \in \mathbb{Q}_p^n}} v(x, t) = v(x_0, t_0) < 0$$

and thus, by formula (3.1), $(\mathbf{W}_{\alpha}v)(x_0,t_0) \geq 0, (\mathbf{W}_{\alpha_k}v)(x_0,t_0) \geq 0$ for all k, and $\frac{\partial v}{\partial t}(x_0,t_0) \leq 0$, hence

$$\frac{\partial v}{\partial t}(x_0, t_0) - a_0(x, t)(\mathbf{W}_{\alpha} v)(x_0, t_0) - \sum_{k=1}^N a_k(x, t)(\mathbf{W}_{\alpha_k} v)(x_0, t_0) + b(x, t)v(x_0, t_0) < 0.$$

Now, by (6.2),

$$\begin{aligned} \frac{\partial v}{\partial t}(x,t) &- a_0(x,t)(\mathbf{W}_{\alpha}v)(x,t) - \sum_{k=1}^N a_k(x,t)(\mathbf{W}_{\alpha_k}v)(x,t) + b(x,t)v(x,t) \\ &= \rho - \sigma \left[a_0(x,t)(\mathbf{W}_{\alpha}\psi)(x) + \sum_{k=1}^N a_k(x,t)(\mathbf{W}_{\alpha_k}\psi)(x) \right] + b(x,t)\left[\rho t + \sigma \psi(x)\right] \\ &\geq \rho - \sigma M > 0 \end{aligned}$$

We have obtained a contradiction, thus $u(x,t) \ge 0$. Finally taking -u(x,t) instead of u(x,t), we conclude that $u(x,t) \equiv 0$.

7. Markov Processes

In this section we show that the fundamental solution $\Lambda(x, t, \xi, \tau)$ of Cauchy problem (5.1) is the transition density of a Markov process. We need some preliminary results.

Lemma 7.1. If the coefficients $a_k(x,t)$ and b(x,t) are nonnegative, then

$$\Lambda(x, t, \xi, \tau) \ge 0$$

Proof. It is sufficient to show that

$$u(x,t) = \int_{\mathbb{Q}_p^n} \Lambda(x,t,\xi,\tau) \varphi(\xi) d^n \xi \ge 0,$$

where u(x,t) is the solution of Cauchy problem (5.1) with $f(x,t) \equiv 0$, and initial condition $u(x,0) = \varphi(x) \ge 0$ with $\varphi \in S(\mathbb{Q}_p^n)$. From (5.12), (5.13), and Lemma 4.3 (iii), it follows that

(7.1)
$$u(x,t) \to 0 \text{ as } ||x||_p \to \infty.$$

Now, if u(x,t) < 0, then there exist $x_0 \in \mathbb{Q}_p^n$ and $t_0 \in (0,T]$ such that

(7.2)
$$\inf_{\substack{0 \le t \le T, \\ x \in \mathbb{Q}_p^n}} u(x,t) = u(x_0,t_0) < 0.$$

This implies that $(\mathbf{W}_{\alpha}u)(x_0, t_0) \ge 0$, $(\mathbf{W}_{\alpha_k}u)(x_0, t_0) \ge 0$ for all k, and $\frac{\partial u}{\partial t}(x_0, t_0) \le 0$. On the other hand,

$$\frac{\partial u}{\partial t}(x,t) - a_0(x,t)(\mathbf{W}_{\alpha}u)(x,t) - \sum_{k=1}^N a_k(x,t)(\mathbf{W}_{\alpha_k}u)(x,t) = 0.$$

By using the uniform parabolicity condition $a_0(x,t) \ge \mu > 0$, we get $(\mathbf{W}_{\alpha}u)(x_0,t_0) = 0$, then by (3.1), $u(x,t_0)$ is constant, and by (7.1), $u(x,t_0) \equiv 0$, which contradicts (7.2).

Lemma 7.2. If $b(x,t) \equiv 0$, then

$$\int_{\mathbb{Q}_p^n} \Lambda(x, t, \xi, \tau) d^n \xi = 1.$$

Proof. By integrating (5.14) in the variable ξ over whole the space \mathbb{Q}_p^n , and by using Lemma 4.9 (iii), we have

$$\frac{\partial}{\partial t} \left(\int\limits_{\mathbb{Q}_p^n} \Lambda(x, t, \xi, \tau) d^n \xi \right) = 0,$$

thus $\int_{\mathbb{Q}_p^n} \Lambda(x, t, \xi, \tau) d^n \xi$ is independent of t. Now, by integrating (5.12) over whole space \mathbb{Q}_p^n in variable ξ and by using Lemma 4.3 (iv), we have

$$\int_{\mathbb{Q}_p^n} \Lambda(x,t,\xi,\tau) d^n \xi = 1 + \int_{\tau}^t \int_{\mathbb{Q}_p^n} \int_{\mathbb{Q}_p^n} Z(x-\eta,t-\theta,\eta,\theta) \phi(\eta,\theta,\xi,\tau) d^n \eta d^n \xi d\theta.$$

The result is obtained by taking $t = \tau$ in the above formula.

Lemma 7.3. If $b(x,t) \equiv 0$ and $f(x,t) \equiv 0$, then the function $\Lambda(x,t,\xi,\tau)$ satisfies the following property:

(7.3)
$$\Lambda(x,t,\xi,\tau) = \int_{\mathbb{Q}_p^n} \Lambda(x,t,y,\sigma) \Lambda(y,\sigma,\xi,\tau) d^n y.$$

Proof. Consider the following initial value problem:

(7.4)
$$\begin{cases} \frac{\partial u}{\partial t}(x,t) - a_0(x,t)(\mathbf{W}_{\alpha}u)(x,t) - (\widetilde{\mathbf{W}}u)(x,t) = 0\\ u(x,\tau) = \varphi(x), \ x \in \mathbb{Q}_p^n \text{ and } t \in (\tau,\sigma], \end{cases}$$

by Theorem 5.5, $u(x,\sigma) = \int_{\mathbb{Q}_n^n} \Lambda(x,\sigma,\xi,\tau) \varphi(\xi) d^n \xi$. Now consider

(7.5)
$$\begin{cases} \frac{\partial u}{\partial t}(x,t) - a_0(x,t)(\mathbf{W}_{\alpha}u)(x,t) - (\widetilde{\mathbf{W}}u)(x,t) = 0\\ u(x,\sigma) = \int_{\mathbb{Q}_p^n} \Lambda(x,\sigma,\xi,\tau)\varphi(\xi)d^n\xi, \ x \in \mathbb{Q}_p^n, \ t \in (\sigma,T], \ \text{with} \ \tau < \sigma < T, \end{cases}$$

by Theorem 5.5 and Fubini's Theorem, the solution of (7.5) is given by

$$u(x,t) = \int_{\mathbb{Q}_p^n} \left(\int_{\mathbb{Q}_p^n} \Lambda(x,t,y,\sigma) \Lambda(y,\sigma,\xi,\tau) d^n y \right) \varphi(\xi) d^n \xi.$$

On the other hand, (7.5) is equivalent to

(7.6)
$$\begin{cases} \frac{\partial u}{\partial t}(x,t) - a_0(x,t)(\mathbf{W}_{\alpha}u)(x,t) - (\widetilde{\mathbf{W}}u)(x,t) = 0\\ u(x,\tau) = \varphi(x), \quad x \in \mathbb{Q}_p^n \ , \ t \in (\tau,T] \ , \end{cases}$$

which has solution given by $u(x,t) = \int_{\mathbb{Q}_p^n} \Lambda(x,t,\xi,\tau) \varphi(\xi) d^n \xi$. Now, by Theorem 6.3,

$$\int_{\mathbb{Q}_p^n} \Lambda(x,t,\xi,\tau)\varphi(\xi)d^n\xi = \int_{\mathbb{Q}_p^n} \left(\int_{\mathbb{Q}_p^n} \Lambda(x,t,y,\sigma)\Lambda(y,\sigma,\xi,\tau)d^ny \right) \varphi(\xi)d^n\xi,$$

for any test function φ , which implies (7.3).

Theorem 7.4. If the coefficients $a_k(x,t)$, $k = 1, \dots, N$ are nonnegative bounded continuous functions, $b(x,t) \equiv 0$, $0 \leq \lambda < \alpha_1 - n$ (if $a_1(x,t) = \dots a_k(x,t) \equiv 0$, we shall suppose that $0 \leq \lambda < \alpha - n$), and $f(x,t) \equiv 0$, then the fundamental solution $\Lambda(x,t,\xi,\tau)$ is the transition density of a bounded right-continuous Markov process without second kind discontinuities.

Proof. The result follows from [10, Theorem 3.6] by using Lemmas (7.1)-(7.2)-(7.3), and (5.12)-(5.13), and Lemma 4.3 (iii).

8. The Cauchy Problem is Well-Posed

In this section, we study the continuity of the solution of Cauchy problem (5.1) with respect to $\varphi(x)$ and f(x,t). We assume that the coefficients $a_k(x,t)$, $k = 0, 1, \ldots, N$ are nonnegative bounded continuous functions, b(x,t) is a bounded continuous function, $0 \leq \lambda < \alpha_1 - n$ (if $a_1(x,t) = \ldots = a_k(x,t) \equiv 0$, we shall suppose that $0 \leq \lambda < \alpha - n$), $\varphi(x) \in \mathfrak{M}_{\lambda}$ and $f(x,t) \in \mathfrak{M}_{\lambda}$, uniformly in t, with $0 \leq \lambda < \alpha_1 - n$.

We identify \mathfrak{M}_{λ} with the \mathbb{R} -vector space of all the functions " $\phi(x,t) \in \mathfrak{M}_{\lambda}$, uniformly in t," and introduce on \mathfrak{M}_{λ} the following norm:

$$\|\phi\|_{\mathfrak{M}_{\lambda}} := \sup_{t \in [0,T]} \sup_{x \in \mathbb{Q}_{p}^{n}} \left| \frac{\phi\left(x,t\right)}{1 + \|x\|_{p}^{\lambda}} \right|.$$

From now on, we consider \mathfrak{M}_{λ} as topological vector space with the topology induced by $\|\cdot\|_{\mathfrak{M}_{\lambda}}$. We also consider $\mathfrak{M}_{\lambda} \times \mathfrak{M}_{\lambda}$ as topological vector space with the topology induced by the norm $\|\cdot\|_{\mathfrak{M}_{\lambda}} + \|\star\|_{\mathfrak{M}_{\lambda}}$.

Theorem 8.1. With the above hypotheses, consider the following operator: $\mathfrak{M}_{\lambda} \times \mathfrak{M}_{\lambda} \qquad \underline{L} \qquad \mathfrak{M}_{\lambda}$

$$(\varphi(x), f(x,t)) \rightarrow u(x,t),$$

where u(x,t) is given by (5.11). Then $||u(x,t)||_{\mathfrak{M}_{\lambda}} \leq C(||\varphi(x)||_{\mathfrak{M}_{\lambda}} + ||f(x,t)||_{\mathfrak{M}_{\lambda}})$, i.e. L is a continuous operator.

Proof. We write $u(x,t) = u_1(x,t) + u_2(x,t)$ where

$$u_1(x,t) = \int_0^t \int_{\mathbb{Q}_p^n} \Lambda(x,t,\xi,\tau) f(\xi,\tau) d^n \xi d\tau \text{ and } u_2(x,t) = \int_{\mathbb{Q}_p^n} \Lambda(x,t,\xi,0) \varphi(\xi) d^n \xi$$

as before. Now

$$\begin{aligned} |u_1(x,t)| &\leq \int_0^t \int_{\mathbb{Q}_p^n} |\Lambda(x,t,\xi,\tau)| \left| f(\xi,\tau) \right| d^n \xi d\tau \\ &\leq \|f(x,t)\|_{\mathfrak{M}_\lambda} \left\{ \int_0^t \int_{\mathbb{Q}_p^n} |\Lambda(x,t,\xi,\tau)| d^n \xi d\tau + \int_0^t \int_{\mathbb{Q}_p^n} |\Lambda(x,t,\xi,\tau)| \left\| \xi \right\|_p^\lambda d^n \xi d\tau \right\}, \end{aligned}$$

by (5.12)-(5.13), (5.3) and Proposition 4.4,

$$\begin{aligned} |u_{1}(x,t)| &\leq C_{0} \, \|f\left(x,t\right)\|_{\mathfrak{M}_{\lambda}} \left\{ \int_{0}^{t} (t-\tau)^{1+\frac{n-\alpha}{\alpha-n}} \, d\tau + \int_{0}^{t} (t-\tau)^{2-\frac{\lambda}{\alpha-n}+\frac{n-\alpha}{\alpha-n}} \, d\tau \right. \\ &+ \sum_{k=1}^{N+1} \int_{0}^{t} (t-\tau)^{1+\frac{n-\alpha_{k}}{\alpha-n}} \, d\tau + \left(1+\|x\|_{p}^{\lambda}\right) \int_{0}^{t} (t-\tau)^{1+\frac{n-\alpha}{\alpha-n}} \, d\tau \\ &+ \left(1+\|x\|_{p}^{\lambda}\right) \int_{0}^{t} (t-\tau)^{2-\frac{\lambda}{\alpha-n}+\frac{n-\alpha}{\alpha-n}} \, d\tau + \left(1+\|x\|_{p}^{\lambda}\right) \times \\ &\sum_{k=1}^{N+1} \int_{0}^{t} (t-\tau)^{1+\frac{n-\alpha_{k}}{\alpha-n}} \, d\tau \right\} \leq \|f\left(x,t\right)\|_{\mathfrak{M}_{\lambda}} \left\{ C_{1}\left(T\right) + C_{2}\left(T\right) \left(1+\|x\|_{p}^{\lambda}\right) \right\}. \end{aligned}$$

Hence,

$$\left|\frac{u_1(x,t)}{1+\|x\|_p^{\lambda}}\right| \le \|f(x,t)\|_{\mathfrak{M}_{\lambda}} \left\{\frac{C_1\left(T\right)}{1+\|x\|_p^{\lambda}} + C_2\left(T\right)\right\}$$

In the same form, one shows that

$$\left|\frac{u_{2}(x,t)}{1+\|x\|_{p}^{\lambda}}\right| \leq \|\varphi(x)\|_{\mathfrak{M}_{\lambda}}\left\{\frac{C_{1}'(T)}{1+\|x\|_{p}^{\lambda}}+C_{2}'(T)\right\},$$

therefore $\|u(x,t)\|_{\mathfrak{M}_{\lambda}} \leq C\left(\|\varphi(x,t)\|_{\mathfrak{M}_{\lambda}} + \|f(x,t)\|_{\mathfrak{M}_{\lambda}}\right).$

Acknowledgement 8.2. The authors wish to thank to Sergii Torba for many useful comments and discussions, which led to an improvement of this work.

References

- [1] Albeverio S., Khrennikov A. Yu., Shelkovich V. M., Theory of *p*-adic distributions: linear and nonlinear models. Cambridge University Press, 2010.
- [2] Albeverio S., Karwoski W., Diffusion in p-adic numbers. In: K. Ito, H. Hida (Eds.), Gaussian Random Fields, pp. 86-99, 1991, World Scientific, Singapore.
- [3] Albeverio S., Karwoski W., A random walk on p-adics: the generator and its spectrum, Stochastic Process. Appl. 53 (1994), 1-22.
- [4] Avetisov V. A., Bikulov A. Kh., Osipov V. A., p-adic models of ultrametric diffusion in the conformational dynamics of macromolecules, Proc. Steklov Inst. Math. 2004, no. 2 (245), 48–57.
- [5] Avetisov V. A., Bikulov A. Kh., Osipov V. A., p-adic description of characteristic relaxation in complex systems, J. Phys. A 36 (2003), no. 15, 4239–4246.
- [6] Avetisov V. A., Bikulov A. H., Kozyrev S. V., Osipov V. A., p-adic models of ultrametric diffusion constrained by hierarchical energy landscapes, J. Phys. A 35 (2002), no. 2, 177–189.
- [7] Casas-Sánchez O.F., Zúñiga-Galindo W. A., p-adic elliptic quadratic forms, parabolic-type pseudodifferential equations with variable coefficients and Markov processes, p-Adic Numbers Ultrametric Anal. Appl. 6 (2014), no. 1, 1–20.
- [8] Chacón-Cortes L.F., Zúñiga-Galindo W. A., Nonlocal Operators, Parabolic-Type Equations, and Ultrametric Random Walks, J. Math. Phys. 54, 113503 (2013).
- [9] Dragovich B., Khrennikov A. Yu., Kozyrev S. V., Volovich, I. V., On p-adic mathematical physics, p-Adic Numbers Ultrametric Anal. Appl. 1 (2009), no. 1, 1–17.
- [10] Dynkin E. B., Markov processes. Vol. I. Springer-Verlag, 1965.
- [11] Karwowski W., Diffusion processes with ultrametric jumps, Rep. Math. Phys. 60 (2007), no. 2, 221–235.
- [12] Kochubei A.N., Parabolic equations over the field of p-adic numbers, Math. USSR-Izv. 39 (1992), no. 3, 1263–1280.
- [13] Kochubei Anatoly N., Pseudo-differential equations and stochastics over non-Archimedean fields. Marcel Dekker, Inc., New York, 2001.
- [14] Khrennikov A. Yu., Kozyrev S. V., p-adic pseudodifferential operators and analytic continuation of replica matrices, Theoret. and Math. Phys. 144 (2005), no. 2, 1166–1170.
- [15] Mézard Marc, Parisi Giorgio, Virasoro Miguel Angel, Spin glass theory and beyond. World Scientific, 1987.
- [16] Rammal R., Toulouse G., Virasoro M. A., Ultrametricity for physicists, Rev. Modern Phys. 58 (1986), no. 3, 765–788.
- [17] Rodríguez-Vega John Jaime, On a general type of p-adic parabolic equations, Rev. Colombiana Mat. 43 (2009), no. 2, 101–114.
- [18] Rodríguez-Vega J. J., Zúñiga-Galindo W. A., Taibleson operators, p-adic parabolic equations and ultrametric diffusion, Pacific J. Math. 237 (2008), no. 2, 327–347.
- [19] Taibleson M. H., Fourier analysis on local fields, Princeton University Press, 1975.
- [20] Torba S. M., Zúñiga-Galindo W. A., Parabolic Type Equations and Markov Stochastic Processes on Adeles, J. Fourier Anal. Appl. 19 (2013), no. 4, 792835.
- [21] Varadarajan V. S., Path integrals for a class of p-adic Schrödinger equations, Lett. Math. Phys. 39 (1997), no. 2, 97–106.
- [22] Vladimirov V. S., Volovich I. V., Zelenov E. I., p-adic analysis and mathematical physics, World Scientific, 1994.
- [23] Zúñiga-Galindo W. A., Parabolic equations and Markov processes over p-adic fields, Potential Anal. 28 (2008), no. 2, 185–200.

Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Departamento de Matemáticas- Unidad Querétaro, Libramiento Norponiente #2000, Fracc. Real de Juriquilla. Santiago de Querétaro, Qro. 76230, México

E-mail address: fchaconc@math.cinvestav.edu.mx, wazuniga@math.cinvestav.edu.mx