Inference for a Special Bilinear Time Series Model

Shiqing Ling¹, Liang Peng² and Fukang Zhu³

Abstract. It is well known that estimating bilinear models is quite challenging. Many different ideas have been proposed to solve this problem. However, there is not a simple way to do inference even for its simple cases. This paper studies the special bilinear model

$$Y_t = \mu + \phi Y_{t-2} + b Y_{t-2} \varepsilon_{t-1} + \varepsilon_t,$$

where $\{\varepsilon_t\}$ is a sequence of i.i.d. random variables with mean zero. We first give a sufficient condition for the existence of a unique stationary solution for the model and then propose a GARCH-type maximum likelihood estimator for estimating the unknown parameters. It is shown that the GMLE is consistent and asymptotically normal under only finite fourth moment of errors. Also a simple consistent estimator for the asymptotic covariance is provided. A simulation study confirms the good finite sample performance. Our estimation approach is novel and nonstandard and it may provide a new insight for future research in this direction. *Key words and phrases*: Asymptotic distribution, Bilinear model, LSE, MLE.

AMS 2010 subject classifications: Primary 62F12, 62M10; secondary 60G10.

1 Introduction

The general bilinear time series model is defined by the equation

$$Y_{t} = \mu + \sum_{i=1}^{p} \phi_{i} Y_{t-i} + \sum_{j=1}^{q} \psi_{j} \varepsilon_{t-j} + \sum_{l=1}^{m} \sum_{l'=0}^{k} b_{ll'} Y_{t-l} \varepsilon_{t-l'} + \varepsilon_{t},$$
(1)

where $\{\varepsilon_t\}$ is a sequence of independent and identically distributed random variables with mean zero and variance σ^2 . It was proposed by Granger and Anderson (1978a) and has been widely ¹Department of Mathematics, Hong Kong University of Science and Technology, Hong Kong, China.

²Department of Risk Management and Insurance, Georgia State University, USA.

³School of Mathematics, Jilin University, Changchun 130012, China.

applied in many areas such as control theory, economics and finance. The structure of model (1) has been studied in the literature especially for some special cases. For example, Subba Rao (1981) considered model (1) with $\psi_1 = \cdots = \psi_q = 0$; Davis and Resnick (1996) studied the asymptotic behavior of the correlation function for the simple bilinear model $Y_t = bY_{t-1}\varepsilon_{t-1} + \varepsilon_t$; Phan and Tran (1981), Turkman and Turkman (1997) and Basrak et al. (1999) studied the model $Y_t = \phi_1 Y_{t-1} + bY_{t-1}\varepsilon_{t-1} + \varepsilon_t$; Zhang and Tong (2001) considered the model $Y_t = bY_{t-1}\varepsilon_t + \varepsilon_t$. A sufficient condition for stationarity of the general model was obtained by Liu and Brockwell (1988), which is far away from the necessary one as pointed out by Liu (1989). A simplified sufficient condition is given by Liu (1990a).

It is known that estimating the general bilinear model is quite challenging. Many different ideas have been proposed to solve this problem for some special cases of (1), see Pham and Tran (1981), Guegan and Pham (1989), Wittwer (1989), Liu (1990b), Kim and Billard (1990), Kim et al. (1990), Sesay and Subba Rao (1992), Gabr (1998) and Hili (2008). Extension to periodic bilinear models is studied by Bibi and Aknouche (2010) and Bibi and Gautier (2010). However, the asymptotic theory is either rarely established or only derived by assuming that ε_t follows a normal distribution in these papers. The Hellinger distance estimation in Hili (2008) even assumes that the density of ε_t is known. To understand this difficulty, let us look at the least squares estimator (LSE) considered by Pham and Tran (1981). The LSE is equivalent to the quasi-maximum likelihood estimator, which is the minimizer of

$$L_n(\theta) = \sum_{t=1}^n \varepsilon_t^2(\theta),$$

where θ is the vector consisting of all parameters in the model and its true value is $\theta_0, \varepsilon_t(\theta_0) = \varepsilon_t$ and

$$\varepsilon_t(\theta) = Y_t - \mu - \sum_{i=1}^p \phi_i Y_{t-i} - \sum_{j=1}^q \psi_j \varepsilon_{t-j}(\theta) - \sum_{l=1}^m \sum_{l'=0}^k b_{ll'} Y_{t-l} \varepsilon_{t-l'}(\theta).$$

Given a sample $\{Y_1, \dots, Y_n\}$, one needs an efficient way to calculate the residual $\varepsilon_t(\theta)$ such that the effect from the initial values $\{Y_0, Y_{-1}, \dots\}$ is ignorable. This is the so-called invertibility of the model. Although Liu (1990a) gave a sufficient condition for invertibility, it still remains unknown on how to use it to derive the asymptotic limit of the above LSE. Another type of invertibility was proposed by Granger and Anderson (1978b). That is, model (1) is said to be invertible if $\lim_{t\to\infty} E(\varepsilon_t - \hat{\varepsilon}_t)^2 = 0$, where $\hat{\varepsilon}_t$ is an estimator of ε_t . Along this direction, the invertibility of a special bilinear model was studied by Subba Rao (1981), Pham and Tran (1981) and Wittwer (1989). This type of invertibility may be useful for forecasting, but it is not useful for proving asymptotic normality of estimators of parameters. This is because we need the property of $\varepsilon_t(\theta)$ at a neighborhood of the true parameter θ_0 for deriving the asymptotic limit of the estimator. For example, to obtain the asymptotic normality of the LSE, we need the score function $\frac{\partial \varepsilon_t(\theta)}{\partial \theta}$ to have a finite second moment, which in general results in some very restrictive requirements for model (1). Let us further illustrate this issue as follows.

For the following simple bilinear model

$$Y_t = bY_{t-2}\varepsilon_{t-1} + \varepsilon_t, \tag{2}$$

one needs $\prod_{i=1}^{m} Y_{t-i}$ has a finite moment for any m in order to have $E\left\{\frac{\partial \varepsilon_t(\theta)}{\partial \theta}\right\}^2 < \infty$. Grahn (1995) showed that $EY_t^{2m} < \infty$ if and only if $b^{2m} E\varepsilon_t^{2m} < 1$. Note that $E|Y_t|^m < \infty$ for any m is equivalent to b = 0 when $\varepsilon_t \sim N(0, \sigma^2)$. Thus, it is almost impossible to establish the asymptotic normality of the LSE for model (2) unless some special conditions are imposed. Instead Grahn (1995) proposed a nonstandard conditional LSE procedure for model (2) by using the facts that $E(Y_t^2|Y_s, s \leq t-2) = \sigma^2 + b^2 \sigma^2 Y_{t-2}^2$ and $E(Y_tY_{t-1}|Y_s, s \leq t-2) = b\sigma^2 Y_{t-2}$. Although Grahn (1995) derived the asymptotic normality for the conditional LSE, the asymptotic variance and its estimator are not given, so some ad hoc method such as bootstrap method is needed to construct confidence intervals for b. Furthermore, the moment condition required is $EY_t^8 < \infty$, which reduces to $b^8 \sigma^8 < 1/105$ when $\varepsilon_t \sim N(0, \sigma^2)$. This is quite restrictive on the parametric space of (b, σ) . When $\varepsilon_t \sim N(0, \sigma^2)$, Giordano (2000) and Giordano and Vitale (2003) obtained the formula of the asymptotic variance for the conditional LSE of b, which can be estimated too. Liu (1990b) considered the LSE estimation for the model

$$Y_t = \phi Y_{t-p} + b Y_{t-p} \varepsilon_{t-q} + \varepsilon_t, \tag{3}$$

with $p \ge 1$, and obtained its asymptotic normality by assuming that $\frac{\partial \varepsilon_t(\theta)}{\partial \theta}$ has a finite second moment. As in model (2), this condition may only hold when b = 0 if $\varepsilon_t \sim N(0, \sigma^2)$. When $|\varepsilon_t| \le c$ (a constant) holds almost surely and $\phi = 0$, Liu (1990b) showed that this condition holds when $|b| \le \frac{1}{2c}$ which is a small parameter space when c is large. In general, one cannot check whether this condition holds when ε_t is not bounded. That is, a general asymptotic theory for LSE or maximum likelihood estimator (MLE) has not been established for model (3) up to now.

In this paper, we first give a sufficient condition for the existence of a unique stationary solution for a slightly more general model than (2), and then propose a GARCH-type MLE (GMLE) for estimating the unknown parameters. It is shown that the GMLE is consistent and asymptotically normal under only finite fourth moment of errors. We organize this paper as follows. Section 2 presents our main results. Section 3 reports some simulation results. Section 4 concludes. All proofs are given in Section 5.

2 Estimation and Asymptotic Results

Throughout we consider the following special bilinear model:

$$Y_t = \mu + \phi Y_{t-2} + b Y_{t-2} \varepsilon_{t-1} + \varepsilon_t, \tag{4}$$

where $\{\varepsilon_t\}$ is a sequence of independent and identically distributed random variables with mean zero and variance $\sigma^2 > 0$. Let $\ln^+ x = \max\{\ln x, 0\}$ be the positive part of the logarithm, and define

$$X_t = \begin{pmatrix} Y_t \\ Y_{t-1}(\phi + b\varepsilon_t) \end{pmatrix}, \quad A_t = \begin{pmatrix} 0 & 1 \\ & \\ \phi + b\varepsilon_t & 0 \end{pmatrix}, \quad B_t = \begin{pmatrix} \mu + \varepsilon_t \\ & \\ 0 \end{pmatrix}$$

Then (4) can be rewritten as $X_t = A_t X_{t-1} + B_t$. It is easy to check that

$$\prod_{i=1}^{2m} A_i = \begin{pmatrix} \prod_{i=1}^m (\phi + b\varepsilon_{2i}) & 0\\ 0 & \prod_{i=1}^m (\phi + b\varepsilon_{2i-1}) \end{pmatrix}, \quad \prod_{i=1}^{2m+1} A_i = \begin{pmatrix} 0 & \prod_{i=1}^m (\phi + b\varepsilon_{2i})\\ \prod_{i=1}^{m+1} (\phi + b\varepsilon_{2i-1}) & 0 \end{pmatrix}$$

for any integer $m \ge 1$. For vector $x = (x_1, x_2)^{\top}$ and 2×2 matrix y, define $|x| = (x_1^2 + x_2^2)^{1/2}$ and

$$|y|| = \max_{|x|=1} |yx|. \text{ Then}$$
$$\ln \left\| \prod_{i=1}^{2m} A_i \right\|^2 = \max \left\{ \sum_{i=1}^m \ln(\phi + b\varepsilon_{2i})^2, \sum_{i=1}^m \ln(\phi + b\varepsilon_{2i-1})^2 \right\}$$

and

$$\ln \left\| \prod_{i=1}^{2m+1} A_i \right\|^2 = \max \left\{ \sum_{i=1}^m \ln(\phi + b\varepsilon_{2i})^2, \sum_{i=1}^{m+1} \ln(\phi + b\varepsilon_{2i-1})^2 \right\},\$$

which imply that

$$\gamma = \lim_{n \to \infty} \frac{1}{n} \ln \left\| \prod_{i=1}^{n} A_i \right\| = E \ln |\phi + b\varepsilon_1|.$$

Note that $E \ln^+ |B_1| = E \ln^+ |\mu + \varepsilon_1|$. Therefore, when $E \ln^+ |\mu + \varepsilon_1| < \infty$ and $E \ln |\phi + b\varepsilon_1| < 0$, it follows from Theorem 3.2.5 in Basrak (2000) that $X_n = B_n + \sum_{m=1}^{\infty} \prod_{i=0}^{m-1} A_{n-i}B_{n-m}$ converges almost surely and is the unique strictly stationary solution of (4). Since we assume that $0 < E\varepsilon_1^2 < \infty$, $E \ln^+ |\mu + \varepsilon_1| < \infty$ holds naturally.

The following theorem summarizes the above arguments.

Theorem 1. Assume $E \ln |\phi + b\varepsilon_1| < 0$. Then there exists a unique strictly stationary solution to model (4), and the solution is ergodic and has the following representation:

$$Y_t = \mu + \varepsilon_t + \sum_{i=1}^{\infty} \prod_{r=0}^{i-1} (\phi + b\varepsilon_{t-2r-1})(\mu + \varepsilon_{t-2i}).$$

Remark 1. If the model (4) is irreducible, then the condition $E \ln |\phi + b\varepsilon_1| < 0$ is a necessary condition for stationarity, which is a direct consequence of Bougerol and Picard (1992, Theorem 2.5). From Theorem 3 in Kristensen (2009) we know that a sufficient condition for irreducibility is that ε_t has a continuous component at zero and $|\phi| < 1$. **Remark 2.** It follows from Jensen's inequality that $2E \ln |\phi + b\varepsilon_1| = E \ln(\phi + b\varepsilon_1)^2 < \ln E(\phi + b\varepsilon_1)^2 = \ln(\phi^2 + \sigma^2 b^2)$ for $b \neq 0$. Hence model (4) is still stationary when $\phi^2 + \sigma^2 b^2 = 1$ and $b \neq 0$.

Remark 3. When $P(\phi + b\varepsilon_1 > 0) = 1$, results in Kesten (1973) can be employed to show that Y_t has a heavy tail. However, it remains unknown on the tail behavior of Y_t when $P(\phi + b\varepsilon_1 > 0) < 1$. This is in contrast to the well-studied simple bilinear model $Y_t = \phi Y_{t-1} + bY_{t-1}\varepsilon_{t-1} + \varepsilon_t$ in the literature, where the tail property has been clear, but statistical inference for parameters remains unsolved when only some moment condition on ε_t is assumed.

Next we estimate the unknown parameters. Let \mathcal{F}_t be the σ -fields generated by $\{\varepsilon_s : s \leq t\}$. Assume that $\{Y_1, Y_2, \dots, Y_n\}$ are generated by model (4). By noting that

$$E[Y_t|\mathcal{F}_{t-2}] = \mu + \phi Y_{t-2},$$

Var $[Y_t|\mathcal{F}_{t-2}] = E[(Y_t - \mu - \phi Y_{t-2})^2|\mathcal{F}_{t-2}] = \sigma^2(1 + b^2 Y_{t-2}^2),$

we propose to estimate parameters by maximizing the following quasi-log-likelihood function:

$$L_n(\theta) = \sum_{t=1}^n \ell_t(\theta) \quad \text{and} \quad \ell_t(\theta) = -\frac{1}{2} \left[\ln[\sigma^2 (1+b^2 Y_{t-2}^2)] + \frac{(Y_t - \mu - \phi Y_{t-2})^2}{\sigma^2 (1+b^2 Y_{t-2}^2)} \right],$$

where $\theta = (\mu, \phi, \sigma^2, b^2)^{\top}$ is the unknown parameter and its true value is denoted by θ_0 . The maximizer $\hat{\theta}_n$ of $L_n(\theta)$ is called the GMLE of θ_0 . Although the estimation idea has appeared in Francq and Zakoïn (2004), Ling (2004) and Truquet and Yao (2012), the challenge is that $\left\{\frac{\partial \ell_t(\theta)}{\partial \theta}\right\}$ is no longer a martingale difference, which complicates the derivation of the asymptotic limit. A straightforward calculation shows that

$$\begin{split} \frac{\partial \ell_t(\theta)}{\partial \mu} &= \frac{Y_t - \mu - \phi Y_{t-2}}{\sigma^2 (1 + b^2 Y_{t-2}^2)}, \\ \frac{\partial \ell_t(\theta)}{\partial \phi} &= \frac{Y_{t-2} (Y_t - \mu - \phi Y_{t-2})}{\sigma^2 (1 + b^2 Y_{t-2}^2)}, \\ \frac{\partial \ell_t(\theta)}{\partial \sigma^2} &= -\frac{1}{2\sigma^2} \left[1 - \frac{(Y_t - \mu - \phi Y_{t-2})^2}{\sigma^2 (1 + b^2 Y_{t-2}^2)} \right], \\ \frac{\partial \ell_t(\theta)}{\partial b^2} &= -\frac{Y_{t-2}^2}{2(1 + b^2 Y_{t-2}^2)} \left[1 - \frac{(Y_t - \mu - \phi Y_{t-2})^2}{\sigma^2 (1 + b^2 Y_{t-2}^2)} \right]. \end{split}$$

By solving

$$\sum_{t=1}^{n} \frac{\partial \ell_t(\theta)}{\partial \mu} = \sum_{t=1}^{n} \frac{\partial \ell_t(\theta)}{\partial \phi} = \sum_{t=1}^{n} \frac{\partial \ell_t(\theta)}{\partial \sigma^2} = 0,$$

we can write the GMLE for μ , ϕ , σ^2 explicitly in terms of b^2 . Hence, using these explicit expressions and the equation $\sum_{t=1}^{n} \frac{\partial \ell_t(\theta)}{\partial b^2} = 0$, we can first obtain the GMLE for b^2 , and then obtain the GMLE for μ , ϕ , σ^2 .

It is easy to check that $E\left[\frac{\partial \ell_t(\theta_0)}{\partial \theta}\Big|\mathcal{F}_{t-2}\right] = 0$, but $\left\{\frac{\partial \ell_t(\theta_0)}{\partial \theta}\right\}_{t=1}^{\infty}$ can not be a martingale difference. Therefore we can not use the central limit theory for martingale difference to derive the asymptotic limit. Instead we will show that $\left\{\frac{\partial \ell_t(\theta_0)}{\partial \theta}\right\}_{t=1}^{\infty}$ is a near-epoch dependent sequence so that the asymptotic limit of the proposed GMLE can be derived. Denote

$$\begin{split} \Omega &= E\left[\frac{\partial\ell_t(\theta_0)}{\partial\theta} + \frac{\partial\ell_{t-1}(\theta_0)}{\partial\theta}\right] \left[\frac{\partial\ell_t(\theta_0)}{\partial\theta} + \frac{\partial\ell_{t-1}(\theta_0)}{\partial\theta}\right]^\top - E\left[\frac{\partial\ell_t(\theta_0)}{\partial\theta}\frac{\partial\ell_t(\theta_0)}{\partial\theta^\top}\right],\\ \Sigma &= \operatorname{diag}\left\{E\left[\frac{1}{\sigma_0^2(1+b_0^2Y_{t-2}^2)} \begin{pmatrix} 1 & Y_{t-2} \\ Y_{t-2} & Y_{t-2}^2 \end{pmatrix}\right], \ E\left(\begin{array}{cc} \frac{1}{2\sigma_0^4} & \frac{Y_{t-2}^2}{2\sigma_0^2(1+b_0^2Y_{t-2}^2)} \\ \frac{Y_{t-2}^2}{2\sigma_0^2(1+b_0^2Y_{t-2}^2)} & \frac{Y_{t-2}^4}{2(1+b_0^2Y_{t-2}^2)^2} \end{array}\right)\right\}. \end{split}$$

The following theorem gives the asymptotic properties of the GMLE.

Theorem 2. Suppose the parameter space Θ is a compact subset of $\{\theta : E \ln |\phi + b\varepsilon_1| < 0, |\mu| \le \overline{\mu}, |\phi| \le \overline{\phi}, \underline{\omega} \le \sigma^2 \le \overline{\omega}, \underline{\alpha} \le b^2 \le \overline{\alpha}\}$, where $\overline{\mu}, \overline{\phi}, \underline{\omega}, \overline{\omega}, \underline{\alpha}$ and $\overline{\alpha}$ are some finite positive constants, and the true parameter value θ_0 is an interior point in Θ . Further assume $E\varepsilon_1^4 < \infty$. Then as $n \to \infty$,

- (a) $\hat{\theta}_n \to \theta_0$ almost surely,
- (b) $\sqrt{n}(\hat{\theta}_n \theta_0) \xrightarrow{d} N(0, \Sigma^{-1}\Omega\Sigma^{-1}).$

Remark 4. To ensure the positive definiteness of Σ in Theorem 2, we only need to show the two sub-matrices are positive definite, which is equivalent to show the determinants of these two submatrices are positive. Obviously Cauchy-Schwarz inequality implies that the determinant of the second sub-matrix in Σ is positive. Put $A = 1 + b_0^2 Y_t^2$, then the determinant of the first sub-matrix
$$\begin{split} &is \; \sigma_0^{-4}\{E(A^{-1})E(1-A^{-1})b_0^{-2}-[E(Y_tA^{-1})]^2\}=\sigma_0^{-4}b_0^{-2}\{E(A^{-1})-[E(A^{-1})]^2-b_0^2[E(Y_tA^{-1})]^2\}>\\ &\sigma_0^{-4}b_0^{-2}\{E(A^{-1})-E(A^{-2})-E(b_0^2Y_t^2A^{-2})\}=0. \end{split}$$



Figure 1: Region of (b, ϕ) such that $E \ln |\phi + b\varepsilon_1| < 0$ when $\varepsilon_1 \sim N(0, 1)$.

Remark 5. Figure 1 gives the region of (b, ϕ) such that $E \ln |\phi + b\varepsilon_1| < 0$ when $\varepsilon_1 \sim N(0, 1)$. So |b| can be greater than 1, i.e., the asymptotic limit of the proposed GMLE holds under some weaker conditions than the condition $|b\sigma| < 105^{-1/8} \approx 0.5589$ required by the conditional LSE in Grahn(1995). Moreover, Ω and Σ can be estimated simply by

$$\begin{split} \hat{\Omega}_{n} &= \frac{1}{n} \sum_{t=1}^{n} \left[\frac{\partial l_{t}(\hat{\theta}_{n})}{\partial \theta} + \frac{\partial l_{t-1}(\hat{\theta}_{n})}{\partial \theta} \right] \left[\frac{\partial l_{t}(\hat{\theta}_{n})}{\partial \theta} + \frac{\partial l_{t-1}(\hat{\theta}_{n})}{\partial \theta} \right]^{\top} - \frac{1}{n} \sum_{t=1}^{n} \left[\frac{\partial l_{t}(\hat{\theta}_{n})}{\partial \theta} \frac{\partial l_{t}(\hat{\theta}_{n})}{\partial \theta^{\top}} \right], \\ \hat{\Sigma}_{n} &= \operatorname{diag} \left\{ \frac{1}{n} \sum_{t=1}^{n} \left[\frac{1}{\hat{\theta}_{n3}(1 + \hat{\theta}_{n4}Y_{t-2}^{2})} \begin{pmatrix} 1 & Y_{t-2} \\ Y_{t-2} & Y_{t-2}^{2} \end{pmatrix} \right], \\ \frac{1}{n} \sum_{t=1}^{n} \left(\frac{\frac{1}{2\hat{\theta}_{n3}^{2}}}{\frac{Y_{t-2}^{2}}{2\hat{\theta}_{n3}(1 + \hat{\theta}_{n4}Y_{t-2}^{2})}} \frac{Y_{t-2}^{4}}{2(1 + \hat{\theta}_{n4}Y_{t-2}^{2})^{2}} \right) \right\}, \end{split}$$

respectively, where $\hat{\theta}_n = (\hat{\theta}_{n1}, \hat{\theta}_{n2}, \hat{\theta}_{n3}, \hat{\theta}_{n4})^{\top}$. The consistency follows from Lemma 2 in Section 5.

Since
$$\sum_{t=1}^{n} \frac{\partial \ell_t(\theta)}{\partial b^2} = 0$$
 is equivalent to $\sum_{t=1}^{n} \frac{\partial \ell_t(\theta)}{\partial b} = 0$, one can not estimate b by the above

GMLE. In order to estimate b, we need a consistent estimator for the sign of b. Write

$$(Y_t - \mu - \phi Y_{t-2})(Y_{t-1} - \mu - \phi Y_{t-3}) = \varepsilon_t (bY_{t-3}\varepsilon_{t-2} + \varepsilon_{t-1}) + b^2 Y_{t-3} Y_{t-2} \varepsilon_{t-2} \varepsilon_{t-1} + bY_{t-2} \varepsilon_{t-1}^2.$$

It is easy to see that $E\{(Y_t - \mu - \phi Y_{t-2})(Y_{t-1} - \mu - \phi Y_{t-3})|\mathcal{F}_{t-2}\} = b\sigma^2 Y_{t-2}$, which motivates to estimate b by minimizing the following least squares

$$\sum_{t=2}^{n} \{ (Y_t - \mu - \phi Y_{t-2})(Y_{t-1} - \mu - \phi Y_{t-3}) - b\sigma^2 Y_{t-2} \}^2$$

with μ , ϕ and σ^2 being replaced by the corresponding GMLE. However, in order to avoid requiring some moment conditions on Y_t , we propose to minimize the weighted least squares

$$\sum_{t=2}^{n} \frac{\{(Y_t - \mu - \phi Y_{t-2})(Y_{t-1} - \mu - \phi Y_{t-3}) - b\sigma^2 Y_{t-2}\}^2}{(1 + Y_{t-2}^2)\sqrt{1 + Y_{t-3}^2}}$$

with μ, ϕ, σ^2 being replaced by the corresponding GMLE. This results in

$$\tilde{b}_n = \left(\hat{\theta}_{n3} \sum_{t=2}^n \frac{Y_{t-2}^2}{(1+Y_{t-2}^2)\sqrt{1+Y_{t-3}^2}}\right)^{-1} \sum_{t=2}^n \frac{(Y_t - \hat{\theta}_{n1} - \hat{\theta}_{n2}Y_{t-2})(Y_{t-1} - \hat{\theta}_{n1} - \hat{\theta}_{n2}Y_{t-3})Y_{t-2}}{(1+Y_{t-2}^2)\sqrt{1+Y_{t-3}^2}}.$$

Like Theorem 2 (a), it is easy to show that $\tilde{b}_n = b + o_p(1)$. Using \tilde{b}_n to estimate the sign of b, we obtain an estimator for b as $\hat{b}_n = \operatorname{sgn}(\tilde{b}_n)\sqrt{\hat{\theta}_{n4}}$. It easily follows from Theorem 2 that $\hat{b}_n = b + o_p(1)$ and the asymptotic limit of $2b\sqrt{n}(\hat{b}_n - b)$ is the same as that of $\sqrt{n}(\hat{\theta}_{n4} - b^2)$ given in Theorem 2. As stated in the simulation study, we propose to use $2b\sqrt{n}(\hat{b}_n - b)$ rather than $2\hat{b}_n\sqrt{n}(\hat{b}_n - b)$ to construct a confidence interval for b although both share the same asymptotic limit. Moreover we do not propose to estimate b directly by \tilde{b}_n . The reason is that like Grahm (1995) we can not derive the formula and a consistent estimator for the asymptotic variance of $\sqrt{n}(\tilde{b}_n - b)$. Moreover, \tilde{b}_n is a less efficient estimator than \hat{b}_n in general.

Theorem 2 excludes the case of b = 0, which reduces the bilinear model to a linear model. Hence testing $H_0: b = 0$ is of interest. Write $\Theta = [-\bar{\mu}, \bar{\mu}] \times [-\bar{\phi}, \bar{\phi}] \times [\underline{\omega}, \overline{\omega}] \times [0, \overline{\alpha}]$, where $\bar{\mu}, \bar{\phi}, \underline{\omega}, \overline{\omega}$ and $\overline{\alpha}$ are some finite positive constants. Then the case of b = 0 means that $\theta = (\mu, \phi, \sigma^2, b^2)^{\top}$ lies at the boundary of the compact set Θ , which implies that the case of b = 0 is the well-known nonstandard situation of maximum likelihood estimation. The following theorem easily follows from Lemmas 1–3 in Section 5 and the same arguments in deriving (2.2) in Self and Liang (1987).

Theorem 3. Suppose the parameter space Θ satisfies $E \ln |\phi + b\varepsilon_1| < 0$, and the true parameter value $\theta_0 = (\mu_0, \phi_0, \sigma_0^2, 0)^\top$ satisfies that $(\mu_0, \phi_0, \sigma_0^2)^\top$ is an interior point of $[-\bar{\mu}, \bar{\mu}] \times [-\bar{\phi}, \bar{\phi}] \times [\underline{\omega}, \overline{\omega}]$. Further assume $E\varepsilon_1^4 < \infty$. Then as $n \to \infty$,

(a) $\hat{\theta}_n \to \theta_0$ almost surely,

 $(b) \sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow{d} (Z_1, Z_2, Z_3, Z_4)^\top I(Z_4 > 0) + (Z_1 - \sigma_{14}\sigma_{44}^{-1}Z_4, Z_2 - \sigma_{24}\sigma_{44}^{-1}Z_4, Z_3 - \sigma_{34}\sigma_{44}^{-1}Z_4, 0)^\top I(Z_4 < 0), \ where \ (Z_1, Z_2, Z_3, Z_4)^\top \sim N(0, \Sigma^{-1}\Omega\Sigma^{-1}), \ \Sigma^{-1}\Omega\Sigma^{-1} = (\sigma_{ij}) \ and \ \Sigma \ and \ \Omega \ are \ given \ in \ Theorem \ 2.$

Remark 6. Using the consistent estimators for Ω and Σ in Remark 5, one can easily simulate the asymptotic limit of $\sqrt{n}(\hat{\theta}_n - \theta_0)$ so that interval estimation is obtained. For testing $H_0: b = 0$ against $H_a: b \neq 0$, we let $\hat{\sigma}_{ij}$ denote the consistent estimator for σ_{ij} given in Remark 5, but with $\hat{\theta}_{n,4}$ being replaced by 0. By Theorem 3 one rejects H_0 at level ξ whenever $\hat{\theta}_{n,4} > \sqrt{\hat{\sigma}_{44}}z_{\xi}/\sqrt{n}$, where $P(N(0,1) > z_{\xi}) = \xi$. We also remark that the likelihood ratio tests in Self and Liang (1987) do not apply to our bilinear model even for the case of $b^2 > 0$. The reason is that $\left\{\frac{\partial \ell_t(\theta)}{\partial \theta}\right\}$ can not be a martingale difference, and so Ω in Theorem 2 is different from the standard one $E\left\{\frac{\partial \ell_t(\theta)}{\partial \theta}\frac{\partial \ell_t(\theta)}{\partial \theta^{\top}}\right\}$, which is necessary to ensure Wilks theorem holds for the likelihood ratio approach.

3 Simulation

We investigate the finite sample performance of the proposed GMLE by drawing 1,000 random samples of size n = 200 and 1,000 from model (4) with $\mu = 0, b = \pm 0.1$ or $\pm 1, \phi = 0$ or 0.9, and $\varepsilon_t \sim N(0, 1)$. We compute the GMLE $\hat{\theta}_n = (\hat{\theta}_{n1}, \dots, \hat{\theta}_{n4})^{\top}$ for $\theta = (\mu, \phi, \sigma^2, b^2)^{\top}$ and \hat{b}_n . For an estimator $\hat{\beta}$, we use $E(\hat{\beta})$, $SD(\hat{\beta})$ and $\widehat{SD}(\hat{\beta})$ to denote the sample mean of $\hat{\beta}$, sample standard deviation of $\hat{\beta}$ and sample mean of the standard deviation estimator given in Remark 5 of $\hat{\beta}$ based on the 1,000 samples.

Tables 1 and 2 report these quantities, which show that the proposed GMLE has a small bias (i.e, $E(\cdot)$ close to the true value) and the proposed variance estimator is accurate too (i.e., $\widehat{SD}(\cdot)$ close to $SD(\cdot)$). From these two tables, we also observe that $SD(\hat{\beta})$ and $\widehat{SD}(\hat{\beta})$ are much smaller when n = 1000 than those when n = 200. Although the proposed estimator for b has a small bias, the proposed variance estimator performs badly when b is small. This is due to some very small values of $\hat{\theta}_{n4}$. However, the variance estimator for $2b\hat{b}_n$ is reasonably well and much accurate than that for \hat{b}_n . Hence, we suggest to use $2b\sqrt{n}(\hat{b}_n - b)$ instead of $\sqrt{n}(\hat{b}_n - b)$ to construct a confidence interval for b in practice.

Next we use Remark 6 to test $H_0: b = 0$ against $H_a: b \neq 0$ at levels 0.1 and 0.05. We draw 10,000 random samples of size n = 200 and 1,000 from model (4) with $\mu = 0, b = b^*/n^{0.25}, \phi = 0.1$ or 0.9, $b^* = 0, 0.5, 1$, and $\varepsilon_t \sim N(0, 1)$. The empirical size and power are reported in Table 3, where $b^* = 0$ corresponds to the size. From Table 3, we observe that the proposed test has a reasonably accurate size and non-trivial power.

4 Conclusions

Many different ideas have been proposed for estimating parameters in bilinear models. Unfortunately asymptotic limit is either missing or derived under some restrictive distribution assumption on errors. By focusing on a simple bilinear model, we first give a sufficient condition for the existence of a unique stationary solution for the model and then propose a GARCH-type maximum likelihood estimator for estimating parameters. It is shown that the proposed estimator is consistent and asymptotically normal under mild conditions. Moreover, the new estimation approach is novel, nonstandard and has good finite sample behavior.

(b,ϕ)	(0.1, 0)	(0.1, 0.9)	(-0.1, 0)	(-0.1, 0.9)	(1, 0)	(1, 0.9)	(-1, 0)	(-1, 0.9)
$E(\hat{\theta}_{n1})$	0.0008	-0.0020	0.0002	0.0021	-0.0014	0.0054	0.0039	-0.0009
$\mathrm{SD}(\hat{\theta}_{n1})$	0.0707	0.0890	0.0720	0.0882	0.1058	0.1315	0.1015	0.1369
$\widehat{\mathrm{SD}}(\widehat{\theta}_{n1})$	0.0701	0.0740	0.0703	0.0740	0.0999	0.1244	0.0997	0.1241
$E(\hat{\theta}_{n2})$	-0.0026	0.8793	-0.0092	0.8797	-0.0113	0.8887	-0.0052	0.8892
$\mathrm{SD}(\hat{\theta}_{n2})$	0.0714	0.0399	0.0706	0.0400	0.1000	0.0878	0.0979	0.0891
$\widehat{\mathrm{SD}}(\widehat{\theta}_{n2})$	0.0698	0.0352	0.0693	0.0350	0.0977	0.0868	0.0974	0.0869
$E(\hat{\theta}_{n3})$	0.9648	0.9896	0.9744	0.9919	1.0209	1.0254	1.0263	1.0280
$\mathrm{SD}(\hat{\theta}_{n3})$	0.1115	0.1161	0.1047	0.1179	0.1976	0.2535	0.1993	0.2524
$\widehat{\mathrm{SD}}(\widehat{\theta}_{n3})$	0.1196	0.1253	0.1209	0.1259	0.1855	0.2245	0.1880	0.2280
$E(\hat{\theta}_{n4})$	0.0355	0.0124	0.0344	0.0117	0.9987	1.0256	0.9882	1.0201
$\mathrm{SD}(\hat{\theta}_{n4})$	0.0615	0.0166	0.0570	0.0168	0.3183	0.3205	0.3246	0.3041
$\widehat{\mathrm{SD}}(\widehat{\theta}_{n4})$	0.0738	0.0174	0.0728	0.0172	0.2786	0.2667	0.2797	0.2671
$E(\hat{b}_n)$	0.0879	0.0780	-0.0986	-0.0736	0.9872	1.0015	-0.9812	-0.9996
$\mathrm{SD}(\hat{b}_n)$	0.1666	0.0793	0.1572	0.0793	0.1555	0.1505	0.1593	0.1446
$\widehat{\mathrm{SD}}(\hat{b}_n)$	4.7444	0.8028	4.5939	0.8511	0.1379	0.1294	0.1393	0.1303
$E(2b\hat{b}_n)$	0.0176	0.0156	0.0197	0.0147	1.9744	2.0030	1.9624	1.9992
$\mathrm{SD}(2b\hat{b}_n)$	0.0333	0.0159	0.0314	0.0159	0.3111	0.3010	0.3187	0.2892
$\widehat{\mathrm{SD}}(2b\hat{b}_n)$	0.0738	0.0174	0.0728	0.0172	0.2786	0.2667	0.2797	0.2671

Table 1: Sample mean and sample standard deviation are reported for the proposed GMLE for $(\mu, \phi, \sigma^2, b^2)^{\top}$ and b with n = 200.

(b,ϕ)	(0.1, 0)	(0.1, 0.9)	(-0.1, 0)	(-0.1, 0.9)	(1, 0)	(1, 0.9)	(-1, 0)	(-1, 0.9)
$E(\hat{\theta}_{n1})$	0.0002	-0.0004	0.0001	0.0007	-0.0005	0.0031	0.0002	-0.0001
$\mathrm{SD}(\hat{\theta}_{n1})$	0.0311	0.0330	0.0310	0.0331	0.0444	0.0563	0.0445	0.0556
$\widehat{\mathrm{SD}}(\widehat{\theta}_{n1})$	0.0317	0.0325	0.0317	0.0325	0.0448	0.0549	0.0447	0.0549
$E(\hat{\theta}_{n2})$	-0.0044	0.8956	-0.0041	0.8958	-0.0036	0.8971	-0.0026	0.8972
$\mathrm{SD}(\hat{\theta}_{n2})$	0.0306	0.0155	0.0311	0.0156	0.0428	0.0382	0.0434	0.0385
$\widehat{\mathrm{SD}}(\hat{\theta}_{n2})$	0.0317	0.0153	0.0376	0.0153	0.0444	0.0392	0.0444	0.0392
$E(\hat{\theta}_{n3})$	0.9907	1.0017	0.9904	1.0005	1.0037	1.0014	1.0024	1.0102
$\mathrm{SD}(\hat{\theta}_{n3})$	0.0492	0.0555	0.0492	0.0555	0.0829	0.1033	0.0844	0.1042
$\widehat{\mathrm{SD}}(\hat{\theta}_{n3})$	0.0550	0.0564	0.0550	0.0563	0.0846	0.1019	0.0849	0.1029
$E(\hat{\theta}_{n4})$	0.0178	0.0095	0.0182	0.0097	0.9969	1.0050	0.9975	0.9949
$\mathrm{SD}(\hat{\theta}_{n4})$	0.0247	0.0070	0.0251	0.0070	0.1365	0.1192	0.1358	0.1214
$\widehat{\mathrm{SD}}(\hat{\theta}_{n4})$	0.0327	0.0072	0.0329	0.0072	0.1284	0.1194	0.1298	0.1185
$E(\hat{b}_n)$	0.0935	0.0871	-0.0935	-0.0887	0.9961	1.0008	-0.9964	-0.9956
$\mathrm{SD}(\hat{b}_n)$	0.0953	0.0434	0.0974	0.0432	0.0680	0.0589	0.0678	0.0606
$\widehat{\mathrm{SD}}(\hat{b}_n)$	1.8724	0.1336	1.9108	0.1349	0.0642	0.0594	0.0649	0.0593
$E(2b\hat{b}_n)$	0.0187	0.0174	0.0187	0.0177	1.9923	2.0015	1.9929	1.9912
$\mathrm{SD}(2b\hat{b}_n)$	0.0191	0.0087	0.0195	0.0086	0.1359	0.1178	0.1356	0.1212
$\widehat{\mathrm{SD}}(2b\hat{b}_n)$	0.0327	0.0072	0.0329	0.0072	0.1284	0.1194	0.1298	0.1185

Table 2: Sample mean and sample standard deviation are reported for the proposed GMLE for $(\mu, \phi, \sigma^2, b^2)^{\top}$ and b with n = 1000.

	level 0.1				level 0.05			
(n,ϕ)	b = 0	$b = 0.5n^{-0.25}$	$b = n^{-0.25}$		b = 0	$b = 0.5n^{-0.25}$	$b = n^{-0.25}$	
(200, 0.1)	0.0790	0.1142	0.2513		0.0449	0.0634	0.1550	
(200, 0.9)	0.0777	0.3151	0.7575		0.0422	0.2025	0.6267	
(1000, 0.1)	0.0815	0.1237	0.3030		0.0375	0.0592	0.1795	
(1000, 0.9)	0.0758	0.4026	0.9604		0.0336	0.2619	0.9052	

Table 3: Empirical size and power are reported for the proposed test in Remark 6 for testing $H_0: b = 0$ against $H_a: b \neq 0$ at levels 0.1 and 0.05.

5 Proofs

We first give one lemma, which plays a key role in the proofs of other lemmas.

Lemma 1. Under assumptions of Theorem 2,

(a)
$$E \sup_{\theta \in \Theta} |\ell_t(\theta)| < \infty;$$

(b) $E\ell_t(\theta)$ achieves its unique maximum value at $\theta = \theta_0.$

Proof. Since $E|\varepsilon_t| < \infty$, similar to the proof of Lemma 1 in Ling (2004), we can show that there exists a $\delta \in (0, 1)$ such that $E|\phi + b\varepsilon_t|^{\delta} < 1$. Using this and the expression of Y_t in Theorem 1, we can show that $E|Y_t|^{\delta} < \infty$. Take $\delta_0 = \delta/2$. Thus, by Jensen's inequality, it follows that

$$\begin{split} E \sup_{\theta \in \Theta} |\ln[\sigma^2(1+b^2Y_{t-2}^2)]| &\leq \sup_{\theta \in \Theta} |\ln\sigma^2| + \delta_0^{-1}E\ln(1+\overline{\alpha}Y_{t-2}^2)^{\delta_0} \\ &\leq |\ln\overline{\omega}| + \delta_0^{-1}\ln(1+\overline{\alpha}^{\delta_0}E|Y_{t-2}|^{\delta}) < \infty, \end{split}$$

where the following elementary relationship is used: $(a_1 + a_2)^s \leq a_1^s + a_2^s$ for all $a_1, a_2 > 0$ and $s \in [0, 1]$. Furthermore, since $Y_t - \mu - \phi Y_{t-2} = \varepsilon_t - (\mu - \mu_0) - (\phi - \phi_0)Y_{t-2} + b_0\varepsilon_{t-1}Y_{t-2}$, it can

be shown that

$$\begin{aligned}
E \sup_{\theta \in \Theta} \frac{(Y_t - \mu - \phi Y_{t-2})^2}{\sigma^2 (1 + b^2 Y_{t-2}^2)} \\
&\leq \frac{4}{\omega} \left[E \sup_{\theta \in \Theta} \frac{\varepsilon_t^2}{1 + b^2 Y_{t-2}^2} + E \sup_{\theta \in \Theta} \frac{(\mu - \mu_0)^2}{1 + b^2 Y_{t-2}^2} + E \sup_{\theta \in \Theta} \frac{(\phi - \phi_0)^2 Y_{t-2}^2}{1 + b^2 Y_{t-2}^2} + E \sup_{\theta \in \Theta} \frac{b_0^2 \varepsilon_{t-1}^2 Y_{t-2}^2}{1 + b^2 Y_{t-2}^2} \right] \\
&\leq \frac{4}{\omega} \left(\overline{\omega} + 4\overline{\mu}^2 + \frac{4\overline{\phi}^2}{\underline{\alpha}} + \frac{\overline{\omega} \,\overline{\alpha}}{\underline{\alpha}} \right) < \infty.
\end{aligned}$$
(5)

Hence, (a) holds.

To prove (b), by noting that

$$E[(Y_t - \mu - \phi Y_{t-2})^2 | \mathcal{F}_{t-2}] = E[(\varepsilon_t - (\mu - \mu_0) - (\phi - \phi_0)Y_{t-2} + b_0\varepsilon_{t-1}Y_{t-2})^2 | \mathcal{F}_{t-2}]$$
$$= [(\mu - \mu_0) + (\phi - \phi_0)Y_{t-2}]^2 + \sigma_0^2(1 + b_0^2Y_{t-2}^2),$$

we have

$$E\ell_{t}(\theta) = -\frac{1}{2}E\left[\ln[\sigma^{2}(1+b^{2}Y_{t-2}^{2})] + \frac{(Y_{t}-\mu-\phi Y_{t-2})^{2}}{\sigma^{2}(1+b^{2}Y_{t-2}^{2})}\right]$$
$$= -\frac{1}{2}\left\{E\ln[\sigma^{2}(1+b^{2}Y_{t-2}^{2})] + E\frac{\sigma_{0}^{2}(1+b_{0}^{2}Y_{t-2}^{2})}{\sigma^{2}(1+b^{2}Y_{t-2}^{2})}\right\} - E\frac{[(\mu-\mu_{0})+(\phi-\phi_{0})Y_{t-2}]^{2}}{2\sigma^{2}(1+b^{2}Y_{t-2}^{2})}.$$
 (6)

The second term in (6) reaches its maximum at zero, and this occurs if and only if $\mu = \mu_0$ and $\phi = \phi_0$. The first term in (6) is equal to

$$-\frac{1}{2}\left[-E(\ln M_t) + EM_t\right] - \frac{1}{2}E\ln[\sigma_0^2(1+b_0^2Y_{t-2}^2)],\tag{7}$$

where $M_t = \frac{\sigma_0^2 (1 + b_0^2 Y_{t-2}^2)}{\sigma^2 (1 + b^2 Y_{t-2}^2)}$. As in Lemma 2 of Ling (2004), (7) reaches its maximum $-1/2 - E \ln[\sigma_0^2 (1 + b_0^2 Y_{t-2}^2)]/2$, and this occurs if and only if $\sigma^2 = \sigma_0^2$ and $b^2 = b_0^2$. Thus, $E\ell_t(\theta)$ is uniquely maximized at θ_0 .

Lemma 2. Under assumptions of Theorem 2,

(a)
$$E \sup_{\theta \in \Theta} \left\| \frac{\partial \ell_t(\theta)}{\partial \theta} \right\|^2 < \infty,$$

(b) $E \sup_{\theta \in \Theta} \left[\frac{\partial^2 \ell_t(\theta)}{\partial \theta \partial \theta^\top} \right] < \infty,$

$$(c) \quad \sup_{\theta \in \Theta} \left| \frac{1}{n} \sum_{t=1}^{n} \left\{ \frac{\partial^2 \ell_t(\theta)}{\partial \theta \partial \theta^{\top}} - E \frac{\partial^2 \ell_t(\theta)}{\partial \theta \partial \theta^{\top}} \right\} \right| = o_p(1) \quad as \quad n \to \infty,$$

$$(d) \quad \sup_{\theta \in \Theta} \left| \frac{1}{n} \sum_{t=1}^{n} \left\{ \left\| \frac{\partial \ell_t(\theta)}{\partial \theta} \right\|^2 - E \left\| \frac{\partial \ell_t(\theta)}{\partial \theta} \right\|^2 \right\} \right| = o_p(1) \quad as \quad n \to \infty.$$

Proof. As in (5), it is easy to show that

$$\begin{split} E \sup_{\theta \in \Theta} \left[\frac{\partial \ell_t(\theta)}{\partial \phi} \right]^2 &= E \sup_{\theta \in \Theta} \frac{Y_{t-2}^2 [\varepsilon_t - (\mu - \mu_0) - (\phi - \phi_0)Y_{t-2} + b_0\varepsilon_{t-1}Y_{t-2}]^2}{\sigma^4 (1 + b^2 Y_{t-2}^2)^2} \\ &\leq \frac{4}{\omega^2} \left[E \sup_{\theta \in \Theta} \frac{\varepsilon_t^2 Y_{t-2}^2}{(1 + b^2 Y_{t-2}^2)^2} + E \sup_{\theta \in \Theta} \frac{(\mu - \mu_0)^2 Y_{t-2}^2}{(1 + b^2 Y_{t-2}^2)^2} \right] \\ &+ E \sup_{\theta \in \Theta} \frac{(\phi - \phi_0)^2 Y_{t-2}^4}{(1 + b^2 Y_{t-2}^2)^2} + E \sup_{\theta \in \Theta} \frac{b_0^2 \varepsilon_{t-1}^2 Y_{t-2}^4}{(1 + b^2 Y_{t-2}^2)^2} \right] < \infty, \\ E \sup_{\theta \in \Theta} \left[\frac{\partial \ell_t(\theta)}{\partial b^2} \right]^2 &\leq \frac{1}{2} E \sup_{\theta \in \Theta} \frac{Y_{t-2}^4}{(1 + b^2 Y_{t-2}^2)^2} + \frac{1}{2} E \sup_{\theta \in \Theta} \frac{Y_{t-2}^4 (Y_t - \mu - \phi Y_{t-2})^4}{\sigma^4 (1 + b^2 Y_{t-2}^2)^4} \\ &= \frac{1}{2} E \sup_{\theta \in \Theta} \frac{Y_{t-2}^4}{(1 + b^2 Y_{t-2}^2)^2} \\ &+ \frac{1}{2} E \sup_{\theta \in \Theta} \frac{Y_{t-2}^4 [\varepsilon_t - (\mu - \mu_0) - (\phi - \phi_0)Y_{t-2} + b_0\varepsilon_{t-1}Y_{t-2}]^4}{\sigma^4 (1 + b^2 Y_{t-2}^2)^4} \\ &\leq \frac{1}{2\underline{\alpha}^2} + 2E \sup_{\theta \in \Theta} \frac{Y_{t-2}^4 [\varepsilon_t^2 + (\mu - \mu_0)^2 + (\phi - \phi_0)^2 Y_{t-2}^2 + b_0^2 \varepsilon_{t-1}^2 Y_{t-2}^2]^2}{\sigma^4 (1 + b^2 Y_{t-2}^2)^4} \\ &\leq \frac{1}{2\underline{\alpha}^2} + \frac{8}{\underline{\omega}^2} \left[E \sup_{\theta \in \Theta} \frac{\varepsilon_t^4 Y_{t-2}^4}{(1 + b^2 Y_{t-2}^2)^4} + E \sup_{\theta \in \Theta} \frac{(\mu - \mu_0)^4 Y_{t-2}^4}{(1 + b^2 Y_{t-2}^2)^4} \\ &+ E \sup_{\theta \in \Theta} \frac{(\phi - \phi_0)^4 Y_{t-2}^8}{(1 + b^2 Y_{t-2}^2)^4} + E \sup_{\theta \in \Theta} \frac{b_0^4 \varepsilon_{t-1}^4 Y_{t-2}^8}{(1 + b^2 Y_{t-2}^2)^4} \right] < \infty. \end{split}$$

Similarly, we can show that other terms in (a) are finite too. Hence, (a) holds.

A straightforward calculation gives that

$$\begin{split} \frac{\partial^2 \ell_t(\theta)}{\partial \mu^2} &= -\frac{1}{\sigma^2 (1 + b^2 Y_{t-2}^2)}, \\ \frac{\partial^2 \ell_t(\theta)}{\partial \phi^2} &= -\frac{Y_{t-2}^2}{\sigma^2 (1 + b^2 Y_{t-2}^2)}, \\ \frac{\partial^2 \ell_t(\theta)}{\partial \sigma^4} &= \frac{1}{2\sigma^4} \left[1 - \frac{2(Y_t - \mu - \phi Y_{t-2})^2}{\sigma^2 (1 + b^2 Y_{t-2}^2)} \right], \\ \frac{\partial^2 \ell_t(\theta)}{\partial b^4} &= \frac{Y_{t-2}^4}{2(1 + b^2 Y_{t-2}^2)^2} \left[1 - \frac{2(Y_t - \mu - \phi Y_{t-2})^2}{\sigma^2 (1 + b^2 Y_{t-2}^2)} \right], \\ \frac{\partial^2 \ell_t(\theta)}{\partial \mu \partial \phi} &= -\frac{Y_{t-2}}{\sigma^2 (1 + b^2 Y_{t-2}^2)}, \\ \frac{\partial^2 \ell_t(\theta)}{\partial \mu \partial \sigma^2} &= -\frac{Y_t - \mu - \phi Y_{t-2}}{\sigma^4 (1 + b^2 Y_{t-2}^2)}, \end{split}$$

$$\begin{split} \frac{\partial^2 \ell_t(\theta)}{\partial \mu \partial b^2} &= -\frac{Y_{t-2}^2(Y_t - \mu - \phi Y_{t-2})}{\sigma^2(1 + b^2 Y_{t-2}^2)^2}, \\ \frac{\partial^2 \ell_t(\theta)}{\partial \phi \partial \sigma^2} &= -\frac{Y_{t-2}(Y_t - \mu - \phi Y_{t-2})}{\sigma^4(1 + b^2 Y_{t-2}^2)}, \\ \frac{\partial^2 \ell_t(\theta)}{\partial \phi \partial b^2} &= -\frac{Y_{t-2}^3(Y_t - \mu - \phi Y_{t-2})}{\sigma^2(1 + b^2 Y_{t-2}^2)^2}, \\ \frac{\partial^2 \ell_t(\theta)}{\partial \sigma^2 \partial b^2} &= -\frac{Y_{t-2}^2(Y_t - \mu - \phi Y_{t-2})^2}{2\sigma^4(1 + b^2 Y_{t-2}^2)^2}. \end{split}$$

Using these formulas and some similar arguments in proving (a), we can show that (b) holds. (c) and (d) follow from Theorem 3.1 in Ling and McAleer (2003). \Box

Lemma 3. Under assumptions of Theorem 2,

$$\frac{1}{\sqrt{n}}\sum_{t=1}^{n}\frac{\partial\ell_t(\theta_0)}{\partial\theta}\overset{d}{\to}N(0,\Omega).$$

Proof. Let $C = (c_1, c_2, c_3, c_4)^\top$ be any constant vector with $C^\top C \neq 0$ and define

$$\begin{split} S_n &\equiv \frac{C^{\top}}{\sqrt{n}} \sum_{t=1}^n \frac{\partial l_t(\theta_0)}{\partial \theta} \\ &= \frac{1}{\sqrt{n}} \sum_{t=1}^n \left[\frac{c_1 \xi_{1t}}{\sqrt{\sigma_0^2 (1 + b_0^2 Y_{t-2}^2)}} + \frac{c_2 Y_{t-2} \xi_{1t}}{\sqrt{\sigma_0^2 (1 + b_0^2 Y_{t-2}^2)}} - \frac{c_3 \xi_{2t}}{2\sigma_0^2} - \frac{c_4 Y_{t-2}^2 \xi_{2t}}{2(1 + b_0^2 Y_{t-2}^2)} \right] \\ &\equiv \frac{1}{\sqrt{n}} \sum_{t=1}^n s_t, \end{split}$$

where

$$\xi_{1t} = \frac{\varepsilon_t + b_0 Y_{t-2} \varepsilon_{t-1}}{\sqrt{\sigma_0^2 (1 + b_0^2 Y_{t-2}^2)}}, \quad \xi_{2t} = 1 - \frac{(\varepsilon_t + b_0 Y_{t-2} \varepsilon_{t-1})^2}{\sigma_0^2 (1 + b_0^2 Y_{t-2}^2)}$$

Since $E(s_t s_{t+k}) = 0$ if $|k| \ge 2$, we have

$$\sigma_n^2 \equiv E\left(\frac{1}{\sqrt{n}}\sum_{t=1}^n s_t\right)^2 \to Es_t^2 + 2Es_ts_{t-1} = C^{\top}\Omega C,$$

as $n \to \infty$. Next we show that $\{s_t\}$ is $L^2(\nu)$ -near-epoch dependent series, that is,

$$E[s_t - E(s_t | \mathcal{F}_t^m)]^2 = O(m^{-\nu}),$$
(8)

for any $\nu > 2$ and large m, where $\mathcal{F}_t^m = \sigma\{\varepsilon_t, \cdots, \varepsilon_{t-m}\}$. Put

$$Y_{m,t} = \mu_0 + \varepsilon_t + \sum_{1 \le i \le m/2 - 1} \prod_{r=0}^{i-1} (\phi_0 + b_0 \varepsilon_{t-2r-1}) (\mu_0 + \varepsilon_{t-2i}).$$

Then $Y_{m,t} \in \mathcal{F}_t^m$. From the proof of Lemma 1, there exists a $\delta \in (0,1)$ such that $E|\phi_0 + b_0\varepsilon_t|^{\delta} < 1$. Thus, by Theorem 1, we have

$$E|Y_{t} - Y_{m,t}|^{\delta} = E \left| \sum_{i \ge m/2} \prod_{r=0}^{i-1} (\phi_{0} + b_{0}\varepsilon_{t-2r-1})(\mu_{0} + \varepsilon_{t-2i}) \right|^{\delta}$$

$$\leq \sum_{i \ge m/2} E \left| \prod_{r=0}^{i-1} (\phi_{0} + b_{0}\varepsilon_{t-2r-1})(\mu_{0} + \varepsilon_{t-2i}) \right|^{\delta}$$

$$= \sum_{i \ge m/2} \prod_{r=0}^{i-1} E |\phi_{0} + b_{0}\varepsilon_{t-2r-1}|^{\delta} E |\mu_{0} + \varepsilon_{t-2i}|^{\delta}$$

$$= O\left(\sum_{i \ge m/2} (E|\phi_{0} + b_{0}\epsilon_{1}|^{\delta})^{i-1}\right)$$

$$= O(\rho^{m}), \qquad (9)$$

where $\rho \in (0, 1)$. It follows from (9) that

$$E \left| \frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} - \frac{Y_{m,t-2}}{1+b_0^2 Y_{m,t-2}^2} \right| \\
 = \frac{2}{|b_0|} E \left| \frac{b_0 Y_{t-2}}{2(1+b_0^2 Y_{t-2}^2)} - \frac{b_0 Y_{m,t-2}}{2(1+b_0^2 Y_{m,t-2}^2)} \right| \\
 \le \frac{2}{|b_0|} E \left| \frac{b_0 Y_{t-2}}{2(1+b_0^2 Y_{t-2}^2)} - \frac{b_0 Y_{m,t-2}}{2(1+b_0^2 Y_{m,t-2}^2)} \right|^{\delta} \\
 \le \frac{2}{|b_0|} \left\{ E \left| \frac{b_0 Y_{t-2}}{2(1+b_0^2 Y_{t-2}^2)} - \frac{b_0 Y_{m,t-2}}{2(1+b_0^2 Y_{t-2}^2)} \right|^{\delta} + E \left| \frac{b_0 Y_{m,t-2}}{2(1+b_0^2 Y_{m,t-2}^2)} - \frac{b_0 Y_{m,t-2}}{2(1+b_0^2 Y_{m,t-2}^2)} \right|^{\delta} \right\} \\
 \le \frac{4|b_0|^{\delta}}{|b_0|} E|Y_{t-2} - Y_{m,t-2}|^{\delta} \\
 = O(\rho^m),
 \tag{10}$$

which implies that

$$\begin{split} E \left| \frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} - E\left(\frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} \Big| \mathcal{F}_t^m\right) \right| \\ &\leq E \left| \frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} - \frac{Y_{m,t-2}}{1+b_0^2 Y_{m,t-2}^2} \right| + E \left[E\left(\left| \frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} - \frac{Y_{m,t-2}}{1+b_0^2 Y_{m,t-2}^2} \right| \Big| \mathcal{F}_t^m\right) \right] \\ &= 2E \left| \frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} - \frac{Y_{m,t-2}}{1+b_0^2 Y_{m,t-2}^2} \right| \end{split}$$

$$= O(\rho^m). \tag{11}$$

Similar to (10), we can show hat

$$E\left|\frac{Y_{t-2}^2}{1+b_0^2 Y_{t-2}^2} - E\left(\frac{Y_{t-2}^2}{1+b_0^2 Y_{t-2}^2}\Big|\mathcal{F}_t^m\right)\right| = O(\rho^m)$$
(12)

for some $\delta \in (0, 1)$. Furthermore, since ε_t and ε_{t-1} are independent of Y_{t-2} and $Y_{t-2}^2/(1+b_0^2Y_{t-2}^2)$ is bounded, it follows from (11) and (12) that

$$\begin{split} & E \left| \frac{Y_{t-2}\xi_{1t}}{\sqrt{1+b_0^2 Y_{t-2}^2}} - E\left(\frac{Y_{t-2}\xi_{1t}}{\sqrt{1+b_0^2 Y_{t-2}^2}} \Big| \mathcal{F}_t^m \right) \right|^2 \\ & \leq 2E\varepsilon_t^2 E \left| \frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} - E\left(\frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} \Big| \mathcal{F}_t^m \right) \right|^2 + 2b_0^2 E\varepsilon_{t-1}^2 E \left| \frac{Y_{t-2}^2}{1+b_0^2 Y_{t-2}^2} - E\left(\frac{Y_{t-2}^2}{1+b_0^2 Y_{t-2}^2} \Big| \mathcal{F}_t^m \right) \right|^2 \\ & = O\left(E \left| \frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} - E\left(\frac{Y_{t-2}}{1+b_0^2 Y_{t-2}^2} \Big| \mathcal{F}_t^m \right) \right| \right) + O\left(E \left| \frac{Y_{t-2}^2}{1+b_0^2 Y_{t-2}^2} - E\left(\frac{Y_{t-2}^2}{1+b_0^2 Y_{t-2}^2} \Big| \mathcal{F}_t^m \right) \right| \right) \\ & = O(\rho^m). \end{split}$$

Similar inequalities hold for other terms in s_t and hence (8) holds. Therefore we conclude that $S_n \xrightarrow{d} N(0, C^{\top}\Omega C)$ by Theorem 21.1 in Billingsley (1968). Furthermore, by the Cramér-Wold device, we complete the proof.

Proof of Theorem 2. Part (a) follows from Theorem 1(a) in Ling and McAleer (2010) and Lemma 1 (Assumption 2(i) in that paper automatically holds since we only need one initial value). First, part (a) of this theorem implies that $\hat{\theta}_n$ converges a.s. to θ_0 . Second, $\frac{1}{n} \sum_{t=1}^n \frac{\partial^2 \ell_t(\theta)}{\partial \theta \partial \theta^{\top}}$ exists and is continuous in Θ . Third, it follows from Lemma 2(b)-(c) that $\frac{1}{n} \sum_{t=1}^n \frac{\partial^2 \ell_t(\hat{\theta}_n)}{\partial \theta \partial \theta^{\top}}$ converges to $-\Sigma$ in probability. Fourth, by Lemma 3, we have $\frac{1}{\sqrt{n}} \sum_{t=1}^n \frac{\partial \ell_t(\theta_0)}{\partial \theta} \xrightarrow{d} N(0, \Omega)$. Thus, all conditions in Theorem 4.1.3 in Amemiya (1985) hold, i.e., $\sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow{d} N(0, \Sigma^{-1}\Omega\Sigma^{-1})$.

Proof of Theorem 3. Note that (10) follows directly from (9) without the involved derivations. Hence, the theorem can be shown by repeating Lemmas 1–3 and using the same arguments in deriving (2.2) in Self and Liang (1987). \Box Acknowledgments. Ling's research was supported by the Hong Kong Research Grants Council (Grant HKUST641912, 603413 and FSGRF12SC12). Peng's research was supported by NSF grant DMS-1005336 and Simons Foundation. Zhu's research was supported by National Natural Science Foundation of China (11371168, 11271155), Specialized Research Fund for the Doctoral Program of Higher Education (20110061110003), Science and Technology Developing Plan of Jilin Province (20130522102JH) and Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry.

References

- [1] Amemiya, T. (1985). Advanced Econometrics. Cambridge: Harvard University Press.
- [2] Basrak, B. (2000). The Sample Autocorrelation Function of Non-linear Time Series. Ph.D. thesis, University of Groningen.
- [3] Basrak, B., Davis, R.A. and Mikosch, T. (1999). The sample ACF of a simple bilinear process. Stochastic Processes and their Applications, 83, 1-14.
- [4] Bibi, A. and Aknouche, A. (2010). Yule-Walker type estimators in periodic bilinear models: strong consistency and asymptotic normality. *Statistical Methods and Applications*, **19**, 1-30.
- [5] Bibi, A. and Gautier, A. (2010). Consistent and asymptotically normal estimators for periodic bilinear models. Bulletin of the Korean Mathematical Society, 47, 889-905.
- [6] Billingsley, P. (1968). Convergence of Probability Measures. New York: John Wiley & Sons Inc.
- Bougerol, P. and Picard, N. (1992). Strict stationarity of generalized autoregressive processes. Annals of Probability, 20, 1714-1730.
- [8] Davis, R.A. and Resnick, S.I. (1996). Limit theory for bilinear processes with heavy-tailed noise. *Annalsof Applied Probability*, 6, 1191-1210.
- [9] Francq, C. and Zakoïn, J.M. (2004). Maximum likelihood estimation of pure GARCH and ARMA-GARCH processes. *Bernoulli*, 10, 605-637.

- [10] Gabr, M.M. (1998). Robust estimation of bilinear time series models. Communications in Statistics-Theory and Methods, 27, 41-53.
- [11] Giordano, F. (2000). The variance of CLS estimators for a simple bilinear model. Quaderni di Statistica, 2, 147-155.
- [12] Giordano, F. and Vitale, C. (2003). CLS asymptotic variance for a particular relevant bilinear time series model. *Statistical Methods and Applications*, **12**, 169-185.
- [13] Grahn, T. (1995). A conditional least squares approach to bilinear time series estimation. Journal of Time Series Analysis, 16, 509-529.
- [14] Granger, C.W.J. and Andersen, A.P. (1978a). An Introduction to Bilinear Time Series Models.Göttingen: Vandenhoeck & Ruprecht.
- [15] Granger, C.W.J. and Andersen, A.P. (1978b). On the invertibility of time series models. Stochastic Processes and their Applications, 8, 87-92.
- [16] Guegan, D. and Pham, D.T. (1989). A note on the estimation of the parameters of the diagonal bilinear model by the method of least squares. Scandinavian Journal of Statistics, 16, 129-136.
- [17] Hili, O. (2008). Hellinger distance estimation of general bilinear time series models. Statistical Methodology, 5, 119-128.
- [18] Kesten, H. (1973). Random difference equations and renewal theory for products of random matrices. Acta Mathematica, 131, 207–248.
- [19] Kim, W.Y. and Billard, L. (1990). Asymptotic properties for the first-order bilinear time series model. Communications in Statistics-Theory and Methods, 19, 1171-1183.
- [20] Kim, W.Y., Billard, L. and Basawa, I.V. (1990). Estimation for the first-order diagonal bilinear time series model. *Journal of Time Series Analysis*, **11**, 215-229.
- [21] Kristensen, D. (2009). On stationarity and ergodicity of the bilinear model with applications to GARCH models. Journal of Time Series Analysis, 30, 125-144.
- [22] Ling, S. (2004). Estimation and testing stationarity for double-autoregressive models. Journal of the Royal Statistical Society Series B, 66, 63-78.

- [23] Ling, S. and McAleer, M. (2003). Asymptotic theory for a vector ARMA-GARCH model. *Econometric Theory*, **19**, 280-310.
- [24] Ling, S. and McAleer, M. (2010). A general asymptotic theory for time-series models. Statistica Neerlandica, 64, 97-111.
- [25] Liu, J. (1989). A simple condition for the existence of some stationary bilinear time series. Journal of Time Series Analysis, 10, 33-39.
- [26] Liu, J. (1990a). A note on causality and invertibility of a general bilinear time series model. Advances in Applied Probability, 22, 247-250.
- [27] Liu, J. (1990b). Estimation for some bilinear time series. Stochastic Models, 6, 649-665.
- [28] Liu, J. and Brockwell, P. J. (1988). On the general bilinear time series model. Journal of Applied Probability, 25, 553-564.
- [29] Pham, D.T. and Tran, L.T. (1981). On the first-order bilinear time series model. Journal of Applied Probability, 18, 617-627.
- [30] Self, S.G. and Liang, K.Y. (1987). Asymptotic properties of maximum likelihood estimators and likelihood ratio tests under nonstandard conditions. *Journal of the American Statistical Association*, 82, 605-610.
- [31] Sesay, S.A.O. and Subba Rao, T. (1992). Frequency-domain estimation of bilinear time series models. Journal of Time Series Analysis, 13, 521-545.
- [32] Subba Rao, T. (1981). On the theory of bilinear time series models. Journal of the Royal Statistical Society Series B, 43, 244-255.
- [33] Truquet, L. and Yao, J. (2012). On the quasi-likelihood estimation for random coefficient autoregressions. Statistics, 46, 505–521.
- [34] Turkman, K.F. and Turkman, M.A.A. (1997). Extremes of bilinear time series models. Journal of Time Series Analysis, 18, 305-319.
- [35] Wittwer, G. (1989). Some remarks on bilinear time series models. *Statistics*, **20**, 521-529.

[36] Zhang, Z. and Tong, H. (2001). On some distributional properties of a first-order nonnegative bilinear time series model. *Journal of Applied Probability*, 38, 659-671.