AN ELEMENTARY CHARACTERISATION OF SIFTED WEIGHTS

MATĚJ DOSTÁL AND JIŘÍ VELEBIL

ABSTRACT. Sifted colimits (those that commute with finite products in sets) play a major rôle in categorical universal algebra. For example, varieties of (many-sorted) algebras are precisely the free cocompletions under sifted colimits of (many-sorted) Lawvere theories. Such a characterisation does not depend on the existence of finite products in algebraic theories, but on the above fact that these products commute with sifted colimits and another condition: finite products form a *sound* class of limits.

In this paper we study the notion of soundness for general classes of weights in enriched category theory. We show that soundness of a given class of weights is equivalent to having a 'nice' characterisation of *flat* weights for that class. As an application, we give an elementary characterisation of sifted weights for the enrichment in categories and in preorders. We also provide a number of examples of sifted weights using our elementary criterion.

1. INTRODUCTION

The classical theory of locally presentable [8] and accessible categories [15], [20] (see also the more recent [2]) has been generalised to locally \mathbb{D} -presentable and \mathbb{D} -accessible categories for a 'good' class \mathbb{D} of small categories in [1]. The classical theory hinges a lot upon the interplay of two classes of categories: the class of λ -small categories for limits and the class of λ -filtered categories for colimits, where λ is a fixed regular cardinal. The precise nature of the interplay is that

 λ -small limits commute with λ -filtered colimits in the category of all sets and mappings.

The idea of [1] was to develop a more general theory of locally presentable and accessible categories based on the fact that one has a fixed class \mathbb{D} of small categories that replaces the class of λ -small categories. The corresponding class of colimits, called \mathbb{D} -filtered, is then defined by the requirement that

D-limits commute with D-filtered colimits in the category of sets and mappings.

It has been showed in [1] that a great deal of the classical theory can be developed for the concept of \mathbb{D} -filteredness, provided that the class \mathbb{D} satisfies a side condition that is called *soundness*.

Roughly speaking, sound classes \mathbb{D} allow for an easier detection of \mathbb{D} -filteredness: for a sound class \mathbb{D} a category \mathscr{E} is \mathbb{D} -filtered if the process of taking colimits over \mathscr{E} commutes with taking \mathbb{D} -limits of *representable functors*.

For example, the class \mathbb{D} consisting of finite discrete categories is sound. The corresponding \mathbb{D} -filtered colimits turn out to be precisely the *sifted* colimits of [16]. Free cocompletions of small categories under sifted colimits generalise the notion of a variety, as shown in [3]. In fact, the notion of a sifted colimit turned out to be a cornerstone notion in the categorical treatment of universal algebra, see [4].

One can pass from categories to categories enriched in a suitable monoidal \mathscr{V} and ask whether the above results can be reproduced. Since (co)limits over a class of categories have to be replaced by *weighted* (co)limits for \mathscr{V} -categories, one is naturally forced to define and study *soundness of classes of weights*. Definitions of soundness of a class of weights have appeared in [14] (Axiom A) and in Section 4 of [7]. The provided definitions of soundness, when specialised to categories enriched in sets, are, however, *weaker* than the definition in [1].

Results of the paper. For categories enriched in a general \mathscr{V} , we give in Section 3 a definition of soundness of a class Ψ of weights that is equivalent to the notion of [1] when \mathscr{V} is the category of sets. We show in Proposition 3.8 that all notions of soundness that have appeared in the literature coincide when the class Ψ satisfies two side conditions that often arise in practice. Namely, the class Ψ has to be *locally small* in the sense of [12] and *saturated* in the sense of [5] (there called a *closed* class).

Date: 9 May 2014.

Key words and phrases. Weighted colimits, sound classes, sifted colimits.

Matěj Dostál acknowledges the support by the grant No. SGS14/186/OHK3/3T/13 of the Grant Agency of the Czech Technical University in Prague. Jiří Velebil acknowledges the support by the grant No. P202/11/1632 of the Czech Science Foundation.

Our definition allows us to give, for a sound class Ψ , a characterisation of Ψ -flat weights (those weights φ such that φ -colimits commute with Ψ -limits in \mathscr{V}) by means of a certain coend. We show that this characterisation, when \mathscr{V} is the category of sets and \mathbb{D} is sound, boils down to the characterisation of \mathbb{D} -filteredness of [1] in terms of cocones for certain diagrams.

In Section 4 we turn the coend characterisation to a useful criterion of Ψ -flat weights when the sound class Ψ is the class of weights for finite products. Thus Ψ -flat weights are precisely the (enriched) *sifted* weights. Specialising to the enrichment in Cat, we can therefore deal with siftedness for 2-categories. We apply the coend criterion to give elementary proofs of siftedness of various weights used in 2-dimensional universal algebra, see, e.g. [6].

Acknowledgement. We thank John Bourke for the discussions concerning various aspects of siftedness.

2. Preliminaries on classes of weights

We introduce now the basic notation and results on weighted limits and colimits that we will need later. The material here is standard; for more details we refer to the book [10] and the paper [12].

We fix a complete and cocomplete symmetric monoidal category $\mathscr{V} = (\mathscr{V}_o, \otimes, I, [-, -])$. All categories, functors and natural transformations are to be understood as \mathscr{V} -categories, \mathscr{V} -functors and \mathscr{V} -natural transformations.

Limits and colimits. A weight is a functor $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$, where \mathscr{E} is small. Given a weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ and a diagram $D : \mathscr{E} \longrightarrow \mathscr{K}$, a colimit of D weighted by φ is an object $\varphi \star D$ together with an isomorphism

$$\mathscr{K}(\varphi \star D, X) \cong [\mathscr{E}^{op}, \mathscr{V}](\varphi, \mathscr{K}(D-, X))$$

natural in X. Dually, a *limit* of a diagram $D : \mathscr{E}^{op} \longrightarrow \mathscr{K}$ weighted by $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ is an object $\{\varphi, D\}$ together with an isomorphism

$$\mathscr{K}(X, \{\varphi, D\}) \cong [\mathscr{E}^{op}, \mathscr{V}](\varphi, \mathscr{K}(X, D-))$$

natural in X.

Given a weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$, a category \mathscr{K} is called φ -cocomplete if it has colimits $\varphi \star D$ of all diagrams $D : \mathscr{E} \longrightarrow \mathscr{K}$. Analogously, we define φ -completeness of \mathscr{K} .

Suppose \mathscr{K} and \mathscr{L} have φ -colimits. A functor $H : \mathscr{K} \longrightarrow \mathscr{L}$ between φ -cocomplete categories *preserves* φ -colimits (or, is φ -*cocontinuous*) if the canonical comparison $\varphi \star HD \longrightarrow H(\varphi \star D)$ is an isomorphism, for all $D : \mathscr{E} \longrightarrow \mathscr{K}$.

Even more generally, given a class Φ of weights, we say that \mathscr{K} is Φ -cocomplete (Φ -complete, resp.) if it has φ -colimits (φ -limits, resp.) for all φ in Φ . Analogously, we define Φ -cocontinuous (Φ -continuous, resp.) functors.

Free cocompletions and saturated classes of weights. Given a class Φ of weights, let us write U_{Φ} : Φ -Cocts \longrightarrow Cat for the forgetful 2-functor from the 2-category Φ -Cocts of Φ -cocomplete categories, Φ continuous functors and natural transformations. Then U_{Φ} has a left adjoint pseudofunctor, yielding a *free* Φ -cocompletion $\Phi(\mathscr{K})$ of any category \mathscr{K} .

For a small category \mathscr{E} , the free Φ -cocompletion $\Phi(\mathscr{E})$ can be computed via a transfinite process, namely the closure of \mathscr{E} in $[\mathscr{E}^{op}, \mathscr{V}]$ under Φ -colimits.¹ We will need the first step $\Phi_1(\mathscr{E})$ of this process: $\Phi_1(\mathscr{E})$ is the full subcategory of $[\mathscr{E}^{op}, \mathscr{V}]$ spanned by Φ -colimits of representables. Hence a weight $\alpha : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ is in $\Phi_1(\mathscr{E})$ iff it is of the form $\operatorname{Lan}_{T^{op}}\varphi$ for some $T : \mathscr{D} \longrightarrow \mathscr{E}$ and some weight $\varphi : \mathscr{D}^{op} \longrightarrow \mathscr{V}$ in Φ .

The class Φ is called *saturated* (the concept introduced in [5], there called *closed*) if, for any small category \mathscr{E} , the free cocompletion $\Phi(\mathscr{E})$ consists precisely of all the weights $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ that belong to Φ . If we put Φ^* to be the largest class such that the 2-categories Φ^* -Cocts and Φ -Cocts coincide, then Φ^* is saturated and it is the least saturated class containing Φ .

Commutation of limits and colimits, flatness. Let Φ and Ψ be classes of weights. We say that Φ colimits commute with Ψ -limits in \mathscr{V} , if for any $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ in Φ , the functor

$$\varphi \star (-) : [\mathscr{E}, \mathscr{V}] \longrightarrow \mathscr{V}$$

preserves Ψ -limits. We denote by Ψ^+ the class of Ψ -flat weights, i.e., all weights φ such that φ -colimits commute with Ψ -limits in \mathscr{V} .

¹In fact, the same transfinite process can be applied to obtain $\Phi(\mathscr{K})$ for any category \mathscr{K} . There is, however, a slight technicality concerning size when \mathscr{K} is not small. Since we will not need $\Phi(\mathscr{K})$ for large \mathscr{K} , we refer to [10] for more details.

Example 2.1. Suppose \mathscr{V} is cartesian closed. By $\text{const}_1 : \mathscr{D}^{op} \longrightarrow \mathscr{V}$ we denote the weight that is constantly the terminal object 1. Such weights will be called *conical*. Any class \mathbb{D} of small categories induces a class

$\Psi_{\mathbb{D}}$

of conical weights $const_1 : \mathscr{D}^{op} \longrightarrow \mathscr{V}$ with \mathscr{D}^{op} in \mathbb{D} .

- (1) Suppose $\mathscr{V} = \mathsf{Set}$. Then to say that a small category \mathscr{E} is \mathbb{D} -filtered in the sense of [1] is to say that the conical weight $\mathsf{const}_1 : \mathscr{E}^{op} \longrightarrow \mathsf{Set}$ is $\Psi_{\mathbb{D}}$ -flat. Indeed: (co)limits of diagrams weighted by conical weights yield the usual notions defined by (co)cones.
- (2) Suppose \mathscr{V} is arbitrary (but still cartesian closed). The class $\Psi_{\mathbb{D}}$ for \mathbb{D} consisting of all finite discrete categories will be denoted by Π . The corresponding class of Π -flat weights is called the class of *sifted* weights. We will say more on sifted weights in Section 4.

3. Sound classes of weights

In this section we generalise the definition of a sound class \mathbb{D} of small (ordinary) categories to soundness of a class Ψ of weights for a general \mathscr{V} .

Soundness of a class \mathbb{D} of small categories was defined (in case $\mathscr{V} = \mathsf{Set}$) in [1] using connectedness of a certain category of cocones. Since cocones are not available for general \mathscr{V} , we use a different phrasing (already implicit in [1]). We prove in Proposition 3.5 below that our definition coincides with that of [1] in case $\mathscr{V} = \mathsf{Set}$.

For any Ψ -flat weight φ , the functor $\varphi \star (-)$ is obliged to preserve all Ψ -limits by the definition of Ψ -flatness (see Section 2). Soundness of Ψ means that we can choose a *smaller* class of Ψ -limits to detect Ψ -flatness.

Definition 3.1. A class Ψ of weights is called *sound* if a weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ is Ψ -flat whenever the functor

$$\varphi \star (-) : [\mathscr{E}, \mathscr{V}] \longrightarrow \mathscr{V}$$

preserves Ψ -limits of representables.

Example 3.2.

(1) In case $\mathscr{V} = \mathsf{Set}$, the list

 \mathbb{D} =finite categories, \mathbb{D} =finite discrete categories, \mathbb{D} =empty class

yields a list of sound classes $\Psi_{\mathbb{D}}$ of weights by Example 2.3 of [1].

By the same example, the one-element class \mathbb{D} consisting of the scheme for pullbacks, or the twoelement class \mathbb{D} consisting of the scheme for pullbacks and terminal objects, yield classes $\Psi_{\mathbb{D}}$ that are not sound.

- (2) It has been proved in [11] that the class Π of weights for finite products is sound, for every cartesian closed \mathscr{V} .
- (3) The class \mathcal{P} of all weights is sound for any \mathscr{V} . Indeed, the class of all \mathcal{P} -flat weights is precisely the class \mathcal{Q} of all small-projective weights by Proposition 6.20 of [12]. By the same proposition, small-projectivity of $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ can be detected by the fact that $\varphi \star (-) : [\mathscr{E}, \mathscr{V}] \longrightarrow \mathscr{V}$ preserves a particular limit of representables, namely the limit $\{\varphi, Y\}$, where $Y : \mathscr{E}^{op} \longrightarrow [\mathscr{E}, \mathscr{V}]$ is the Yoneda embedding. Thus, \mathcal{P} is sound.
- (4) The class Ω of all small-projective weights is sound for any \mathscr{V} . By Remark 8.17 of [12], the class of Ω -flat weights coincides with the class \mathcal{P} of all weights. Hence the condition on soundness is vacuous.

The following easy result shows that the 'testing weights' for Ψ -flatness can be taken in a special form:

Proposition 3.3. For a class Ψ the following are equivalent:

- (1) Ψ is sound.
- (2) The weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ is Ψ -flat, whenever $\varphi \star (-)$ preserves $\Psi_1(\mathscr{E})$ -limits of representables, i.e., whenever the canonical morphism

$$\operatorname{can}: \varphi \star \{\psi, Y-\} \longrightarrow \{\psi, \varphi\} \tag{3.1}$$

is an isomorphism, for every $\psi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ in $\Psi_1(\mathscr{E})$.

Proof. Let $Y : \mathscr{E}^{op} \longrightarrow [\mathscr{E}, \mathscr{V}]$ be the Yoneda embedding. Definition 3.1 requires the canonical morphism

$$\varphi \star \{\psi, YT^{op}-\} \longrightarrow \{\psi, \varphi \cdot T^{op}\}$$

to be an isomorphism, for every $\psi : \mathscr{D}^{op} \longrightarrow \mathscr{V}$ in Ψ and every $T : \mathscr{D} \longrightarrow \mathscr{E}$.

The weight $\operatorname{Lan}_{T^{op}}\psi: \mathscr{E}^{op} \longrightarrow \mathscr{V}$ is in $\Psi_1(\mathscr{E})$ and every weight in $\Psi_1(\mathscr{E})$ has this form, for some $\psi: \mathscr{D}^{op} \longrightarrow \mathscr{V}$ in Ψ and some $T: \mathscr{D} \longrightarrow \mathscr{E}$.

Since there are isomorphisms

$$\psi, YT^{op}\} \cong \{\operatorname{Lan}_{T^{op}}\psi, Y\}, \quad \{\psi, \varphi \cdot T^{op}\} \cong \{\operatorname{Lan}_{T^{op}}\psi, \varphi\}$$

the equivalence of (1) and (2) follows.

ł

The canonical morphism in (3.1) can be rewritten using coends and Yoneda Lemma as the morphism

$$\operatorname{can}: \int^{e} [\mathscr{E}^{op}, \mathscr{V}](Ye, \varphi) \otimes [\mathscr{E}^{op}, \mathscr{V}](\psi, Ye) \longrightarrow [\mathscr{E}^{op}, \mathscr{V}](\psi, \varphi)$$
(3.2)

that is given by composition in $[\mathscr{E}^{op}, \mathscr{V}]$. We illustrate now on two well-known classes that this coend description yields precisely the 'classical' description of flatness by means of the category of cocones.

Example 3.4 (Sifted weights and flat weights for $\mathscr{V} = \mathsf{Set}$). Suppose $\mathscr{V} = \mathsf{Set}$. Recall that every $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Set}$ has a *category of elements* $\mathsf{elts}(\varphi)$: the objects are pairs (x, e) with $x \in \varphi e$ and a morphism from (x, e) to (x', e') is a morphism $t : e \longrightarrow e'$ in \mathscr{E} such that $\varphi t(e') = e$ holds.

- (1) Let Π be the sound class of weights for finite products. The category $\Pi_1(\mathscr{E})$ is spanned by finite coproducts of representables in $[\mathscr{E}^{op}, \mathsf{Set}]$. Hence a general testing weight $\psi : \mathscr{E}^{op} \longrightarrow \mathsf{Set}$ for Π -flatness by Proposition 3.3 has the form $\coprod_{i \in I} Ye_i$ where I is a finite set.
 - We show now that (3.2) yields the well-known characterisation of *sifted* weights, see [16]. Indeed, given a general weight $\varphi : \mathcal{E}^{op} \longrightarrow \mathsf{Set}$, the mapping can has the form

$$\mathsf{can}: \int \ \varphi e \times \prod_{i \in I} \mathscr{E}(e_i, e) \longrightarrow \prod_{i \in I} \varphi e_i, \quad [(x, (t_i))] \mapsto (\varphi t_i(x))$$

Hence can is a bijection if and only if the following two conditions hold:

(a) The mapping can is surjective, i.e., for every element of $\prod_{i \in I} \varphi e_i$, i.e., for every *I*-tuple (x_i) of elements of φ there is an e, an element $x \in \varphi e$ and an *I*-tuple $t_i : e_i \longrightarrow e$ of morphisms in \mathscr{E} such that $\varphi t_i(x) = x_i$.

Briefly: on every *I*-tuple of objects of $elts(\varphi)$ there is a cocone.

(b) The mapping can is injective, i.e., for any pair (x, (t_i)), (x', (t'_i)) such that φt_i(x) = φt'_i(x') holds for all i, i.e., for any two cocones of the same I-tuple of objects of elts(φ), there is a zig-zag in & that connects these cocones in elts(φ).

To summarise: a weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Set}$ is Π -flat iff its category of elements is sifted (every finite family of elements has a cocone and every two cocones for the same finite family are connected by a zig-zag).

(2) Let Ψ be the sound class of finite (conical) limits, i.e., let Ψ = Ψ_D for the class D of finite categories. The category Ψ₁(𝔅) is spanned by finite colimits of representable functors in [𝔅^{op}, Set]. Thus, a general testing weight ψ : 𝔅^{op} → Set for Ψ-flatness has the form ψ = colim YC for a diagram C : 𝔅 → 𝔅 with 𝔅 finite.

Given a general weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Set}$, the mapping can is a bijection iff two conditions hold:

- (a) The mapping can is surjective, i.e., every finite diagram in $elts(\varphi)$ has a cocone.
- (b) The mapping can is injective, i.e., any two cocones for the same finite diagram in elts(φ) are connected by a zig-zag in elts(φ).

The above two conditions together state that the category of cocones of finite diagrams in $\mathsf{elts}(\varphi)$ is nonempty and connected. This means that the category $\mathsf{elts}(\varphi)$ is filtered. As expected, Ψ -flat weights are precisely the flat ones.

In both cases above, the classes of testing weights can be simplified. For example, for siftedness, one can choose only nullary coproduct of representables and binary coproducts of representables as the testing weights. We will use this fact in Section 4 below.

We prove now that Definition 3.1 coincides with the definition of soundness from [1] (this definition is condition (2) of the proposition).

Proposition 3.5. Suppose $\mathscr{V} = \mathsf{Set.}$ For a class \mathbb{D} of small categories, the following conditions are equivalent:

- (1) The class $\Psi_{\mathbb{D}}$ of conical weights $\text{const}_1 : \mathscr{D}^{op} \longrightarrow \text{Set with } \mathscr{D}^{op}$ in \mathbb{D} is sound.
- (2) A category \mathscr{E} is \mathbb{D} -filtered whenever the category of cocones for any functor $T : \mathscr{D} \longrightarrow \mathscr{E}$ with \mathscr{D}^{op} in \mathbb{D} is nonempty and connected.

Proof. We will use the canonical morphism (3.2). Observe first that $[\mathscr{E}^{op}, \mathsf{Set}](\psi, \mathsf{const}_1)$ is a one-element set for any small category \mathscr{E} and any $\psi : \mathscr{E}^{op} \longrightarrow \mathsf{Set}$, since const_1 is a terminal object in $[\mathscr{E}^{op}, \mathsf{Set}]$.

By Proposition 3.3 any testing weight $\psi : \mathscr{E}^{op} \longrightarrow \mathsf{Set}$ for $\Psi_{\mathbb{D}}$ -flatness of $\mathsf{const}_1 : \mathscr{E}^{op} \longrightarrow \mathsf{Set}$ has the form $\operatorname{Lan}_{T^{op}}\mathsf{const}_1$ for some $T : \mathscr{D} \longrightarrow \mathscr{E}$, where $\mathsf{const}_1 : \mathscr{D}^{op} \longrightarrow \mathsf{Set}$ is in $\Psi_{\mathbb{D}}$. The left-hand side of (3.2) therefore has the form

$$\int^{e} [\mathscr{E}^{op}, \mathsf{Set}](\mathrm{Lan}_{T^{op}}\mathsf{const}_{1}, Ye) \cong \int^{e} [\mathscr{D}^{op}, \mathsf{Set}](\mathsf{const}_{1}, Ye \cdot T^{op}) \cong \int^{e} [\mathscr{D}^{op}, \mathsf{Set}](\mathsf{const}_{1}, \mathscr{E}(T-, e))$$

Observe that the category of elements of $[\mathscr{D}^{op}, \mathsf{Set}](\mathsf{const}_1, \mathscr{E}(T-, e))$ is precisely the category of cocones for T that have e as a vertex.

Thus (3.2) is a bijection iff

$$\int^{e} [\mathscr{D}^{op}, \mathsf{Set}](\mathsf{const}_1, \mathscr{E}(T-, e)) \cong 1$$

holds. From this, the equivalence of (1) and (2) follows immediately.

There is another important issue related to sound classes of weights. Adapting freely the terminology of [4], we may call a small Ψ -complete category \mathscr{T} a Ψ -theory. The category Ψ -Alg(\mathscr{T}) of Ψ -algebras for \mathscr{T} is the full subcategory of [\mathscr{T}, \mathscr{V}] spanned by functors that preserve Ψ -limits.

By definition, Ψ -flat colimits commute with Ψ -limits. Hence the category Ψ -Alg(\mathscr{T}) is closed in $[\mathscr{T}, \mathscr{V}]$ under Ψ -flat colimits and it contains the representables, for any Ψ -theory \mathscr{T} . Therefore, for any class Ψ and any Ψ -theory \mathscr{T} , there is an inclusion

$$\Psi^+(\mathscr{T}^{op}) \subseteq \Psi\text{-}\mathsf{Alg}(\mathscr{T})$$

since $\Psi^+(\mathscr{T}^{op})$ is the closure in $[\mathscr{T}, \mathscr{V}]$ of the representables under Ψ -flat colimits. We discuss now the case when the above inclusion is an equality.

Lemma 3.6. Suppose the class Ψ is sound. Then $\Psi^+(\mathscr{T}^{op}) = \Psi$ -Alg (\mathscr{T}) holds for any Ψ -theory \mathscr{T} .

Proof. Suppose $\varphi : \mathscr{T} \longrightarrow \mathscr{V}$ preserves Ψ -limits. Then $\varphi \star (-) : [\mathscr{T}^{op}, \mathscr{V}] \longrightarrow \mathscr{V}$ preserves Ψ -limits of representables. By soundness of Ψ , this means that φ is Ψ -flat. Hence φ is in $\Psi^+(\mathscr{T}^{op})$.

Remark 3.7. The equality $\Psi^+(\mathscr{T}^{op}) = \Psi - \mathsf{Alg}(\mathscr{T})$ for any Ψ -theory \mathscr{T} is an important fact that allows for the development of an abstract theory of 'algebras' for Ψ .

For example, when $\mathscr{V} = \mathsf{Set}$ and $\Psi = \Pi$, much of the theory of varieties of algebras hinges upon the above equality, see [4]. Namely, a Π -theory \mathscr{T} is then precisely a (many-sorted) Lawvere theory, and a functor $\varphi : \mathscr{T} \longrightarrow \mathsf{Set}$ is an algebra for \mathscr{T} iff φ is a sifted weight.

Another instance of the above is the equality, for any category \mathscr{T} with finite limits, of the free cocompletion $\operatorname{Ind}(\mathscr{T}^{op})$ of \mathscr{T}^{op} under filtered colimits and the category $\operatorname{Lex}(\mathscr{T},\operatorname{Set})$ of functors preserving finite limits. This coincidence is vital in interpreting locally finitely presentable categories as categories of algebras for essentially algebraic theories, see [2].

In Remark 2.6 of [1], the authors present the class $\Psi_{\mathbb{D}}$ for $\mathbb{D}=(\text{pullbacks+terminal object})$ as the example of a class that is *not sound*, yet the equality $\Psi_{\mathbb{D}}^+(\mathscr{T}^{op}) = \Psi_{\mathbb{D}}\text{-}\mathsf{Alg}(\mathscr{T})$ holds. We show now that such a counterexample is essentially due to the fact that $\Psi_{\mathbb{D}}$ is not saturated.

In what follows we require a free Ψ -theory to exist on every small category. More precisely, we require the class Ψ to be *locally small* (see [12]): the category $\Psi(\mathscr{D})$ is small for every small \mathscr{D} .

Proposition 3.8. Suppose the class Ψ is locally small and saturated. Then the following are equivalent:

- (1) Ψ is sound.
- (2) $\Psi^+(\mathscr{T}^{op}) = \Psi\text{-}\mathsf{Alg}(\mathscr{T})$ holds for any $\Psi\text{-}theory \mathscr{T}$.

Proof. It suffices to prove that (2) implies (1). Consider any weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathscr{V}$ such that the functor

$$F \equiv \varphi \star (-) : [\mathscr{E}, \mathscr{V}] \longrightarrow \mathscr{V}$$

preserves Ψ -limits of representables. We prove that F preserves all Ψ -limits.

Denote by

$$\overset{Y_{\mathscr{E}^{op}}}{\overbrace{Z^{op}}} \mathscr{C}^{op} \xrightarrow{W} [\mathscr{E}, \mathscr{V}]$$

the factorisation of the Yoneda embedding where $Z^{op} : \mathscr{E}^{op} \longrightarrow \mathscr{C}^{op}$ is the closure of \mathscr{E}^{op} in $[\mathscr{E}, \mathscr{V}]$ under Ψ -limits. Notice that \mathscr{C}^{op} is small since Ψ is locally small.

 \square

By the construction, \mathscr{C}^{op} is a Ψ -theory and the functor $W : \mathscr{C}^{op} \longrightarrow [\mathscr{E}, \mathscr{V}]$ preserves Ψ -limits. Furthermore, every object of \mathscr{C}^{op} is a Ψ -limit of representables, since Ψ is saturated. Hence the composite

$$\overline{\varphi} \equiv \mathscr{C}^{op} \xrightarrow{W} [\mathscr{E}, \mathscr{V}] \xrightarrow{F} \mathscr{V}$$

preserves Ψ -limits. By our assumption (2), the equality $\Psi^+(\mathscr{C}) = \Psi$ -Alg (\mathscr{C}^{op}) holds. Hence the functor $\overline{\varphi} \star (-) : [\mathscr{C}, \mathscr{V}] \longrightarrow \mathscr{V}$ preserves Ψ -limits.

Moreover,

$$\overline{\varphi} \star (-) \cong (-) \star FW$$

holds, since $\overline{\varphi} \star (-) \cong \operatorname{Lan}_{Y_{\mathscr{C}^{op}}} \overline{\varphi}$, and $\operatorname{Lan}_{Y_{\mathscr{C}^{op}}} W \cong (-) \star W$, and F preserves $\operatorname{Lan}_{Y_{\mathscr{C}^{op}}} W$ since $F = \varphi \star (-)$ preserves colimits. See the diagram



In the adjunction $(-) \star W \dashv \widetilde{W} : [\mathscr{E}, \mathscr{V}] \longrightarrow [\mathscr{C}, \mathscr{V}]$ the functor \widetilde{W} is fully faithful, since W is dense. Thus, the composite

$$[\mathscr{E},\mathscr{V}] \xrightarrow{\widetilde{W}} [\mathscr{C},\mathscr{V}] \xrightarrow{(-)\star W} [\mathscr{E},\mathscr{V}] \xrightarrow{F} \mathscr{V}$$

is isomorphic to $F = \varphi \star (-)$ and it preserves all Ψ -limits. We proved that φ is Ψ -flat; the proof is finished. \Box

Example 3.9. Local smallness of a class Ψ is a nontrivial property.

- (1) Every small class is locally small. Example 15 of [21] may be modified to show that, when $\mathscr{V} = \mathsf{Set}$, there exists a locally small class Ψ such that Ψ is not contained in the saturation of any small class of weights.
- (2) Every subclass of a locally small class is locally small again. The saturation of a locally small class is locally small again.
- (3) The class \mathcal{P} of all weights is typically not locally small. For example, $\mathcal{P}(\mathscr{D}) = [\mathscr{D}^{op}, \mathsf{Set}]$ in case $\mathscr{V} = \mathsf{Set}$. In fact, \mathcal{P} is locally small iff \mathscr{V} is small, i.e., iff \mathscr{V} is a commutative quantale. If \mathscr{V} is a commutative quantale, then every class of weights is locally small.
- (4) Even classes substantially smaller than \mathcal{P} may not be locally small. For example, consider the class \mathcal{Q} of small-projective weights that yields the Cauchy completion $\mathcal{Q}(\mathcal{D})$ of the category \mathcal{D} , see [12].
 - (a) The class Ω is locally small, whenever \mathscr{V} is locally presentable as a monoidal category by Theorem 6 of [13].
 - (b) However, in case \mathscr{V} is the monoidal closed category Sup of complete join-semilattices and joinpreserving maps, the class \mathcal{Q} is not locally small by Section 1 of [13]. In this enrichment, the Cauchy completion $\mathcal{Q}(\mathscr{D})$ of a small category \mathscr{D} always contains all small coproducts of representables.

Remark 3.10. We do not know whether local smallness of Ψ can be omitted from the assumptions of Proposition 3.8.

We can combine the above with Theorem 8.11 of [12] to obtain further characterisation of soundness of locally small saturated classes.

Corollary 3.11. For a locally small saturated class Ψ , the following conditions are equivalent:

- (1) Ψ is sound.
- (2) $\Psi^+(\mathscr{T}^{op}) = \Psi \operatorname{-Alg}(\mathscr{T})$ holds for any Ψ -theory \mathscr{T} .
- (3) For any small \mathscr{D} , every weight $\varphi : \mathscr{D}^{op} \longrightarrow \mathscr{V}$ is a Ψ -flat colimit of a diagram in $\Psi(\mathscr{D})$.
- (4) For any small \mathscr{D} , the closure of $\Psi(\mathscr{D})$ in $[\mathscr{D}^{op}, \mathscr{V}]$ under Ψ -flat colimits is all of $[\mathscr{D}^{op}, \mathscr{V}]$.

Remark 3.12. The class $\Psi_{\mathbb{D}}$ for $\mathbb{D}=(\text{pullbacks}+\text{terminal object})$ is locally small and not saturated. It is also not a sound class by Remark 2.6 of [1]. The saturation $\Psi_{\mathbb{D}}^*$ is the class of finitely presentable weights: this is a locally small, saturated and sound class.

In fact, one can always assume that Ψ is a locally small saturated class, since it is easy to prove the following:

If Ψ is a locally small and sound class, so is Ψ^* .

Indeed, if Ψ is locally small, then so is Ψ^* . Suppose Ψ is sound. For proving soundness of Ψ^* , suppose \mathscr{T} is a Ψ^* -theory. By Lemma 3.6 we have the equality $\Psi^+(\mathscr{T}^{op}) = \Psi$ -Alg (\mathscr{T}) . Moreover, the equality Ψ -Alg $(\mathscr{T}) = \Psi^*$ -Alg (\mathscr{T}) holds by the definition of Ψ^* . Furthermore, the equality $\Psi^+ = \Psi^{*+}$ holds by Proposition 5.4 of [12]. Hence $\Psi^{*+}(\mathscr{T}^{op}) = \Psi^*$ -Alg (\mathscr{T}) holds for every Ψ^* -theory \mathscr{T} . The class Ψ^* , being locally small and saturated, is sound by Proposition 3.8.

4. An elementary characterisation of sifted weights

In this section we analyse the isomorphism (3.2) in more detail for the enrichment in Cat. We then turn the analysis into a useful elementary criterion of *siftedness* of weights enriched in Cat. Finally, we comment on similarities and differences in using the criterion for siftedness in another '2-dimensional enrichment', namely that in Pre (the category of preorders and monotone maps).

An analysis of the coend in (3.2). Suppose that $\mathscr{V} = \mathsf{Cat}$. Let $\psi, \varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ be any weights. Then the coend

$$\int^{e} [\mathscr{E}^{op}, \mathsf{Cat}](Ye, \varphi) \times [\mathscr{E}^{op}, \mathsf{Cat}](\psi, Ye)$$
(4.1)

is a category that can be computed as a coequaliser in Cat of the parallel pair

$$\coprod_{e,e'} [\mathscr{E}^{op}, \mathsf{Cat}](Ye', \varphi) \times \mathscr{E}(e, e') \times [\mathscr{E}^{op}, \mathsf{Cat}](\psi, Ye) \xrightarrow{L}_{R} \coprod_{e} [\mathscr{E}^{op}, \mathsf{Cat}](Ye, \varphi) \times [\mathscr{E}^{op}, \mathsf{Cat}](\psi, Ye)$$
(4.2)

of functors

$$\begin{array}{l} L: (\widehat{x}: Ye' \longrightarrow \varphi, f: e \longrightarrow e', \tau: \psi \longrightarrow Ye) \mapsto (\widehat{x} \cdot Yf: Ye \longrightarrow \varphi, \tau: \psi \longrightarrow Ye) \\ R: (\widehat{x}: Ye' \longrightarrow \varphi, f: e \longrightarrow e', \tau: \psi \longrightarrow Ye) \mapsto (\widehat{x}: Ye' \longrightarrow \varphi, Yf \cdot \tau: \psi \longrightarrow Ye') \end{array}$$

Thus the coend (4.1) has the following description (see, e.g., [17]):

(1) The objects are equivalence classes

$$(\widehat{x}, \tau)]_{\sim}$$

where $\hat{x}: Ye \longrightarrow \varphi$ and $\tau: \psi \longrightarrow Ye$ are natural transformations. The equivalence is generated by

$$(\widehat{x}, Yf \cdot \tau) \sim (\widehat{x} \cdot Yf, \tau)$$

for all $\widehat{x}: Ye \longrightarrow \varphi, \, \tau: \psi \longrightarrow Ye' \, f: e' \longrightarrow e \text{ in } \mathscr{D}.$

(2) The morphisms are equivalence classes

$$[((u_1,v_1),\ldots,(u_n,v_n))]_{\approx}$$

of finite sequences $((u_1, v_1), \ldots, (u_n, v_n))$ such that every pair (u_i, v_i) is a morphism in the category $\coprod_e [\mathscr{E}^{op}, \mathsf{Cat}](Ye, \varphi) \times [\mathscr{E}^{op}, \mathsf{Cat}](\psi, Ye)$ and

$$cod(u_1, v_1) \sim dom(u_2, v_2), \quad cod(u_2, v_2) \sim dom(u_3, v_3), \quad \dots, \quad cod(u_{n-1}, v_{n-1}) \sim dom(u_n, v_n)$$

The equivalence relation \approx is generated from the following two conditions

 $(u * Yw, v) \approx (u, Yw * v), \quad ((u_1, v_1), (u_2, v_2)) \approx (u_2 \cdot u_1, v_2 \cdot v_1)$

by reflexivity, symmetry, transitivity and composition (concatenation). Above, by * we denote the horizontal composition of natural transformations.

It will be useful to work with the following graphical representation. The sequence $((u_1, v_1), \ldots, (u_n, v_n))$ as above is going to be depicted as

The above picture is called a hammock from (\hat{x}, τ) to (\hat{x}', τ') . The wiggly arrow in the above hammock, for example from (\hat{x}, τ) to (\hat{x}_1, τ_1) , represents a zig-zag connecting e and e_1 in \mathscr{E} that witnesses the equivalence $(\hat{x}, \tau) \sim (\hat{x}_1, \tau_1)$.

The whole hammock (4.3) gets evaluated to the composite modification

$$(u_n * v_n) \cdot (u_{n-1} * v_{n-1}) \cdot \cdots \cdot (u_1 * v_1) : \widehat{x} \cdot \tau \longrightarrow \widehat{x}' \cdot \tau'$$

in $[\mathscr{E}^{op}, \mathsf{Cat}]$. Up to the equivalence \approx , this is how the evaluation functor can works.

The functor **can** is an isomorphism of categories iff it is bijective on objects and fully faithful. Hence, the following two conditions have to hold:

- (1) The 1-dimensional aspect. To give $\alpha: \psi \longrightarrow \varphi$ is to give a unique $[(\hat{x}, \tau)]_{\sim}$ such that $\hat{x} \cdot \tau = \alpha$ holds.
- (2) The 2-dimensional aspect. To give a modification $\Xi : \alpha \longrightarrow \alpha'$ is to give a unique equivalence class $[((u_1, v_1), \dots, (u_n, v_u))]_{\approx}$ such that Ξ is the composite $(u_n * v_n) \cdots (u_1 * v_1)$.

Siftedness for enrichment in categories. We are going to fix the class Π of (conical) weights for finite products, see Example 2.1. Recall, by Example 2.1 again, that the Π -flat weights are called sifted.

It is proved in [11] that the class $\Pi_1(\mathscr{E})$ of testing weights for siftedness of a weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ can be reduced further to the empty coproduct $\mathsf{const}_0 : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ of representables and to binary coproducts $\mathscr{E}(-, e_1) + \mathscr{E}(-, e_2) : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$. Hence, using (3.2), the following result holds:

Lemma 4.1. A weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ is sifted iff the following two conditions hold:

- (1) The unique functor from $\int^e \varphi e$ to the one-morphism category 1 is an isomorphism.
- (2) For any e_1 , e_2 in \mathscr{E} , the canonical morphism

$$\mathsf{can}: \int^e \varphi e \times \mathscr{E}(e_1, e) \times \mathscr{E}(e_2, e) \longrightarrow \varphi e_1 \times \varphi e_2$$

is an isomorphism.

Remark 4.2. By analogy to the case $\mathscr{V} = \mathsf{Set}$, we may call the first condition above *connectedness* of the weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ and the second condition expresses that the diagonal 2-functor $\Delta : \mathscr{E} \longrightarrow \mathscr{E} \times \mathscr{E}$ is *cofinal* in the sense that the 2-cell

$$\mathcal{E}^{op} \xrightarrow{\Delta^{op}} \mathcal{E}^{op} \times \mathcal{E}^{op}$$

$$\varphi \xrightarrow{\delta} \mathcal{E}^{op} \times \mathcal{E}^{op}$$

$$\varphi \xrightarrow{\delta} \mathcal{E}^{op} \times \mathcal{E}^{op}$$

$$\varphi \xrightarrow{\delta} \mathcal{E}^{op} \times \mathcal{E}^{op}$$

where $\delta_e : \varphi e \longrightarrow \varphi e \times \varphi e$ is the diagonal functor, is a left Kan extension. Indeed, it suffices to consider the isomorphism

$$\int^{e} \varphi e \times \mathscr{E}(e_1, e) \times \mathscr{E}(e_2, e) \cong \int^{e} \varphi e \times (\mathscr{E}^{op} \times \mathscr{E}^{op})(\Delta^{op} e, (e_1, e_2))$$

We apply the criteria of Lemma 4.1, together with the analysis of (3.2) using hammocks, for giving elementary proofs of siftedness of various weights.

Example 4.3 (A weight that is not sifted). We start with an example of a weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ that is not sifted, although the 'underlying' ordinary functor

$$\mathscr{E}_{o}^{op} \xrightarrow{\varphi_{o}} \operatorname{Cat}_{o} \xrightarrow{\operatorname{ob}} \operatorname{Set}$$

is sifted.

Consider the one-morphism category \mathscr{S} with the only object s. Denote by \mathscr{E}^{op} the free completion of \mathscr{S} under finite products. It follows immediately that the only 2-cells in \mathscr{E}^{op} are identities.

Let $\chi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ be the product-preserving functor defined by $\chi(s) = 2$, where 2 is the two-element chain, considered as a category. We define φ to be the following modification of χ : where $\chi(s^n) = 2^n$, we let $\varphi(s^n) = 2^n$ for every n > 1. The structure on 2^n is that of an almost discrete preorder with the only nontrivial inequality being $(0, \ldots, 0) \leq (1, \ldots, 1)$. The action of φ on morphisms is defined as for χ . Of course, φ does *not* preserve products, but the composite

$$\mathscr{E}_{o}^{op} \xrightarrow{\varphi_{o}} \mathsf{Cat}_{o} \xrightarrow{\mathsf{ob}} \mathsf{Set}$$

does; in fact, it is not hard to see that this ordinary functor constitutes an algebra for the ordinary algebraic theory \mathscr{E}_{o}^{op} and thus it is a sifted weight by [4].

It is enough now to find pairs (x_1, x_2) and (y_1, y_2) from $\varphi(s) \times \varphi(s)$ such that $(x_1, x_2) \leq (y_1, y_2)$ holds but there is no hammock to witness this inequality. Consider $(x_1, x_2) = (0, 1)$ and $(y_1, y_2) = (1, 1)$. Firstly, we make use of the fact that there are no nontrivial 2-cells in \mathscr{E}^{op} . This implies that the 'lax' parts of the hammock consist only of inequalities between the elements of $\varphi(s^n) = 2^n$ for some s^n . But these are precisely the diagonal inequalities $(0, \ldots, 0) \leq (1, \ldots, 1)$. Together with the fact that the only morphisms of the form $s^n \longrightarrow s$ in \mathscr{E}^{op} are the product projections, it is easy to see that there is no way how any hammock could evaluate its right-hand side to (1, 1) and its left-hand side to (0, 1).

Remark 4.4. Siftedness of the composite $\mathbf{ob} \cdot \varphi_o : \mathscr{E}_o^{op} \longrightarrow \mathsf{Set}$ establishes precisely the 1-dimensional aspect of siftedness: the functor can is bijective on objects iff $\mathbf{ob} \cdot \varphi_o$ is sifted. From this it immediately follows that a weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ with \mathscr{E} locally discrete (i.e., with only the identity 2-cells) and such that every φe is a discrete category is sifted iff the composite $\mathbf{ob} \cdot \varphi_o : \mathscr{E}_o^{op} \longrightarrow \mathsf{Set}$ is sifted in the ordinary sense.

The 2-dimensional aspect of siftedness of $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ has to be verified in general. Example 4.3 exhibits such a situation when \mathscr{E} is locally discrete and Example 4.6 shows a conical weight $\mathsf{const}_1 : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ that is not sifted although the underlying ordinary category \mathscr{E}_o is sifted in the ordinary sense.

Example 4.5 (Siftedness for weights based on the simplicial category). Recall from, e.g., [19], that the simplicial category Δ has finite ordinals as objects and monotone maps as morphisms. It can be proved rather easily that the morphisms of Δ can be obtained from $id_1 : 1 \longrightarrow 1$, $\eta : 0 \longrightarrow 1$ and $\mu : 2 \longrightarrow 1$ by ordinal sums subject to monad axioms. Hence we will draw the morphisms of Δ as string diagrams that are generated from the following strings



that represent $id_1 : 1 \longrightarrow 1$, $\eta : 0 \longrightarrow 1$ and $\mu : 2 \longrightarrow 1$, respectively, by vertical concatenation that is subject to the unit axioms



We show that both the conical weight on Δ and the weight given by inclusion of Δ into Cat are sifted weights. In fact, from our reasoning it will be clear that the same holds of almost any truncation Δ_n . The truncated category Δ_n is just the full subcategory of Δ spanned by finite ordinals up to n.

- It is known that const₁ : Δ → Set is an ordinary sifted weight, and therefore even the conical weight const₁ : Δ → Cat is sifted due to the fact that there are no non-trivial 2-cells in Δ, see Remark 4.4. Every truncation Δ_n (for n ≥ 1) of the simplicial category Δ gives rise to a conical sifted weight as well.
- (2) Suppose the weight $\varphi : \Delta \longrightarrow \mathsf{Cat}$ is given by inclusion. Here

and the associativity axiom

$$\Delta_{a} \xrightarrow{\varphi_{o}} \mathsf{Cat}_{a} \xrightarrow{\mathsf{ob}} \mathsf{Set}$$

is a representable weight $\Delta_o(1, -)$. For each object n of Δ the category $\varphi(n)$ is the free linearly ordered category on an n-element chain. We will show an elementary proof that φ is a sifted weight. First of all, let us check that the coend $\int^n \varphi n$ is isomorphic to the one-morphism category 1. Of course, the category $\int^n \varphi n$ has precisely one object: given any two objects $x \in \varphi(n)$ and $y \in \varphi(m)$, they are equivalent by \sim if there exists a string diagram $\sigma : \varphi(n) \longrightarrow \varphi(m)$ such that x gets mapped to y by σ . A diagram like this always exists; we illustrate this on an example situation with n = 4and m = 3:



Now given any morphism $f: x \longrightarrow x'$ in $\varphi(n)$, we show that $f \approx id_*$, where id_* is the identity morphism on the only object * of $\varphi(1)$. This is again immediate when using the string diagrams:

consider the only string diagram $!: \varphi(n) \longrightarrow \varphi(1)$. It maps all morphisms in $\varphi(n)$ to the identity morphism, see for example the diagram below.



So the category $\int_{-\infty}^{n} \varphi n$ indeed has only one morphism. Now we show the isomorphism

$$\int^{n} \varphi n \times \Delta(n, n_1) \times \Delta(n, n_2) \cong \varphi n_1 \times \varphi n_2$$

by showing that the canonical morphism is bijective on objects and fully faithful. On objects, the canonical morphism takes an object $x \in \varphi(n)$, two string diagrams $\sigma : \varphi(n) \longrightarrow \varphi(n_1)$ and $\tau : \varphi(n) \longrightarrow \varphi(n_2)$, and computes the pair $(\sigma(x), \tau(x))$. It is immediate that for any pair (y, z) in $\varphi n_1 \times \varphi n_2$ there exists a tuple (x, σ, τ) that is mapped to (y, z). More is true: we can always choose $x = * \in \varphi(1)$ and the string diagrams σ, τ are the obvious diagrams choosing y and z, respectively.



This proves that can is bijective on objects. In order to prove that can is full, we will show that given any pair of morphisms $g: y \longrightarrow y'$ and $h: z \longrightarrow z'$ in $\varphi(m)$ and $\varphi(p)$ respectively, there is a morphism $f: x \longrightarrow x'$ in $\varphi(n)$ and two string diagrams sending the morphism f to g and h, respectively. But there is again a canonical such $f: x \longrightarrow x'$ in $\varphi(2)$ with the obvious inclusions, as is shown in the example diagram below.



Thus we have proved fullness and faithfulness of the canonical functor can. The weight φ is sifted. We have actually proved that any truncation $\varphi_n : \Delta_n \longrightarrow \mathsf{Cat}$ of the inclusion weight is also sifted for $n \ge 2$.

The 2-dimensional aspect of siftedness is crucial for Cat-enriched weights even in the case of conical weights, as we show in the following easy example.

Example 4.6 (A conical weight that is not sifted). Consider the diagram scheme for reflexive coequalisers satisfying $\delta_0 \cdot \sigma = \delta_1 \cdot \sigma = id_1$, and adjoin freely a 2-cell α to it:



The resulting 2-category \mathscr{E} , when considered as a conical weight, is *not* sifted, although the underlying ordinary category \mathscr{E}_o is sifted in the ordinary sense (see, e.g., Chapter 3 of [4]).

Example 4.7 (Siftedness for the weight for Kleisli objects). The weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ such that φ -colimits yield Kleisli objects is described in [18]. We will recall the definition of the weight φ and prove that it is sifted. That the weight φ is sifted is known from Proposition 8.43 in [6]: in this example we show an elementary proof of this fact.

The 2-category \mathscr{E} is the suspension $\Sigma\Delta$ of the simplicial category Δ . This means that \mathscr{E} has a unique object, say e_0 , and that the hom-category $\mathscr{E}(e_0, e_0)$ is the category Δ . Morphisms in \mathscr{E} are finite ordinals, and the 2-cells are 'monad-like' string diagrams as described in Example 4.5.

The category $\varphi(e_0)$ is defined as follows: the objects are finite *non-zero* ordinals, that is, objects of the form 1 + n for some $n < \omega$. Every object 1 + n is understood as a (n + 1)-element chain with a distinguished bottom element. The morphisms in $\varphi(e_0)$ are precisely the monotone maps that preserve the distinguished

bottom element. This definition of $\varphi(e_0)$ again allows a pictorial description in terms of string diagrams. The morphisms in $\varphi(e_0)$ are string diagrams generated by the basic diagrams



that express the fact that the diagram $\overline{}$ is an algebra for the monad given by the unit and multiplication $\overline{}$.

The 2-functor $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$ is defined on the morphisms and 2-cells of \mathscr{E}^{op} by concatenation: for a given morphism $n : e_0 \longrightarrow e_0$, the functor $\varphi(n) : \varphi(e_0) \longrightarrow \varphi(e_0)$ maps an object $1 + m \in \varphi(e_0)$ to the object 1 + m + n. A string diagram s in $\varphi(e_0)$ is mapped to the diagram $\varphi(n)(s)$, defined as the diagram s concatenated n identity strings. We show an example of this assignment for n = 1:



Likewise, given a 2-cell $\theta : m \longrightarrow n$ in \mathscr{E} , the natural transformation $\varphi(\theta)$ is defined componentwise: for an object 1 + m in $\varphi(e_0)$, the morphism $\varphi(\theta)_{1+m}$ is the concatenation of the identity diagram on 1 + m with the diagram θ . For example, given the diagram \blacktriangleleft as θ and m = 2, the component $\varphi(\theta)_3$ is the following string diagram in $\varphi(e_0)$:



Now to prove that φ is a sifted weight, we need to verify that there are canonical isomorphisms

$$\int^{e} \varphi e \cong \mathbb{1}, \qquad \int^{e} \varphi e \times \mathscr{E}(e_0, e) \times \mathscr{E}(e_0, e) \cong \varphi e_0 \times \varphi e_0 \tag{4.4}$$

proving that $\varphi \star (-)$ preserves nullary and binary products. We first analyse parts of a general hammock (4.3) for the weight φ with the testing weight $\psi = \prod_{i \in I} \mathscr{E}(-, e_i)$. The left-hand side rectangle on the diagram below



represents the information that for each $i \in I$ and the morphisms given in the diagram we have that equalities $f + s_i = t_i$ and x = y + f hold in natural numbers. This situation is depicted on the right-hand side of the above diagram. In general, the tuples (x, s_i) and (y, t_i) are related by the equivalence relation \sim if and only if $x + s_i = y + t_i$ holds for all $i \in I$.

The rectangle of the form



is represented by the concatenation of two string diagrams u and v_i for each $i \in I$.



The above diagram is an example of string diagrams that are equivalent: the 'sliding' of the division between the string diagrams generates the equivalence relation \approx . Observe moreover that morphisms in the coend are *n*-tuples of composable string diagrams. Any such *n*-tuple is equivalent to a 1-tuple, but the fact that we are allowed to vertically 'decompose' any string diagram to *n* parts is important in the proof of siftedness for φ . In the following diagram



we can see such a decomposition of a string diagram into a 2-tuple of shorter string diagrams.

With the complete description of the weight φ and of the hammocks, we can conclude that we have the canonical isomorphisms in (4.4):

(1) The weight φ satisfies the isomorphism

$$\int^e \varphi(e) \cong \mathbb{1}.$$

Indeed, the coend $\int^{e} \varphi(e)$ has precisely one object: any pair 1 + n and 1 + m of objects in $\varphi(e_0)$ is related by a hammock of length 2:

To show that $\int^e \varphi(e)$ has a unique morphism, we will prove that any string diagram $\sigma: 1+m \longrightarrow 1+n$ is congruent by the equivalence relation \approx to an identity string diagram $\mathrm{id}_k: k \longrightarrow k$ for some natural number k. We have to distinguish two cases. If the diagram σ does not contain as a subdiagram, then it is trivially a concatenation of two string diagrams $\sigma_0 = \mathrm{id}_1: 1 \longrightarrow 1$ and $\sigma_1: m \longrightarrow n$, and therefore $\sigma \approx \mathrm{id}_1$ holds. If σ contains σ , then it is necessary to factor it into a composition of two diagrams (and denote the red part of the diagram by ω):



This decomposition is unique. Take the identity morphism id_n and decompose it in the same way into a concatenation of ω with (τ_1, id_n) . By the first case we have that $\tau_1 \approx \sigma_1$, and the equivalence $\mathrm{id}_n \approx \mathrm{id}_n$ is trivial. This decomposition thus witnesses the equivalence $\sigma \approx \mathrm{id}_n$.

(2) The second isomorphism

$$\int^{e} \varphi e \times \mathscr{E}(e_0, e) \times \mathscr{E}(e_0, e) \cong \varphi e_0 \times \varphi e_0$$

is proved similarly. Given two objects 1 + m and 1 + n from $\varphi(e_0)$, there is a triple (1, m, n) that gets mapped exactly to (1 + m, 1 + n) by the canonical functor. For any other triple (k, m', n') that is mapped to (1 + m, 1 + n) we have the equalities k + m' = 1 + m and k + n' = 1 + n. Therefore $(1, m, n) \sim (k, m', n')$ holds and the canonical functor is bijective on objects.

To prove that the canonical functor is full, we show that for any two string diagrams $\sigma : 1 + m \longrightarrow 1 + n$ and $\tau : 1 + p \longrightarrow 1 + q$ there is a triple (ω, α, β) getting mapped to (σ, τ) . But again, as in the case of the first isomorphism, take ω to be the diagram



and factor the diagrams σ and τ into pairs $\alpha = (\sigma_1, \mathrm{id}_n)$ and $\beta = (\tau_1, \mathrm{id}_q)$ in a way that $\omega * \alpha = \sigma$ and $\omega * \beta = \tau$, where * denotes the horizontal composition. Faithfulness of the canonical functor then comes easily from the fact that the morphisms in the coend have the above mentioned 'normal form'.

Siftedness for enrichment in preorders. The enrichment in the category Pre of preorders and monotone maps is in many aspects similar to the enrichment in Cat, but the computations are much simpler. In fact, we will be able to give a full characterisation of sifted *conical* weights $const_1 : \mathscr{E}^{op} \longrightarrow Pre$, see Example 4.8.

The crucial coend

$$\int^e \varphi e \times [\mathscr{E}^{op}, \mathsf{Pre}](\psi, Ye)$$

is computed as a coequaliser in Pre of two monotone maps L and R that are defined in the same way as for $\mathscr{V} = \mathsf{Cat}$, see (4.2). Moreover, the coequaliser of L and R can be computed in two steps. First we compute the coequaliser on the level of underlying sets. This yields a set of equivalence classes of the form $[(\widehat{x}, \tau)]_{\sim}$ w.r.t. the equivalence \sim generated by L and R. The set of equivalence classes is then equipped with a least preorder \sqsubseteq satisfying the following condition:

If
$$(\widehat{x}, \tau) \leq (\widehat{y}, \sigma)$$
, then $[(\widehat{x}, \tau)]_{\sim} \sqsubseteq [(\widehat{y}, \sigma)]_{\sim}$

where \leq denotes the preorder of the coproduct $\coprod_e \varphi e \times [\mathscr{E}^{op}, \mathsf{Pre}](\psi, Ye)$.

Below, we will also use harmocks for the enrichment in Pre. These are pictures like (4.3) but the 2-cells u_i , v_i are replaced by mere inequality signs.

We show now that for *conical* weights $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Pre}$ the 2-dimensional aspect of siftedness is vacuous. That this is not true for *general* weights $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Pre}$ is demonstrated by the weight of Example 4.3: all categories there are in fact enriched in Pre .

Example 4.8 (Sifted conical weights). The reasoning is similar to Example 3.4 above. Elements of $\Psi_1(\mathscr{E})$ are finite coproducts $\coprod_{i \in I} Ye_i$ of representables in $[\mathscr{E}^{op}, \operatorname{Pre}]$. By Yoneda Lemma, every $\tau : \psi \longrightarrow Ye$ can be identified with a cocone $t_i : e_i \longrightarrow e$. Then the requirement that for any two natural transformations $\tau : \psi \longrightarrow Ye$ and $\sigma : \psi \longrightarrow Ye$ the equivalence $\tau \sim \sigma$ has to hold, corresponds to the fact that the cocones $t_i : e_i \longrightarrow e$ (corresponding to τ and σ respectively) have to be connected by a zig-zag. The 2-dimensional aspect of siftedness is vacuous in this case.

Thus a weight $const_1 : \mathscr{E}^{op} \longrightarrow Pre$ is sifted if and only if the *ordinary* functor

$$\mathscr{E}_{o}^{oop} \xrightarrow{(\mathsf{const}_1)_o} \mathsf{Pre}_o \xrightarrow{\mathsf{ob}} \mathsf{Set}$$

is sifted in the *ordinary* sense.

Example 4.9 (Sifted weights in general). Consider a general weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Pre.}$ To establish the isomorphism

$$\mathsf{can}: \int^e \varphi e \times \prod_{i \in I} \mathscr{E}(e_i, e) \longrightarrow \prod_{i \in I} \varphi e_i,$$

of preorders we need the monotone map can to be bijective and order-reflecting. As we noticed earlier, the coend is computed as a coequaliser in Set equipped with a freely generated preorder. More precisely, there are two conditions for a weight to be sifted:

(1) To obtain bijectivity of the can mapping we demand that

$$\mathscr{E}_{o}^{op} \xrightarrow{\varphi_{o}} \operatorname{Pre}_{o} \xrightarrow{\operatorname{ob}} \operatorname{Set}$$

be an ordinary sifted weight.

(2) Order-reflectivity of can means that given any two tuples $(x_i) \leq (x'_i)$ from $\prod_{i \in I} \varphi e_i$ we can form a hammock



such that its left-hand vertical side evaluates to (x_i) , and its right-hand vertical side evaluates to (x'_i) .

Remark 4.10. Observe that the characterisations of sifted weights for enrichments in Cat and Pre are strongly related. This is because the computations of coequalisers are essentially the same.

In fact, the requirements for a weight $\varphi : \mathscr{E}^{op} \longrightarrow \mathsf{Pre}$ to be sifted (as enriched in Pre) are exactly the requirements of siftedness for the weight $\varphi' : \mathscr{E}^{op} \longrightarrow \mathsf{Cat}$, with φ' being the weight φ considered as enriched in Cat .

The situation is rather different when considering sifted weights for the enrichment in the category Pos of all posets and monotone maps. The computation of a coequaliser in Pos runs in two steps: one computes the coequaliser in preorders *and then* performs the poset-reflection. It is the second step that brings in additional identifications and makes the characterisation of siftedness quite complex.

5. Conclusions and future work

We gave a generalisation of the concept of a sound class of small categories [1] to a sound class Ψ of weights in the context of enriched category theory. When passing to classes of weights, we showed that an easy analysis of flatness w.r.t. a sound class of weights is possible, providing us, for example, with known characterisations of siftedness and filteredness of ordinary categories. The same result yields elementary proofs of siftedness of various weights for the enrichment in categories or preorders.

A further characterisation of soundness is essentially contained in [7]. Namely, a class Ψ of weights is sound iff the KZ-monad $\mathscr{K} \mapsto \mathscr{P}(\mathscr{K})$ of free cocompletions under all colimits lifts to the 2-category Ψ -Cont of all Ψ -complete categories, all Ψ -continuous functors and all natural transformations. We thank John Bourke for pointing this out to us.

The theory of lex colimits of [9] works precisely due to the lifting of $\mathscr{K} \mapsto \mathscr{P}(\mathscr{K})$ to Ψ -Cont for Ψ being the class of weights for finite limits. Hence a theory of 'colimits in the Ψ -world' can be developed for any sound class Ψ . Since lex colimits are used to understand exactness of enriched categories, one can expect a theory of 'exactness in the Ψ -world' for any sound class Ψ . This is the matter of future research.

References

- J. Adámek, F. Borceux, S. Lack and J. Rosický, A classification of accessible categories, J. Pure Appl. Algebra 175 (2002), 7–30. 1, 2, 3, 4, 5, 6, 14
- [2] J. Adámek and J. Rosický, Locally presentable and accessible categories, Cambridge University Press 1994. 1, 5
- [3] J. Adámek and J. Rosický, On sifted colimits and generalized varieties, Theory Appl. Categ. 8 (2001), 33–53.
- [4] J. Adámek, J. Rosický and E. Vitale, Algebraic theories, Cambridge Tracts in Mathematics 184, 2011. 1, 5, 8, 10
- [5] M. H. Albert and G. M. Kelly, The closure of a class of colimits, J. Pure Appl. Algebra 51 (1988), 1–17. 1, 2
- [6] J. Bourke, Codescent objects in 2-dimensional universal algebra, PhD Thesis, University of Sydney, 2010. 2, 10
- [7] B. J. Day and S. Lack, Limits of small functors, J. Pure Appl. Algebra 210 (2007), 651-663. 1, 14
- [8] P. Gabriel and F. Ulmer, Lokal präsentierbare Kategorien, Lecture Notes in Mathematics 221, Springer 1971. 1
- [9] R. Garner and S. Lack, Lex colimits, J. Pure Appl. Algebra 216 (2012), 1372–1396. 14
- [10] G. M. Kelly, Basic concepts of enriched category theory, London Math. Soc. Lecture Notes Series 64, Cambridge Univ. Press, 1982, also available as Repr. Theory Appl. Categ. 10 (2005). 2
- [11] G. M. Kelly and S. Lack, Finite product-preserving-functors, Kan extensions and strongly-finitary 2-monads, Appl. Categ. Structures 1 (1993), 85–94. 3, 8
- [12] G. M. Kelly and V. Schmitt, Notes on enriched categories with colimits of some class, Theory Appl. Categ. 14.17 (2005), 399–423. 1, 2, 3, 5, 6, 7
- [13] S. R. Johnson, Small Cauchy completions, J. Pure Appl. Algebra 62 (1989), 35-45. 6
- [14] S. Lack and J. Rosický, Notions of Lawvere theory, Appl. Categ. Structures 19.1 (2011), 363–391. 1
- [15] C. Lair, Catégories modelables et catégories esquissables, *Diagrammes* 6 (1981), 1–20. 1
- [16] C. Lair, Sur le genre d'esquissabilité des catégories modelables (accessibles) possédant les produits de deux, Diagrammes 35 (1996), 25–52. 1, 4

- [17] F. W. Lawvere, Functorial semantics of algebraic theories, PhD Thesis, Columbia University 1963, available as Repr. Theory Appl. Categ. 5 (2004), 1–121.
- [18] F. W. Lawvere, Ordinal sums and equational doctrines, in Seminar on triples and categorical homology theory (B. Eckmann and M. Tierney, eds.), LNM 80, Springer 1969, available as Repr. Theory Appl. Categ. 18 (2008), 1–303. 10
- [19] S. Mac Lane, Categories for the working mathematician, 2nd ed., Springer 1998. 9
- [20] M. Makkai and R. Paré, Accessible categories, American Mathematical Society, Providence 1989. 1
- [21] J. Velebil and J. Adámek, A remark on conservative cocompletion of categories, J. Pure Appl. Algebra 168 (2002), 107–124. 6

DEPARTMENT OF MATHEMATICS, FACULTY OF ELECTRICAL ENGINEERING, CZECH TECHNICAL UNIVERSITY IN PRAGUE, CZECH REPUBLIC

E-mail address: dostamat@math.feld.cvut.cz

DEPARTMENT OF MATHEMATICS, FACULTY OF ELECTRICAL ENGINEERING, CZECH TECHNICAL UNIVERSITY IN PRAGUE, CZECH REPUBLIC

E-mail address: velebil@math.feld.cvut.cz