

EQUIISINGULARITY OF FAMILIES OF ISOLATED DETERMINANTAL SINGULARITIES

J. J. NUÑO-BALLESTEROS, B. ORÉFICE-OKAMOTO, J. N. TOMAZELLA

ABSTRACT. We study the topological triviality and the Whitney equisingularity of a family of isolated determinantal singularities. On one hand, we give a Lê-Ramanujam type theorem for this kind of singularities by using the vanishing Euler characteristic. On the other hand, we extend the results of Teissier and Gaffney about the Whitney equisingularity of hypersurfaces and complete intersections, respectively, in terms of the constancy of the polar multiplicities.

1. INTRODUCTION

The celebrated Lê-Ramanujan theorem [10] states that any 1-parameter family of isolated hypersurface singularities $\{(X_t, 0)\}$ of dimension $d \neq 2$ has constant topological type if the Milnor number $\mu(X_t, 0)$ is constant on t . The same result is also known to be true for isolated complete intersection singularities (ICIS) of dimension $d \neq 2$ (see [15]). The condition $d \neq 2$ is needed because of the use of the h -cobordism theorem in the proof. It is worth to mention that the case $d = 2$ remains open until now. For space curves (not necessarily ICIS), the topological triviality of the family was obtained by Buchweitz-Greuel, by using an appropriate version of Milnor number (see [1]).

For plane curves, Zariski proved [19] that the μ -constant condition is also equivalent to the fact that the family is Whitney equisingular. However, the Briançon-Speder example [2] shows that this is not true anymore for hypersurfaces of dimension $d \geq 2$. Teissier solved completely in [18] the Whitney equisingularity question for hypersurfaces. He showed that the family is Whitney equisingular if and only if the μ^* sequence is constant on the family. This result was generalized by Gaffney [5, 6] for ICIS of any dimension d , with the use of the polar multiplicities $m_i(X_0, 0)$, $i = 0, \dots, d$. Note that the polar multiplicities $m_i(X_0, 0)$, for $i = 0, \dots, d - 1$ are defined for any variety $(X_0, 0)$ (see [11]), but $m_d(X_0, 0)$ is defined only for ICIS in [5]. For space curves, a family is Whitney equisingular if and only if $m_1(X_t, 0)$ is constant on t , where $m_1(X_t, 0)$ is defined as the Milnor number of a generic linear projection (see [13]).

Here we study the topological triviality and the Whitney equisingularity, for families of isolated determinantal singularities (IDS). Determinantal varieties are those defined by the zeros of an ideal generated by the minors of a given size of a matrix. We assume that the variety has the expected dimension according to the sizes of the matrix and of the minors. Moreover, we restrict ourselves to the case that the determinantal variety has isolated singularity (in some sense related to the determinantal structure). Any ICIS (and hence any hypersurface) is a particular case of an IDS.

2000 *Mathematics Subject Classification*. Primary 32S15; Secondary 58K60, 32S30.

Key words and phrases. Determinantal singularity, topological triviality, Whitney equisingularity.

The first author was partially supported by DGICYT Grant MTM2012-33073 and FAPESP Grant 2012/18805-2. The second author was partially supported by FAPESP Grant 2013/14014-3. The third author was partially supported by FAPESP Grant 2013/10856-0.

On one hand, we show that a family $\{(X_t, 0)\}$ of IDS of dimension $d \neq 2$ has constant topological type provided that:

- (1) the family is good (i.e., it has isolated singularity uniformly in the family),
- (2) the determinantal smoothing of any member $(X_t, 0)$ has the homotopy type of a bouquet of d -spheres,
- (3) the vanishing Euler characteristic $\nu(X_t, 0)$ is constant on t .

The number $\nu(X_0, 0)$ was introduced in a previous paper [14] for an IDS $(X_0, 0)$ and it is defined as

$$\nu(X_0, 0) = (-1)^d(\chi(X_A) - 1),$$

where X_A is the determinantal smoothing of $(X_0, 0)$ and $\chi(\cdot)$ denotes the Euler characteristic. In the particular case of an ICIS, the determinantal smoothing coincides with the Milnor fibre and hence, $\nu(X_0, 0) = \mu(X_0, 0)$. Moreover, for ICIS the condition (2) is always satisfied and the condition (1) follows from (3). Thus, our result generalizes the ICIS version of the Lê-Ramanujam theorem.

On the other hand, we also show that a good family $\{(X_t, 0)\}$ of IDS of any dimension d is Whitney equisingular if and only if all the polar multiplicities $m_i(X_t, 0)$, $i = 0, \dots, d$ are constant on the family. Again, we find in [14] an adapted version of the top polar multiplicity $m_d(X_0, 0)$ of an IDS $(X_0, 0)$, which generalizes that of [5] for ICIS. Hence, our theorem generalizes the results of Teissier or Gaffney in the particular cases of hypersurfaces or ICIS respectively.

2. ISOLATED DETERMINANTAL SINGULARITIES

We recall here from [14] the main definitions and properties of IDS. Let $0 < s \leq m \leq n$ be integer numbers, $M_{m,n}$ be the set of complex matrices of size $m \times n$ and let $M_{m,n}^s$ be the subset of matrices with rank less than s .

We consider a holomorphic map germ

$$f : (\mathbb{C}^N, 0) \rightarrow M_{m,n}$$

and we say that the germ

$$(X_0, 0) = f^{-1}(M_{m,n}^s)$$

is a determinantal germ of type $(m, n; s)$ if the dimension of $(X_0, 0)$ is equal to

$$d = N - (m - s + 1)(n - s + 1).$$

In [14], we introduced a new concept of isolated determinantal singularity. We defined the germ $(X_0, 0)$ to be an isolated determinantal singularity, shortening IDS, if $s = 1$ or $N < (m - s + 2)(n - s + 2)$, X_0 is smooth at x and $\text{rank}(f(x)) = s - 1$, for all $x \neq 0$ in a neighborhood of the origin.

We fix a small enough representative $f : B_\epsilon \rightarrow M_{m,n}$, where B_ϵ is the open ball centered at the origin and of radius $\epsilon > 0$. Given a matrix A in $M_{m,n}$, let $f_A : B_\epsilon \rightarrow M_{m,n}$ be the map defined by

$$f_A(x) = f(x) + A$$

and let

$$X_A = f_A^{-1}(M_{m,n}^s).$$

We see in [14] that there exists a non-empty Zariski open set W in $M_{m,n}$ such that X_A is smooth if A belongs to W and the Euler characteristic of X_A , $\chi(X_A)$ does not depend on

A in W . The vanishing Euler characteristic of the IDS $(X_0, 0)$ was, then, defined by

$$\nu(X_0, 0) = (-1)^d(\chi(X_A) - 1),$$

with A in the Zariski open set W . We also call X_A the special determinantal smoothing of $(X_0, 0)$. In the case of an ICIS ($s = 1$), X_A can be seen as the Milnor fibre of $(X_0, 0)$ and hence, $\nu(X_0, 0)$ coincides with the Milnor number $\mu(X_0, 0)$.

We recall now the notion of determinantal deformation. We consider a map germ $F : (\mathbb{C}^N \times \mathbb{C}, 0) \rightarrow M_{m,n}$ such that $F(x, 0) = f(x)$ for all $x \in \mathbb{C}^N$. Then we denote $\mathcal{X} = F^{-1}(M_{m,n}^s)$ and we say that the projection

$$\begin{aligned} \pi : (\mathcal{X}, 0) &\rightarrow (\mathbb{C}, 0) \\ (x, t) &\mapsto t \end{aligned}$$

is a determinantal deformation of $(X_0, 0)$. If we fix again a representative $F : B_\epsilon \times D \rightarrow M_{m,n}$, then we set $f_t(x) = F(x, t)$ and $X_t = f_t^{-1}(M_{m,n}^s)$. With this notation, it is common to write simply X_t for the deformation, instead of π .

A determinantal smoothing of $(X_0, 0)$ is a determinantal deformation X_t such that X_t is smooth and $\text{rank}(f_t(x)) = s - 1$ for all $x \in X_t$ and for all $t \in D - \{0\}$. In this case, we showed in [14]:

$$\nu(X_0, 0) = (-1)^d(\chi(X_t) - 1).$$

If X_t is a determinantal deformation of $(X_0, 0)$ as above, we say that:

- (1) X_t is origin preserving if $0 \in S(X_t)$, for all t in D , where $S(X_t)$ denotes the singular set of X_t . Then we say that $\{(X_t, 0)\}_{t \in D}$ is a 1-parameter family of IDS;
- (2) X_t is good if there exists $\epsilon > 0$ such that $S(X_t) = \{0\}$ on B_ϵ , for all t in D ;
- (3) X_t has constant topological type if for each $t \in D$ there exists $\epsilon_t > 0$ such that (B_{ϵ_t}, X_t) is homeomorphic to (B_ϵ, X_0) .
- (4) X_t is Whitney equisingular if it is good and $\{\mathcal{X} - T, T\}$ satisfies the Whitney conditions, where $T = \{0\} \times D$.

Finally, we also recall the definition of polar multiplicities. They were defined by Lê-Teissier [11] for any d -dimensional variety $(X_0, 0) \subset (\mathbb{C}^N, 0)$ and for each $i = 0, \dots, d - 1$. We choose a generic linear projection $p : \mathbb{C}^N \rightarrow \mathbb{C}^{d-i+1}$, then the i -th polar multiplicity is equal to

$$m_i(X_0, 0) = m_0(\overline{S(p|_{X_0^0})}, 0),$$

where X_0^0 is the smooth part of X_0 and $m_0(Z, 0)$ denotes the usual multiplicity of any variety $(Z, 0)$.

In the case that $(X_0, 0)$ is an IDS, we introduced in [14] the d -th polar multiplicity as

$$m_d(X_0, 0) = \#S(p|_{X_A}),$$

where now X_A is a determinantal smoothing and $p : \mathbb{C}^N \rightarrow \mathbb{C}$ is a generic linear function.

3. THE TOPOLOGY OF THE DETERMINANTAL SMOOTHING

In this section, we describe the topology of the determinantal smoothing X_A of an IDS $(X_0, 0)$ of dimension d . We know that X_A has the homotopy type of a CW-complex of real dimension d . We start by showing that the polar multiplicities of $(X_0, 0)$ are related to the cell structure of the smoothing X_A .

Lemma 3.1. *Let $M = X_A \cap B_{\epsilon'}$ for some $0 < \epsilon' < \epsilon$. Let $g : X_A \rightarrow \mathbb{C}$ be a holomorphic Morse function with no critical points on $X_A \cap S_{\epsilon'}$. Then,*

$$M \cong (g^{-1}(c) \cap M) \text{ with } m \text{ } d\text{-cells attached,}$$

where $m = \#S(g|_M)$ and c is a regular value of g .

Proof. See the proof of Theorem A.5 of [14]. \square

Theorem 3.2. *Let $(X_0, 0)$ be a d -dimensional IDS and let X_A be a determinantal smoothing. Then X_A has the homotopy type of CW-complex with $m_i(X_0, 0)$ cells of dimension i , for $i = 0, \dots, d$.*

Proof. Let's proceed by finite induction over d . If $d = 0$, X_A is a finite set with $\chi(X_A)$ points. By [14, 3.5],

$$\chi(X_A) = \nu(X_0, 0) + 1 = \dim_{\mathbb{C}} \mathcal{O}_{X_0, 0} = m_0(X_0, 0).$$

Assume that the result is true for any IDS with dimension $d - 1$. We choose a generic linear map $p : \mathbb{C}^N \rightarrow \mathbb{C}$. Then by Lemma 3.1,

$$X_A \cong (p^{-1}(c) \cap X_A) \text{ with } \#S(p|_{X_A}) \text{ } d\text{-cells attached.}$$

We have that $p^{-1}(c) \cap X_A$ is the smoothing of the $(d-1)$ -dimensional IDS $(X_0 \cap p^{-1}(0), 0)$. By induction hypothesis, $p^{-1}(c) \cap X_A$ has a CW-complex structure with $m_i(X_0 \cap p^{-1}(0), 0)$ cells of dimension i , for $i = 0, \dots, d-1$. But it follows from [11] that $m_i(X_0, 0) = m_i(X_0 \cap p^{-1}(0), 0)$ for $i = 0, \dots, d-1$, and $m_d(X_0, 0) = \#S(p|_{X_A})$, by definition of the d -th polar multiplicity. \square

Corollary 3.3. *If $(X_0, 0)$ is a d -dimensional IDS, then*

$$\nu(X_0, 0) = (-1)^d \left(\sum_{i=0}^d (-1)^i m_i(X_0, 0) - 1 \right).$$

Proof. By definition $\nu(X_0, 0) = (-1)^d (\chi(X_A) - 1)$, then

$$\begin{aligned} \nu(X_0, 0) &= (-1)^d (\chi(X_A) - 1) \\ &= (-1)^d \left(\sum_{i=0}^d (-1)^i m_i(X_0, 0) - 1 \right). \end{aligned}$$

\square

Example 3.4. Let $(X_0, 0)$ be the determinantal surface in $(\mathbb{C}^4, 0)$ given by the 2×2 minors of the matrix:

$$\begin{pmatrix} x & y & z \\ y & z & w \end{pmatrix}$$

We showed in [14] that $m_0(X_0, 0) = 3$, $m_1(X_0, 0) = 4$ and $m_2(X_0, 0) = 3$. Thus, any determinantal smoothing X_A has the homotopy type of a 2-dimensional CW-complex with 3 0-cells, 4 1-cells and 3 2-cells.

Another interesting topological property is that any determinantal smoothing is always connected (unless it is 0-dimensional).

Lemma 3.5. *If $(X_0, 0)$ is an IDS of dimension $d \geq 1$, then X_A is connected.*

Proof. From the long exact homotopy sequence of the pair $(\overline{X_A}, \partial\overline{X_A})$, the canonical mapping

$$\pi_i(\partial\overline{X_A}) \rightarrow \pi_i(\overline{X_A})$$

is surjective if $i = d - 1$ and is an isomorphism if $i \leq d - 2$ (see [4], proof of Theorem 1). Moreover, we have $\pi_i(\partial\overline{X_0}) = \pi_i(\partial\overline{X_A})$ and $\pi_i(\overline{X_A}) = \pi_i(X_A)$.

If $d \geq 2$, $(X_0, 0)$ is normal, therefore irreducible and $\partial\overline{X_0}$ is connected. Hence, $\pi_0(X_A) \cong \pi_0(\partial\overline{X_0})$ and X_A is connected. If $d = 1$, X_A is connected by Theorem 4.2.2 of [1]. \square

It was shown by Milnor [12], that the Milnor fibre of a d -dimensional isolated hypersurface singularity has the homotopy type of a bouquet of spheres of dimension d . This was generalized later by Hamm [7] for ICIS. For the purpose of the next section, it would be interesting to know whether this is also true for general IDS.

When $d = 1$, this is always true, since any connected 1-dimensional CW-complex has the homotopy type of a bouquet of 1-spheres. For $d = 2$, we know from [4] that $\beta_1(X_A) = 0$, but this is not enough to ensure that X_A has the homotopy type of a bouquet of 2-spheres, since we should need that X_A is simply connected. We do not know at this moment if this is true or not for a 2-dimensional IDS. For $d \geq 3$, the result is definitely false in general: Damon and Pike [3] give examples of 3-dimensional IDS for which $\beta_2(X_A) \neq 0$.

In general, we will use the following well known criterion for a d -dimensional CW-complex to be a bouquet of spheres of dimension d . The proof follows easily by using Hurewicz and Whitehead theorems (see also the proof of [12, 6.5]).

Lemma 3.6. *Let X be a connected d -dimensional CW-complex, with $d \geq 2$. The following statements are equivalent:*

- (1) $H_2(X) = \cdots = H_{d-1}(X) = \pi_1(X) = 1$;
- (2) X is $(d - 1)$ -connected;
- (3) X has the homotopy type of a bouquet of spheres of dimension d .

Finally, we deduce the following direct consequence of the previous lemma and the results of [4].

Corollary 3.7. *If $(X_0, 0)$ is a d -dimensional IDS in \mathbb{C}^{d+k} with $k \leq d - 1$ and such that $H_{d-k+1}(X_A) = \cdots = H_{d-1}(X_A) = 0$, then X_A has the homotopy type of a bouquet of $\nu(X_0, 0)$ spheres of dimension d .*

4. THE TOPOLOGY OF A DETERMINANTAL DEFORMATION

We consider here a determinantal deformation $(\mathcal{X}, 0)$ of the IDS $(X_0, 0)$. If the deformation is given by $(\mathcal{X}, 0) = F^{-1}(M_{m,n}^s)$, we fix a representative $F : B \times D \rightarrow M_{m,n}$, where $B \subset \mathbb{C}^N$ and $D \subset \mathbb{C}$ are small enough open balls centered at the origin. We set the following notation for each $t \in D$ and $A \in M_{m,n}$:

- $f_t : B \rightarrow M_{m,n}$ the map defined by $f_t(x) = F(x, t)$ and $X_t = f_t^{-1}(M_{m,n}^s)$;
- $f_{t,A} : B \rightarrow M_{m,n}$ the map defined by $f_{t,A}(x) = f_t(x) + A$ and $X_{t,A} = f_{t,A}^{-1}(M_{m,n}^s)$.

Lemma 4.1. *The subset $W \subset D \times M_{m,n}$ of pairs (t, A) such that $X_{t,A}$ is smooth and $\text{rank } f_{t,A}(x) = s - 1$ for any $x \in X_{t,A}$ is non-empty Zariski open. Moreover, the topological type of $X_{t,A}$ does not depend on (t, A) in W .*

Proof. Just follow the arguments of [14, Theorem 3.4]. \square

We fix some notations which will be useful for some of the results of this section. We choose the ball B_ϵ such that the only singular point of X_0 in B_ϵ is the origin. Now we fix a generic matrix $A \in M_{m,n}$ such that X_A is a determinantal smoothing of $(X_0, 0)$. Moreover, since the subset W of Lemma 4.1 is open, we can shrink the disk D if necessary, in such a way that for any $t \in D$, $(t, A) \in W$. Hence, $X_{t,A}$ is smooth, $\text{rank } f_{t,A}(x) = s - 1$ for any $x \in X_{t,A}$ and $X_{t,A}$ is homeomorphic to X_A .

Fix $t \in D$ and let $\{x_1, \dots, x_k\}$ be the singular set of $X_t \cap B_\epsilon$. For each $i = 1, \dots, k$, choose an open disc $D_i \subset B_\epsilon$ centered at x_i and such that x_i is the only critical point of $X_t \cap D_i$ (see fig. 1). It follows that $X_{t,A} \cap D_i$ is a determinantal smoothing of (X_t, x_i) . Moreover, we can choose a continuous map $r : X_{t,A} \rightarrow X_t$ such that $r(X_{t,A} \cap D_i) = \{x_i\}$ and r is a homeomorphism outside D_i .

Finally, we set:

$$\begin{aligned} U &= X_t \cap (\cup_{i=1}^k \overline{D}_i), \\ V &= X_t \cap (B_\epsilon - \cup_{i=1}^k D_i), \\ \tilde{U} &= X_{t,A} \cap (\cup_{i=1}^k \overline{D}_i), \\ \tilde{V} &= X_{t,A} \cap (B_\epsilon - \cup_{i=1}^k D_i). \end{aligned}$$




FIGURE 1.

The following theorem states that the difference between the global invariants in the fibres of a deformation measures exactly the vanishing Euler characteristic of the deformation. This fact is well known for curve singularities (see [1]), for hypersurfaces (see [16]) and probably for ICIS, although we could not find any good reference in the literature. We prove it here for general IDS.

Theorem 4.2. *Let $(X_0, 0)$ be a d -dimensional IDS and let X_t be any determinantal deformation. Then,*

$$\nu(X_0, 0) - \sum_{x \in S(X_t)} \nu(X_t, x) = (-1)^d (\chi(X_t) - 1)$$

Proof. With the above notation we have for X_t :

$$\chi(X_t) = \chi(U \cup V) = \chi(U) + \chi(V) - \chi(U \cap V) = \chi(V) + k,$$

because each $D_i \cap X_t$ is contractible and $\chi(U \cap V) = 0$ (since $U \cap V$ is a closed manifold of odd dimension).

Analogously, for $X_{t,A}$ we get:

$$\chi(X_t) = \chi(\tilde{U} \cup \tilde{V}) = \chi(\tilde{U}) + \chi(\tilde{V}) - \chi(\tilde{U} \cap \tilde{V}) = \chi(\tilde{V}) + \sum_{i=1}^k \chi(X_{t,A} \cap D_i).$$

On the other hand,

$$\begin{aligned} (-1)^d \nu(X_0, 0) + 1 &= \chi(X_A) \\ &= \chi(X_{t,A}) \\ &= \chi(\tilde{V}) + \sum_{i=1}^k \chi(X_{t,A} \cap D_i) \\ &= \chi(V) + \sum_{i=1}^k ((-1)^d \nu(X_t, x_i) + 1) \\ &= \chi(X_t) - k + \sum_{i=1}^k (-1)^d \nu(X_t, x_i) + k \\ &= \chi(X_t) + (-1)^d \sum_{i=1}^k \nu(X_t, x_i), \end{aligned}$$

which gives the desired equality. □

Corollary 4.3. *Let X_t be a determinantal deformation of an IDS $(X_0, 0)$. Then, the sum $\sum_{x \in S(X_t)} \nu(X_t, x)$ is constant on t if and only if $\chi(X_t) = 1$.*

We recall that a deformation $(\mathcal{X}, 0)$ of $(X_0, 0)$ is topologically trivial if there exists a homeomorphism $\Phi : (\mathcal{X}, 0) \rightarrow (X_0 \times \mathbb{C}, 0)$ which commutes with the projection, that is, $\pi \circ \Phi = \pi$, where $\pi(x, t) = t$. In particular, this implies that X_t is homeomorphic to X_0 , for t small enough.

Corollary 4.4. *If X_t is a topologically trivial determinantal deformation of an IDS $(X_0, 0)$, then*

$$\sum_{x \in S(X_t)} \nu(X_t, x) = \nu(X_0, 0).$$

Remark 4.5. We know from [12] that the Milnor number of a hypersurface with isolated singularity is a topological invariant, that is, if $(X_0, 0)$ and $(X'_0, 0)$ are hypersurfaces with isolated singularity and have the same topological type (i.e., (B_ϵ, X_0) is homeomorphic to $(B_{\epsilon'}, X'_0)$), then $\mu(X_0, 0) = \mu(X'_0, 0)$. However, this is not true in general for higher codimension. For instance, take $(X_0, 0) \subset (\mathbb{C}^3, 0)$ as the determinantal curve given by the 2×2 -minors of

$$\begin{pmatrix} x & y & z \\ y & z & x^2 \end{pmatrix}$$

and take $(X'_0, 0)$ as the x -axis $y = z = 0$. Both curves have the same topological type, since both are irreducible, $(X_0, 0)$ is the curve parametrized by $t \mapsto (t^3, t^4, t^5)$ and $(X'_0, 0)$ is smooth. However, $\mu(X_0, 0) = 4$ and $\mu(X'_0, 0) = 0$.

In order to deduce some other consequences of Theorem 4.2, we need some connection properties of the general fibre X_t of a determinantal deformation of an IDS $(X_0, 0)$.

Lemma 4.6. *If $(X_0, 0)$ is an IDS with dimension $d \geq 1$ then any determinantal deformation X_t is connected for t small enough.*

Proof. We consider the Mayer-Vietoris sequences of the pairs (U, V) in X_t and (\tilde{U}, \tilde{V}) in $X_{t,A}$ where the vertical arrows are induced by the continuous map $r : X_{t,A} \rightarrow X_t$:

$$\begin{array}{ccccccc} \dots & \longrightarrow & H_0(U \cap V) & \longrightarrow & H_0(U) \oplus H_0(V) & \longrightarrow & H_0(X_t) \longrightarrow 0 \\ & & \uparrow \alpha_0 & & \uparrow \beta_0 & & \uparrow \gamma_0 \\ \dots & \longrightarrow & H_0(\tilde{U} \cap \tilde{V}) & \longrightarrow & H_0(\tilde{U}) \oplus H_0(\tilde{V}) & \longrightarrow & H_0(X_{t,A}) \longrightarrow 0 \end{array}$$

Note that β_0 is an isomorphism, since r is a homeomorphism between \tilde{V} and V and it is also a bijection between the connected components of \tilde{U} and U . Then, γ_0 is an epimorphism by elementary calculations. On the other hand, since $X_A \simeq X_{t,A}$ is connected by Lemma 3.5, we have that X_t is also connected. \square

Lemma 4.7. *If $(X_0, 0)$ is an IDS with dimension $d \geq 2$, then $\beta_1(X_t) = 0$ for any determinantal deformation X_t and for t small enough.*

Proof. We consider the Mayer-Vietoris sequence of the pair (U, V) in X_t for the homology with rational coefficients:

$$\begin{array}{ccccccc} \dots & \longrightarrow & H_1(U; \mathbb{Q}) \oplus H_1(V; \mathbb{Q}) & \xrightarrow{\phi_1} & H_1(X_t; \mathbb{Q}) & \xrightarrow{\phi_2} & \\ & & \xrightarrow{\phi_2} & H_0(U \cap V; \mathbb{Q}) & \xrightarrow{\phi_3} & H_0(U; \mathbb{Q}) \oplus H_0(V; \mathbb{Q}) & \longrightarrow \dots \end{array}$$

Since $d \geq 2$, all the singularities (X_t, x_i) are normal and hence, irreducible. Thus, $U \cap V$ has exactly k connected components (one for each singular point) and the induced morphism $H_0(U \cap V; \mathbb{Q}) \rightarrow H_0(U; \mathbb{Q})$ is an isomorphism. This implies that ϕ_3 is a monomorphism, $\phi_2 = 0$ and hence, ϕ_1 is an epimorphism. In fact, since $H_1(U; \mathbb{Q}) = 0$, we can consider $\phi_1 : H_1(V; \mathbb{Q}) \rightarrow H_1(X_t; \mathbb{Q})$.

Thus, we have

$$\begin{array}{ccc} H_1(V; \mathbb{Q}) & \xrightarrow{\phi_1} & H_1(X_t; \mathbb{Q}) \\ \uparrow \beta_1 & & \uparrow \gamma_1 \\ H_1(\tilde{V}; \mathbb{Q}) & \longrightarrow & H_1(X_{t,A}; \mathbb{Q}) \end{array}$$

where ϕ_1 is epimorphism and β_1 is isomorphism. Then, γ_1 is also epimorphism. But $(X_0, 0)$ is normal and by [4], $H_1(X_{t,A}; \mathbb{Q}) = H_1(X_A; \mathbb{Q}) = 0$, therefore $H_1(X_t; \mathbb{Q}) = 0$. \square

As a consequence of the two previous lemmas and Theorem 4.6, we can conclude that the vanishing Euler characteristic of 2-dimensional IDS is upper semi-continuous.

Corollary 4.8. *If $(X_0, 0)$ is a 2-dimensional IDS, then $\nu(X_t, x) \leq \nu(X_0, 0)$ for any determinantal deformation X_t and for any $x \in S(X_t)$.*

Proof. We know that $\nu(X_t, x) = \beta_2(X_{t,A}) \geq 0$, for any $x \in S(X_t)$. Hence, by 4.2, 4.6 and 4.7:

$$\nu(X_t, x) \leq \sum_{x \in S(X_t)} \nu(X_t, x) = \nu(X_0, 0) - \chi(X_t) + 1 = \nu(X_0, 0) - \beta_2(X_t) \leq \nu(X_0, 0).$$

\square

In order to proof the main theorem of this section (Theorem 4.11), we will need to use the fact that X_t is contractible. This could be concluded from Theorem 4.6 if we knew that X_t was $(d-1)$ -connected. In the two following lemmas, we show that this is true if the determinantal smoothing of (X_t, x) is assumed to be $(d-1)$ -connected, for any singular point x .

Lemma 4.9. *Let $(X_0, 0)$ be an IDS of dimension $d \geq 2$ and let \mathcal{X} be a determinantal deformation of $(X_0, 0)$. If there is a determinantal smoothing X_A of $(X_0, 0)$ such that $\pi_1(X_A) = 1$, then $\pi_1(X_t) = 1$ for t small enough.*

Proof. In this proof we change slightly the notation for the sets $U, V, \tilde{U}, \tilde{V}$ from the beginning of this section. Since we are going to use Van Kampen theorem, we need that these sets are open.

Let $D'_i \subset D_i$ be another ball with center x_i . Let us denote

$$V = X_t - (\cup_{i=1}^n D'_i), \quad \tilde{V} = X_A - (\cup_{i=1}^n D'_i),$$

and, for each $i = 1, \dots, k$,

$$U_i = X_t \cap D_i, \quad U = U_1 \cup \dots \cup U_k.$$

We observe that U_1 is contractible (because D_1 is a Milnor ball for (X_t, x_1)) and, therefore, connected and $U_1 \cap V$ is also connected (because it has the same homotopy type of the link of (X_t, x_1) , see the proof of Lemma 3.5). We will show that V is also connected. In fact, we consider the following Mayer-Vietoris sequence in the reduced homology:

$$\dots \longrightarrow \tilde{H}_0(U \cap V) \longrightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \longrightarrow \tilde{H}_0(X_t) .$$

Since $\tilde{H}_0(X_t) = 0$ by Lemma 4.6, the map

$$\tilde{H}_0(U \cap V) \longrightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V)$$

is surjective. However, the number of connected components of $U \cap V$ is equal to the number of connected components of U , so $\tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U)$ is an isomorphism and therefore, $\tilde{H}_0(V) = 0$.

By the Van Kampen theorem, the induced map

$$\pi_1(U_1) *_{\pi_1(U_1 \cap V)} \pi_1(V) \longrightarrow \pi_1(U_1 \cup V)$$

is an isomorphism. But $\pi_1(U_1) = 1$, hence

$$\pi_1(V) \longrightarrow \pi_1(U_1 \cup V)$$

is surjective.

Now, we denote $V_2 = V \cup U_1$. We know, similarly, that U_2, V_2 e $U_2 \cap V_2$ are connected. By the Van Kampen theorem, the induced map

$$\pi_1(U_2) *_{\pi_1(U_2 \cap V_2)} \pi_1(V) \longrightarrow \pi_1(U_2 \cup V_2)$$

is an isomorphism. But again $\pi_1(U_2) = 1$, so $\pi_1(V_2) \rightarrow \pi_1(U_2 \cup V_2)$ is surjective. By taking composition with the above map, we have that

$$\pi_1(V) \longrightarrow \pi_1(U_1 \cup U_2 \cup V)$$

is also surjective.

Proceeding in this way, we conclude that there the following map is surjective:

$$\pi_1(V) \longrightarrow \pi_1(U_1 \cup \cdots \cup U_n \cup V) = \pi_1(X_t).$$

We have, then, the following commutative diagram

$$\begin{array}{ccc} \pi_1(V) & \longrightarrow & \pi_1(X_t) \\ \uparrow & & \uparrow \\ \pi_1(\tilde{V}) & \longrightarrow & \pi_1(X_A) = 1 \end{array}$$

where the top line is an epimorphism and the left column is an isomorphism. Therefore, $\pi_1(X_t) = 1$. \square

Lemma 4.10. *Let X_0 be an IDS of dimension $d \geq 1$ and let X_t be a determinantal deformation of it such that the determinantal smoothing of (X_t, x) is $(d-1)$ -connected, for all $x \in S(X_t)$ and t small enough. Then X_t is also $(d-1)$ -connected, for all t small enough.*

Proof. If $d = 1$, the result follows from Lemma 4.6. If $d \geq 2$, from Lemma 4.9, we know that $\pi_1(X_t) = 1$ for t small enough. We will show that $H_i(X_t) = 0$ for any $i = 2, \dots, k$.

We keep the notation of the beginning of this section for the sets $U, V, \tilde{U}, \tilde{V}$. We consider again the Mayer-Vietoris sequences as in the proof of Lemma 4.6:

$$\begin{array}{ccccccc} \dots & \longrightarrow & H_i(U \cap V) & \longrightarrow & H_i(U) \oplus H_i(V) & \longrightarrow & H_i(X_t) & \longrightarrow \\ & & \uparrow \alpha_i & & \uparrow \beta_i & & \uparrow \gamma_i & \\ \dots & \longrightarrow & H_i(\tilde{U} \cap \tilde{V}) & \longrightarrow & H_i(\tilde{U}) \oplus H_i(\tilde{V}) & \longrightarrow & H_i(X_{t,A}) & \longrightarrow \\ & & \uparrow \alpha_{i-1} & & \uparrow \beta_{i-1} & & \uparrow \gamma_{i-1} & \\ & \longrightarrow & H_{i-1}(U \cap V) & \longrightarrow & H_{i-1}(U) \oplus H_{i-1}(V) & \longrightarrow & H_{i-1}(X_t) & \longrightarrow \dots \\ & & \uparrow \alpha_{i-1} & & \uparrow \beta_{i-1} & & \uparrow \gamma_{i-1} & \\ & \longrightarrow & H_{i-1}(\tilde{U} \cap \tilde{V}) & \longrightarrow & H_{i-1}(\tilde{U}) \oplus H_{i-1}(\tilde{V}) & \longrightarrow & H_{i-1}(X_{t,A}) & \longrightarrow \dots \end{array}$$

We use an appropriate version of the five lemma (see [8]). We have:

- (1) β_i is an epimorphism, since r is a homeomorphism between \tilde{V} and V and $H_i(U) = 0$ (each connected component $U \cap D_j$ is contractible).
- (2) α_{i-1} is an isomorphism since r is a homeomorphism between $\tilde{U} \cap \tilde{V}$ and $U \cap V$.
- (3) β_{i-1} is a monomorphism, since $H_{i-1}(\tilde{U}) = 0$ from the hypothesis and β_{i-1} is an isomorphism between $H_{i-1}(\tilde{V})$ and $H_{i-1}(V)$.

Therefore, γ_i is an epimorphism and since $H_i(X_{t,A}) = 0$, we also have $H_i(X_t) = 0$. \square

It is known that any 1-parameter family of hypersurfaces, or more generally ICIS, with dimension different of two has constant topological type if and only if it has constant Milnor number (see [10, 15]). In the following theorem, which is the main result of this section, we generalize this fact for a family of IDS. Our proof follows the arguments given in [1, 10, 15] for space curves, hypersurfaces or ICIS, respectively.

Theorem 4.11. *Let $\{(X_t, 0)\}$ be a good family of IDS of dimension $d \neq 2$ such that the determinantal smoothing of $(X_t, 0)$ is $(d-1)$ -connected and $\nu(X_t, 0)$ is constant for t small enough. Then the family has constant topological type.*

Proof. We observe that we have three cases to consider:

Case 1: If $(X, 0)$ is a curve germ. In this case the theorem is showed in [1].

Case 2: If $(X, 0)$ is an hypersurface germ. This result is showed in [10].

Case 3: If $d > 2$ and $\text{codim} X_0 > 1$. We will proceed to the proof of this case.

Choose an $\epsilon > 0$ small enough such that the origin is the only critical point in $X_0 \cap B_\epsilon$ and such that X_0 is transversal to S_η for any η such that $0 < \eta \leq \epsilon$. If t is small enough, X_t is also transversal to S_ϵ . Consider the map

$$\begin{aligned} \pi : \mathcal{X} \cap (S_\epsilon \times \mathbb{C}) &\rightarrow D \subset \mathbb{C} \\ (x, t) &\mapsto t \end{aligned}$$

Since π is a proper submersion, by the Ehresmann fibration theorem, π is a locally trivial fibration. Therefore for t small enough $(S_\epsilon, X_t \cap S_\epsilon)$ is diffeomorphic to $(S_\epsilon, X_0 \cap S_\epsilon)$.

We know that $(B_\epsilon, X_0 \cap B_\epsilon)$ is homeomorphic to $(C(S_\epsilon), C(X_0 \cap S_\epsilon))$, where $C(M)$ denotes the cone of M . For each t , let $0 < \epsilon_t < \epsilon$ such that 0 is the only critical point of $X_t \cap B_{\epsilon_t}$ and X_t is transversal to S_η for any $0 < \eta \leq \epsilon_t$. Then $(B_{\epsilon_t}, X_t \cap B_{\epsilon_t})$ is homeomorphic to $(C(S_{\epsilon_t}), C(X_t \cap S_{\epsilon_t}))$. We will show that $(S_{\epsilon_t}, X_t \cap S_{\epsilon_t})$ is homeomorphic to $(S_\epsilon, X_t \cap S_\epsilon)$ and, therefore,

$$(B_\epsilon, X_0) \cong (C(S_\epsilon), C(X_0 \cap S_\epsilon)) \cong (C(S_\epsilon), C(X_t \cap S_\epsilon)) \cong (C(S_{\epsilon_t}), C(X_t \cap S_{\epsilon_t})) \cong (B_{\epsilon_t}, X_t).$$

Let $M = B_\epsilon - \overset{\circ}{B}_{\epsilon_t}$ and $V = M \cap X_t$. The boundary of M is $\partial M = S_\epsilon \sqcup S_{\epsilon_t}$ and the boundary of V is $\partial V = (X_t \cap S_\epsilon) \sqcup (X_t \cap S_{\epsilon_t})$ (see fig. 2). Obviously, the inclusions $S_\epsilon, S_{\epsilon_t} \hookrightarrow M$ are homotopy equivalences, we show that $X_t \cap S_\epsilon, X_t \cap S_{\epsilon_t} \hookrightarrow V$ are also homotopy equivalences. By Whitehead and Hurewicz theorems, we only need to prove that the induced maps between the fundamental groups and the homology groups are isomorphisms.

We have that $X_t \cap B_{\epsilon_t}$ and $X_t \cap B_\epsilon$ are contractible (in the first one B_{ϵ_t} is a Milnor ball for $(X_t, 0)$ and the second one follows from Lemma 4.10 and Theorem 4.2). By Van Kampen theorem,

$$1 = \pi_1(X_t \cap B_\epsilon) = \pi_1(X_t \cap B_{\epsilon_t}) *_{\pi_1(X_t \cap S_{\epsilon_t})} \pi_1(V),$$

but $\pi_1(X_t \cap B_{\epsilon_t}) = 1$, $\pi_1(X_t \cap S_{\epsilon_t}) = \pi_1(X_{t,A}) = 1$, so necessarily $\pi_1(V) = 1$ and the map $\pi_1(X_t \cap S_{\epsilon_t}) \rightarrow \pi_1(V)$ is an isomorphism. On the other hand, $\pi_1(X_t \cap S_\epsilon) = \pi_1(X_0 \cap S_\epsilon) = \pi_1(X_A) = 1$, hence the map $\pi_1(X_t \cap S_\epsilon) \rightarrow \pi_1(V)$ is also an isomorphism.

For the homology, we use the Mayer-Vietoris sequence:

$$\begin{aligned} \dots &\longrightarrow H_{i+1}(X_t \cap B_\epsilon) \longrightarrow H_i(X_t \cap S_{\epsilon_t}) \longrightarrow H_i(X_t \cap B_{\epsilon_t}) \oplus H_i(V) \longrightarrow \\ &\longrightarrow H_i(X_t \cap B_\epsilon) \longrightarrow \dots \end{aligned}$$

Since $H_i(X_t \cap B_\epsilon) = 0$ and $H_i(X_t \cap B_{\epsilon_t}) = 0$, for all $i > 0$, we deduce that the map $H_i(X_t \cap S_{\epsilon_t}) \rightarrow H_i(V)$ is an isomorphism for all $i > 0$. Finally, the map $H_i(X_t \cap S_\epsilon) \rightarrow H_i(V)$ is also an isomorphism by the Poincaré duality theorem for cobordism (see [9]).

Therefore, the pairs $(S_\epsilon, X_t \cap S_\epsilon)$ and $(S_{\epsilon_t}, X_t \cap S_{\epsilon_t})$ are h -cobordant (see [17]). Furthermore, $\dim X_t \cap S_\epsilon = 2d - 1 \geq 5$ and $\text{codim}_{S_\epsilon}(X_t \cap S_\epsilon) = 2N - 1 - (2d - 1) = 2(N - d) \geq 4$, (where now the dimensions are taken as real manifolds). This implies that $\pi_1(S_\epsilon - S_\epsilon \cap X_t) = 1$. Hence, by the relative h -cobordism theorem of Smale [17, Theorem 1.4], $(S_{\epsilon_t}, X_t \cap S_{\epsilon_t})$ is homeomorphic to $(S_\epsilon, X_t \cap S_\epsilon)$. □




FIGURE 2.

We remark that the condition that the determinantal smoothing is $(d - 1)$ -connected is satisfied in case that X_t is a curve ([1]); $d = N - 1$, that is, X_t is a hypersurface ([12]) and $s = 1$, that is, X_t is an ICIS ([7]). Moreover, it is also known that the constancy of $\nu(X_t, 0)$ implies that the family is good in such cases. Therefore, Theorem 4.11 indeed extends the results of [10, 15].

5. WHITNEY EQUISINGULARITY

It is well known that any Whitney equisingular family is topologically trivial by the Thom first isotopy lemma. However, the converse is not true in general. In fact, the Briançon-Speder example [2] shows that the μ -constant condition is not a sufficient condition for the Whitney equisingularity of a family (even for a family of hypersurfaces). However, it is known that a family of ICIS $(X_t, 0)$ of dimension d is Whitney equisingular if and only if the polar multiplicities $m_i(X_t, 0)$, $i = 0, \dots, d$ are constant on t (see Gaffney and Teissier [5, 18]). In this section, we use our definition of d -th polar multiplicity to generalize the results of Gaffney and Teissier in [5, 18].

We start by giving the relationship between the d -th polar multiplicity of an IDS of dimension d and of a determinantal deformation of it.

Lemma 5.1. *Let $(X_0, 0)$ be a d -dimensional IDS and let X_t be a determinantal deformation. For a generic linear projection $p : \mathbb{C}^N \rightarrow \mathbb{C}$, we have:*

$$m_d(X_0, 0) = \sum_{y \in X_t^0} \mu(p|_{X_t}, y) + \sum_{x \in S(X_t)} m_d(X_t, x),$$

where X_t^0 denotes the smooth part of X_t .

Proof. We choose a ball B_ϵ such that the origin is the only singular point of X_0 in B_ϵ . Let $\{x_1, \dots, x_k\}$ be the singular set of X_t and $\{y_1, \dots, y_l\}$ be the singular set of $p|_{X_t^0}$.

For each $i = 1, \dots, k$, choose a Milnor disc D_i for (X_t, x_i) and for each $j = 1, \dots, l$, choose a Milnor disc E_j around y_j for $p|_{X_t^0}$ (see fig. 3). Let p_a be a generic linear deformation of p and let $X_{t,A}$ be a determinantal smoothing such that $p_a|_{X_{t,A}}$ is a Morse

function. We have

$$\begin{aligned}
m_d(X_0, 0) &= \#S(p_a|_{X_{t,A}}) \\
&= \sum_{j=1}^l \#S(p_a|_{X_{t,A} \cap E_j}) + \sum_{i=1}^k \#S(p_a|_{X_{t,A} \cap D_i}) \\
&= \sum_{j=1}^l \mu(p|_{X_t, y_j}) + \sum_{i=1}^k m_d(X_t, x_i).
\end{aligned}$$

□




FIGURE 3.

We recall from Teissier [18] the definition of the relative polar multiplicities $m_i(\mathcal{X}, \pi, 0)$, with $i = 0, \dots, d$, of a $(d+1)$ -dimensional variety $(\mathcal{X}, 0)$ with respect to $\pi : (\mathcal{X}, 0) \rightarrow (\mathbb{C}, 0)$. Given a generic linear projection $P : \mathbb{C}^{N+1} \rightarrow \mathbb{C}^{d-i+1}$, then $m_i(\mathcal{X}, \pi, 0) = m_0(\overline{S(P|_{\mathcal{X}^0})}, 0)$.

Lemma 5.2. *Let $\{(X_t, 0)\}_{t \in D}$ be a good family of d -dimensional IDS. If $m_d(X_t, 0)$ is constant on $t \in D$, then $m_d(\mathcal{X}, \pi, 0) = 0$.*

Proof. Choose $\epsilon > 0$ such that the single critical point of $X_0 \cap B_\epsilon$ is the origin. From the previous lemma, we have

$$m_d(X_0, 0) = \sum_{y \in X_t^0} \mu(p|_{X_t, y}) + m_d(X_t, 0),$$

where $p : \mathbb{C}^N \rightarrow \mathbb{C}$ is any generic linear projection. Since $m_d(X_t, 0)$ is constant and all the terms in the right hand side of the equation are bigger or equal to 0, $\mu(p|_{X_t, y}) = 0$ for all $y \in X_t^0$ and hence $p|_{X_t^0}$ is regular. We denote $P(x, t) = (p(x), t)$, then

$$m_d(\mathcal{X}, \pi, 0) = m_0(\overline{S(P|_{\mathcal{X}^0})}, 0) = m_0(\overline{\{(x, t) : x \in S(p|_{X_t^0})\}}, 0) = m(\emptyset, 0) = 0.$$

□

We will present now the main result of this section. The theorem bellow generalizes the results of Gaffney [5] and Teissier [18] by showing necessary and sufficient conditions for a family of IDS to be Whitney equisingular.

Theorem 5.3. *A good family of d -dimensional IDS $\{(X_t, 0)\}_{t \in D}$ is Whitney equisingular if and only if all the polar multiplicities $m_i(X_t, 0)$, $i = 0, \dots, d$ are constant on $t \in D$.*

Proof. Assume, first, that the family is Whitney-equisingular. From the results of Tessier in [18], $m_i(\mathcal{X}, \pi, (0, t))$ is constant on $t \in D$ for $i = 1, \dots, d$. In particular, $m_d(\mathcal{X}, \pi, (0, t))$ is constant on $t \in D$.

We have that $m_d(\mathcal{X}, \pi, 0) = 0$. In fact, if it was not true, then $m_d(\mathcal{X}, \pi, (0, t)) = m_d(\mathcal{X}, \pi, 0) \neq 0$, for $t \in D$. Therefore, since $(0, t)$ does not belong to $S(P|_{\mathcal{X}^0})$, we would have

$$T = \{0\} \times D \subset \overline{S(P|_{\mathcal{X}^0})} - S(P|_{\mathcal{X}^0}),$$

but this is not possible because $\overline{S(P|_{\mathcal{X}^0})}$ is analytic of dimension 1 and hence $\overline{S(P|_{\mathcal{X}^0})} - S(P|_{\mathcal{X}^0})$ has dimension 0.

Now if $m_d(\mathcal{X}, \pi, 0) = 0$, we have $\overline{S(P|_{\mathcal{X}^0})} = \emptyset$ and thus, $S(p|_{\mathcal{X}^0}) = \emptyset$. This implies that

$$\sum \mu(p|_{\mathcal{X}_t^0}, y_i) = 0,$$

and from Lemma 5.2, $m_d(X_t, 0) = m_d(X_0, 0)$.

The fact that the other polar multiplicities are constant follows from Corollary 5.6 of [5].

Assume, now, that $m_i(X_t, 0)$ is constant for all $i = 0, \dots, d$. In particular, $m_d(X_t, 0)$ is constant and from Lemma 5.2,

$$m_d(\mathcal{X}, \pi, 0) = 0.$$

Hence, the family X_t is Whitney equisingular by Corollary 5.12 of [5]. \square

We remark that in the case of a family of ICIS $\{(X_t, 0)\}_{t \in D}$ of dimension d , the condition that $m_d(X_t, 0)$ is constant on t implies that the family is good (see [5, 5.9]).

REFERENCES

- [1] R. O. Buchweitz, G. M. Greuel, *The Milnor number and deformations of complex curve singularities*, Invent. Math. **58** (1980), No. 3, 241–248.
- [2] J. Briançon, J. P. Speder, *La trivialité topologique n'implique pas les conditions de Whitney*, C. R. Acad. Sci. Paris Sér. A-B **280** (1975), No. 6, 365–367.
- [3] J. Damon, B. Pike, *Solvable groups, free divisors and nonisolated matrix singularities II: Vanishing topology*, Geom. Topol. **18** (2014), No. 2, 911–962.
- [4] G. M. Greuel, J. Steenbrik, *On the topology of smoothable singularities. Singularities*, Part 1 (Arcata, Calif., 1981), Proc. Sympos. Pure Math. **40**, Amer. Math. Soc., Providence, R.I. (1983), 535–545.
- [5] T. Gaffney, *Polar multiplicities and equisingularity of map germs*, Topology **32** (1993), No. 2, 185–223.
- [6] T. Gaffney, *Multiplicities and equisingularity of ICIS germs*, Invent. Math. **123** (1996), No. 1, 209–220.
- [7] H. A. Hamm, *Lokale topologische Eigenschaften komplexer Räume*, Math. Ann. **191** (1971), 235–252.
- [8] T. W. Hungerford, *Algebra*, Reprint of the 1974 original. Graduate Texts in Mathematics **73**, Springer-Verlag, New York-Berlin (1980).
- [9] A. A. Kosinski, *Differential Manifolds*, Pure and Applied Mathematics **138**, Academic Press, Inc., Boston MA (1993).
- [10] Lê Dung Tráng, C. P. Ramanujam, *The invariance of Milnor's number implies the invariance of the topological type*, Amer. J. Math. **98** (1976), No. 1, 67–78.
- [11] Lê Dung Tráng, B. Teissier, *Variétés polaires locales et classes de Chern de variétés singulières*, Ann. Math. **114** (1981), No. 3, 457–491.

- [12] J. Milnor, *Singular points of complex hypersurfaces*, Ann. of Math. Studies no. 61, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo (1968).
- [13] J. J. Nuño-Ballesteros, J. N. Tomazella, *The Milnor number of a function on a space curve germ*, Bull. London Math. Soc. **40** (2008), No. 1, 129–138.
- [14] J. J. Nuño-Ballesteros, B. Oréface-Okamoto, J. N. Tomazella, *The vanishing Euler characteristic of an isolated determinantal singularity*, Israel J. Math. **197** (2013), No. 1, 475–495.
- [15] A. J. Parameswaran, *Topological equisingularity for isolated complete intersection singularities*. Compositio Math. **80** (1991), No. 3, 323–336.
- [16] D. Siersma, *Vanishing cycles and special fibres*. Singularity theory and its applications, Part I (Coventry, 1988/1989), Lecture Notes in Math. **1462**, Springer, Berlin, (1991), 292–301.
- [17] S. Smale, *On the structure of manifolds*, American J. of Math. **84**, (1962), No. 3, 387–399.
- [18] B. Teissier, *Cycles évanescents, sections planes et conditions de Whitney*, Singularités à Cargèse (Rencontre Singularités Géom. Anal., Inst. Études Sci., Cargèse, 1972), Asterisque, Soc. Math. France, Paris (1973), Nos. 7 et 8, 285–362.
- [19] O. Zariski, *Contributions to the problem of equisingularity. 1970 Questions on Algebraic Varieties*, Edizioni Cremonese, Rome (1969), 261–343.

DEPARTAMENT DE GEOMETRIA I TOPOLOGIA, UNIVERSITAT DE VALÈNCIA, CAMPUS DE BURJASSOT,
46100 BURJASSOT SPAIN

E-mail address: Juan.Nuno@uv.es

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE SÃO CARLOS, CAIXA POSTAL 676,
13560-905, SÃO CARLOS, SP, BRAZIL

E-mail address: bruna@dm.ufscar.br

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE SÃO CARLOS, CAIXA POSTAL 676,
13560-905, SÃO CARLOS, SP, BRAZIL

E-mail address: tomazella@dm.ufscar.br