

BOUNDARY VALUES OF FUNCTIONS OF DIRICHLET SPACES L_2^1 ON CAPACITARY BOUNDARIES

V. GOL'DSHTEIN AND A. UKHLOV

ABSTRACT. We prove that any weakly differentiable function with square integrable gradient can be extended to a capacitary boundary of any simply connected plane domain $\Omega \neq \mathbb{R}^2$ except a set of a conformal capacity zero. For locally connected at boundary points domains the capacitary boundary coincides with the Euclidean one. A concept of a capacitary boundary was proposed by V. Gol'dshtein and S. K. Vodop'yanov in 1978 for a study of boundary behavior of quasi-conformal homeomorphisms. We prove in details the main properties of the capacitary boundary. An abstract version of the extension property for more general classes of plane domains is discussed also.

1. INTRODUCTION

Let Ω be a domain in \mathbb{R}^2 . We consider a Dirichlet space (a uniform Sobolev space) $L_2^1(\Omega)$ of locally integrable functions with the square integrable weak gradient $\nabla u \in L_2(\Omega)$ equipped with the seminorm

$$\|u\|_{L_2^1(\Omega)} = \|\nabla u\|_{L_2(\Omega)}.$$

The paper is devoted to study of the boundary behavior for functions $u \in L_2^1(\Omega)$.

By the standard definition functions of $L_2^1(\Omega)$ are defined only up to a set of measure zero, but they can be redefined quasieverywhere i. e. up to a set of conformal capacity zero. Indeed, every function $u \in L_2^1(\Omega)$ has a unique quasicontinuous representation $\tilde{u} \in L_2^1(\Omega)$. A function \tilde{u} is termed quasicontinuous if for any $\varepsilon > 0$ there is an open set U_ε such that the conformal capacity of U_ε is less then ε and the function \tilde{u} is continuous on the set $\Omega \setminus U_\varepsilon$ (see, for example [15, 23]). The concept of quasicontinuity can be obviously extended to the closure $\overline{\Omega}$ of Ω .

In this paper we deals with quasicontinuous representations of functions $u \in L_2^1(\Omega)$.

One of main results of the paper is:

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^2$, $\Omega \neq \mathbb{R}^2$, be a simply connected domain which is locally connected at any boundary point $x \in \partial\Omega$. Then for any function $u \in L_2^1(\Omega)$ there exists a quasicontinuous function $\tilde{u} : \overline{\Omega} \rightarrow \mathbb{R}$ such that $\tilde{u}|_\Omega = u$.*

Remark 1.2. The quasicontinuous function $\tilde{u} : \overline{\Omega} \rightarrow \mathbb{R}$ is defined at any point of $\partial\Omega$ except a set of conformal capacity zero (i.e. quasieverywhere).

The main ingredient of our method is a well-known concept of the conformal capacity and a less known concept of the conformal capacitary boundary introduced by V. Gol'dshtein and S. K. Vodop'yanov [14] for quasiconformal homeomorphisms.

⁰**Key words and phrases:** Sobolev Spaces, Conformal Mappings.

⁰**2000 Mathematics Subject Classification:** 46E35, 30C65, 30C85.

In the plane case "points" of the conformal capacity boundary coincide with the Caratheodory prime ends.

Main properties of the space $L_2^1(\mathbb{D})$ where $\mathbb{D} \subset \mathbb{R}^2$ is the unit disc are well known. Dirichlet spaces $L_2^1(\Omega)$ are conformal invariants. Therefore the Riemann Mapping Theorem permits us to transfer necessary information about boundary behavior of spaces $L_2^1(\Omega)$ from $L_2^1(\mathbb{D})$ in the case of simply connected domains Ω .

More precisely, we extend the concept of quasicontinuity to a "capacity" completion of a domain Ω . We construct a conformal capacity boundary as a completion $\{\widetilde{\Omega}_\rho, \rho\}$ of a metric space $\{\Omega_\rho, \rho\}$ for a conformal capacity metric ρ (see section 1). Roughly speaking, an "ideal" capacity boundary point is a boundary continuum of the conformal capacity zero.

Our method allow us to treat the general case of simply connected plane domains $\Omega \subset \mathbb{R}^2$. We prove that any function $u \in L_2^1(\Omega)$ has a quasicontinuous extension onto the conformal capacity boundary $H_\rho = \widetilde{\Omega}_\rho \setminus \Omega_\rho$. The main result is:

Theorem 1.3. *Let $\Omega \subset \mathbb{R}^2$ be a simply connected domain, $\Omega \neq \mathbb{R}^2$. Then for any function $u \in L_2^1(\Omega)$ there exists a quasicontinuous function $\tilde{u} : \widetilde{\Omega}_\rho \rightarrow \mathbb{R}$ such that $\tilde{u}|_\Omega = u$.*

Remark 1.4. A concept of the conformal capacity metric ρ and the conformal capacity boundary was proposed in [14]. By a quasi-invariance of the conformal capacity under (quasi)conformal homeomorphisms any such homeomorphism $\varphi : \Omega \rightarrow \Omega'$ is a bi-Lipschitz homeomorphism $\varphi : (\Omega, \rho) \rightarrow (\Omega', \rho)$ for corresponding conformal metrics and can be extended to a homeomorphism $\tilde{\varphi} : (\widetilde{\Omega}, \rho) \rightarrow (\widetilde{\Omega}', \rho)$ of the capacity completions [14]. Recall that the paper [14] is a short note and contains only sketches of proofs.

There is the vast literature concerning of "ideal" boundaries of plane domains in the context of conformal homeomorphisms. We discuss few such concepts in the last section.

The paper is organized as follows:

Main properties of the conformal capacity metric are proved in Section 2. The focus is on the local properties of the metric at boundary points and its dependence on the local topological properties of the boundary. In section 3 we discuss an analog of the Luzin property for the capacity metric. In section 4 we discuss a sufficient condition for existence of an extension of functions $u \in L_2^1(\Omega)$ to the capacity boundary. We call this condition as a strong Luzin property for the capacity metric. We prove this condition for comparatively large classes of domains that include extension domains for $L_2^1(\Omega)$. In Section 5 we apply the abstract construction of Section 4 to simply connected plane domains and we prove main results about extension of functions $u \in L_2^1(\Omega)$ to the capacity boundary.

In terminology of the theory of Sobolev spaces we solved the classical trace problem for $L_2^1(\Omega)$ in simply connected plane domains.

Remark 1.5. The classical trace problem for Sobolev spaces is of essential interest, mainly due to its important applications to boundary-value problems for partial differential equations. Boundary value problems can be specified with the help of traces to $\partial\Omega$ of Sobolev functions.

There is an extensive literature devoted to the trace problem of Sobolev spaces. Among the multitude of results we mention the monographs of P. Grisvard [7],

J. L. Lions and E. Magenes [21], V. G. Maz'ya and S. Poborchi [23], [28], and the papers [2], [3], [6], [16], [17], [18], [24], [25], [26], [27], [31], [32], [37], [40], [41].

For smooth domains the traces of Sobolev functions are Besov spaces. In the case of Lipschitz domains the traces can be described also in terms of Besov spaces. For arbitrary non Lipschitz domain the trace problem is open. For cusp type singularities a description of traces can be found in [12] in terms of weighted Sobolev spaces.

2. CONFORMAL CAPACITARY METRIC

Let Ω be a plane domain and F_0, F_1 two disjoint compact subset of Ω . We call the triple $E = (F_0, F_1; \Omega)$ a condenser.

The value

$$\text{cap}(E) = \text{cap}(F_0, F_1; \Omega) = \inf \int_{\Omega} |\nabla v|^2 dx,$$

where the infimum is taken over all nonnegative functions $v \in C(\Omega) \cap L_2^1(\Omega)$, such that $v = 0$ in a neighborhood of the set F_0 , and $v \geq 1$ in a neighborhood of the set F_1 , is called the conformal capacity of the condenser $E = (F_0, F_1; \Omega)$.

For finite values of capacity $0 < \text{cap}(F_0, F_1; \Omega) < +\infty$ there exists a unique function u_0 (an extremal function) such that:

$$\text{cap}(F_0, F_1; \Omega) = \int_{\Omega} |\nabla u_0|^2 dx.$$

An extremal function is continuous in Ω , monotone in the domain $\Omega \setminus (F_0 \cup F_1)$, equal to zero on F_0 and is equal to one on F_1 [15, 39].

Definition 2.1. A homeomorphism $\varphi : \Omega \rightarrow \Omega'$ between plane domains is called K -quasiconformal if it preserves orientation, belongs to the Sobolev class $L_{2,loc}^1(\Omega)$ and the distortion inequality

$$\max_{|\xi|=1} |D\varphi(x) \cdot \xi| \leq K \min_{|\xi|=1} |D\varphi(x) \cdot \xi|$$

holds for almost all $x \in \Omega$.

Infinitesimally, quasiconformal homeomorphisms carry circles to ellipses with eccentricity uniformly bounded by K . If $K = 1$ we recover conformal homeomorphisms, while for $K > 1$ plane quasiconformal mappings need not be smooth. The theory of quasiconformal mappings can be find, for example, in [38].

It is well known that the conformal capacity is quasi-invariant under action of plane quasiconformal homeomorphisms.

2.1. Definition of the conformal capacity metrics. A connected closed (with respect to Ω) set is called a continuum. Fix a continuum F in the domain $\Omega \subset \mathbb{R}^2$ and a compact domain V such that $F \subset V \subset \overline{V} \subset \Omega$, and the boundary ∂V is an image of the unit circle $S(0, 1)$ under some quasiconformal homeomorphism of \mathbb{R}^2 .

Definition 2.2. Choose arbitrarily points $x, y \in \Omega \subset \mathbb{R}^n$ and joint x, y by a rectifiable curve $l(x, y)$. Define the conformal capacity distance between x and y in Ω with respect to pair (F, V) as the following quantity

$$\rho_{(F,V)}(x, y) = \inf_{l(x,y)} \{ \text{cap}^{\frac{1}{2}}(F, l(x, y) \setminus V; \Omega) + \text{cap}^{\frac{1}{2}}(\partial\Omega, l(x, y) \cap V; \Omega) \}$$

where the infimum is taken over all curves $l(x, y)$ satisfying the above conditions.

This definition firstly was introduced in [14], where was claimed that the distance $\rho_{(F,V)}(x, y)$ is a metric in Ω which is quasi-invariant under quasiconformal homeomorphisms and is invariant under conformal ones.

Denote by $\{\tilde{\Omega}, \rho_{(F,V)}\}$ the standard completion of the metric space $\{\Omega, \rho_{(F,V)}\}$ and by H_ρ the set $\{\tilde{\Omega}, \rho_{(F,V)}\} \setminus \{\Omega, \rho_{(F,V)}\}$. We call H_ρ a conformal capacity boundary of Ω . It will be proved that the topology of H_ρ does not depends on choice of a pair (F, V) . Moreover two conformal capacity metrics are equivalent for any different choice of pairs (F_1, V_1) and (F_2, V_2) . This is a justification of the notation H_ρ for the conformal capacity boundary.

2.2. Equivalence of different conformal capacity distances. We start from an important technical observation:

Lemma 2.3. *Let Ω be a domain in \mathbb{R}^2 and F_1 be a compact in Ω . Then there exists a constant $0 < K < \infty$ such that*

$$\frac{1}{K} \text{cap}(F_{02}, F_1; \Omega) \leq \text{cap}(F_{01}, F_1; \Omega) \leq K \text{cap}(F_{02}, F_1; \Omega)$$

for every compacts $F_{01} \subset \Omega$, $F_{02} \subset \Omega$ such that compacts F_{01} , F_{02} , F_1 are mutually disjoint.

Proof. Let $U_1 \supset F_{01}$ be a ε -neighborhood of a set F_{01} in Ω and $U_2 \supset F_{02}$ be a ε -neighborhood of the set F_{02} in Ω such that $\overline{U_1} \subset \Omega$ and $\overline{U_2} \subset \Omega$. Because the set F_{01} is a compact set there exist a finite covering $\{B_i\}_{i=1, \dots, N}$ of F_{01} by balls $B_i \in \Omega$ of radius ε . It means that $\Omega \supset \bigcup_{i=1}^N B_i \supset F_1$.

Then

$$\text{cap}(F_{01}, F_1; \Omega) \leq \text{cap}\left(\bigcup_{i=1}^N \overline{B_i}, F_1; \Omega\right).$$

For every ball B_i , $i = 1, \dots, N$, we can construct a bi-Lipschitz homeomorphism ψ_i of Ω onto itself that maps B_i onto B_1 . Using quasi-invariance of the conformal capacity under bi-Lipschitz homeomorphisms we have

$$\text{cap}(F_{01}, F_1; \Omega) \leq \text{cap}\left(\bigcup_{i=1}^N \overline{B_i}, F_1; \Omega\right) \leq C_1 \cdot \text{cap}(\overline{B_1}, F_1; \Omega)$$

where a constant C_1 depends only on the multiplicity of the covering and F_1 .

Applying the same construction to F_{02} we can construct a finite covering $\{\tilde{B}_i\}_{i=1}^N$ of F_{02} by balls $\tilde{B}_i \in \Omega$ of radius ε such that $\Omega \supset \bigcup_{i=1}^N \tilde{B}_i \supset F_{02}$. Then

$$\text{cap}(F_{02}, F_1; \Omega) \leq C_2 \cdot \text{cap}(\overline{\tilde{B}_1}, F_1; \Omega)$$

where a constant C_2 depends only on the multiplicity of the covering of F_2 and F_2 itself.

To combine both estimate we construct a bi-Lipschitz homeomorphism φ of Ω that maps the ball $B_1 \subset U_1$ onto a ball $\tilde{B}_1 \subset U_2$. Using quasi invariance of the conformal capacity under bi-Lipschitz homeomorphisms we obtain

$$\text{cap}(\overline{B_1}, F_1; \Omega) \leq M \text{cap}(\overline{\tilde{B}_1}, F_1; \Omega) \leq M \text{cap}(\overline{U_2}, F_1; \Omega)$$

where M is the Lipschitz constant of φ . Hence

$$\text{cap}_p(F_{01}, F_1; \Omega) \leq C_1 M \text{cap}_p(\overline{U}_2, F_1; \Omega).$$

Therefore

$$(2.1) \quad \text{cap}(F_{01}, F_1; \Omega) \leq \lim_{\varepsilon \rightarrow 0} K \text{cap}(\overline{U}_2, F_1; \Omega) = K \text{cap}(F_{02}, F_1; \Omega).$$

Using the same construction and the inverse bi-Lipschitz homeomorphisms φ^{-1} we obtain

$$(2.2) \quad \text{cap}(F_{02}, F_1; \Omega) \leq \lim_{\varepsilon \rightarrow 0} C_2 M \text{cap}(\overline{U}_1, F_1; \Omega) = C_2 M \text{cap}(F_{01}, F_1; \Omega).$$

Using inequalities 2.1 and 2.2 we obtain

$$\frac{1}{K} \text{cap}(F_{02}, F_1; \Omega) \leq \text{cap}(F_{01}, F_1; \Omega) \leq K \text{cap}(F_{02}, F_1; \Omega).$$

□

Using this lemma we prove

Theorem 2.4. *Let Ω be a domain in \mathbb{R}^2 . Suppose that $\rho_{(F_1, V_1)}$ and $\rho_{(F_2, V_2)}$ are two conformal capacity distances on Ω . Then there exists a constants $0 < K < \infty$ such that*

$$\frac{1}{K} \rho_{(F_2, V_2)}(x, y) \leq \rho_{(F_1, V_1)}(x, y) \leq K \rho_{(F_1, V_1)}(x, y) \text{ for any } x, y \in \Omega.$$

Proof. Consider two continuums F_1 and F_2 in the domain Ω . Without loss of generality we can suppose that any admissible for the conformal capacity metric curve $l(x, y)$ does not intersect V_1 and V_2 . Then by Lemma 2.3

$$\frac{1}{K} \text{cap}(F_2, l(x, y); \Omega) \leq \text{cap}(F_1, l(x, y); \Omega) \leq K \text{cap}(F_2, l(x, y); \Omega)$$

for any admissible for the conformal capacity metric curve $l(x, y)$. Hence

$$\frac{1}{K} \rho_{(F_2, V_2)}(x, y) \leq \rho_{(F_1, V_1)}(x, y) \leq K \rho_{(F_1, V_1)}(x, y)$$

for any $x, y \in \Omega$. □

2.3. Conformal capacity distance is a metric. The sketch of the proof can be found in [14]. For readers convenience we prove this fact in details.

Lemma 2.5. *Let Ω be a domain in \mathbb{R}^2 and F be a compact subdomain of Ω . Suppose that $x \in \Omega \setminus \overline{F}$, $B(x, 2r) \subset \Omega \setminus \overline{F}$, $r > 0$ and curve γ joints x and $S(x, 2r)$. Then $\text{cap}(F, \gamma; \Omega) > c(r) > 0$.*

Proof. Choose a continuum F_r , such that $F_r \cap \gamma = \emptyset$, which connects spheres $S(x, r)$ and $S(x, 2r)$. By Lemma 2.3

$$\text{cap}(F, \gamma; \Omega) \geq Q \text{cap}(F_r, \gamma; \Omega).$$

Using the properties of capacity and Proposition 4.6 from [8] we obtain

$$\text{cap}(F_r, \gamma; \Omega) \geq \text{cap}(F_r, \gamma \cap \{z : r \leq |x - z| \leq 2r\}; \Omega) \geq c(r) > 0.$$

Hence $\text{cap}(F, \gamma; \Omega) \geq c(r) > 0$. □

Theorem 2.6. *Let Ω be a domain in \mathbb{R}^2 . Then the $\rho_{(F, V)}(x, y)$ is a metric in Ω .*

Proof. By standard properties of the conformal capacity [23] $\rho_{(F,V)}(x, x) = 0$.

Let $\rho_{(F,V)}(x, y) = 0$. Assume by contradiction that $x \neq y$ and denote the Euclidean distance $\text{dist}(x, y)$ as r . Then there exists a sequence of curves $\{l_k(x, y)\}$, $k = 1, 2, \dots$, such that

$$\text{cap}(F, l_k(x, y) \setminus V; \Omega) \rightarrow 0 \text{ while } k \rightarrow \infty$$

and

$$\text{cap}(\partial\Omega, l_k(x, y) \cap V; \Omega) \rightarrow 0 \text{ while } k \rightarrow \infty.$$

Hence a sequence of extremal functions $u_k \in L_2^1(\Omega)$ for the capacities

$$\text{cap}(F, l_k(x, y) \setminus V; \Omega)$$

tends to zero in the space $L_2^1(\Omega)$ and a sequence of extremal functions $v_k \in L_2^1(\Omega)$ for the capacities

$$\text{cap}(\partial\Omega, l_k(x, y) \cap V; \Omega)$$

tends to zero in the space $L_2^1(\Omega)$.

So, the sequence u_k tends to zero except a set of the conformal capacity zero and the sequence v_k tends to zero except for a set of the conformal capacity zero. But by Lemma 2.5

$$\|u_k|_{L_2^1(\Omega)}\| + \|v_k|_{L_2^1(\Omega)}\| \geq c(r) > 0 \text{ for all } k.$$

Contradiction. Therefore $x = y$.

By the definition of the conformal capacity we have

$$\begin{aligned} \text{cap}^{\frac{1}{2}}(F, l(x, y) \setminus V; \Omega) + \text{cap}^{\frac{1}{2}}(\partial\Omega, l(x, y) \cap V; \Omega) \\ = \text{cap}^{\frac{1}{2}}(F, l(y, x) \setminus V; \Omega) + \text{cap}^{\frac{1}{2}}(\partial\Omega, l(y, x) \cap V; \Omega). \end{aligned}$$

Hence $\rho_{(F,V)}(x, y) = \rho_{(F,V)}(y, x)$.

Prove the triangle inequality. Choose arbitrary points $x, y, z \in \Omega$. By the sub-additive property of the capacity [23] we obtain

$$\begin{aligned} \text{cap}^{\frac{1}{2}}(F, (l(x, z) \setminus V) \cup (l(z, y) \setminus V); \Omega) \\ \leq \text{cap}^{\frac{1}{2}}(F, (l(x, z) \setminus V); \Omega) + \text{cap}^{\frac{1}{2}}(F, (l(z, y) \setminus V); \Omega) \end{aligned}$$

and

$$\begin{aligned} \text{cap}^{\frac{1}{2}}(\partial\Omega, (l(x, z) \cap V) \cup (l(z, y) \cap V); \Omega) \\ \leq \text{cap}^{\frac{1}{2}}(\partial\Omega, (l(x, z) \cap V); \Omega) + \text{cap}^{\frac{1}{2}}(\partial\Omega, (l(z, y) \cap V); \Omega). \end{aligned}$$

Hence $\rho_{(F,V)}(x, y) \leq \rho_{(F,V)}(x, z) + \rho_{(F,V)}(y, z)$. Therefore $\rho_{p;(F,V)}$ is a metric. \square

Theorem 2.7. *The topology induced by the conformal capacity metric $\rho_{(F,V)}$ into the domain $\Omega \subset \mathbb{R}^2$ coincides with the Euclidean topology.*

Proof. Let $U \subset \Omega$ be an open set with respect to Euclidean metric. Let $x_0 \in U$ such that

$$\overline{B(x_0, 3r)} = \overline{\{x \in \Omega : |x - x_0| < 3r\}} \subset U.$$

Then for every point $y \in \partial B(x_0, r)$ we have that

$$\begin{aligned} \rho_{(F,V)}(x_0, y) = \\ \inf_{l(x_0 y)} \{ \text{cap}^{\frac{1}{2}}(F, l(x_0 y) \setminus V; \Omega) + \text{cap}^{\frac{1}{2}}(\partial\Omega, l(x_0 y) \cap V; \Omega) \} > 0, \end{aligned}$$

since H^1 -Hausdorff measure of the sets $l(x_0y) \setminus V$ (or $l(x_0y) \cap V$ is positive). Hence U is the open set with respect to the conformal capacity metric in the domain Ω . The inverse inclusion can be proved similarly. \square

We complete by the standard manner the metric space $\Omega_\rho = \Omega_{\rho_{(F,V)}}$. In the completion $\tilde{\Omega}_\rho$ we define the conformal capacity boundary of Ω as $H_\rho = \tilde{\Omega}_\rho \setminus \Omega_\rho$. It means that a boundary element $h \in H_\rho$ is a class of fundamental (in the metric ρ) sequences $\{x_n\}_{n=1}^\infty$.

Theorem 2.8. *Suppose that $\rho_{(F_1,V_1)}$ and $\rho_{(F_2,V_2)}$ are two conformal capacity metrics on Ω . The metric spaces $\{H_\rho, \rho_{(F_1,V_1)}\}$ and $\{H_\rho, \rho_{(F_2,V_2)}\}$ are quasi-isometric, i.e there exist a constant $0 < K < \infty$ such that*

$$\frac{1}{K} \rho_{(F_2,V_2)}(x, y) \leq \rho_{(F_1,V_1)}(x, y) \leq K \rho_{(F_1,V_1)}(x, y)$$

for any $x, y \in H_\rho$.

Proof. Consider two continuums F_1 and F_2 in the domain Ω and suppose that sequences $\{x_k\}$ and $\{y_k\}$ are fundamental sequences in the metric $\rho_{(F_2,V_2)}$. If the sequences $\{x_k\}$ and $\{y_k\}$ have limit point $x, y \in \Omega$, then the sequences are fundamental in the metric $\rho_{(F_1,V_1)}$ because in the domain Ω conformal capacity topologies are equivalent to the Euclidean topology. Let fundamental sequences $\{x_k \in \Omega\}$ and $\{y_k \in \Omega\}$ has limit points $x, y \in \{H_\rho, \rho_{(F_2,V_2)}\}$.

By Theorem 2.4 for any $x_k, y_k \in \Omega$ there exist a constant $0 < K < \infty$ such that

$$\frac{1}{K} \rho_{(F_2,V_2)}(x_k, y_k) \leq \rho_{(F_1,V_1)}(x_k, y_k) \leq K \rho_{(F_1,V_1)}(x_k, y_k)$$

Passing to limit by $k \rightarrow \infty$ we conclude the proof. \square

Remark 2.9. Both metrics are equivalent in completions of Ω . It can be proved by the same way but it is more technical. For our main aims this fact is not important.

2.4. Asymptotic behavior of the conformal capacity metric. For study of an asymptotic behavior of the conformal capacity metric we need a few well known estimates of the conformal capacity (see, for example [23]). For readers convenience we reproduce simple proofs of these facts adapted to the conformal capacity metric study:

Lemma 2.10. *Consider the unit disc $\mathbb{D}(0, 1) \subset \mathbb{R}^2$ and continuums*

$$F_0 = (-1, -\frac{1}{2}] \subset \mathbb{D} \text{ and } F_1 = [0, \varepsilon] \subset \mathbb{D}, \quad 0 < \varepsilon < \frac{1}{4}.$$

Then

$$\text{cap}(F_0, F_1; \mathbb{R}^2) \geq c_1 \ln(1 + \varepsilon), \quad c_1 = \text{const.}$$

Proof. Consider the conformal mapping

$$\varphi : \mathbb{C} \rightarrow \mathbb{C}, \quad \varphi(z) = \frac{1}{z}.$$

Then by the capacity estimates [10]

$$\begin{aligned} \text{cap}(F_0, F_1; \mathbb{R}^2) &= \text{cap}(\varphi(F_0), \varphi(F_1); \mathbb{R}^2) \geq \text{cap}([-2, -1], [\frac{1}{\varepsilon}, 1 + \frac{1}{\varepsilon}]; \mathbb{R}^2) \\ &= c_1 \ln \frac{1 + \frac{1}{\varepsilon}}{\frac{1}{\varepsilon}} = c_1 \ln(1 + \varepsilon). \end{aligned}$$

□

Lemma 2.11. *Consider the unit disc $\mathbb{D}(0, 1) \subset \mathbb{R}^2$ and continuums $F_0 = \{z : |z| \geq 1\}$ and $F_1 = [0, \varepsilon] \subset \mathbb{D}$, $0 < \varepsilon < \frac{1}{4}$. Then*

$$\text{cap}(F_0, F_1; \mathbb{R}^2) \leq c_2 \left(\ln \frac{1}{\varepsilon} \right)^{-1}, \quad c_2 = \text{const.}$$

Proof. By the capacity estimates [10]

$$\text{cap}(F_0, F_1; \mathbb{R}^2) \leq \text{cap}(F_0, \overline{D(0, \varepsilon)}, \mathbb{R}^2) = c_2 \left(\ln \frac{1}{\varepsilon} \right)^{-1}.$$

□

From these lemmas immediately follows:

Proposition 2.12. *Let $x = (0, 0)$ and $y = (\varepsilon, 0)$ are points of the unit disc $\mathbb{D}(0, 1) \subset \mathbb{R}^2$. Then we have the following asymptotic behavior of the conformal metric in the unit disk $D(0, 1)$:*

$$\lim_{x \rightarrow y} \frac{\rho(x, y)}{|x - y|} \geq \lim_{\varepsilon \rightarrow 0} c_1 \frac{\ln(1 + \varepsilon)}{\varepsilon} = c_1.$$

and

$$\lim_{x \rightarrow y} \frac{\rho(x, y)}{|x - y|} \leq \lim_{\varepsilon \rightarrow 0} c_2 \frac{\left(\ln \frac{1}{\varepsilon} \right)^{-1}}{\varepsilon} < +\infty.$$

Now we study topological properties of the boundary H_ρ .

Definition 2.13. For arbitrary point $h \in H_\rho$ we consider disks $D(h, \varepsilon)$, $\varepsilon > 0$, defined in terms of the conformal capacity metric $\rho_{(F, V)}$.

Call the set

$$s_h = \bigcap_{\varepsilon > 0} \overline{D(h, \varepsilon) \cap \Omega} \subset \overline{\mathbb{R}^2}$$

the realization (impression) of a boundary element $h \in H_\rho$.

Recall that a domain Ω is called locally connected at a point $z_0 \in \partial\Omega$ if z_0 has arbitrarily small connected neighborhoods in Ω . By C. Caratheodory [4] the domain Ω is locally connected at every boundary point if and only if every prime end has trivial realization.

Lemma 2.14. *Let a realization s_h of a boundary element $h \in H_\rho$ is one-point. Then for every sequence $\{x_m \in \Omega\}$ from $\rho_{(F, V)}(x_m, h) \rightarrow 0$ follows that $|x_m - s_h| \rightarrow 0$ (while $m \rightarrow \infty$).*

Proof. Suppose that $\rho_{(F,V)}(x_m, h) \rightarrow 0$ while $m \rightarrow \infty$. Because the realization s_h of a boundary element $h \in H_\rho$ is a point, then

$$\text{diam} \left(\overline{D(h, \varepsilon) \cap \Omega} \right) = \sup_{x, y \in \overline{D(h, \varepsilon) \cap \Omega}} |x - y| \rightarrow 0 \text{ while } \varepsilon \rightarrow 0.$$

The sequence $\{x_m\}$ belongs to a boundary element $h \in H_\rho$ and so

$$|x_m - x_n| \rightarrow 0 \text{ while } m, n \rightarrow \infty.$$

Hence, the sequence $\{x_n\}$ is a sequence Cauchy in the Euclidean metric, and we have that $|x_m - s_h| \rightarrow 0$ while $m \rightarrow \infty$. \square

Lemma 2.15. *Let a domain Ω is locally connected at a point $x \in \partial\Omega$ and $x \in s_h$ for some and $h \in H_\rho$. Then for every sequence $\{x_m \in \Omega\}$ such that $|x_m - x| \rightarrow 0$ we have $\rho_{(F,V)}(x_m, h) \rightarrow 0$ (while $m \rightarrow \infty$).*

Proof. Since the domain Ω be locally connected at the point $x \in \partial\Omega$ then any two points x_k, x_m from the sequence $\{x_n\}$ can be connected by a geodesic path $l(x_k, x_m)$ such that its length tends to zero for $k, m \rightarrow \infty$. Without loss of generality we can suppose that $l(x_k, x_m) \cap V = \emptyset$. Hence $\text{cap}_\rho(F_1, l(x_k, x_m); \Omega)$ tends to zero for $k, m \rightarrow \infty$ and therefore

$$\lim_{n \rightarrow \infty} \rho_{(F,V)}(x_n, h) = 0.$$

\square

From these lemmas follows:

Theorem 2.16. *Let a domain Ω is locally connected at any point $x \in \partial\Omega$. Then the identical mapping $i : \Omega \rightarrow \Omega$ can be extend to a homeomorphism $\tilde{i}_\rho : \tilde{\Omega} \rightarrow \tilde{\Omega}_\rho$ if and only if all realizations s_h of boundary elements $h \in H_\rho$ are one-points.*

Proof. Suppose that an identical mapping $i : \Omega \rightarrow \Omega$ can be extended to a homeomorphism $\tilde{i}_\rho : \tilde{\Omega} \rightarrow \tilde{\Omega}_\rho$. Then every boundary element $h \in H_\rho$ coincides with a point $x \in \partial\Omega$ and so has an one-point realization.

Inversely, let all realizations s_h of a boundary elements $h \in H_\rho$ are one-points. Then extending an identical mapping $i : \Omega \rightarrow \Omega$ to the mapping $\tilde{i}_\rho : \tilde{\Omega}_\rho \rightarrow \tilde{\Omega}$ by the rule $\tilde{i}_\rho(h) = s_h$ we obtain a one-to-one correspondence $\tilde{i}_\rho : \tilde{\Omega}_\rho \rightarrow \tilde{\Omega}$. Let us check continuity of \tilde{i}_ρ and \tilde{i}_ρ^{-1} .

Suppose that $x_k \rightarrow x$ in $\tilde{\Omega}$. Because all realizations s_h of boundary elements $h \in H_\rho$ are one-points and $x \in s_h$ then by Lemma 2.15 follows $\rho_{(F,V)}(x_m, h) \rightarrow 0$, while $m \rightarrow \infty$. We prove that \tilde{i}_ρ is continuous.

Suppose $h_k \rightarrow h_0$ in $\tilde{\Omega}_\rho$. Because Ω is locally connected then realizations of h_k and h_0 are one point sets and we can identify h_k and h_0 with their realizations. By Lemma 2.14 $h_k \rightarrow h_0$ in Ω . Therefore \tilde{i}_ρ^{-1} is also continuous. Therefore \tilde{i}_ρ is a homeomorphism. \square

2.5. Conformal capacitary boundary and Carathéodory prime ends. The notion of the ideal boundary in the terms of prime end was introduced by Carathéodory [4]. The Cartheodory prime ends represent a compactification of plane domains in the relative distance introduced by Lavrentiev [19]. (A detailed historical sketch can be found in [29]). We prove that the capacitary boundary is homeomorphic to the Carathéodory boundary.

Theorem 2.17. *Let $\Omega \subset \mathbb{R}^2$ be a simply connected domain, $\Omega \neq \mathbb{R}^2$. Then the capacity boundary H_ρ is homeomorphic to the Carathéodory boundary $\partial_C \Omega$.*

Proof. The Carathéodory boundary $\partial_C \Omega$ is homeomorphic to the boundary of the unit disc $\partial \mathbb{D}$. The capacity boundary is homeomorphic to the boundary of the unit disc $\partial \mathbb{D}$ also. Hence the capacity boundary H_ρ is conformally equivalent to the Carathéodory boundary $\partial_C \Omega$. \square

On the base of this theorem we give some examples [5] of boundary elements $h \in H_\rho$ of the conformal capacity boundary.

Example 2.18. *Let*

$$X = \{(x, y) : y = 1/3^n \text{ for some } n \geq 1 \text{ and } -1 \leq x \leq 2\}$$

and

$$Y = \{(x, y) : y = 2/3^n \text{ for some } n \geq 1 \text{ and } -2 \leq x \leq 1\}.$$

Let $\Omega = (-2, 2) \times (0, 1) \setminus (X \cup Y)$. The boundary element of this domain is $h = \{(x, 0) : -1 \leq x \leq 1\}$.

Example 2.19. *Let $\Omega = \mathbb{R}^2 \setminus K$, where K is given in polar coordinates by*

$$K = \{(r, \theta) : \theta = 2\pi p \text{ for some integer } n \geq 1 \text{ and some odd integer } p \\ \text{such that } 0 < p < 2^n, 0 \leq r \leq 1/2^n\}.$$

The boundary element $h \in H_\rho$ of this domain at the origin is homeomorphic to a Cantor set.

By C. Caratheodory [4] the domain Ω is called locally connected at boundary points point if and only if every boundary element has trivial realization. Hence we have the following corollary of Theorem 2.16:

Corollary 2.20. *Let a simply connected domain Ω is locally connected at any point $x \in \partial \Omega$. Then the identical mapping $i : \Omega \rightarrow \Omega$ can be extend to a homeomorphism $\tilde{i}_\rho : \tilde{\Omega}_\rho \rightarrow \tilde{\Omega}$.*

3. STRONG LUZIN PROPERTY FOR THE CAPACITARY METRIC AND BOUNDARY VALUES OF SOBOLEV FUNCTIONS

Recall the notion of the conformal capacity of a set $E \subset \Omega$. Let Ω be a domain in \mathbb{R}^2 and a compact $F \subset \Omega$. The conformal capacity of the compact F is defined by

$$\text{cap}(F; \Omega) = \inf\{\|u\|_{L_2^1(\Omega)}^2, u \geq 1 \text{ on } F, u \in C_0(\Omega)\}.$$

By the similar way we can define the conformal capacity of open sets.

For arbitrary set $E \subset \Omega$ we define a inner conformal capacity as

$$\underline{\text{cap}}(E; \Omega) = \sup\{\text{cap}(e; \Omega), e \subset E \subset \Omega, e \text{ is a compact}\},$$

and a outer conformal capacity as

$$\overline{\text{cap}}(E; \Omega) = \inf\{\text{cap}(U; \Omega), E \subset U \subset \Omega, U \text{ is an open set}\}.$$

A set $E \subset \Omega$ is called conformal capacity measurable, if $\underline{\text{cap}}(E; \Omega) = \overline{\text{cap}}(E; \Omega)$. The value

$$\text{cap}(E; \Omega) = \underline{\text{cap}}(E; \Omega) = \overline{\text{cap}}(E; \Omega)$$

is called the conformal capacity of the set $E \subset \Omega$.

The classical Luzin theorem asserts that every measurable function is **uniformly** continuous if it is restricted to the complement of an open set of arbitrary small measure. It is reasonable to conjecture that every function $u \in L_2^1(\Omega)$ is **uniformly** continuous if it is restricted to the complement of an open subset of $\Omega \subset \mathbb{R}^2$ of arbitrary small conformal capacity. Unfortunately this conjecture is wrong for an arbitrary domain and is correct only under additional conditions on Ω . The weak version of The Luzin theorem is correct for the capacity:

Theorem 3.1. (*Weak Luzin theorem for p -capacity [23]*) *Let $\Omega \subset \mathbb{R}^2$ be an open set. For any function $u \in L_2^1(\Omega)$ and for any $\varepsilon > 0$ there exists an open set $U_\varepsilon \in \Omega$, $\text{cap}(U_\varepsilon; \Omega) < \varepsilon$, such that $u|_{\Omega \setminus U_\varepsilon}$ is continuous.*

We discuss here the strong version of the Luzin property for the capacity:

Definition 3.2. Strong Luzin capacitary property. A domain $\Omega \subset \mathbb{R}^2$ possesses a strong Luzin capacitary property if for every function $u \in L_2^1(\Omega)$ and for any $\varepsilon > 0$ there exists an open set U_ε of the conformal capacity less then ε and such that the restriction of the function u on $\Omega \setminus U_\varepsilon$ is uniformly continuous for the conformal capacitary metric.

This property looks very restrictive, but, in reality, it is correct for a large set of domains. We prove in this section that any extension domain possess the Luzin capacitary property and we prove in next section that any quasiconformal homeomorphism preserves the strong Luzin capacitary property.

Our main motivation for a study of this property is the following result:

Theorem 3.3. *Let a domain $\Omega \subset \mathbb{R}^2$ possesses the strong Luzin capacitary property. Then for any function $u \in L_2^1(\Omega)$ there exists a quasicontinuous function $\tilde{u} : \tilde{\Omega}_\rho \rightarrow \mathbb{R}$ defined quasieverywhere on H_ρ such that $\tilde{u}|_\Omega = u$.*

Proof. Because Ω possesses the strong Luzin capacitary property then for every $\varepsilon > 0$ there exists an open set $U_\varepsilon \subset \Omega$ such that $\text{cap}(U_\varepsilon) < \varepsilon$ and any function $u \in L_2^1(\Omega)$ is uniformly continuous for the conformal capacitary metric on the closed (with respect to Ω) set $\Omega^\varepsilon = \Omega \setminus U_\varepsilon$. Consider the completion $\tilde{\Omega}^\varepsilon$ of the set Ω^ε in the complete metric space $(\tilde{\Omega}_\rho, \rho)$. The function $u \in L_p^1(\Omega)$ will be uniformly continuous on the metric space $(\Omega_\rho^\varepsilon, \rho)$. Hence by the Tietz theorem there exists an extension \tilde{u}_ε of u to $\tilde{\Omega}^\varepsilon$. Put $\tilde{\Omega}^0 = \cup_{\varepsilon > 0} \tilde{\Omega}^\varepsilon$. Then the function u possesses an extension \tilde{u} to the metric space $(\tilde{\Omega}^0, \rho)$ and $\text{cap}(\tilde{\Omega}_\rho \setminus \tilde{\Omega}^0) = 0$ because $\Omega_{\varepsilon_1} \supset \Omega_{\varepsilon_2}$ if $\varepsilon_1 < \varepsilon_2$. Therefore $\tilde{u}|_{H_\rho}$ defined quasi-everywhere on H_ρ and represents the boundary value of the function $u \in L_2^1(\Omega)$ on the capacitary boundary H_ρ . \square

Remark 3.4. The function \tilde{u} defined quasi-everywhere on H_ρ in the following sense. For any $\varepsilon > 0$ there exists an open set $U_\varepsilon \subset \Omega$ such that the function u is uniformly continuous on $\Omega \setminus U_\varepsilon$, $\text{cap}(\Omega_\rho \setminus U_\varepsilon) < \varepsilon$, and the continuous extension of $\tilde{u} : \Omega \setminus U_\varepsilon \rightarrow \mathbb{R}$ to its completion $(\widetilde{\Omega \setminus U_\varepsilon}, \rho)$ coincides with \tilde{u} on $H_\rho \cap (\widetilde{\Omega \setminus U_\varepsilon}, \rho)$.

Combining the previous Theorem and Theorem 2.20 we obtain immediately

Theorem 3.5. *Let a domain $\Omega \subset \mathbb{R}^2$ possesses the strong Luzin capacitary property and be locally connected at any boundary point. Then for any function $u \in L_2^1(\Omega)$ there exists a quasicontinuous function $\tilde{u} : \tilde{\Omega} \rightarrow \mathbb{R}$ defined quasieverywhere on $\partial\Omega$ such that $\tilde{u}|_\Omega = u$.*

The strong Luzin capacity property is valid for the large class of domains, namely extension domains. The class of extension domains includes domains with smooth or Lipschitz boundaries (see for example [23]).

Definition 3.6. A domain $\Omega \subset \mathbb{R}^2$ is said to be a Sobolev L_2^1 -extension domain if there exists a bounded linear operator $E : L_2^1(\Omega) \rightarrow L_2^1(\mathbb{R}^n)$ such that for any $u \in L_2^1(\Omega)$ the condition $E(u)|_\Omega = u$ holds.

We call the operator E an extension operator. It is known that a simply connected domain $\Omega \subset \mathbb{R}^2$ is a L_2^1 -extension domain if and only if Ω is a quasidisc [9].

Recall that a domain $\Omega \subset \mathbb{R}^2$ is called a quasidisc if there exists a quasiconformal homeomorphism $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $\Omega = \varphi(\mathbb{D})$

Theorem 3.7. *If a bounded domain $\Omega \subset \mathbb{R}^2$ is a L_2^1 -extension domain then the identity mapping $id : H_\rho \rightarrow \partial\Omega$ is a homeomorphism.*

Proof. Because Ω is a L_2^1 -extension domain then there exists an extension operator

$$E : L_2^1(\Omega) \rightarrow L_2^1(\mathbb{R}^n)$$

such that for any $u \in L_2^1(\Omega)$ we have $E(u)|_\Omega = u$. Hence

$$\frac{1}{\|E\|} \|E(u)\|_{L_2^1(\mathbb{R}^n)} \leq \|u\|_{L_2^1(\Omega)} \leq \|E(u)\|_{L_2^1(\mathbb{R}^n)}.$$

By the definition of the conformal capacity for any condenser $(F_0, F_1; \Omega)$ the following inequality

$$\frac{1}{\|E\|^2} \text{cap}(F_0, F_1; \mathbb{R}^n) \leq \text{cap}(F_0, F_1; \Omega) \leq \text{cap}(F_0, F_1; \mathbb{R}^n)$$

holds

So, by the definition of the conformal capacity metric for any points $x, y \in \Omega$ and any pair (F, V) from the previous inequality follows

$$\frac{1}{\|E\|^2} \hat{\rho}_{(F,V)}(x, y) \leq \rho_{(F,V)}(x, y) \leq \hat{\rho}_{(F,V)}(x, y)$$

where $\hat{\rho}_{(F,V)}(x, y)$ is the conformal capacity metric in \mathbb{R}^2 and $\rho_{(F,V)}(x, y)$ is the conformal capacity metric in Ω . It means that the metric $\rho_{(F,V)}(x, y)$ is equivalent to the metric $\hat{\rho}_{(F,V)}(x, y)$ on Ω . By Theorem 2.7 the topology induced by the metric $\hat{\rho}_{(F,V)}(x, y)$ on \mathbb{R}^2 coincides with the Euclidean topology and so the topology of H_ρ coincides with the Euclidean topology of $\partial\Omega$. Because metrics $\rho_{(F,V)}(x, y)$ and $\hat{\rho}_{(F,V)}(x, y)$ are equivalent on Ω the theorem proved. \square

Theorem 3.8. *Let $\Omega \subset \mathbb{R}^2$ be a bounded L_2^1 -extension domain. Then Ω possesses the strong Luzin capacity property.*

Proof. Choose arbitrarily a function $u \in L_2^1$. Because Ω is an extension domain there exists an extension $\hat{u} \in L_2^1(\mathbb{R}^2)$ of u . By the Theorem 3.1 for any $\varepsilon > 0$ there exist such open set $U_\varepsilon \in \mathbb{R}^2$ of conformal capacity less than ε such that the function \hat{u} is continuous on $\mathbb{R}^2 \setminus U_\varepsilon$. Because the domain Ω is bounded the function $\hat{u}|_{\Omega \setminus U_\varepsilon}$ is uniformly continuous. Hence the function u is uniformly continuous on $\Omega \setminus U_\varepsilon$. By monotonicity of conformal capacity $\text{cap}(U_\varepsilon \cap \Omega) < \text{cap}(U_\varepsilon) < \varepsilon$. By the previous Theorem 3.7 the function u is uniformly continuous for p -capacity metric in $\Omega \setminus U_\varepsilon$ also. \square

Combining Theorem 3.3, Theorem 3.8 and 3.7 we obtain

Theorem 3.9. *Let a domain $\Omega \subset \mathbb{R}^2$ be a bounded L_2^1 -extension domain. Then for any function $u \in L_2^1(\Omega)$ there exists a quasicontinuous function $\tilde{u} : \overline{\Omega} \rightarrow \mathbb{R}$ defined quasieverywhere on $\partial\Omega$ such that $\tilde{u}|_\Omega = u$.*

Theorem 3.3, Theorem 3.8 and 3.7 can be easily extended to a more flexible class of so-called quasi-extension domains:

Definition 3.10. A domain $\Omega \subset \mathbb{R}^2$ is said to be a Sobolev L_2^1 -quasi-extension domain if for any $\varepsilon > 0$ there exist such open set U_ε of conformal capacity less than ε that $\Omega \setminus \overline{U_\varepsilon}$ is a L_2^1 -extension domain.

Typical examples of such domains are domains with boundary singularities of conformal capacity zero.

Theorem 3.11. *If a bounded domain Ω is a L_2^1 quasi-extension domain then the identity mapping $id : \partial\Omega \rightarrow H_\rho$ is a homeomorphism.*

Proof. Follows directly from Theorem 3.7 and the countable subadditivity of capacity. \square

Theorem 3.12. *Let $\Omega \subset \mathbb{R}^2$ be a bounded L_2^1 quasi-extension domain. Then Ω possesses the strong Luzin capacitary property.*

Proof. Follows directly from Theorem 3.7 and the countable subadditivity of capacity. \square

Theorem 3.13. *Let a domain $\Omega \subset \mathbb{R}^2$ be a bounded L_2^1 quasi-extension domain. Then for any function $u \in L_2^1(\Omega)$ there exists a quasicontinuous function $\tilde{u} : \overline{\Omega} \rightarrow \mathbb{R}$ defined quasieverywhere on $\partial\Omega$ such that $\tilde{u}|_\Omega = u$.*

4. BOUNDARY VALUES OF SOBOLEV FUNCTIONS FOR SIMPLY CONNECTED DOMAINS

Using the Riemann Mapping Theorem we prove that any simply connected domain possess the strong Luzin capacitary property, that permits to extend main results to any simply connected domain.

The unit disk $\mathbb{D}(0,1) \subset \mathbb{R}^2$ is the L_2^1 -extension domain and possess the strong Luzin capacitary property. Remember that the conformal capacity of condensors is a quasi-invariant for quasiconformal homeomorphisms $\varphi : \Omega \rightarrow \Omega'$ between two plane domains Ω and Ω' . Hence the conformal capacitary metric is also a quasi-invariant for quasiconformal homeomorphisms. Moreover, from this remark immediately follows

Proposition 4.1. ([14]) *Any quasiconformal homeomorphism $\varphi : \Omega \rightarrow \Omega'$ between two plane domains Ω and Ω' induces a quasi-isometry of $\tilde{\Omega}_\rho$ and $\tilde{\Omega}'_\rho$.*

Corollary 4.2. *Let $\varphi : \mathbb{D} \rightarrow \Omega$ be a quasiconformal homeomorphism of the unit disc \mathbb{D} onto a domain $\Omega \in \mathbb{R}^2$. Then Ω possesses the strong Luzin capacitary property.*

Proof. Choose a function $u \in L_2^1(\Omega)$. Because $\varphi : \mathbb{D} \rightarrow \Omega$ is a quasiconformal homeomorphism, then the composition $v := u \circ \varphi$ belongs to $L_2^1(\mathbb{D})$ (see, for example [10]). Because \mathbb{D} possesses the Luzin capacitary property, then for any $\varepsilon > 0$ there exist an open set V_ε of the conformal capacity less than ε such that $v|_{\mathbb{D} \setminus V_\varepsilon}$ is

uniformly continuous. The conformal capacity is a quasiinvariant for a quasiconformal homeomorphism φ . It means that there exist a constant Q which depends only on the quasiconformal distortion of φ and such that the conformal capacity of $U_\varepsilon := \varphi(V_\varepsilon)$ is less than $Q\varepsilon$. By the previous proposition φ^{-1} induces a quasisisometry of $\tilde{\Omega}_\rho$ and \tilde{D}_ρ . Therefore $u = v \circ \varphi^{-1}$ is uniformly continuous on $G \setminus U_\varepsilon$. We proved that Ω possesses the strong Luzin capacity property \square

From the previous proposition and Theorem 3.3 immediately follows

Theorem 4.3. *Let $\Omega \subset \mathbb{R}^2$ be a simply connected domain, $\Omega \neq \mathbb{R}^2$. Then for any function $u \in L_2^1(\Omega)$ there exists a quasicontinuous function $\tilde{u} : \tilde{\Omega}_\rho \rightarrow \mathbb{R}$ defined quasi-everywhere on the capacity boundary H_ρ such that $\tilde{u}|_\Omega = u$.*

And its version for simply connected domains locally connected at any boundary point

Theorem 4.4. *Let $\Omega \subset \mathbb{R}^2$, $\Omega \neq \mathbb{R}^2$ be a simply connected domain locally connected at any boundary point. Then for any function $u \in L_2^1(\Omega)$ there exists a quasicontinuous function $\tilde{u} : \bar{\Omega} \rightarrow \mathbb{R}$ defined quasi-everywhere on the boundary $\partial\Omega$ such that $\tilde{u}|_\Omega = u$.*

For readers convenience we repeat some basic facts about quasidisks.

Definition 4.5. A domain Ω is called a K -quasidisk if it is an image of the unit disc \mathbb{D} of a K -quasiconformal homeomorphism of the plane onto itself.

It is well known that the boundary of any K -quasidisk Ω admits a K^2 -quasiconformal reflections and thus, for example, any conformal homeomorphism $\varphi : \mathbb{D} \rightarrow \Omega$ can be extended to a K^2 -quasiconformal homeomorphism of the whole plane to itself.

Boundaries of quasidisk are called quasicircles. It is known that there are quasicircles for which no segment has finite length. The Hausdorff dimension of quasicircles was first investigated by Gehring and Vaisala (1973) [13], who proved that it can take all values in the interval $[1, 2)$. S. Smirnov proved recently [35] that the Hausdorff dimension of any K -quasicircle is at most $1 + k^2$, where $k = (K - 1)/(K + 1)$.

Ahlfors's 3-point condition [1] gives a complete geometric characterization: a Jordan curve γ in the plane is a quasicircle if and only if for each two points a, b on γ the (smaller) arc between them has diameter comparable to $|a - b|$. This condition is easily checked for the snowflake. On the other hand, every quasicircle can be obtained by an explicit snowflake-type construction (see [33]).

Because any quasidisk is an L_2^1 -extension domain we can reformulate previous results in the terms of quasidisks.

Proposition 4.6. *Let a domain $\Omega \subset \mathbb{R}^2$ is a quasidisk. Then the identity mapping $id : H_\rho \rightarrow \partial\Omega$ is a homeomorphism.*

Proposition 4.7. *Let a domain $\Omega \subset \mathbb{R}^2$ is a quasidisk. Then Ω possesses the strong Luzin capacity property.*

5. HISTORICAL SKETCH AND CONCLUSIONS

The concept of the ideal boundaries is common for the geometry and the analysis. The Poincaré disc is a model of the hyperbolic plane that provides a geometrical realization of the ideal boundary of the hyperbolic plane with help of a conformal homeomorphism.

By the Riemann Mapping Theorem any simply connected plane domain $\Omega \neq \mathbb{R}^2$ is conformally equivalent to the unit disc. However the boundary behavior of plane conformal homeomorphisms can not be described in terms of Euclidean boundaries but it can be described in terms of ideal boundary elements (prime ends) that was introduced by C. Caratheodory. By the Caratheodory Theorem any conformal homeomorphism $\varphi : \mathbb{D} \rightarrow \Omega$ induces one to one correspondence of prime ends.

M. A. Lavrentiev [19] introduced a metric (a relative distance) for prime ends. G. D. Suvorov [36] constructed a counterexample that demonstrates an absence of the triangle inequality for the Lavrentiev relative distance and proposed a more accurate concept of relative distance that support the triangle inequality. In terms of this metric the Caratheodory prime ends are a geometric representation of "ideal" compactification "boundary points". There exists a number of different conformally invariant intrinsic metrics. A detailed survey can be found in the paper of V. M. Miklyukov [29].

For dimension more than two by the Liouville theorem the class of conformal homeomorphisms coincides with the Mobius transformations. Even for quasiconformal homeomorphisms nothing similar to the Riemann mapping theorem is not correct.

By our opinion two main constructions of a quasiconformally invariant "ideal" boundary were proposed. The first one was in the spirit of Banach algebras. Recall that Royden algebra $\mathbb{R}(\Omega)$ is a quasiconformal invariant by M. Nakai [30] for dimension two and by L. G. Lewis [20] for arbitrary dimension. As any Banach algebra the Royden algebra produces a compactification of Ω and any quasiconformal homeomorphism induces a homeomorphism of such compactifications.

The second one is so-called capacity boundary proposed by V. Gol'dshtein and S. K. Vodop'janov [14]. Its construction is based on a notion of the conformal capacity. Remember that the conformal capacity is a quasiinvariant of quasiconformal homeomorphisms. By [14] quasiconformal homeomorphism can be extended to a homeomorphism of domains with capacity boundaries.

The Royden compactification does not coincide with the Caratheodory compactification. The "ideal" elements of the capacity boundary are Caratheodory prime ends.

Necessary and sufficient condition for existence of continuous traces of $L_p^1(\Omega)$, $p > 2$ were obtained by Shvartsman [34] in terms of quasi-hyperbolic metrics.

REFERENCES

- [1] L. V. Ahlfors. Quasiconformal reflections, Acta Math., 109, (1963), 291-301,
- [2] N. Aronszajn, P. Szeptycki, Theory of Bessel potentials. IV, Ann. Inst. Fourier 25 (1975), 27-69.
- [3] O. V. Besov, The behavior of differentiable functions on a non-smooth surface. Studies in the theory of differentiable functions of several variables and its applications, IV, Tr. Mat. Inst. Steklova 117 (1972), 3-10.
- [4] C. Caratheodory, Über die Begrenzung einfach zusammenhangender Gebiete, Math. Ann. 73 (1913), 323-370.
- [5] D. B. A. Epstein, Prime ends, Proc. London Math. Soc. 42 (1981), 385-414.
- [6] E. Gagliardo, Caratterizzazioni delle tracce sulla frontiera relative ad alcune classi di funzioni in n -variabili, Rend. Semin. Mat. Univ. Padova 27 (1957), 284-305.
- [7] P. Grisvard, Elliptic Problems in Non-Smooth Domains, Monogr. Stud. Math., Pitman, Boston (1985)
- [8] V. Gol'dshtein, L. Gurov and A. S. Romanov, Homeomorphisms that induce monomorphisms of Sobolev spaces, Israel J. Math. 91 (1995), 31-60.

- [9] V.M. Gol'dshtein, S.K. Vodop'janov. (1980). Prolongement des fonctions de classe L_2^1 et applications quasiconformes. C.R.Ac.Sc., Paris, 290, 10, A453-456.
- [10] V. Gol'dshtein, Yu. G. Reshetnyak, Quasiconformal mappings and Sobolev spaces, Kluwer Academic Publishers Group, Dordrecht (1990).
- [11] V. Gol'dshtein, A. Ukhlov, Sobolev homeomorphisms and composition operators, Sobolev spaces in mathematics. Int. Math. Ser (N. Y.) 11 (2010).
- [12] V. Gol'dshtein, M. Ju. Vasilchik, Embedding theorems and boundary-value problems for cusp domains, Trans. Amer. Math. Soc. 362 (2010), 1963–1979.
- [13] F. W. Gehring, J. Väisälä, Hausdorff dimension and quasiconformal mappings, Journal of the London Mathematical Society 6 (1973), 504-512.
- [14] V. Gol'dshtein, S. K. Vodop'yanov, Metric completion of a domain by means of a conformal capacity that is invariant under quasiconformal mappings, Dokl. Akad. Nauk SSSR. 238 (1978), 1040–1042.
- [15] J. Heinonen, T. Kilpeläinen, O. Martio, Nonlinear Potential Theory of Degenerate Elliptic Equations, Oxford Math. Monographs, Oxford Univ. Press, 1993.
- [16] A. Jonsson, The trace of potentials on general sets, Ark. Mat. 17 (1979), 1–18.
- [17] A. Jonsson, Besov spaces on surfaces with singularities, Manuscripta Math. 71 (1991), 121–152.
- [18] A. Jonsson, H. Wallin, Function Spaces on Subsets of \mathbb{R}^n , Math. Rep., vol. 2, part 1, Harwood Acad. Publ., London, (1984).
- [19] M. A. Lavrentiev, On continuity of univalent functions in closed domains, C. R. (Doklady) Acad. Sci. URSS (N. S.) 4 (1936), 207–210.
- [20] L. G. Lewis, Quasiconformal mappings and Royden algebras in space, Trans. Amer. Math. Soc. 158 (1971), 481–492.
- [21] J. L. Lions, E. Magenes, Non-Homogeneous Boundary Value Problems and Applications, Springer-Verlag, Berlin, New York, (1972).
- [22] J. Maly, Hölder type quasicontinuity, Potential Analysis 2 (1993), 249–254.
- [23] V. G. Maz'ya, Sobolev Spaces, Springer-Verlag, Berlin, (1985).
- [24] V. G. Maz'ya, S. Poborchi, Traces of functions from S.L. Sobolev spaces on small and large components of the boundary, Math. Notes 45 (1989), 312–317.
- [25] V. G. Maz'ya, S. Poborchi, Traces of functions with a summable gradient in a domain with a cusp at the boundary, Math. Notes 45 (1989), 39–44.
- [26] V. G. Maz'ya, S. Poborchi, Boundary traces of functions from Sobolev spaces on a domain with a cusp, Siberian Adv. Math. 1 (1991), 75–107.
- [27] V. G. Maz'ya, S. Poborchi, Yu. Netrusov, Boundary values of functions from Sobolev spaces in some non-Lipschitzian domains, Algebra i Analiz 11 (1999), 141–170.
- [28] V. G. Maz'ya, S. Poborchi, Differentiable Functions on Bad Domains, World Scientific, River Edge, NJ, (1997).
- [29] V. M. Miklyukov, The relative distance of M. A. Lavrent'ev and prime ends on nonparametric surfaces, Ukr. Math. Bull 1 (2004), 353–376.
- [30] M. Nakai, Algebraic criterion on quasiconformal equivalence of Riemann surfaces, Nagoya Math. J. 16 (1960), 157–184.
- [31] S. M. Nikol'skii, Boundary properties of functions defined on a region with angular points, Amer. Math. Soc. Transl. Ser. 2 83 (1958), 101–158.
- [32] S. Poborchi, Continuity of the boundary trace operator $W_p^1(\partial\Omega) \rightarrow L_Q(\partial\Omega)$ for a domain with outward peak, Vestnik St. Petersburg Univ. Math. 38 (2005), 37–44.
- [33] S. Rohde, Quasircles modulo bilipschitz maps, Rev. Mat. Iberoamericana, 17(3), (2001), 643-659.
- [34] P. Shvartsman, On the boundary values of Sobolev W_p^1 -functions, Adv. in Math. 225 (2010), 2162–2221.
- [35] S. Smirnov, Dimension of quasircles, Acta Math., 205(1), (2010), 189-197.
- [36] G. D. Suvorov, Remarks on a theorem of Lavrent'ev, Tomskii Gos. Univ. Uc. Zap. Mat. Meh. 25 (1956), 3–8.
- [37] M. Yu. Vasil'chik, The boundary behavior of functions of Sobolev spaces defined on a planar domain with a peak vertex on the boundary, Siberian Adv. Math. 14 (2004), 92–115.
- [38] J. Väisälä J., Lectures on n -dimensional quasiconformal mappings, Springer-Verlag (Berlin, 1971).

- [39] S. K. Vodop'yanov, V. M. Gol'dshtein, Yu. G. Reshetnyak, On geometric properties of functions with generalized first derivatives, *Uspekhi Mat. Nauk* 34 (1979), 17–65.
- [40] G. N. Yakovlev, Boundary properties of functions of the class $W_p^{(l)}$ in regions with corners, *Soviet Math.* 2 (1961), 1177–1180.
- [41] G. N. Yakovlev, Dirichlet problem for a region with a non-Lipschitz boundary, *Differ. Equ.* 1 (1965), 847–858.