

KR-THEORY OF COMPACT LIE GROUPS WITH GROUP ANTI-INVOLUTIONS

CHI-KWONG FOK

ABSTRACT. Let G be a compact, connected, and simply-connected Lie group, equipped with an anti-involution a_G which is the composition of a Lie group involutive automorphism σ_G and the group inversion. We view (G, a_G) as a Real (G, σ_G) -space via the conjugation action. In this note, we exploit the notion of Real equivariant formality discussed in [Fo] to compute the ring structure of the equivariant KR -theory of G . In particular, we show that when G does not have Real representations of complex type, the equivariant KR -theory is the ring of Grothendieck differentials of the coefficient ring of equivariant KR -theory over the coefficient ring of ordinary KR -theory, thereby generalizing a result of Brylinski-Zhang's ([BZ]) for the complex K -theory case.

Keywords: KR -theory, compact Lie group, Real equivariant formality, Real representation ring

2010 Mathematics Subject Classification: 19L47; 57T10

1. INTRODUCTION

Let G be a compact, connected and simply-connected Lie group, viewed as a G -space via the conjugation action. According to the main result of [BZ], the equivariant K -theory ring $K_G^*(G)$ is isomorphic to $\Omega_{R(G)/\mathbb{Z}}$, the ring of Grothendieck differentials of the complex representation ring of G over the integers (in fact, Brylinski-Zhang proved that this is true for $\pi_1(G)$ being torsion-free). Assuming further that G is equipped with an involutive automorphism σ_G , the author gave in [Fo] an explicit description of the ring structure of the equivariant KR -theory (cf. [At2], [At3] and [AS] for definition of KR -theory) $KR_G^*(G)$ by drawing on Brylinski-Zhang's result, Seymour's result on the module structure of $KR^*(G)$ (cf. [Se]) and the notion of Real equivariant formality. $KR_G^*(G)$ in general has far more complicated ring structure and, among other things, is not a ring of Grothendieck differentials, as one would expect from Brylinski-Zhang's theorem. This is because in general the algebra generators of the equivariant KR -theory ring do not simply square to 0.

In this note, we equip G instead with an anti-involution $a_G := \sigma_G \circ \text{inv}$. Denoting the (G, σ_G) -space (G, a_G) by G^- for brevity, we compute the ring structure of $KR_G^*(G^-)$ following the idea of [Fo]. We find that there exists a derivation of the graded ring $KR_G^0(\text{pt}) \oplus KR_G^{-4}(\text{pt})$ taking values in $KR_G^1(G^-) \oplus KR_G^{-3}(G^-)$ (cf. Proposition 3.5) and

Date: August 16, 2015.

that any element in the image of the derivation squares to 0 (see Propositions 3.3, 3.5 and 4.8(1), and compare with [Fo, Theorem 4.30, Proposition 4.31]). In particular,

Theorem 1.1. *If G does not have any Real representation of complex type with respect to σ_G , then the derivation in Proposition 3.5 induces the following ring isomorphism*

$$KR_G^*(G^-) \cong \Omega_{KR_G^*(pt)/KR^*(pt)}$$

Hence an anti-involution is the ‘right’ involution needed to generalize Brylinski-Zhang’s result in the context of KR -theory. As a by-product, we also obtain the following

Corollary 1.2. *If G is a compact connected Real Lie group (not necessarily simply-connected) and X a compact Real G -space, then for any x in $KR_G^1(X)$ or $KR_G^{-3}(X)$, $x^2 = 0$.*

Note that graded commutativity only implies that x^2 is 2-torsion.

Throughout this note, G is a compact, connected and simply-connected Lie group unless otherwise specified. We sometimes omit the notation for the involution when it is clear from the context that a Real structure is implicitly assumed.

2. BACKGROUND

In this section, we recall some relevant definitions and results from [BZ] and [Fo] needed in this note. We refer the reader to [At2], [At3] and [AS] for the basic definition of (equivariant) KR -theory, which we shall omit here.

Definition 2.1. Let G be a compact Lie group equipped with an involutive automorphism σ_G , i.e. a Real compact Lie group, and X a finite CW -complex equipped with an involution.

- (1) (cf. [Fo, Proposition 2.29]) Let $c : KR_G^*(X) \rightarrow K_G^*(X)$ be the *complexification map* which forgets the Real structure of Real vector bundles, and $r : K_G^*(X) \rightarrow KR_G^*(X)$ be the *realification map* defined by

$$[E] \mapsto [E \oplus \sigma_X^* \sigma_G^* \overline{E}]$$

where σ_G^* means twisting the original G -action on E by σ_G .

- (2) (cf. [Fo, Definitions 2.1 and 2.5]) Let $\delta : R(G) \rightarrow K^{-1}(G)$ be the derivation of $R(G)$ taking values in the $R(G)$ -module $K^{-1}(G)$ (the module structure is realized by the augmentation homomorphism), where $\delta(\rho)$ is represented by the complex of vector bundles

$$\begin{aligned} 0 \rightarrow G \times \mathbb{R} \times V \rightarrow G \times \mathbb{R} \times V \rightarrow 0, \\ (g, t, v) \mapsto (g, t, -t\rho(g)v) \text{ if } t \geq 0, \\ (g, t, v) \mapsto (g, t, tv) \text{ if } t \leq 0. \end{aligned}$$

We define $\delta_G : R(G) \rightarrow K_G^{-1}(G)$ similarly. $\delta_G(\rho)$ is represented by the same complex of vector bundles where G acts on $G \times \mathbb{R} \times V$ by $g_0 \cdot (g_1, t, v) = (g_0 g_1 g_0^{-1}, t, \rho_V(g_0)v)$.

- (3) Let σ_n be the class of the standard representation of $U(n)$ in $R(U(n))$. Let T be the standard maximal torus of $U(n)$. Let $\sigma_{\mathbb{R}}$ be the complex conjugation on $U(n)$, T , $U(n)/T$ or $U(\infty)$. Let $\sigma_{\mathbb{H}}$ be the symplectic type involution on $U(2m)$ (given

- by $g \mapsto J_m \bar{g} J_m^{-1}$, $U(2\infty)$ or the involution $gT \mapsto J_m \bar{g} T$ on $U(2m)/T$. Let $a_{\mathbb{R}}$ and $a_{\mathbb{H}}$ be the corresponding anti-involutions on the unitary groups.
- (4) (cf. [Fo, Definitions 2.9 and 2.20]) A Real representation V of G is a finite-dimensional complex representation of G equipped with an anti-linear involution σ_V such that $\sigma_V(g \cdot v) = \sigma_G(g) \cdot \sigma_V(v)$. Similarly a Quaternionic representation is one equipped with an anti-linear endomorphism J_V such that $J_V^2 = -\text{Id}_V$ and $J_V(g \cdot v) = \sigma_G(g) \cdot J_V(v)$. For $\mathbb{F} = \mathbb{R}$ or \mathbb{H} , a morphism between V and $W \in \mathcal{R}ep_{\mathbb{F}}(G)$ is a linear transformation from V to W which commutes with G and respects both σ_V and σ_W . Let $\mathcal{R}ep_{\mathbb{R}}(G)$ (resp. $\mathcal{R}ep_{\mathbb{H}}(G)$) be the category of Real (resp. Quaternionic) representations of G . The Real (resp. Quaternionic) representation group of G , denoted by $RR(G)$ (resp. $RH(G)$) is the Grothendieck group of $\mathcal{R}ep_{\mathbb{R}}(G)$ (resp. $\mathcal{R}ep_{\mathbb{H}}(G)$).
- (5) (cf. [Fo, Definitions 2.11 and 2.20]) Let V be an irreducible Real (resp. Quaternionic) representation of G . Its *commuting field* is defined to be $\text{Hom}_G(V, V)^{\sigma_V}$, which is isomorphic to either \mathbb{R} , \mathbb{C} or \mathbb{H} . Let $RR(G, \mathbb{F})$ (resp. $RH(G, \mathbb{F})$) be the abelian group generated by the isomorphism classes of irreducible Real (resp. Quaternionic) representations with \mathbb{F} as the commuting field.
- (6) (cf. [Fo, Definition 2.19]) Let $R(G, \mathbb{C})$ be the abelian group generated by the isomorphism classes of those irreducible complex representations V satisfying $V \not\cong \sigma_G^* \bar{V}$.

Theorem 2.2. *Let G be a compact connected Lie group with torsion-free fundamental group.*

- (1) [Ho] $K^*(G)$ is a free \mathbb{Z} -module. If P is the image of the map $\delta : R(G) \rightarrow K^{-1}(G)$, then

$$K^*(G) \cong \bigwedge_{\mathbb{Z}}^* P$$

- (2) [BZ] Let $\varphi : \Omega_{R(G)/\mathbb{Z}} \rightarrow K_G^*(G)$ be the $R(G)$ -algebra homomorphism defined by the following

- (a) $\varphi(\rho_V) := [G \times V] \in K_G^*(G)$, where G acts on $G \times V$ by $g_0 \cdot (g_1, v) = (g_0 g_1 g_0^{-1}, \rho_V(g_0)v)$,
and
(b) $\varphi(d\rho_V) = \delta_G(\rho)$.

Then φ is an $R(G)$ -algebra isomorphism.

Definition 2.3. (1) ([HL, Definition 4.1]) A G -space X is *weakly equivariantly formal* if the map $K_G^*(X) \otimes_{R(G)} \mathbb{Z} \rightarrow K^*(X)$ induced by the forgetful map is a ring isomorphism, where \mathbb{Z} is viewed as an $R(G)$ -module through the augmentation homomorphism.

- (2) ([Fo, Definition 4.2]) A Real G -space X is a *Real equivariantly formal* space if
- G is a Real compact Lie group,
 - X is a weakly equivariantly formal G -space, and
 - the forgetful map $KR_G^*(X) \rightarrow KR^*(X)$ admits a section $s_R : KR^*(X) \rightarrow KR_G^*(X)$ which is a $KR^*(\text{pt})$ -module homomorphism.
- (3) ([Fo, Definition 4.4]) For a section $s : K^*(X) \rightarrow K_G^*(X)$ (resp. $s_R : KR^*(X) \rightarrow KR_G^*(X)$) and $a \in K^*(X)$ (resp. $a \in KR^*(X)$), we call $s(a)$ (resp. $s_R(a)$) a (*Real*) *equivariant lift* of a , with respect to s (resp. s_R).

Remark 2.4. By Theorem 2.2 and noting that the forgetful map takes $\delta_G(\rho) \in K_G^{-1}(G)$ to $\delta(\rho) \in K^{-1}(G)$, we have that G is weakly equivariantly formal if it is a compact connected Lie group with torsion-free fundamental group.

The following result is a structure theorem for the equivariant KR -theory of Real equivariantly formal spaces.

Theorem 2.5. ([Fo, Theorem 4.5]) *Let X be a Real equivariantly formal space. For any element $a \in K^*(X)$ (resp. $a \in KR^*(X)$), let $a_G \in K_G^*(X)$ (resp. $a_G \in KR_G^*(X)$) be a (Real) equivariant lift of a with respect to a group homomorphic section s (resp. s_R which is a $KR^*(pt)$ -module homomorphism). Then the map*

$$f : (RR(G, \mathbb{R}) \oplus RH(G, \mathbb{R})) \otimes KR^*(X) \oplus r(R(G, \mathbb{C}) \otimes K^*(X)) \rightarrow KR_G^*(X),$$

$$\rho_1 \otimes a_1 \oplus r(\rho_2 \otimes a_2) \mapsto \rho_1 \cdot (a_1)_G \oplus r(\rho_2 \cdot (a_2)_G).$$

is a group isomorphism. In particular, if $R(G, \mathbb{C}) = 0$, then f is a $KR_G^*(pt)$ -module isomorphism.

3. A PRELIMINARY DESCRIPTION OF $KR_G^*(G^-)$

Lemma 3.1. *Let X be a finite CW-complex equipped with an involution. We have that*

$$KR^1(X) \cong [X, (U(\infty), a_{\mathbb{R}})]_{\mathbb{R}}$$

$$KR^{-3}(X) \cong [X, (U(2\infty), a_{\mathbb{H}})]_{\mathbb{R}}$$

where $[X, Y]_{\mathbb{R}}$ means the set of Real homotopy equivalence classes of Real maps from X to Y . Here Real homotopy equivalence is the one witnessed by a family of Real maps.

Proof. The representing space for KR^{-q} is a Real space which is homeomorphic to the representing space for K^{-q} and whose subspace fixed under the Real structure is homeomorphic to that for KO^{-q} . By the definition of Real vector bundles and KR -theory, and noting that the representing spaces for K -theory and KO -theory are homogeneous spaces constructed out of various infinite dimensional matrix groups (orthogonal, unitary, and symplectic), the representing spaces for KR -theory are obtained from those for KO -theory by ‘complexifying’ the relevant matrix groups (to be distinguished from the usual complexification of Lie groups). For example, $O(\infty)$ is transformed to $(U(\infty), \sigma_{\mathbb{R}})$ (analogous to the fact that \mathbb{R} is complexified to \mathbb{C}), $Sp(\infty)$ to $(U(2\infty), \sigma_{\mathbb{H}})$ (analogous to the fact that $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{C} \cong M_2(\mathbb{C})$), and $U(\infty)$ to $(U(\infty) \times U(\infty), (g_1, g_2) \mapsto (\bar{g}_2, \bar{g}_1))$ (analogous to the fact that $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} \cong \mathbb{C} \oplus \mathbb{C}$). We shall show the case for KR^1 . The other case is similar. The representing space for KR^1 is $(U(\infty) \times U(\infty)/U(\infty)_{\Delta}, (g_1, g_2)U(\infty)_{\Delta} \mapsto (\bar{g}_2, \bar{g}_1)U(\infty)_{\Delta})$ (here Δ means diagonal subgroup), which is obtained from that for KO^1 , which in turn is $U(\infty)/O(\infty)$ (cf. [Bot]). We have that $(U(\infty), a_{\mathbb{R}})$ is Real diffeomorphic to $(U(\infty) \times U(\infty)/U(\infty)_{\Delta}, (g_1, g_2)U(\infty)_{\Delta} \mapsto (\bar{g}_2, \bar{g}_1)U(\infty)_{\Delta})$ by the map $g \mapsto (g, e)U(\infty)$. \square

Definition 3.2. Let $\delta_{\mathbb{R}}^{\text{inv}} : RR(G) \rightarrow KR^1(G^-)$ send ρ to the Real homotopy class of it, viewed as the Real map $G^- \rightarrow (U(\infty), a_{\mathbb{R}})$. Define $\delta_{\mathbb{H}}^{\text{inv}} : RH(G) \rightarrow KR^{-3}(G^-)$ similarly.

Proposition 3.3. *If ρ be in $RR(G)$ with (V, σ_V) being the underlying finite dimensional Real vector space of the Real unitary representation, then $\delta_{\mathbb{R}}(\rho)$ is represented by the following complex of Real vector bundles*

$$0 \longrightarrow G \times \mathbb{R} \times \mathbb{C} \times (V \oplus V) \longrightarrow G \times \mathbb{R} \times \mathbb{C} \times (V \oplus V) \longrightarrow 0$$

$$(g, t, z, v_1, v_2) \mapsto \begin{cases} \left(g, t, z, \begin{pmatrix} -t\rho(g) & \bar{z}I_V \\ zI_V & t\rho(g)^* \end{pmatrix} \begin{pmatrix} iv_1 \\ iv_2 \end{pmatrix} \right) & \text{if } t \geq 0 \\ \left(g, t, z, \begin{pmatrix} tI_V & \bar{z}I_V \\ zI_V & -tI_V \end{pmatrix} \begin{pmatrix} iv_1 \\ iv_2 \end{pmatrix} \right) & \text{if } t \leq 0 \end{cases}$$

where the Real structure on $G \times \mathbb{R} \times \mathbb{C} \times (V \oplus V)$ is given by

$$(g, t, z, v_1, v_2) \mapsto (\sigma_G(g)^{-1}, t, -z, \sigma_V(v_2), \sigma_V(v_1))$$

Similarly, if $\rho \in RH(G)$ with (V, J_V) being the underlying finite dimensional Quaternionic vector space of the Quaternionic unitary representation, then $\delta_{\mathbb{H}}(\rho)$ is represented by the same complex of Real vector bundles except that the Real structure on $G \times \mathbb{R} \times \mathbb{C} \times (V \oplus V)$ is given by

$$(g, t, z, v_1, v_2) \mapsto (\sigma_G(g)^{-1}, t, z, -J_V(v_2), J_V(v_1))$$

Proof. It is straightforward to verify that the given Real structures indeed commute with the middle maps of the complex of vector bundles, and that they are canonical. The complex of vector bundles, with the Real structures forgotten, is the tensor product of the following two complexes

$$0 \longrightarrow G \times \mathbb{C} \times \mathbb{C} \longrightarrow G \times \mathbb{C} \times \mathbb{C} \longrightarrow 0$$

$$(g, z, w) \mapsto (g, z, izw)$$

$$0 \longrightarrow G \times \mathbb{R} \times V \longrightarrow G \times \mathbb{R} \times V \longrightarrow 0$$

$$(g, t, v) \mapsto \begin{cases} (g, t, -it\rho(g)v) & \text{if } t \geq 0 \\ (g, t, itv) & \text{if } t \leq 0 \end{cases}$$

(cf. [ABS, Proposition 10.4]) which represent the Bott class $\beta \in K^{-2}(G)$ and $\delta(\rho) \in K^{-1}(G)$ as defined in [BZ] respectively (the middle maps of the above two complexes differ from the ones conventionally used to define β and $\delta(\rho)$ by multiplication by i , which is homotopy equivalent to the identity map). Besides, the KR -theory classes represented by the complexes of Real vector bundles live in degree 1 and -3 pieces respectively because of the type of the involution of the middle maps restricted to $\mathbb{R} \times \mathbb{C}$. In sum, the two complexes of Real vector bundles represent canonical Real lifts of $\delta(\rho)$. Therefore they must represent $\delta_{\mathbb{R}}^{\text{inv}}(\rho)$ (resp. $\delta_{\mathbb{H}}^{\text{inv}}(\rho)$). \square

Definition 3.4. Let $\delta_{\mathbb{R}}^{G, \text{inv}} : RR(G) \rightarrow KR_G^1(G^-)$ send ρ to the complex of Real vector bundles as in Proposition 3.3 equipped with the equivariant structure given by

$$(g, t, z, v_1, v_2) \mapsto (\sigma_G(g)^{-1}, t, z, \rho(g)v_1, \rho(g)v_2)$$

Define $\delta_{\mathbb{H}}^{G, \text{inv}} : RH(G) \rightarrow KR_G^{-3}(G^-)$ similarly.

Proposition 3.5. *Identifying $RR(G)$ with $KR_G^0(\text{pt})$ and $RH(G)$ with $KR_G^{-4}(\text{pt})$ (cf. [AS, Sect. 8]), $\delta_{\mathbb{R}}^{G,\text{inv}} \oplus \delta_{\mathbb{H}}^{G,\text{inv}}$ is a derivation of the graded ring $KR_G^0(\text{pt}) \oplus KR_G^{-4}(\text{pt})$ taking values in $KR_G^1(G^-) \oplus KR_G^{-3}(G^-)$.*

Proof. The proof can be easily adapted from the one of [BZ, Proposition 3.1] by straightforwardly modifying the homotopy ρ_s and replacing the definition of the map δ_G given there (which is incorrect) with the one in [Fo, Definition 2.5]. The modified homotopy can be easily seen to intertwines with both Real structures of the complex of Real vector bundles as in Proposition 3.3. \square

Proposition 3.6. $\delta_G(\overline{a_G^*}\rho) = -\delta_G(\overline{\sigma_G^*}\rho)$

Proof. Viewing $\overline{\sigma_G^*}\rho$ and $\overline{a_G^*}\rho$ as maps from G to $U(\infty)$, $\overline{\sigma_G^*}\rho \cdot \overline{a_G^*}\rho$ is the constant map with image being the identity. It follows that

$$0 = \delta_G(\overline{\sigma_G^*}\rho \cdot \overline{a_G^*}\rho) = \delta_G(\overline{\sigma_G^*}\rho) + \delta_G(\overline{a_G^*}\rho)$$

The last equality is the equivariant analogue of [At, Lemma 2.4.6]. \square

The fundamental representations of G are permuted by $\overline{\sigma_G^*}$ (cf. [Se, Lemma 5.5]). Following the notations in [Se] and [Fo], we let $\varphi_1, \dots, \varphi_r, \theta_1, \dots, \theta_s, \gamma_1, \dots, \gamma_t, \overline{\sigma_G^*}\gamma_1, \dots, \overline{\sigma_G^*}\gamma_t$ be the fundamental representations of G , where $\varphi_i \in RR(G, \mathbb{R})$, $\theta_j \in RH(G, \mathbb{R})$ and $\gamma_k \in R(G, \mathbb{C})$.

Definition 3.7. Let λ_k^{inv} be the element in $KR^0(G^-)$ constructed as in the proof of [Se, Proposition 4.6] such that $c(\lambda_k^{\text{inv}}) = \beta^3 \delta(\gamma_k) \delta(\overline{a_G^*}\gamma_k) = -\beta^3 \delta(\gamma_k) \delta(\overline{\sigma_G^*}\gamma_k)$. Let $\lambda_k^{G,\text{inv}}$ be the Real equivariant lift of λ_k^{inv} in $KR_G^0(G^-)$ constructed by adding the natural equivariant structure throughout the construction of λ_k^{inv} such that $c(\lambda_k^{G,\text{inv}}) = \beta^3 \delta_G(\gamma_k) \delta_G(\overline{a_G^*}\gamma_k) = -\beta^3 \delta_G(\gamma_k) \delta_G(\overline{\sigma_G^*}\gamma_k)$.

Applying [Se, Theorem 4.2], one can get the $KR^*(\text{pt})$ -module structure of $KR^*(G^-)$, which is generated, as a $KR^*(\text{pt})$ -algebra, by $\delta_{\mathbb{R}}^{\text{inv}}(\varphi_1), \dots, \delta_{\mathbb{R}}^{\text{inv}}(\varphi_r), \delta_{\mathbb{H}}^{\text{inv}}(\theta_1), \dots, \delta_{\mathbb{H}}^{\text{inv}}(\theta_s), \lambda_1^{\text{inv}}, \dots, \lambda_t^{\text{inv}}$, and realifications of certain products of β^i and $\delta(\gamma_k)$, $k = 1, \dots, t$ (compare with [Se, Theorem 5.6]). Noting that all these generators admit Real equivariant lifts, and that G is a weakly equivariantly formal space by Remark 2.4, we have that G^- is a Real equivariantly formal space by Definition 2.3 (2). Now Theorem 2.5 applies and one can further obtain the $KR_G^*(\text{pt})$ -module structure of $KR_G^*(G^-)$. We shall state the following description of $KR_G^*(G^-)$ without proof. We refer the reader to [Fo, Corollaries 4.10, 4.11, Proposition 4.13, Theorem 4.33] for comparison.

Theorem 3.8. (1) *The map*

$$f : (RR(G, \mathbb{R}) \oplus RH(G, \mathbb{R})) \otimes KR^*(G^-) \oplus r(R(G, \mathbb{C}) \otimes K^*(G)) \rightarrow KR_G^*(G^-)$$

$$\rho_1 \otimes x_1 \oplus r(\rho_2 \otimes x_2) \mapsto \rho_1 \cdot (x_1)_G \oplus r(\rho_2 \cdot (x_2)_G)$$

is a group isomorphism, where $x_G \in KR_G^(G^-)$ is a Real equivariant lift of $x \in KR^*(G^-)$. If $R(G, \mathbb{C}) = 0$, then f is an isomorphism of $KR_G^*(\text{pt})$ -modules.*

(2) $KR_G^*(G^-)$ is generated as an algebra over $KR_G^*(pt)$ (for descriptions of the coefficient ring see [Fo, Section 3]) by $\delta_{\mathbb{R}}^{G,inv}(\varphi_1), \dots, \delta_{\mathbb{R}}^{G,inv}(\varphi_r), \delta_{\mathbb{H}}^{G,inv}(\theta_1), \dots, \delta_{\mathbb{H}}^{G,inv}(\theta_s), \lambda_1^{G,inv}, \dots, \lambda_t^{G,inv}$ and

$$\{r_{\rho,i,\varepsilon_1,\dots,\varepsilon_t,\nu_1,\dots,\nu_t}^{G,inv} := r(\beta^i \cdot \rho \delta_G(\gamma_1)^{\varepsilon_1} \dots \delta_G(\gamma_t)^{\varepsilon_t} \delta_G(\overline{a_G^*} \gamma_1)^{\nu_1} \dots \delta_G(\overline{a_G^*} \gamma_t)^{\nu_t})\}$$

where $\rho \in R(G, \mathbb{C}) \oplus \mathbb{Z} \cdot \rho_{triv}$, $\varepsilon_1, \dots, \varepsilon_t, \nu_1, \dots, \nu_t$ are either 0 or 1, ε_k and ν_k are not equal to 1 at the same time for $1 \leq k \leq t$, and the first index k_0 where $\varepsilon_{k_0} = 1$ is less than the first index k_1 where $\nu_{k_1} = 1$. Moreover,

(a) $(\lambda_k^{G,inv})^2 = 0$.

(b) Let $\omega_t := \delta_{\varepsilon_t, 1-\nu_t}$. Then

$$(r_{\rho,i,\varepsilon_1,\dots,\varepsilon_t,\nu_1,\dots,\nu_t}^{G,inv})^2 = \begin{cases} \eta^2(\rho \cdot \overline{\sigma_G^*} \rho)(\lambda_1^{G,inv})^{\omega_1} \dots (\lambda_t^{G,inv})^{\omega_t} & \text{if } r_{\rho,i,\varepsilon_1,\dots,\varepsilon_t,\nu_1,\dots,\nu_t}^{G,inv} \\ & \text{is of degree } -1 \text{ or } -5 \\ \pm \mu(\rho \cdot \overline{\sigma_G^*} \rho)(\lambda_1^{G,inv})^{\omega_1} \dots (\lambda_t^{G,inv})^{\omega_t} & \text{if } r_{\rho,i,\varepsilon_1,\dots,\varepsilon_t,\nu_1,\dots,\nu_t}^{G,inv} \\ & \text{is of degree } -2 \text{ or } -6 \\ 0 & \text{otherwise} \end{cases}$$

The sign can be determined using formulae in [Fo, Proposition 2.29 (2)].

(c) $r_{\rho,i,\varepsilon_1,\dots,\varepsilon_t,\nu_1,\dots,\nu_t}^{G,inv} \eta = 0$, and $r_{\rho,i,\varepsilon_1,\dots,\varepsilon_t,\nu_1,\dots,\nu_t}^{G,inv} \mu = 2r_{\rho,i+2,\varepsilon_1,\dots,\varepsilon_t,\nu_1,\dots,\nu_t}^{G,inv}$.

Corollary 3.9. *In particular, if $R(G, \mathbb{C}) = 0$, then*

$$\begin{aligned} KR_G^*(G^-) &= \bigwedge_{KR_G^*(pt)} (\delta_{\mathbb{R}}^{G,inv}(\varphi_1), \dots, \delta_{\mathbb{R}}^{G,inv}(\varphi_r), \delta_{\mathbb{H}}^{G,inv}(\theta_1), \dots, \delta_{\mathbb{H}}^{G,inv}(\theta_s)) \\ &\cong \Omega_{KR_G^*(pt)/KR^*(pt)} \end{aligned}$$

as $KR_G^*(pt)$ -modules.

As we can see, the module structure of $KR_G^*(G^-)$ is very similar to that of $KR_G^*(G)$, except that the degrees of the generators are different. Now it remains to find $\delta_{\mathbb{R}}^{G,inv}(\varphi_i)^2$ and $\delta_{\mathbb{H}}^{G,inv}(\theta_j)^2$ so as to complete the description of the ring structure of $KR_G^*(G^-)$. As it turns out, these squares are all zero, in stark contrast to the involutive automorphism case.

4. SQUARES OF THE REAL AND QUATERNIONIC TYPE GENERATORS

This section is devoted to proving that the squares of the real and quaternionic generators are zero, following the strategy outlined in [Fo, Section 4]. We use T to denote the maximal torus of $U(n)$ consisting of diagonal matrices throughout this section.

Applying Brylinski-Zhang's result on the equivariant K -theory of compact connected Lie group G with $\pi_1(G)$ torsion-free and Theorem 2.5, we have

Proposition 4.1. *For $\mathbb{F} = \mathbb{R}$ or \mathbb{H} , we have the following $KR_{(U(n), \sigma_{\mathbb{F}})}^*(pt)$ -module isomorphism*

$$KR_{(U(n), \sigma_{\mathbb{F}})}^*(U(n), a_{\mathbb{F}}) \cong \Omega_{KR_{(U(n), \sigma_{\mathbb{F}})}^*(pt)/KR^*(pt)}$$

The set $\{\delta_{\mathbb{R}}^{G,inv}(\sigma_n), \delta_{\mathbb{R}}^{G,inv}(\bigwedge^2 \sigma_n), \dots, \delta_{\mathbb{R}}^{G,inv}(\bigwedge^n \sigma_n)\}$ is a set of primitive generators for the case $\mathbb{F} = \mathbb{R}$, while $\{\delta_{\mathbb{H}}^{G,inv}(\sigma_{2m}), \delta_{\mathbb{R}}^{G,inv}(\bigwedge^2 \sigma_{2m}), \dots, \delta_{\mathbb{R}}^{G,inv}(\bigwedge^{2m} \sigma_{2m})\}$ is a set of primitive generators for the case $\mathbb{F} = \mathbb{H}$.

Corollary 4.2. *We have the following isomorphism*

$$KR_{(U(n), \sigma_{\mathbb{F}})}^*(U(n), a_{\mathbb{F}}) \cong \Omega_{R(U(n))/\mathbb{Z}} \otimes KR^*(pt)$$

as ungraded $KR^*(pt)$ -modules.

Definition 4.3. Let

$$p_{G,inv}^* : KR_{(U(n), \sigma_{\mathbb{R}})}^*(U(n), a_{\mathbb{R}}) \rightarrow KR_{(T, \sigma_{\mathbb{R}})}^*(T, Id)$$

be the restriction map and the map

$$q_{G,inv}^* : KR_{(U(2m), \sigma_{\mathbb{H}})}^*(U(2m), a_{\mathbb{H}}) \rightarrow KR_{(U(2m)/T \times T, \sigma_{\mathbb{H}})}^*(U(2m)/T \times T, \sigma_{\mathbb{H}} \times Id)$$

induced by the Weyl covering map

$$\begin{aligned} q_G : U(2m)/T \times T &\rightarrow U(2m) \\ (gT, t) &\mapsto gtg^{-1} \end{aligned}$$

Proposition 4.4. *Identifying $KR_{(T, \sigma_{\mathbb{R}})}^*(T, Id)$ with $RR(T, \sigma_{\mathbb{R}}) \otimes KR^*(T, Id)$, we have*

$$p_{G,inv}^*(\delta_{\mathbb{R}}^{G,inv}(\bigwedge^k \sigma_n)) = \sum_{1 \leq j_1 < \dots < j_k \leq n} e_{j_1} \cdots e_{j_k} \otimes \delta_{\mathbb{R}}^{inv}(e_{j_1} + \dots + e_{j_k})$$

where e_i is the 1-dimensional Real representation of $(T, \sigma_{\mathbb{R}})$ with weight being the i -th standard basis vector of the weight lattice. Similarly, identifying $KR_{(U(2m), \sigma_{\mathbb{H}})}^*(U(2m)/T \times T, \sigma_{\mathbb{H}} \times Id)$ with $\mathbb{Z}[e_1^{\mathbb{H}}, \dots, e_{2m}^{\mathbb{H}}, (e_1^{\mathbb{H}} \cdots e_{2m}^{\mathbb{H}})^{-1}] \otimes KR^*(T, Id)$ (cf. [Fo, Proposition 4.25]), where $e_i^{\mathbb{H}}$ is the degree -4 class in $KR_{(U(2m), \sigma_{\mathbb{H}})}^*(U(2m)/T, \sigma_{\mathbb{H}})$ represented by the Quaternionic line bundle $U(2m) \times_T \mathbb{C}_{e_i}$, we have, for $\mathbb{F} = \mathbb{R}$ or \mathbb{H} (depending on the parity of k),

$$q_{G,inv}^*(\delta_{\mathbb{F}}^{G,inv}(\bigwedge^k \sigma_{2m})) = \sum_{1 \leq j_1 < \dots < j_k \leq 2m} e_{j_1}^{\mathbb{H}} \cdots e_{j_k}^{\mathbb{H}} \otimes \delta_{\mathbb{R}}^{inv}(e_{j_1} + \dots + e_{j_k})$$

Proof. The proof is similar to [Fo, Lemma 4.19]. The Proposition follows from the fact that the complex of $U(n)$ -equivariant Real vector bundles representing $\delta_{\mathbb{F}}^{G,inv}(\bigwedge^k \sigma_n)$, as in Proposition 3.3, is decomposed into a direct sum of complexes of T -equivariant Real vector bundles, each of which corresponds to a weight of $\bigwedge^k \sigma_n$. \square

Proposition 4.5. *Both $p_{G,inv}^*$ and $q_{G,inv}^*$ are injective.*

Proof. By [Fo, Lemma 4.19 and Proposition 4.25], and Corollary 4.2 and Proposition 4.4, we can identify both $p_{G,inv}^*$ and $q_{G,inv}^*$ with the map

$$i^* \otimes \text{Id}_{KR^*(pt)} : K_{U(n)}^*(U(n)) \otimes KR^*(pt) \rightarrow K_T^*(T) \otimes KR^*(pt)$$

where the restriction map i^* can factor through $K_T^*(U(n))$ as

$$K_{U(n)}^*(U(n)) \xrightarrow{i_1^*} K_T^*(U(n)) \xrightarrow{i_2^*} K_T^*(T)$$

$i_1^* \otimes \text{Id}_{KR^*(\text{pt})}$ is injective because i_1^* is split injective by [At3, Proposition 4.9]. By adapting [Fo, Lemma 4.20] to the case $G = U(n)$, we have

$$i_2^* \left(\prod_{i=1}^n \delta_T(\bigwedge^i \sigma_n) \right) = d_{U(n)} \otimes \prod_{i=1}^n \delta(e_i)$$

where $d_{U(n)}$ is the Weyl denominator for $U(n)$. By [Fo, Lemma 4.21] and the fact that $rd_{U(n)} \otimes \prod_{i=1}^n \delta(e_i) \neq 0$ for all $r \in KR^*(\text{pt}) \setminus \{0\}$, $i_2^* \otimes \text{Id}_{KR^*(\text{pt})}$ is injective as well. Thus $i^* \otimes \text{Id}_{KR^*(\text{pt})}$, as well as $p_{G,\text{inv}}^*$ and $q_{G,\text{inv}}^*$, are injective. \square

Lemma 4.6. *Let e be the standard representation of S^1 . Then $\delta_{\mathbb{R}}^{\text{inv}}(e)^2 = 0$ in $KR^*(S^1, \text{Id})$.*

Proof. Note that $\delta_{\mathbb{R}}^{\text{inv}}(e) \in KR^{-7}(S^1, \text{Id})$. So $\delta_{\mathbb{R}}^{\text{inv}}(e)^2 \in KR^{-6}(S^1, \text{Id}) \cong KR^{-7}(\text{pt}) = 0$. \square

Proposition 4.7. *For $\mathbb{F} = \mathbb{R}$ or \mathbb{H} , $\delta_{\mathbb{F}}^{G,\text{inv}}(\sigma_n)^2 = 0$ in $KR_{(U(n), \sigma_{\mathbb{F}})}^*(U(n), a_{\mathbb{F}})$.*

Proof. This follows from Propositions 4.4 and 4.5 and Lemma 4.6. \square

The above results finally culminate in the main theorem of this note.

Theorem 4.8. (1) *Let G be a Real compact Lie group, and ρ a Real (resp. Quaternionic) unitary representation of G . Then $\delta_{\mathbb{F}}^{G,\text{inv}}(\rho)^2 = 0$ in $KR_G^*(G^-)$ for $\mathbb{F} = \mathbb{R}$ (resp. $\mathbb{F} = \mathbb{H}$).*

(2) *In particular, if G is connected and simply-connected and $R(G, \mathbb{C}) = 0$, then $\delta_{\mathbb{R}}^{G,\text{inv}} \oplus \delta_{\mathbb{H}}^{G,\text{inv}}$ induces the following ring isomorphism*

$$KR_G^*(G^-) \cong \Omega_{KR_G^*(\text{pt})/KR^*(\text{pt})}$$

Proof. Note that the induced map $\rho^* : KR_{(U(n), \sigma_{\mathbb{F}})}^*(U(n), a_{\mathbb{F}}) \rightarrow KR_G^*(G^-)$ sends $\delta_{\mathbb{F}}^{G,\text{inv}}(\sigma_n)$ to $\delta_{\mathbb{F}}^{G,\text{inv}}(\rho)$ by the interpretation of $\delta_{\mathbb{F}}^{G,\text{inv}}$ in Proposition 3.3. Now part (1) follows from Proposition 4.7. Part (2) follows from part (1), Proposition 3.5 and Corollary 3.9. \square

Note that Theorems 3.8 and 4.8 give a complete description of the ring structure of $KR_G^*(G^-)$. Part (2) of Theorem 4.8 should be viewed as a generalization of Brylinski-Zhang's result in the context of KR-theory.

Last but not least, we obtain, as a by-product, the following

Corollary 4.9. *If G is a compact connected Real Lie group (not necessarily simply-connected) and X a compact Real G -space, then for any x in $KR_G^1(X)$ or $KR_G^{-3}(X)$, $x^2 = 0$.*

Proof. Let EG^n be the join of n copies of G , with the Real structure induced by σ_G and G -action by the left-translation of G . Let $\pi_n^* : KR_G^*(X) \rightarrow KR_G^*(X \times EG^n)$ be the map induced by projection onto X . Consider the map

$$\pi^* := \lim_{\leftarrow n} \pi_n^* : KR_G^*(X) \rightarrow \lim_{\leftarrow n} KR_G^*(X \times EG^n)$$

By adapting the proof of [AS, Corollary 2.3] to the Real case, $\ker(\pi^*) = \bigcap_{n \in \mathbb{N}} IR^n \cdot KR_G^*(X)$, where IR is the augmentation ideal of $RR(G)$. Moreover, $\bigcap_{n \in \mathbb{N}} I^n = (0)$ where I is the augmentation ideal of $R(G)$ (cf. the Note immediately preceding [AH, Section 4.5]), and the map $RR(G) \rightarrow R(G)$ forgetting the Real structure is injective (cf. [Fo, Proposition 2.17(1)]). It follows that $\bigcap_{n \in \mathbb{N}} IR^n = (0)$ and thus π^* is injective. Now it suffices to show that $\pi^*(x)^2 = 0$. Using Lemma 3.1 and compactness of $X \times EG^n/G$, $\pi_n^*(x) \in KR_G^*(X \times EG^n) = KR^*(X \times EG^n/G)$ can be represented by a Real map $f_n : X \times EG^n/G \rightarrow (U(k_n), a_{\mathbb{F}})$ for some k_n . So $\pi_n^*(x) = f_n^* \delta_{\mathbb{F}}^{\text{inv}}(\sigma_{k_n})$, and $\pi_n^*(x)^2 = f_n^* \delta_{\mathbb{F}}^{\text{inv}}(\sigma_{k_n})^2 = 0$ because we have that $\delta_{\mathbb{F}}^{\text{inv}}(\sigma_{k_n})^2 = 0$ by applying the forgetful map $KR_{(U(n), \sigma_{\mathbb{F}})}^*(U(n), a_{\mathbb{F}}) \rightarrow KR^*(U(n), a_{\mathbb{F}})$ to the equation $\delta_{\mathbb{F}}^{G, \text{inv}}(\sigma_{k_n})^2 = 0$ (see Proposition 4.7). Since $\pi^*(x)^2$ is the inverse limit of $\pi_n^*(x)^2 = 0$, $\pi^*(x)^2 = 0$ as desired. \square

Remark 4.10. Corollary 4.9 has the following consequences as a result of forgetting some structures in the equivariant KR -theory.

- (1) When G is trivial, Corollary 4.9 then says that $x^2 = 0$ for any x in $KR^i(X)$, where i is 1 or -3 . This special case can be proved more directly using Lemma 3.1, and noting that $\delta_{\mathbb{F}}^{\text{inv}}(\sigma_n)^2 = 0$ in $KR^*(U(n), a_{\mathbb{F}})$, which can be obtained by applying the map $KR_{(U(n), \sigma_{\mathbb{F}})}^*(U(n), a_{\mathbb{F}}) \rightarrow KR^*(U(n), a_{\mathbb{F}})$ forgetting the equivariant structure to the equation in Proposition 4.7.
- (2) Applying the map $KR^*(U(n), a_{\mathbb{F}}) \rightarrow K^*(U(n))$, which further forgets the Real structure, to the equation $\delta_{\mathbb{F}}^{\text{inv}}(\sigma_n)^2 = 0$, yields $\delta(\sigma_n)^2 = 0$ in $K^*(U(n))$. Pulling back from this ‘universal’ equation, we then get the well-known fact that $x^2 = 0$ for $x \in K^{-1}(X)$.
- (3) If the involution on X is trivial, then $KR^*(X) \cong KO^*(X)$, and we also have that $x^2 = 0$ for $x \in KO^i(X)$, where i is 1 or -3 .

However, it is not true that $x^2 = 0$ if $x \in KR_G^i(X)$ for $i = -1$ or -5 . For example, if η is the non-zero element in $KR^{-1}(\text{pt})$, then $\eta^2 \neq 0$. An example where $x \in KR_G^{-5}(X)$ but $x^2 \neq 0$ can be found in [Fo, Corollary 4.32].

REFERENCES

- [At] M. F. Atiyah, *K-theory*, W. A. Benjamin, Inc., 1964.
- [At2] M. F. Atiyah, *K-theory and reality*, Quart. J. Math. Oxford (2), 17, 367-86, 1966.
- [At3] M. F. Atiyah, *Bott periodicity and the index of elliptic operators*, Quart. J. Math. Oxford (2), 19, 113-40, 1968.
- [ABS] M. F. Atiyah, R. Bott, A. Shapiro, *Clifford modules*, Topology, Vol. 3, Suppl. 1, 3-38, Pergamon Press, 1964.
- [AH] M. F. Atiyah, F. Hirzebruch, *Vector bundles and homogeneous spaces*, Proc. Sympos. Pure Math., Vol. III, 7-38, American Mathematical Society, Providence, R.I., 1961.
- [AS] M. F. Atiyah, G. Segal, *Equivariant K-theory and completion*, J. Differential Geometry 3, 1-18, 1969.
- [Bot] R. Bott, *The periodicity theorem for the classical groups and some of its applications*, Advances in Mathematics, Vol. 4, 353-411, 1970.
- [BZ] J.-L. Brylinski, B. Zhang, *Equivariant K-theory of compact connected Lie groups*, K-Theory, Volume 20, Issue 1, pp. 23-36, May 2000.
- [Fo] C.-K. Fok, *The Real K-theory of compact Lie groups*, Symmetry, Integrability and Geometry: Methods and Applications, Vol. 10(022), 2014.

- [HL] M. Harada, G. Landweber, *Surjectivity for Hamiltonian G -spaces in K -theory*, Trans. Amer. Math. Soc., Vol. 359, No. 12, pp. 6001-6025, Dec. 2007.
- [Ho] L. Hodgkin, On the K -theory of Lie groups, *Topology* **6**, 1-36, 1967.
- [Se] R. M. Seymour, *The real K -theory of Lie groups and homogeneous spaces*, Quart. J. Math. Oxford (2), 24, 7-30, 1973.

NATIONAL CENTER FOR THEORETICAL SCIENCES,
MATHEMATICS DIVISION,
NATIONAL TSING HUA UNIVERSITY
HSINCHU 30013, TAIWAN
E-MAIL: ckfok@ntu.edu.tw
URL: <http://www.math.cornell.edu/~ckfok>