

2-LOCAL TRIPLE HOMOMORPHISMS ON VON NEUMANN ALGEBRAS AND JBW*-TRIPLES

MARIA BURGOS, FRANCISCO J. FERNÁNDEZ-POLO, JORGE J. GARCÉS,
AND ANTONIO M. PERALTA

ABSTRACT. We prove that every (not necessarily linear nor continuous) 2-local triple homomorphism from a JBW*-triple into a JB*-triple is linear and a triple homomorphism. Consequently, every 2-local triple homomorphism from a von Neumann algebra (respectively, from a JBW*-algebra) into a C*-algebra (respectively, into a JB*-algebra) is linear and a triple homomorphism.

1. INTRODUCTION

It is known that the Gleason-Kahane-Żelazko theorem (cf. [23, 28, 45]) admits a reinterpretation affirming that every unital linear local homomorphism from a unital complex Banach algebra A into \mathbb{C} is multiplicative. Formally speaking, the notions of local homomorphisms and local derivations were introduced in 1990, in papers due to Larson and Sourour [34] and Kadison [27]. We recall that given two Banach algebras A and B , a linear mapping $T : A \rightarrow B$ (respectively, $T : A \rightarrow A$) is said to be a *local homomorphism* (respectively, a *local derivation*) if for every a in A there exists a homomorphism $\Phi_a : A \rightarrow B$ (respectively, a derivation $D_a : A \rightarrow A$), depending on a , satisfying $T(a) = \Phi_a(a)$ (respectively, $T(a) = D_a(a)$). A flourishing research on linear local homomorphisms and derivations was built upon the results of Kadison, Larson and Sourour (compare, for example, [1, 5, 7, 8, 13, 15, 17, 18, 20, 26, 31, 32], [36]–[44] and [46], among the over 100 references on the subject).

If in the definition of *local homomorphism*, we relax the assumption concerning linearity with a 2-local behavior, we are led to the notion of (not necessarily linear) *2-local homomorphism*. Let A and B be two C*-algebras, a not necessarily linear nor continuous mapping $T : A \rightarrow B$ is said to be a *2-local homomorphism* (respectively, *2-local *-homomorphism*) if for

1991 *Mathematics Subject Classification*. Primary 46L05; 46L40.

Key words and phrases. Local triple homomorphism, triple homomorphism; 2-local triple homomorphism.

Authors partially supported by the Spanish Ministry of Science and Innovation, D.G.I. project no. MTM2011-23843, and Junta de Andalucía grant FQM375. The fourth author extends his appreciation to the Deanship of Scientific Research at King Saud University (Saudi Arabia) for funding the work through research group no. RGP-361.

every $a, b \in A$ there exists a bounded (linear) homomorphism (respectively, $*$ -homomorphism) $\Phi_{a,b} : A \rightarrow B$, depending on a and b , such that $\Phi_{a,b}(a) = T(a)$ and $\Phi_{a,b}(b) = T(b)$ (see [43], [14]).

In a recent contribution, we establish a generalization of the Kowalski-Słodkowski theorem for 2-local $*$ -homomorphisms on von Neumann algebras, showing that every (not necessarily linear nor continuous) 2-local $*$ -homomorphism from a von Neumann algebra or from a compact C^* -algebra into a C^* -algebra is linear and a $*$ -homomorphism. In the Jordan setting, it is proved that every 2-local Jordan $*$ -homomorphism from a JBW^* -algebra into a JB^* -algebra is linear and a Jordan $*$ -homomorphism (cf. [14]).

Every C^* -algebra A admits a ternary product given by

$$\{a, b, c\} := \frac{1}{2}(ab^*c + cb^*a) \quad (a, b, c \in A).$$

A linear map Φ between C^* -algebras A and B satisfying $\Phi(\{a, b, c\}) = \{\Phi(a), \Phi(b), \Phi(c)\}$, is called a *triple homomorphism*. A *2-local triple homomorphism* between A and B is a not necessarily linear nor continuous map $T : A \rightarrow B$ such that for every $a, b \in A$, there exists a triple homomorphism $\Phi_{a,b} : A \rightarrow B$ with $\Phi_{a,b}(a) = T(a)$ and $\Phi_{a,b}(b) = T(b)$. Motivated by the above commented Kowalski-Słodkowski theorem for von Neumann algebras, it seems natural to consider the following independent problem:

Problem 1.1. *Is every 2-local triple homomorphism between C^* -algebras (automatically) linear?*

It should be noted here that, even in the case of von Neumann algebras, the proofs and arguments given in the study of 2-local $*$ -homomorphisms [14], are no longer valid when considering Problem 1.1, because triple homomorphisms between C^* -algebras do not preserve the natural partial order given by the positive cone in a C^* -algebra.

Problem 1.1 can be posed in the more general setting of JB^* -triples. Let E and F be two JB^* -triples (see subsection 1.1 for definitions). A linear map $\Phi : E \rightarrow F$ which preserves the triple products is called a *triple homomorphism*. A (not necessarily linear nor continuous) mapping $T : E \rightarrow F$ is said to be a *2-local triple homomorphism* if for every $a, b \in E$ there exists a bounded (linear) triple homomorphism $\Phi_{a,b} : E \rightarrow F$, depending on a and b , such that $\Phi_{a,b}(a) = T(a)$ and $\Phi_{a,b}(b) = T(b)$. According to these definitions, we consider the following generalization of Problem 1.1:

Problem 1.2. *Is every 2-local triple homomorphism between JB^* -triples (automatically) linear?*

In this paper we solve Problems 1.1 and 1.2 when the domain is a von Neumann algebra or a JBW^* -triple, respectively. Our main result (Theorem 3.8) asserts that every (not necessarily linear nor continuous) 2-local triple homomorphism from a JBW^* -triple into a JB^* -triple is linear and

a triple homomorphism, and consequently, every 2-local triple homomorphism from a von Neumann algebra (respectively, from a JBW*-algebra) into a C*-algebra (respectively, into a JB*-algebra) is linear and a triple homomorphism (cf. Theorem 3.5 and Corollary 3.6). Our proofs heavily rely on the Bunce-Wright-Mackey-Gleason theorem for JBW*-algebras [9] and deep geometric arguments and techniques, developed in the setting of JB*-triples by R. Braun, W. Kaup and H. Upmeyer [6, 30], B. Russo and Y. Friedman [22], and G. Horn [25].

1.1. Preliminaries. A *JB*-triple* is a complex Banach space, E , together with a continuous triple product $\{.,.,.\} : E \times E \times E \rightarrow E$, $(a, b, c) \mapsto \{a, b, c\}$, which is conjugate-linear in b and symmetric and bilinear in (a, c) and satisfies:

(1) The *Jordan identity*:

$$L(a, b)L(x, y) - L(x, y)L(a, b) = L(L(a, b)x, y) - L(x, L(b, a)y),$$

where $L(a, b)$ denotes the operator given by $L(a, b)x = \{a, b, x\}$;

(2) $L(a, a)$ is an hermitian operator with non-negative spectrum;

(3) $\|\{a, a, a\}\| = \|a\|^3$,

every a, b, x and y in E .

The notion of JB*-triples was introduced by Kaup in the holomorphic classification of bounded symmetric domains in [29]. One of the many kindness exhibited by the class of JB*-triples is that every C*-algebra (respectively, every JB*-algebra) is a JB*-triple with respect to

$$\{a, b, c\} := \frac{1}{2}(ab^*c + cb^*a)$$

(respectively, $\{a, b, c\} := (a \circ b^*) \circ c + (c \circ b^*) \circ a - (a \circ c) \circ b^*$).

A *JBW*-triple* is a JB*-triple which is also a dual Banach space (with a unique isometric predual [4]). It is known that the triple product of a JBW*-triple is separately weak* continuous (cf. [4]).

We recall that an element e in a JB*-triple E is said to be a *tripotent* if $\{e, e, e\} = e$. It is known that for each tripotent e in E we have a decomposition (called the *Peirce decomposition*)

$$E = E_2(e) \oplus E_1(e) \oplus E_0(e),$$

where for $j = 0, 1, 2$, $E_j(e)$ is the $\frac{j}{2}$ -eigenspace of $L(e, e)$. The Peirce subspaces $E_j(e)$ satisfy the following multiplication rules:

$$\{E_i(e), E_j(e), E_k(e)\} \subseteq E_{i-j+k}(e),$$

if $i - j + k \in \{0, 1, 2\}$ and is zero otherwise, and

$$\{E_2(e), E_0(e), E\} = \{E_0(e), E_2(e), E\} = 0.$$

These multiplication rules are called the *Peirce rules*. The natural projection $P_j(e) : E \rightarrow E_j(e)$ of E onto $E_j(e)$ is called the *Peirce- j projection*. The

Peirce projections are contractive and satisfy

$$P_2(e) = L(e, e)(2L(e, e) - Id), \quad P_1(e) = 4L(e, e)(Id - L(e, e)),$$

$$\text{and } P_0(e) = (Id - L(e, e))(Id - 2L(e, e)),$$

where Id denotes the identity map on E (compare [22]). It is also known that for each $x_0 \in E_0(e)$ and $x_2 \in E_2(e)$ we have $\|x_0 + x_2\| = \max\{\|x_0\|, \|x_2\|\}$ (c.f. [22, Lemma 1.3]). The tripotent e is called *complete* when $E_0(e) = \{0\}$.

Another interesting property of the Peirce decomposition asserts that $E_2(e)$ is a unital JB*-algebra with unit e , product $a \circ_e b = \{a, e, b\}$ and involution $a^{\sharp_e} = \{e, a, e\}$ (c.f. [6, Theorem 2.2] and [30, Theorem 3.7]).

Accordingly to the standard terminology, for each element a in a JB*-triple E , we denote $a^{[1]} = a$ and $a^{[2n+1]} := \{a, a^{[2n-1]}, a\}$ ($\forall n \in \mathbb{N}$). It follows from the Jordan identity that JB*-triples are power associative, that is, $\{a^{[2k-1]}, a^{[2l-1]}, a^{[2m-1]}\} = a^{[2(k+l+m)-3]}$. In this paper, the symbol E_a will denote the JB*-subtriple of E generated by a . It is known that E_a is JB*-triple isomorphic (and hence isometric) to $C_0(L)$ for some locally compact Hausdorff space $L \subseteq (0, \|a\|]$, such that $L \cup \{0\}$ is compact and $\|a\| \in L$. It is further known that there exists a triple isomorphism Ψ from E_a onto $C_0(L)$, satisfying $\Psi(a)(t) = t$ ($t \in L$) (compare [29, Lemma 1.14]). In particular, for each natural n , there exists (a unique) element $a^{[1/(2n-1)]}$ in E_a satisfying $(a^{[1/(2n-1)]})^{[2n-1]} = a$.

When a is a norm one element in a JBW*-triple E , the sequence $(a^{[1/(2n-1)]})$ converges in the weak* topology of E to a tripotent in E , which is denoted by $r(a)$ and is called the *range tripotent* of a . The tripotent $r(a)$ is the smallest tripotent e in E satisfying that a is positive in the JBW*-algebra $E_2(e)$ (cf. [19, Lemma 3.3]).

We refer to [16] for a recent monograph on JB*-triples and JB*-algebras.

Throughout the paper, when A is a C*-algebra or a JB*-algebra, the symbol A_{sa} will stand for the set of all self-adjoint elements in A .

2. GENERALITIES ON 2-LOCAL TRIPLE HOMOMORPHISMS

We recall that elements a and b in a JB*-triple E are said to be *orthogonal* (written $a \perp b$) when $L(a, b) = 0$. It is known that $a \perp b$ if and only if $\{a, a, b\} = 0$, if and only if $\{a, b, b\} = 0$ (cf. [11, Lemma 1.1]). A pair of subsets $M, N \subset E$ are called orthogonal ($M \perp N$) if for every $a \in M, b \in N$, we have $a \perp b$.

Throughout the paper, given a 2-local triple homomorphism T between JB*-triples E and F , for each $a, b \in E$, $\Phi_{a,b}$ will denote a (linear) triple homomorphism satisfying $T(a) = \Phi_{a,b}(a)$ and $T(b) = \Phi_{a,b}(b)$.

We begin with some basic properties of 2-local triple homomorphisms.

Lemma 2.1. *Let $T : E \rightarrow F$ be a (not necessarily linear nor continuous) 2-local triple homomorphism between JB^* -triples. The following statements hold:*

- (a) T is 1-homogeneous, that is, $T(\lambda a) = \lambda T(a)$ for every $a \in E$, $\lambda \in \mathbb{C}$;
- (b) T is orthogonality preserving;
- (c) $\{T(a), T(a), T(a)\} = T(\{a, a, a\})$, for every $a \in E$. In particular, every linear 2-local triple homomorphism between JB^* -triples is a triple homomorphism;
- (d) T maps tripotents in E to tripotents in F ;
- (e) For each $a, b \in E$, $\|T(a) - T(b)\| \leq \|a - b\|$, that is, T is 1-lipschitzian and hence continuous;
- (f) For each tripotent e in E with $T(e) \neq 0$, we have $T(E_j(e)) \subseteq F_j(T(e))$, for every $j = 0, 1, 2$, $T(E_2(e) + E_1(e)) \subseteq F_2(T(e)) + F_1(T(e))$, and $T(E_0(e) + E_1(e)) \subseteq F_0(T(e)) + F_1(T(e))$. Furthermore, $T(E_2(e)_{sa}) \subseteq F_2(T(e))_{sa}$;
- (g) For each tripotent $e \in E$ with $T(e) = 0$, the mapping T is zero on $E_2(e) \oplus E_1(e)$.

Proof. The proof of (a) is standard (compare [14, Lemma 2.1]). For the statement (b), we recall that $a \perp b$ if and only if $\{a, a, b\} = 0$ [11, Lemma 1.1]. Let us consider the triple homomorphism $\Phi_{a,b} : E \rightarrow F$. Then

$$\{T(a), T(a), T(b)\} = \{\Phi_{a,b}(a), \Phi_{a,b}(a), \Phi_{a,b}(b)\} = \Phi_{a,b}\{a, a, b\} = 0,$$

which proves $T(a) \perp T(b)$.

- (c) Considering the triple homomorphism $\Phi_{a,a^{[3]}}$, we have

$$\begin{aligned} \{T(a), T(a), T(a)\} &= \{\Phi_{a,a^{[3]}}(a), \Phi_{a,a^{[3]}}(a), \Phi_{a,a^{[3]}}(a)\} \\ &= \Phi_{a,a^{[3]}}(a^{[3]}) = T(\{a, a, a\}). \end{aligned}$$

The second statement follows from the polarization formula

$$8\{x, y, z\} = \sum_{k=0}^3 \sum_{j=1}^2 i^k (-1)^j \left(x + i^k y + (-1)^j z \right)^{[3]}.$$

(d) is clear from (c), and (e) follows from the fact that every triple homomorphism between JB^* -triples is contractive (cf. [3, Lemma 1] and the proof of [14, Lemma 2.1]).

(f) Let us take a tripotent $e \in E$ with $T(e)$ a non-zero tripotent in F . For each $a \in E_j(e)$ we have $L(e, e)(a) = \frac{j}{2}a$. Therefore,

$$\begin{aligned} L(T(e), T(e))T(a) &= \{T(e), T(e), T(a)\} = \{\Phi_{e,a}(e), \Phi_{e,a}(e), \Phi_{e,a}(a)\} \\ &= \Phi_{e,a}(\{e, e, a\}) = \frac{j}{2}\Phi_{e,a}(a) = \frac{j}{2}T(a), \end{aligned}$$

witnessing that $T(E_j(e)) \subseteq F_j(T(e))$, for every $j = 0, 1, 2$. We can similarly show that $T(a) \in \ker(Q(T(e))) = F_0(T(e)) + F_1(T(e))$ for every $a \in \ker(Q(e)) = E_0(e) + E_1(e)$ which shows that $T(E_0(e) + E_1(e)) \subseteq F_0(T(e)) + F_1(T(e))$.

Since $F_2(T(e)) + F_1(T(e)) = \ker(P_0(T(e)))$ and

$$P_0(T(e)) = (Id_F - L(T(e), T(e)))(Id_F - 2L(T(e), T(e))),$$

we can show, applying the triple homomorphism $\Phi_{a,e}$, that, for each element $a \in \ker(P_0(e)) = E_2(e) + E_1(e)$, we have $T(a) \in \ker(P_0(T(e)))$, which gives the other inclusion.

Suppose $a \in E_2(e)_{sa} = \{x \in E_2(e) : x = x^{\sharp e} = \{e, x, e\}\}$. Since

$$\begin{aligned} \{T(e), T(a), T(e)\} &= \{\Phi_{e,a}(e), \Phi_{e,a}(a), \Phi_{e,a}(e)\} \\ &= \Phi_{e,a}(\{e, a, e\}) = \Phi_{e,a}(a) = T(a), \end{aligned}$$

we deduce that $T(a) \in F_2(T(e))_{sa}$.

(g) Suppose $T(e) = 0$ and $a = a_1 + a_2$, where $a_j \in E_j(e)$ for $j = 1, 2$. In such a case

$$\begin{aligned} T(a) &= \Phi_{a,e}(a) = \Phi_{a,e}(\{e, e, a_2\}) + 2\Phi_{a,e}(\{e, e, a_1\}) \\ &= \{\Phi_{a,e}(e), \Phi_{a,e}(e), \Phi_{a,e}(a_2)\} + 2\{\Phi_{a,e}(e), \Phi_{a,e}(e), \Phi_{a,e}(a_1)\} \\ &= \{T(e), T(e), \Phi_{a,e}(a_2)\} + 2\{T(e), T(e), \Phi_{a,e}(a_1)\} = 0. \end{aligned}$$

□

We shall establish next a triple version of [14, Lemma 3.1].

Lemma 2.2. *Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism between JB^* -triples. Then, for each $a \in E$, $T|_{E_a} : E_a \rightarrow F$ is a linear mapping.*

Proof. Let us consider an element $b \in E_a$ of the form $b = \sum_{k=1}^m \alpha_k a^{[2k-1]}$ and the triple homomorphism $\Phi_{a,b}$. The identity

$$T(b) = \Phi_{a,b} \left(\sum_{k=1}^m \alpha_k a^{[2k-1]} \right) = \sum_{k=1}^m \alpha_k \Phi_{a,b}(a)^{[2k-1]} = \sum_{k=1}^m \alpha_k T(a)^{[2k-1]},$$

proves that T is linear on the linear span of the set $\{a^{[2k-1]} : k \in \mathbb{N}\}$. The continuity of T shows that $T|_{E_a}$ is linear. □

Our next technical result establishes that every (not necessarily linear) 2-local triple homomorphism between JB^* -triples is additive on every couple of orthogonal tripotents. The result is a generalization of [14, Lemma 2.2] to the setting of JB^* -triples; it should be noted that, in this more general setting, we need new and independent geometric arguments.

Lemma 2.3. *Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism between JB^* -triples. Let e and f be two orthogonal tripotents in E . Then $T(e + f) = T(e) + T(f)$.*

Proof. Take a real number $\lambda \in (0, 1]$. In this case we have

$$T(e + \lambda f) = \Phi_{e+\lambda f, e}(e + \lambda f) = T(e) + \lambda \Phi_{e+\lambda f, e}(f)$$

with $\Phi_{e+\lambda f, e}(f) \perp T(e)$. We similarly have

$$T(e + \lambda f) = \Phi_{e+\lambda f, f}(e + \lambda f) = \Phi_{e+\lambda f, f}(e) + \lambda T(f).$$

Combining the above identities we have that

$$\begin{aligned} \Phi_{e+\lambda f, f}(e) &= T(e) + \lambda(\Phi_{e+\lambda f, e}(f) - T(f)) \\ &= T(e) + P_0(T(e))(T(e + \lambda f)) - \lambda T(f). \end{aligned}$$

Since $\Phi_{e+\lambda f, f}(e)$ and $T(e)$ are tripotents and

$$T(e) \perp P_0(T(e))(T(e + \lambda f)) - \lambda T(f),$$

it follows that $P_0(T(e))(T(e + \lambda f)) - \lambda T(f)$ also is a tripotent for every $\lambda \in (0, 1]$.

Clearly, the function $f : [0, 1] \rightarrow \{0, 1\}$ defined by

$$f(\lambda) := \|P_0(T(e))T(e + \lambda f) - \lambda T(f)\|,$$

is continuous with $f(0) = 0$, thus $f(\lambda) = 0 \quad \forall \lambda \in [0, 1]$. This implies, in particular, that $f(1) = 0$, or equivalently, $P_0(T(e))T(e + f) = T(f)$, which finishes the proof. \square

The linearity of every (not necessarily linear) 2-local triple homomorphism on finite linear combinations of mutually orthogonal tripotents follows next.

Lemma 2.4. *Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism between JB^* -triples. Let e_1, \dots, e_n be mutually orthogonal tripotents in E . Then*

- (a) $T\left(\sum_{i=1}^n e_i\right) = \sum_{i=1}^n T(e_i)$;
- (b) $T\left(\sum_{i=1}^n \lambda_i e_i\right) = \sum_{i=1}^n \lambda_i T(e_i)$, for every $\lambda_1, \dots, \lambda_n \in \mathbb{C}$.

Proof. (a) We shall argue by induction on n . The case $n = 1$ is clear, while the case $n = 2$ is established in Lemma 2.3. Let us suppose that e_1, \dots, e_n, e_{n+1} are mutually orthogonal tripotents in E . Since $e = e_1 + \dots + e_n$ and e_{n+1} are orthogonal tripotents in E , Lemma 2.3 and the induction hypothesis prove that

$$T\left(\sum_{i=1}^{n+1} e_i\right) = T(e + e_{n+1}) = T(e) + T(e_{n+1}) = \sum_{i=1}^n T(e_i) + T(e_{n+1}).$$

- (b) Fix $j \in \{1, \dots, n\}$ and set $z = \sum_{i=1}^n \lambda_i e_i$ and $e = \sum_{i=1}^n e_i$. The identity

$$\left\{T(e_j), T(e_j), T\left(\sum_{i=1}^n \lambda_i e_i\right)\right\} = \left\{\Phi_{z, e_j}(e_j), \Phi_{z, e_j}(e_j), \Phi_{z, e_j}\left(\sum_{i=1}^n \lambda_i e_i\right)\right\}$$

$$= \Phi_{z, e_j} \left(\left(e_j, e_j, \sum_{i=1}^n \lambda_i e_i \right) \right) = \Phi_{z, e_j}(\lambda_j e_j) = \lambda_j T(e_j),$$

implies that

$$\begin{aligned} T \left(\sum_{i=1}^n \lambda_i e_i \right) &= \Phi_{z, e}(z) = \Phi_{z, e}(\{e, e, z\}) = \{\Phi_{z, e}(e), \Phi_{z, e}(e), \Phi_{z, e}(z)\} \\ &= \{T(e), T(e), T(z)\} = \left\{ T \left(\sum_{j=1}^n e_j \right), T \left(\sum_{j=1}^n e_j \right), T \left(\sum_{i=1}^n \lambda_i e_i \right) \right\} \\ &= (\text{by (a)}) = \left\{ \sum_{j=1}^n T(e_j), \sum_{j=1}^n T(e_j), T \left(\sum_{i=1}^n \lambda_i e_i \right) \right\} = (\text{by orthogonality}) \\ &= \sum_{j=1}^n \left\{ T(e_j), T(e_j), T \left(\sum_{i=1}^n \lambda_i e_i \right) \right\} = \sum_{j=1}^n \lambda_j T(e_j). \end{aligned}$$

□

Let E and F be JB*-triples. We recall that a (not necessarily linear) mapping $f : E \rightarrow F$ is called *orthogonally additive* if $f(a + b) = f(a) + f(b)$ for every $a \perp b$ in E .

Proposition 2.5. *Let E be a JBW*-triple, F a JB*-triple, and suppose that $T : E \rightarrow F$ is a (not necessarily linear) 2-local triple homomorphism. Then T is orthogonally additive.*

Proof. Let a and b be two orthogonal elements in E . The range tripotents $r(a)$ and $r(b)$ are orthogonal, and the JBW*-subtriples $E_2(r(a))$ and $E_2(r(b))$ are also orthogonal (cf. [11, Lemma 1.1]).

For each $\varepsilon > 0$, there exists two algebraic elements $a_\varepsilon = \sum_{k=1}^{m_1} \lambda_k e_k$ and $b_\varepsilon = \sum_{j=1}^{m_2} \mu_j v_j$, where $\lambda_k, \mu_j \in \mathbb{R}$, e_1, \dots, e_{m_1} and v_1, \dots, v_{m_2} are tripotents in $E_2(r(a))$ and $E_2(r(b))$, respectively, and $e_j \perp e_k$, $v_j \perp v_k$ for every $j \neq k$, such that $\|a - a_\varepsilon\| < \frac{\varepsilon}{4}$, and $\|b - b_\varepsilon\| < \frac{\varepsilon}{4}$ (cf. [25, lemma 3.11]). It is clear that $a_\varepsilon + b_\varepsilon$ is a linear combination of mutually orthogonal tripotents. Then, by Lemma 2.1(e) and Lemma 2.4(b),

$$\begin{aligned} \|T(a+b) - T(a) - T(b)\| &= \|T(a+b) - T(a_\varepsilon + b_\varepsilon) + T(a_\varepsilon) + T(b_\varepsilon) - T(a) - T(b)\| \\ &\leq \|T(a+b) - T(a_\varepsilon + b_\varepsilon)\| + \|T(a_\varepsilon) - T(a)\| \\ &\quad + \|T(b_\varepsilon) - T(b)\| < \|(a+b) - (a_\varepsilon + b_\varepsilon)\| + \|a_\varepsilon - a\| + \|b_\varepsilon - b\| < \varepsilon. \end{aligned}$$

Since ε was arbitrarily chosen, we get $T(a+b) = T(a) + T(b)$. □

A simple induction argument, combined with Proposition 2.5, shows:

Corollary 2.6. *Let $(E_i)_{i=1}^n$ be a finite family of JBW^* -triples and let F be a JB^* -triple. Suppose that, for every i , every (not necessarily linear) 2-local triple homomorphism $T : E_i \rightarrow F$ is linear. Then every (not necessarily linear) 2-local triple homomorphism $T : \bigoplus_{i=1, \dots, n}^{\ell_\infty} E_i \rightarrow F$ is linear. \square*

We recall now a result, due to Friedmann and Russo, which has been borrowed from [22, Lemma 1.6].

Lemma 2.7. [22, Lemma 1.6] *Let e be a tripotent in a JB^* -triple E . Then, for each norm-one element $x \in E$ satisfying $P_2(e)x = e$, we have $P_1(e)x = 0$. \square*

In order to make the results more accessible, we have splitted the technical arguments needed in the proofs of our main results into a series of lemmas and propositions, which assure certain almost-linearity properties of 2-local triple homomorphisms.

Lemma 2.8. *Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism between JB^* -triples. Suppose e is a tripotent in E and $z \in E_0(e)$ with $\|z\| < 1$. Then, for each $w \in E$ and each triple homomorphism $\Phi_{w, e+z} : E \rightarrow F$ satisfying $\Phi_{w, e+z}(w) = T(w)$ and $\Phi_{w, e+z}(e+z) = T(e+z)$, we have $\Phi_{w, e+z}(e) = T(e)$, and consequently, $\Phi_{w, e+z}(E_j(e)) \subseteq F_j(T(e))$, for every $j = 0, 1, 2$.*

Proof. By considering the triple homomorphism $\Phi_{e, e+z}$, we obtain that

$$T(e+z) = \Phi_{e, e+z}(e+z) = T(e) + \Phi_{e, e+z}(z),$$

where $\Phi_{e, e+z}(z) \in F_0(\Phi_{e, e+z}(e)) = F_0(T(e))$ and $\|\Phi_{e, e+z}(z)\| \leq \|z\| < 1$. Since $\|z\| < 1$, $\|\Phi_{e, e+z}(z)\| < 1$ and $T(z) \in F_0(T(e))$ (cf. Lemma 2.1), we have,

$$\begin{aligned} \Phi_{w, e+z}(e) &= \Phi_{w, e+z} \left(\lim_{n \rightarrow \infty} (e+z)^{[3^n]} \right) = \lim_{n \rightarrow \infty} (\Phi_{w, e+z}(e+z))^{[3^n]} \\ &= \lim_{n \rightarrow \infty} (T(e+z))^{[3^n]} = \lim_{n \rightarrow \infty} (T(e) + \Phi_{e, e+z}(z))^{[3^n]} = T(e), \end{aligned}$$

where all the above limits are in the norm-topology. \square

Lemma 2.9. *Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism between JB^* -triples. Then the following statements hold:*

(a) *For each tripotent e in E , and each $y \in E_1(e)$, we have*

$$T(e+y) = T(e) + T(y);$$

(b) *Suppose e_1, e_2 , and g are tripotents in E satisfying $e_1 \perp e_2$, $e_1, e_2 \in E_2(g)$, $g \in E_1(e_1) \cap E_1(e_2)$. Then, the identity*

$$T(\lambda_1 e_1 + \mu g + \lambda_2 e_2) = \lambda_1 T(e_1) + \mu T(g) + \lambda_2 T(e_2),$$

holds for every $\lambda_1, \lambda_2, \mu \in \mathbb{C}$.

Proof. (a) Let e be a tripotent in E , and let $y \in E_1(e)$. By Lemma 2.1(g), the desired statement is clear when $T(e) = 0$, so we assume that $T(e) \neq 0$. In this case,

$$T(e + y) = \Phi_{e+y,e}(e + y) = T(e) + \Phi_{e+y,e}(y),$$

where $\Phi_{e+y,e}(y) \in F_1(\Phi_{e+y,e}(e)) = F_1(T(e))$, and we also have

$$T(e + y) = \Phi_{e+y,y}(e + y) = \Phi_{e+y,y}(e) + T(y),$$

with $\Phi_{e+y,y}(e)$ being a tripotent. Therefore,

$$\Phi_{e+y,y}(e) = T(e) + \Phi_{e+y,e}(y) - T(y),$$

with $\Phi_{e+y,e}(y) - T(y) \in F_1(T(e))$. It follows from Lemma 2.7 that

$$0 = P_1(T(e))(\Phi_{e+y,y}(e)) = \Phi_{e+y,e}(y) - T(y),$$

witnessing the desired statement.

(b) We can assume that $\lambda_1, \lambda_2, \mu \neq 0$, otherwise the statement is clear from (a) or from Lemma 2.4. To simplify notation, we set $z = \lambda_1 e_1 + \mu g + \lambda_2 e_2$. Applying (a) we get

$$T(z) = \Phi_{z,\lambda_1 e_1 + \mu g}(z) = \lambda_1 T(e_1) + \mu T(g) + \lambda_2 \Phi_{z,\lambda_1 e_1 + \mu g}(e_2).$$

We also have

$$(2.1) \quad T(z) = \Phi_{z,e_2}(z) = \lambda_1 \Phi_{z,e_2}(e_1) + \mu \Phi_{z,e_2}(g) + \lambda_2 T(e_2).$$

Combining these two equalities we have

$$\Phi_{z,\lambda_1 e_1 + \mu g}(e_2) = T(e_2) + \frac{\mu}{\lambda_2}(\Phi_{z,e_2}(g) - T(g)) + \frac{\lambda_1}{\lambda_2}(\Phi_{z,e_2}(e_1) - T(e_1)),$$

where $\Phi_{z,e_2}(e_1) \in F_0(\Phi_{z,e_2}(e_2)) = F_0(T(e_2))$, $T(e_1) \in F_0(T(e_2))$, $\Phi_{z,e_2}(g) \in F_1(\Phi_{z,e_2}(e_2)) = F_1(T(e_2))$, and $T(g) \in F_1(T(e_2))$ (cf. Lemma 2.1(g)). Lemma 2.7 implies that $T(g) = \Phi_{z,e_2}(g)$, and hence (2.1) writes in the form

$$T(z) = \Phi_{z,e_2}(z) = \lambda_1 \Phi_{z,e_2}(e_1) + \mu T(g) + \lambda_2 T(e_2).$$

The last identity implies that $P_2(T(e_2))T(z) = \lambda_2 T(e_2)$, $P_1(T(e_2))T(z) = \mu T(g)$, and $P_0(T(e_2))T(z) = \lambda_1 \Phi_{z,e_2}(e_1)$.

The identity

$$T(z) = \Phi_{z,e_1}(z) = \lambda_1 T(e_1) + \mu \Phi_{z,e_1}(g) + \lambda_2 \Phi_{z,e_1}(e_2),$$

shows that $P_2(T(e_1))T(z) = \lambda_1 T(e_1)$.

Finally, having in mind that $T(z) \in F_2(T(e_1 + e_2)) = F_2(T(e_1) + T(e_2))$, and $F_0(T(e_2)) \cap F_2(T(e_1 + e_2)) = F_2(T(e_1))$ (cf. [25, 1.12]), we have $\Phi_{z,e_2}(e_1) = T(e_1)$. \square

Lemma 2.10. *Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism from a JBW^* -triple into a JB^* -triple, and let e be a tripotent in E . Then the following statements hold:*

- (a) $T(e + y + z) = T(e) + T(y) + T(z)$, for every $y \in E_1(e)$, and every $z \in E_0(e)$ with $\|z\| < 1$;

- (b) $T(y + z) = T(y) + T(z)$, for every $y \in E_1(e)$, and every $z \in E_0(e)$;
(c) $T(e + y + z) = T(e) + T(y) + T(z)$, for every $y \in E_1(e)$, and every $z \in E_0(e)$. Consequently, $T(\lambda e + y + z) = \lambda T(e) + T(y) + T(z)$, for every $y \in E_1(e)$, every $z \in E_0(e)$, and every $\lambda \in \mathbb{C}$.

Proof. Throughout the proof we set $w = e + y + z$.

(a) We assume first that $T(e) = 0$. By Lemma 2.1(g), $T(y) = 0$. By Lemma 2.8, $\Phi_{w,e+z}(e) = T(e) = 0$ and hence $\Phi_{w,e+z}(y) \in F_1(\Phi_{w,e+z}(e)) = \{0\}$. If we write

$$\begin{aligned} T(e + y + z) &= T(w) = \Phi_{w,e+z}(w) = T(e + z) + \Phi_{w,e+z}(y) \\ &= T(e + z) = (\text{by Proposition 2.5}) = T(e) + T(z) = 0. \end{aligned}$$

Suppose now that $T(e) \neq 0$. Proposition 2.5 implies that

$$T(w) = \Phi_{w,e+z}(w) = T(e + z) + \Phi_{w,e+z}(y) = T(e) + T(z) + \Phi_{w,e+z}(y).$$

By Lemma 2.8, $\Phi_{w,e+z}(e) = T(e)$, and in particular $\Phi_{w,e+z}(y) \in F_1(T(e))$. We also have

$$T(w) = \Phi_{w,y}(w) = T(y) + \Phi_{w,y}(e + z),$$

and hence

$$\Phi_{w,y}(e + z) = T(e) + \Phi_{w,e+z}(y) - T(y) + T(z).$$

Having in mind that $\|\Phi_{w,y}(e + z)\| \leq 1$, Lemma 2.7 implies that

$$0 = P_1(T(e))\Phi_{w,y}(e + z) = \Phi_{w,e+z}(y) - T(y).$$

(b) Since T is 1-homogeneous, we may assume without loss of generality that $\|z\| < 1$. As in the previous case, let us assume that $T(e) = 0$. Under these assumptions, Lemma 2.8 implies that $\Phi_{y+z,e+z}(e + y + z - e)(e) = T(e) = 0$ and hence $\Phi_{y+z,e+z}(E_2(e) \oplus E_1(e)) = \{0\}$. Then

$$\begin{aligned} T(y + z) &= \Phi_{y+z,e+z}(e + y + z - e) = T(e + z) + \Phi_{y+z,e+z}(y - e) \\ &= T(e + z) = (\text{by Proposition 2.5}) = T(e) + T(z) = T(z). \end{aligned}$$

We consider now the case $T(e) \neq 0$. Since we are assuming $\|z\| < 1$, it follows from (a) that

$$\begin{aligned} T(y + z) &= \Phi_{y+z,w}(e + y + z - e) = \Phi_{y+z,w}(e + y + z) - \Phi_{y+z,w}(e) \\ &= T(e) + T(y) + T(z) - \Phi_{y+z,w}(e). \end{aligned}$$

Considering that $T(y+z) \in F_1(T(e)) + F_0(T(e))$ (see Lemma 2.1(f)) we have $P_2(T(e))\Phi_{y+z,w}(e) = T(e)$. Lemma 2.7, applied in the identity $\Phi_{y+z,w}(e) = T(e) + T(y) - T(y + z) + T(z)$, shows that $P_1(T(e))T(y + z) = T(y)$.

By Corollary 2.5, we get

$$\begin{aligned} T(y + z) &= \Phi_{y+z,e+z}(e + z) + \Phi_{y+z,e+z}(y - e) = T(e + z) + \Phi_{y+z,e+z}(y - e) \\ &= T(e) + T(z) + \Phi_{y+z,e+z}(y) - \Phi_{y+z,e+z}(e). \end{aligned}$$

By assumptions $\|z\| < 1$. Thus, applying Lemma 2.8 we show $\Phi_{y+z, e+z}(e) = T(e)$. Therefore, $\Phi_{y+z, e+z}(y) \in F_1(T(e))$ and $P_0(T(e))T(y+z) = T(z)$, which proves that

$$T(y+z) = P_1(T(e))T(y+z) + P_0(T(e))T(y+z) = T(y) + T(z).$$

(c) We begin with the case $T(e) \neq 0$. For each real number $\lambda \in [0, 1]$, we denote $w_\lambda := e + y + \lambda z$. By the assumptions on T

$$T(w_\lambda) = \Phi_{w_\lambda, e}(w_\lambda) = T(e) + \Phi_{w_\lambda, e}(y) + \lambda\Phi_{w_\lambda, e}(z),$$

where $\Phi_{w_\lambda, e}(y) \in F_1(T(e))$, and $\Phi_{w_\lambda, e}(z) \in F_0(T(e))$. Applying (b) we deduce that

$$T(w_\lambda) = \Phi_{w_\lambda, y+\lambda z}(w_\lambda) = T(y) + \lambda T(z) + \Phi_{w_\lambda, y+\lambda z}(e).$$

The above identities show that

$$\Phi_{w_\lambda, y+\lambda z}(e) = T(e) + (\Phi_{w_\lambda, e}(y) - T(y)) + \lambda(\Phi_{w_\lambda, e}(z) - T(z)),$$

and Lemma 2.7 applies to assure that $\Phi_{w_\lambda, e}(y) = T(y)$. Therefore,

$$\Phi_{w_\lambda, y+\lambda z}(e) = T(e) + \lambda(\Phi_{w_\lambda, e}(z) - T(z)).$$

Since $\Phi_{w_\lambda, y+\lambda z}(e)$ is a tripotent, we deduce that

$$P_0(T(e))\Phi_{w_\lambda, y+\lambda z}(e) = \lambda(\Phi_{w_\lambda, e}(z) - T(z)) = P_0(T(e))(T(w_\lambda)) - \lambda T(z)$$

is a tripotent.

Finally the mapping $f : [0, 1] \rightarrow \{0, 1\}$ given by

$$f(\lambda) = \|P_0(T(e))(T(w_\lambda)) - \lambda T(z)\|$$

is (norm) continuous and $f(0) = 0$, then $f(\lambda) = 0$ for every $\lambda \in [0, 1]$, and hence $\Phi_{w_\lambda, e}(z) = T(z)$, which gives the desired statement.

Suppose, finally, that $T(e) = 0$. Lemma 2.1(g) implies that $T(y) = 0$. Let us observe that $\Phi_{w_\lambda, e}(e) = T(e) = 0$. The identities

$$T(w_\lambda) = \Phi_{w_\lambda, e}(w_\lambda) = T(e) + \Phi_{w_\lambda, e}(y) + \lambda\Phi_{w_\lambda, e}(z) = \lambda\Phi_{w_\lambda, e}(z),$$

$$T(w_\lambda) = \Phi_{w_\lambda, y+\lambda z}(w_\lambda) = \Phi_{w_\lambda, y+\lambda z}(e) + T(y) + \lambda T(z) = \Phi_{w_\lambda, y+\lambda z}(e) + \lambda T(z),$$

show that

$$\Phi_{w_\lambda, y+\lambda z}(e) = \lambda\Phi_{w_\lambda, e}(z) - \lambda T(z) = T(w_\lambda) - \lambda T(z),$$

for every $\lambda \in [0, 1]$. Since, for every $0 \leq \lambda \leq 1$, $\Phi_{w_\lambda, y+\lambda z}(e)$ is a tripotent, the function $f : [0, 1] \rightarrow \mathbb{R}$, $f(\lambda) := \|\Phi_{w_\lambda, y+\lambda z}(e)\| = \|T(w_\lambda) - \lambda T(z)\|$ is continuous and takes only the values 0 and 1. Since $f(0) = 0$, we conclude that $f(\lambda) = 0$, for every $\lambda \in [0, 1]$, which proves $T(e+y+z) - T(z) = 0$. \square

We recall that, given a conjugation (conjugate linear isometry of period 2), σ , on a complex Hilbert space H with $\dim(H) = n \in \mathbb{N} \cup \{\infty\}$, the mapping $x \mapsto x^t := \sigma x^* \sigma$ defines a linear involution on $L(H)$. The type-3 Cartan factor, denoted by III_n , is the subtriple of $L(H)$ formed by the t -symmetric operators. Following standard notation, $S_2(\mathbb{C})$ will denote III_2 .

Corollary 2.11. *Let F be a JB^* -triple and let $T : C \rightarrow F$ be a (not necessarily linear) 2-local triple-homomorphism, where C is $M_2(\mathbb{C})$ or $S_2(\mathbb{C})$. Then T is linear and a triple homomorphism.*

Proof. Suppose first that $C = M_2(\mathbb{C})$. We set $e_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $e_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, $y_1 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, and $y_2 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$. Lemma 2.10(c) implies that

$$\begin{aligned} T(\lambda_1 e_1 + \mu_1 y_1 + \mu_2 y_2 + \lambda_2 e_2) &= \lambda_1 T(e_1) + T(\mu_1 y_1 + \mu_2 y_2) + \lambda_2 T(e_2) \\ &= (\text{Proposition 2.5 applied to } y_1 \perp y_2) = \\ &= \lambda_1 T(e_1) + \mu_1 T(y_1) + \mu_2 T(y_2) + \lambda_2 T(e_2). \end{aligned}$$

The linearity follows from the fact that $\{e_1, y_1, y_2, e_2\}$ is a basis of $M_2(\mathbb{C})$.

For the statement concerning $S_2(\mathbb{C})$, we observe that we can assume that $\sigma : H = \ell_2^2 \rightarrow H = \ell_2^2$ is the mapping given by $\sigma(t_1, t_2) = (\bar{t}_1, \bar{t}_2)$. Considering $e_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $e_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, and $y = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, Lemma 2.10(c) implies that

$$T(\lambda_1 e_1 + \mu y + \lambda_2 e_2) = \lambda_1 T(e_1) + \mu T(y) + \lambda_2 T(e_2),$$

which proves that T is linear. \square

Proposition 2.12. *Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism from a JBW^* -triple into a JB^* -triple, and let e be a tripotent in E . Then $T(x + y) = T(x) + T(y)$, for all $x \in E_2(e)$, $y \in E_1(e)$.*

Proof. Let us observe that by Lemma 2.1(g), we may assume that $T(e) \neq 0$. By the norm density of algebraic elements in $E_2(e)$ (cf. [25, lemma 3.11]), together with the continuity of T , it is enough to prove that, for

every algebraic element a in $E_2(e)$ (i.e. $a = \sum_{k=1}^m \lambda_k e_k$, where $\lambda_k \in \mathbb{R}$, and e_1, \dots, e_{m_1} are mutually orthogonal tripotents in $E_2(e)$), we have $T(a + y) = T(a) + T(y)$. We shall prove this statement by induction on the number m of mutually orthogonal tripotents whose linear combination coincides with a .

For the case $m = 1$, we may assume that $a = \lambda_1 e_1 \in E_2(e)$ with $\lambda_1 \neq 0$. Since $y \in E_1(e)$, it follows from Peirce rules that y writes in the form $y = y_1 + y_0$, where $y_k = P_k(e_1)y$, $k = 1, 0$. By Lemma 2.10(c),

$$T(a + y) = T(\lambda_1 e_1 + y_1 + y_0) = T(\lambda_1 e_1) + T(y_1) + T(y_0),$$

and by Lemma 2.10(b), $T(y_1) + T(y_0) = T(y_1 + y_0) = T(y)$, then $T(a + y) = T(a) + T(y)$.

Suppose, by the induction hypothesis, that for every algebraic element b in $E_2(e)$ which is a linear combination of m mutually orthogonal tripotents in $E_2(e)$, we have

$$T(b + y) = T(b) + T(y),$$

for every $y \in E_1(e)$. Let $a = \sum_{i=1}^{m+1} \lambda_i e_i$ be an algebraic element in $E_2(e)$,

and denote by f the tripotent $\sum_{i=1}^{m+1} e_i$. Applying the Peirce decompositions of $a + y$ associated with f and e_1 , we have $a + y = a + y_1 + y_0$, where $y_1 = P_1(f)y$ and $y_0 = P_0(f)y$, and

$$a + y = \lambda_1 e_1 + P_1(e_1)y_1 + \left(\sum_{i=2}^{m+1} \lambda_i e_i + P_0(e_1)y_1 + y_0 \right),$$

where $\left(\sum_{i=2}^{m+1} \lambda_i e_i + P_0(e_1)y_1 + y_0 \right) \in E_0(e_1)$.

Lemma 2.10(c) implies that

$$(2.2) \quad T(a + y) = T(\lambda_1 e_1) + T(P_1(e_1)y_1) + T\left(\sum_{i=2}^{m+1} \lambda_i e_i + P_0(e_1)y_1 + y_0 \right).$$

We observe that $e_1, f \in E_2(e)$, therefore $P_j(e_1)P_k(e) = P_k(e)P_j(e_1)$ and $P_j(f)P_k(e) = P_k(e)P_j(f)$, for every $j, k \in \{0, 1, 2\}$ (cf. [22, Lemma 1.10]).

The induction hypothesis, applied to $\sum_{i=2}^{m+1} \lambda_i e_i \in E_2(e)$ and $P_0(e_1)y_1 + y_0 = P_0(e_1)P_1(f)(y) + P_0(f)(y) = P_0(e_1)P_1(f)P_1(e)(y) + P_0(f)P_1(e)(y) = P_1(e)(P_0(e_1)P_1(f)(y) + P_0(f)(y)) \in E_1(e)$, assures that

$$(2.3) \quad T\left(\sum_{i=2}^{m+1} \lambda_i e_i + P_0(e_1)y_1 + y_0 \right) = T\left(\sum_{i=2}^{m+1} \lambda_i e_i \right) + T(P_0(e_1)y_1 + y_0).$$

Finally, by Lemma 2.4

$$(2.4) \quad T(\lambda_1 e_1) + T\left(\sum_{i=2}^{m+1} \lambda_i e_i \right) = T(a).$$

Since $P_1(e_1)y_1 \in E_1(e_1)$, $P_0(e_1)y_1 + y_0 \in E_0(e_1)$, Lemma 2.10 (b) implies that

$$\begin{aligned} T(P_1(e_1)y_1) + T(P_0(e_1)y_1 + y_0) &= T(P_1(e_1)y_1 + P_0(e_1)y_1 + y_0) \\ &= T(y_1 + y_0) = T(y), \end{aligned}$$

which combined with (2.2), (2.3), and (2.4) prove $T(a+y) = T(a)+T(y)$. \square

Our series of technical results on 2-local triple homomorphisms concludes with an strengthened version of Lemma 2.10.

Lemma 2.13. *Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism from a JBW^* -triple into a JB^* -triple, and let e be a tripotent in E . Then the following statements hold:*

- (a) $T(e + ih + y + z) = T(e) + iT(h) + T(y) + T(z)$, for every $h \in E_2(e)_{sa}$, $y \in E_1(e)$, and every $z \in E_0(e)$ with $\|z\| < 1$;
- (b) $T(ih + y + z) = iT(h) + T(y) + T(z)$, for every $h \in E_2(e)_{sa}$, $y \in E_1(e)$, and every $z \in E_0(e)$;
- (c) $T(e + ih + y + z) = T(e) + iT(h) + T(y) + T(z)$, for every $h \in E_2(e)_{sa}$, $y \in E_1(e)$, and every $z \in E_0(e)$. Consequently,

$$T(\lambda e + ih + y + z) = \lambda T(e) + iT(h) + T(y) + T(z),$$

for every $\lambda \in \mathbb{C}$, $h \in E_2(e)_{sa}$, $y \in E_1(e)$, and every $z \in E_0(e)$.

Proof. Along this proof we set $w = e + ih + y + z$.

We shall assume first that $T(e) = 0$. By Lemma 2.1(g), we have $T(ih) = T(y) = 0$.

(a) It follows from Lemma 2.8 that $\Phi_{w,e+z}(e) = T(e) = 0$, and hence $\Phi_{w,e+z}(E_2(e) \oplus E_1(e)) = \{0\}$. Therefore,

$$\begin{aligned} T(e + ih + y + z) &= \Phi_{w,e+z}(w) = T(e + z) + i\Phi_{w,e+z}(h) + \Phi_{w,e+z}(y) \\ &= (\text{by Proposition 2.5}) = T(e) + T(z) + i\Phi_{w,e+z}(h) + \Phi_{w,e+z}(y) = T(z). \end{aligned}$$

(b) Since T is 1-homogeneous, we may assume, without losing generality, that $\|z\| < 1$. Lemma 2.8 implies that $\Phi_{w-e,e+z}(e) = T(e) = 0$. We write

$$\begin{aligned} T(ih + y + z) &= \Phi_{w-e,e+z}(w - e) = T(e + z) + i\Phi_{w-e,e+z}(h) + \Phi_{w-e,e+z}(y) \\ &= T(e + z) = (\text{by Proposition 2.5}) = T(e) + T(z) = T(z). \end{aligned}$$

(c) Given $\lambda \in [0, 1]$, we set $w_\lambda := e + ih + y + \lambda z$. The identities:

$$T(w_\lambda) = \Phi_{w_\lambda,e}(w_\lambda) = T(e) + i\Phi_{w_\lambda,e}(h) + \Phi_{w_\lambda,e}(y) + \lambda\Phi_{w_\lambda,e}(z) = \lambda\Phi_{w_\lambda,e}(z),$$

and

$$\begin{aligned} T(w_\lambda) &= \Phi_{w_\lambda,ih+y+\lambda z}(w_\lambda) = \Phi_{w_\lambda,ih+y+\lambda z}(e) + T(ih + y + \lambda z) \\ &= (\text{by (b)}) = \Phi_{w_\lambda,ih+y+\lambda z}(e) + iT(h) + T(y) + \lambda T(z) = \Phi_{w_\lambda,ih+y+\lambda z}(e) + \lambda T(z) \end{aligned}$$

assure that

$$\Phi_{w_\lambda,ih+y+\lambda z}(e) = T(w_\lambda) - \lambda T(z).$$

Arguing as in the proof of Lemma 2.10(c) (case $T(e) = 0$), we deduce that $T(w_\lambda) = \lambda T(z)$, for every $0 \leq \lambda \leq 1$.

We suppose from this moment that $T(e) \neq 0$.

(a) We begin with the identity

$$\begin{aligned} T(w) &= \Phi_{w,e+z}(w) = T(e + z) + i\Phi_{w,e+z}(h) + \Phi_{w,e+z}(y) \\ &= (\text{by Proposition 2.5}) = T(e) + T(z) + i\Phi_{w,e+z}(h) + \Phi_{w,e+z}(y). \end{aligned}$$

We deduce, by Lemma 2.8, that $\Phi_{w,e+z}(e) = T(e)$, which, in particular, gives $\Phi_{w,e+z}(y) \in F_1(T(e))$ and $\Phi_{w,e+z}(h) \in F_2(T(e))_{sa}$.

On the other hand,

$$\begin{aligned} T(w) &= \Phi_{w,ih+y}(w) = T(ih+y) + \Phi_{w,ih+y}(e+z) \\ &= (\text{by Proposition 2.12}) = iT(h) + T(y) + \Phi_{w,ih+y}(e+z), \end{aligned}$$

and hence

$$\Phi_{w,ih+y}(e+z) = T(e) + i(\Phi_{w,e+z}(h) - T(h)) + (\Phi_{w,e+z}(y) - T(y)) + T(z).$$

The element $T(e) + i(\Phi_{w,e+z}(h) - T(h))$ lies in the JB*-algebra $F_2(T(e))$ and $(\Phi_{w,e+z}(h) - T(h)) \in F_2(T(e))_{sa}$ (cf. Lemma 2.1(f)) with

$$\|T(e) + i(\Phi_{w,e+z}(h) - T(h))\| \leq \|\Phi_{w,ih+y}(e+z)\| \leq 1.$$

Therefore

$$1 \geq \|T(e) + i(\Phi_{w,e+z}(h) - T(h))\|^2 \geq 1 + \|\Phi_{w,e+z}(h) - T(h)\|^2,$$

witnessing that $\Phi_{w,e+z}(h) = T(h)$. Having in mind that $\|\Phi_{w,ih+y}(e+z)\| \leq 1$, and

$$\Phi_{w,ih+y}(e+z) = T(e) + (\Phi_{w,e+z}(y) - T(y)) + T(z),$$

Lemma 2.7 implies that

$$0 = P_1(T(e))\Phi_{w,ih+y}(e+z) = \Phi_{w,e+z}(y) - T(y).$$

(b) Since T is 1-homogeneous, we may assume, without loss of generality, that $\|z\| < 1$. Denoting $a = ih + y + z$, it follows that

$$\begin{aligned} T(ih+y+z) &= \Phi_{a,w}(e+ih+y+z-e) = \Phi_{a,w}(e+ih+y+z) - \Phi_{a,w}(e) \\ &= T(e+ih+y+z) - \Phi_{a,w}(e) = (\text{by (a)}) = T(e) + iT(h) + T(y) + T(z) - \Phi_{a,w}(e). \end{aligned}$$

On the other hand,

$$\begin{aligned} T(ih+y+z) &= \Phi_{a,e+z}(e+ih+y+z-e) \\ &= T(e+z) + i\Phi_{a,e+z}(h) + \Phi_{a,e+z}(y) - \Phi_{a,e+z}(e) = (\text{by Proposition 2.5}) \\ &= T(e) + T(z) + i\Phi_{a,e+z}(h) + \Phi_{a,e+z}(y) - \Phi_{a,e+z}(e). \end{aligned}$$

We conclude from Lemma 2.8 that $\Phi_{a,e+z}(e) = T(e)$. Therefore $\Phi_{a,e+z}(h) \in F_2(T(e))_{sa}$, $\Phi_{a,e+z}(y) \in F_1(T(e))$,

$$T(ih+y+z) = i\Phi_{a,e+z}(h) + \Phi_{a,e+z}(y) + T(z),$$

and hence

$$\Phi_{a,w}(e) = T(e) + i(T(h) - \Phi_{a,e+z}(h)) + (T(y) - \Phi_{a,e+z}(y)).$$

The arguments given in the final part of the proof of (a) show that $T(h) = \Phi_{a,e+z}(h)$ and $T(y) = \Phi_{a,e+z}(y)$.

(c) For each real number $\lambda \in [0, 1]$, we denote $w_\lambda := e + ih + y + \lambda z$. By the assumptions

$$T(w_\lambda) = \Phi_{w_\lambda,e}(w_\lambda) = T(e) + i\Phi_{w_\lambda,e}(h) + \Phi_{w_\lambda,e}(y) + \lambda\Phi_{w_\lambda,e}(z),$$

where $\Phi_{w_\lambda,e}(h) \in F_2(T(e))_{sa}$, $\Phi_{w_\lambda,e}(y) \in F_1(T(e))$, and $\Phi_{w_\lambda,e}(z) \in F_0(T(e))$. Applying (b) we deduce that

$$T(w_\lambda) = \Phi_{w_\lambda,ih+y+\lambda z}(w_\lambda) = iT(h) + T(y) + \lambda T(z) + \Phi_{w_\lambda,ih+y+\lambda z}(e).$$

The above identities show that

$$\begin{aligned}\Phi_{w_\lambda, ih+y+\lambda z}(e) &= T(e) + i(\Phi_{w_\lambda, e}(h) - T(h)) + (\Phi_{w_\lambda, e}(y) - T(y)) \\ &\quad + \lambda(\Phi_{w_\lambda, e}(z) - T(z)).\end{aligned}$$

Repeating again the arguments given in the final part of the proof of (a) we obtain $T(h) = \Phi_{w_\lambda, e}(h)$ and $T(y) = \Phi_{w_\lambda, e}(y)$. Therefore,

$$\Phi_{w_\lambda, ih+y+\lambda z}(e) = T(e) + \lambda(\Phi_{w_\lambda, e}(z) - T(z)).$$

Since $\Phi_{w_\lambda, ih+y+\lambda z}(e)$ is a tripotent, we deduce that

$$P_0(T(e))\Phi_{w_\lambda, ih+y+\lambda z}(e) = \lambda(\Phi_{w_\lambda, e}(z) - T(z)) = P_0(T(e))(T(w_\lambda)) - \lambda T(z)$$

is a tripotent.

Finally the mapping $f : [0, 1] \rightarrow \{0, 1\}$ given by

$$f(\lambda) = \|P_0(T(e))(T(w_\lambda)) - \lambda T(z)\|$$

is (norm) continuous and $f(0) = 0$. Then $f(\lambda) = 0$ for every $\lambda \in [0, 1]$, and hence $\Phi_{w_1, e}(z) = T(z)$, which gives the desired statement. \square

3. 2-LOCAL TRIPLE HOMOMORPHISMS ON A JBW*-ALGEBRA OR ON A JBW*-TRIPLE

In this section we establish the main results of the paper. Our study on 2-local triple homomorphisms will culminate in a result asserting that every (not necessarily linear) 2-local triple homomorphism from a JBW*-triple into a JB*-triple is linear and a triple homomorphism. In a first step we consider 2-local triple homomorphisms whose domains are JBW*-algebras.

3.1. 2-local triple homomorphisms on a JBW*-algebra. The aim of this subsection is to study 2-local triple homomorphisms from a JBW*-algebra or from a von Neumann algebra into a JB*-triple. The results in these settings are interesting by themselves but also play a crucial role in the proof of our main result for JBW*-triples.

Let $\Phi : \mathcal{J} \rightarrow F$ be a triple homomorphism from a unital JB*-algebra into a JB*-triple. Clearly $\Phi(1)$ is a tripotent in F and $F_2(\Phi(1))$ is a JB*-algebra with unit $\Phi(1)$. Given a in \mathcal{J} , the identities

$$\{\Phi(1), \Phi(1), \Phi(a)\} = \Phi\{1, 1, a\} = \Phi(a),$$

and

$$\Phi(a)^{\sharp_{\Phi(1)}} = \{\Phi(1), \Phi(a), \Phi(1)\} = \Phi\{1, a, 1\} = \Phi(a^*),$$

prove that $\Phi(\mathcal{J}) \subseteq F_2(\Phi(1))$ and $\Phi(\mathcal{J}_{sa}) \subseteq F_2(\Phi(1))_{sa}$. More precisely, Φ is $F_2(\Phi(1))$ -valued and $\Phi : \mathcal{J} \rightarrow F_2(\Phi(1))$ is a unital Jordan *-homomorphism between unital JB*-algebras. For 2-local triple homomorphisms we have:

Lemma 3.1. *Let $T : \mathcal{J} \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism from a unital JB*-algebra into a JB*-triple. Then the following statements hold:*

(a) $T(\mathcal{J}) \subseteq F_2(T(1))$;

- (b) $T(\mathcal{J}_{sa}) \subseteq F_2(T(1))_{sa}$;
(c) $T(a)$ is positive in $F_2(T(1))$ whenever a is positive in \mathcal{J} .

Proof. For each $a \in \mathcal{J}$, and $b \in \mathcal{J}_{sa}$, the comments preceding this lemma assure that $T(a) = \Phi_{1,a}(a) \in F_2(\Phi_{1,a}(1)) = F_2(T(1))$, and $T(b) = \Phi_{1,b}(b) \in F_2(\Phi_{1,b}(1))_{sa} = F_2(T(1))_{sa}$, which proves (a) and (b).

To prove (c), suppose a is a positive element in \mathcal{J} . Since the triple homomorphism $\Phi_{a,1} : \mathcal{J} \rightarrow F_2(\Phi_{a,1}(1)) = F_2(T(1))$ is a unital Jordan *-homomorphism between unital JB*-algebras, $T(a) = \Phi_{1,a}(a)$ is positive in $F_2(\Phi_{a,1}(1)) = F_2(T(1))$. \square

Let \mathcal{J} be a JB*-algebra and let E be a JB*-triple. Following the notation employed in [2] and [10], a *quasi-linear functional* on \mathcal{J} is a function $\rho : \mathcal{J} \rightarrow \mathbb{C}$ such that

- (i) $\rho|_{\mathcal{J}_{\langle h \rangle}} : \mathcal{J}_{\langle h \rangle} \rightarrow \mathbb{C}$ is a linear functional for each $h \in \mathcal{J}_{sa}$, where $\mathcal{J}_{\langle h \rangle}$ denotes the JB*-subalgebra generated by h ;
(ii) $\rho(a + ib) = \rho(a) + i\rho(b)$, for every $a, b \in \mathcal{J}_{sa}$.

If we also assume that, for each $h \in \mathcal{J}_{sa}$, $\rho|_{\mathcal{J}_{\langle h \rangle}}$ is a positive linear functional, we shall say that ρ is *positive quasi-linear functional* on \mathcal{J} . A mapping $\rho : E \rightarrow \mathbb{C}$ is said to be a *quasi-linear functional* on E if for every a in E , the restriction of ρ to the JB*-subtriple, E_a , of E generated by a is linear.

Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism between JB*-triples. For each $\phi \in F^*$, Lemma 2.2 assures that $\phi \circ T : E \rightarrow \mathbb{C}$ is a quasi-linear functional on E in the triple sense. Our next proposition shows that a stronger property holds for 2-local triple homomorphisms from a JBW*-algebra into a JB*-triple.

Proposition 3.2. *Let $T : \mathcal{J} \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism from a JBW*-algebra into a JB*-triple. Then*

$$T(a + ib) = T(a) + iT(b),$$

for every $a, b \in \mathcal{J}_{sa}$.

Proof. It is known that every $a \in \mathcal{J}_{sa}$, can be approximated in norm by a finite (real) linear combination of mutually orthogonal (non-zero) projections in \mathcal{J} ([24, Proposition 4.2.3]). Since T is continuous, it is enough to prove that

$$(3.1) \quad T(a + ib) = T(a) + iT(b),$$

for every $b \in \mathcal{J}_{sa}$ and $a = \sum_{k=1}^m \lambda_k p_k$, where $\lambda_k \in \mathbb{R} \setminus \{0\}$ and p_1, \dots, p_m are mutually orthogonal projections in \mathcal{J} . We shall prove (3.1) by induction on m .

For the case $m = 1$, we assume that $a = \lambda p$ for a non-zero projection p and $\lambda \in \mathbb{R} \setminus \{0\}$. The element b writes in the form $b = P_2(p)(b) + P_1(p)(b) +$

$P_0(p)(b)$, and since $b = b^*$ and p is a projection, $P_2(p)(b) \in \mathcal{J}_{sa}$. Applying Lemma 2.13(c), we have

$$\begin{aligned} T(a + ib) &= \lambda T(p) + iT(P_2(p)(b)) + iT(P_1(p)(b)) + iT(P_0(p)(b)) \\ &= T(a) + iT(P_2(p)(b)) + iT(P_1(p)(b)) + iT(P_0(p)(b)), \end{aligned}$$

by Lemma 2.13(b),

$$= T(a) + T(iP_2(p)(b) + iP_1(p)(b) + iP_0(p)(b)) = T(a + ib).$$

Suppose, by the induction hypothesis, that (3.1) is true for every algebraic element $\sum_{k=1}^{m_1} \mu_k q_k$, with $m_1 \leq m$, $\lambda_k \in \mathbb{R} \setminus \{0\}$ and q_1, \dots, q_{m_1} mutually orthogonal projections in \mathcal{J} . Let us take an algebraic element of the form $a = \sum_{k=1}^{m+1} \lambda_k p_k$. Let us write $b = P_2(p_1)(b) + P_1(p_1)(b) + P_0(p_1)(b)$, and since $b = b^*$ and p_1 is a projection, $P_2(p_1)(b), P_0(p_1)(b) \in \mathcal{J}_{sa}$. Let us observe that

$$T(a + ib) = T\left(\lambda_1 p_1 + iP_2(p_1)(b) + iP_1(p_1)(b) + \sum_{k=2}^{m+1} \lambda_k p_k + iP_0(p_1)(b)\right),$$

where $P_2(p_1)(b) \in \mathcal{J}_2(p_1)_{sa}$, $iP_1(p_1)(b) \in \mathcal{J}_1(p_1)$, and $\sum_{k=2}^{m+1} \lambda_k p_k + iP_0(p_1)(b)$ lies in $\mathcal{J}_0(p_1)$. Lemma 2.13(c) implies that

$$\begin{aligned} T(a + ib) &= \lambda_1 T(p_1) + iT(P_2(p_1)(b)) + T(iP_1(p_1)(b)) \\ &+ T\left(\sum_{k=2}^{m+1} \lambda_k p_k + iP_0(p_1)(b)\right) = (\text{by the induction hypothesis}) = \\ &= \lambda_1 T(p_1) + iT(P_2(p_1)(b)) + T(iP_1(p_1)(b)) + T\left(\sum_{k=2}^{m+1} \lambda_k p_k\right) + T(iP_0(p_1)(b)) \\ &= (\text{by Proposition 2.5}) = T\left(\lambda_1 p_1 + \sum_{k=2}^{m+1} \lambda_k p_k\right) + iT(P_2(p_1)(b)) \\ &+ T(iP_1(p_1)(b)) + T(iP_0(p_1)(b)) = (\text{by Lemma 2.13(b) with } e = p_1) \\ &= T(a) + iT(P_2(p_1)(b) + P_1(p_1)(b) + P_0(p_1)(b)) = T(a) + iT(b). \end{aligned}$$

□

We can establish now a generalization of [14, Theorem 4.1] for 2-local triple homomorphisms.

Theorem 3.3. *Let \mathcal{J} be a JBW^* -algebra with no Type I_2 direct summand and let F be a JB^* -triple. Suppose $T : \mathcal{J} \rightarrow F$ is a (not necessarily linear) 2-local triple homomorphism. Then T is linear and a (continuous) triple homomorphism. More concretely, $T : \mathcal{J} \rightarrow F_2(T(1))$ is a linear unital Jordan $*$ -homomorphism between JB^* -algebras.*

Proof. We know that $T(1)$ is a tripotent in F (cf. Lemma 2.1). By Lemma 3.1 $T(\mathcal{J}) \subseteq F_2(T(1))$ and $T(\mathcal{J}_{sa}) \subseteq F_2(T(1))_{sa}$.

Furthermore, given a projection p in \mathcal{J} , since $p \leq 1$, Lemma 2.1(b) and (d) assure that $T(p)$ is a tripotent in $F_2(T(1))$ with $T(p) \leq T(1)$, which implies that $T(p)$ is a projection in $F_2(T(1))$.

Fix an arbitrary norm-one positive functional φ in $F_2(T(1))^*$. Let $\mathcal{P}(\mathcal{J})$ denote the lattice of projections in \mathcal{J} . The mapping

$$\begin{aligned} \mu_\varphi : \mathcal{P}(\mathcal{J}) &\rightarrow \mathbb{R} \\ \mu_\varphi(p) &:= \varphi(T(p)), \end{aligned}$$

is a finitely additive quantum measure on $\mathcal{P}(\mathcal{J})$ in the terminology employed in [9], i.e. $\mu_\varphi(1) = 1$ and $\mu_\varphi(p_1 + \dots + p_m) = \mu_\varphi(p_1) + \dots + \mu_\varphi(p_m)$, whenever p_1, \dots, p_m are mutually orthogonal projections in \mathcal{J} (this statement follows from Lemma 2.4(a) and the fact that $T(1)$ is the unit element in $F_2(T(1))$). By the Bunce-Wright-Mackey-Gleason theorem [9, Theorem 2.1], there exists a positive linear functional $\phi_\varphi \in \mathcal{J}_{sa}^*$ such that

$$\varphi(T(p)) = \mu_\varphi(p) = \phi_\varphi(p),$$

for every $p \in \mathcal{P}(\mathcal{J})$. It follows from Lemma 2.4(b), the continuity of T , φ , and ϕ_φ , and the norm density of algebraic elements in \mathcal{J}_{sa} that

$$\varphi(T(a)) = \phi_\varphi(a),$$

for every $a \in \mathcal{J}_{sa}$. Therefore,

$$\begin{aligned} \varphi(T(a+b)) &= \phi_\varphi(a+b) = \phi_\varphi(a) + \phi_\varphi(b) \\ &= \varphi(T(a)) + \varphi(T(b)) = \varphi(T(a) + T(b)), \end{aligned}$$

for every $a, b \in \mathcal{J}_{sa}$. Since the positive norm-one functionals in $F_2(T(1))^*$ separate the points of $F_2(T(1))_{sa}$ (cf. [24, Lemma 3.6.8]), we deduce that $T(a+b) = T(a) + T(b)$, for every $a, b \in \mathcal{J}_{sa}$, that is, the restricted mapping $T|_{\mathcal{J}_{sa}} : \mathcal{J}_{sa} \rightarrow F_2(T(1))_{sa} \subseteq F$ is linear.

Finally, Proposition 3.2 shows that $T(a+ib) = T(a) + iT(b)$, for every $a, b \in \mathcal{J}_{sa}$, which gives the desired statement. \square

When in the proof of [14, Corollary 4.4] (respectively, [14, Corollary 2.11]), [14, Proposition 4.2 and Corollary 4.3] (respectively, [14, Proposition 2.7 and Corollary 2.10]) are replaced with Corollary 2.11 and Corollary 2.6, respectively, and having in mind Proposition 3.2, the arguments in those results remain valid to prove:

Corollary 3.4. *Every (not necessarily linear) 2-local triple homomorphism from a Type I_2 JBW*-algebra into a JB*-triple is linear and a triple homomorphism.* \square

The first main result of this note is a consequence of Theorem 3.3, Corollary 3.4 and Corollary 2.6.

Theorem 3.5. *Every (not necessarily linear) 2-local triple homomorphism from a JBW^* -algebra into a JB^* -triple is linear and a triple homomorphism. \square*

The next corollary is interesting by itself.

Corollary 3.6. *Every (not necessarily linear) 2-local triple homomorphism from a von Neumann algebra into a JB^* -triple is linear and a triple homomorphism.*

Since every $*$ -homomorphism between C^* -algebras (respectively, every Jordan $*$ -homomorphism between JB^* -algebras) is a triple homomorphism, Theorems 2.12 and 4.5 and Corollary 4.6 in [14] are direct consequences of the previous Theorem 3.5 and Corollary 3.6.

3.2. 2-local triple homomorphisms on a JBW^* -triple. The rest of the note is devoted to prove the second main result of the paper, in which we shall show that every (not necessarily linear) 2-local triple homomorphism from a JBW^* -triple into a JB^* -triple is linear and a triple homomorphism. The first step toward our goal is the following corollary.

Corollary 3.7. *Let $T : E \rightarrow F$ be a (not necessarily linear) 2-local triple homomorphism from a JBW^* -triple into a JB^* -triple. Then, for each tripotent e in E , $T|_{E_2(e)} : E_2(e) \rightarrow F$ is linear and a triple homomorphism.*

Proof. Clear from Theorem 3.5. \square

We are now in a position to establish the goal of this section.

Theorem 3.8. *Every (not necessarily linear) 2-local triple homomorphism from a JBW^* -triple into a JB^* -triple is linear and a triple homomorphism.*

Proof. Let x, y be two (arbitrary) elements in E . Find a complete tripotent e in E such that $x \in E_2(e)$ (the existence of such a tripotent is guaranteed by [25, Lemma 3.12(1)]). If we write $y = P_2(e)y + P_1(e)y$, then Proposition 2.12 and Corollary 3.7 prove that

$$\begin{aligned} T(x + y) &= T(x + P_2(e)y + P_1(e)y) = T(x + P_2(e)y) + T(P_1(e)y) \\ &= T(x) + T(P_2(e)y) + T(P_1(e)y). \end{aligned}$$

A new application of Proposition 2.12 shows that $T(P_2(e)y) + T(P_1(e)y) = T(P_2(e)y + P_1(e)y)$, and hence $T(x + y) = T(x) + T(y)$. \square

REFERENCES

- [1] S. Albeverio, Sh. Ayupov, K.K. Kudaybergenov, B.O. Nurjanov, Local derivations on algebras of measurable operators, *Commun. Contemp. Math.* **13**, no. 4, 643-657 (2011).
- [2] J.F. Aarnes, Quasi-states on C^* -algebras, *Trans. Amer. Math. Soc.* **149**, 601-625 (1970).
- [3] T.J. Barton, T. Dang, G. Horn, Normal representations of Banach Jordan triple systems, *Proc. Amer. Math. Soc.* **102**, 551-555 (1988).
- [4] T.J. Barton, R.M. Timoney, Weak*-continuity of Jordan triple products and its applications, *Math. Scand.* **59**, 177-191 (1986).

- [5] C. Batty, L. Molnar, On topological reflexivity of the groups of $*$ -automorphisms and surjective isometries of $B(H)$, *Arch. Math.* **67** 415-421 (1996).
- [6] R. Braun, W. Kaup, H. Upmeyer, A holomorphic characterization of Jordan C^* -algebras, *Math. Z.* **161**, no. 3, 277-290 (1978).
- [7] M. Brešar, P. Šemrl, Mappings which preserve idempotents, local automorphisms, and local derivations, *Canad. J. Math.* **45**, no. 3, 483-496 (1993).
- [8] M. Brešar, P. Šemrl, On local automorphisms and mappings that preserve idempotents, *Studia Math.* **113**, no. 2, 101-108 (1995).
- [9] L.J. Bunce, J.D.M. Wright, Continuity and linear extensions of quantum measures on Jordan operator algebras, *Math. Scand.* **64**, 300-306 (1989).
- [10] L.J. Bunce, J.D.M. Wright, The quasi-linearity problem for C^* -algebras, *Pacific J. Math.* **172**, no. 1, 41-47 (1996).
- [11] M. Burgos, F.J. Fernández-Polo, J.J. Garcés, J. Martínez Moreno, A.M. Peralta, Orthogonality preservers in C^* -algebras, JB^* -algebras and JB^* -triples, *J. Math. Anal. Appl.*, **348**, 220-233 (2008).
- [12] M. Burgos, F.J. Fernández-Polo, J.J. Garcés, A.M. Peralta, Orthogonality preservers Revisited, *Asian-European Journal of Mathematics* **2**, No. 3, 387-405 (2009).
- [13] M. Burgos, F.J. Fernández-Polo, J.J. Garcés, A.M. Peralta, Local triple derivations on C^* -algebras, *Communications in Algebra* **42**, 1276-1286 (2014).
- [14] M. Burgos, F.J. Fernández-Polo, J.J. Garcés, A.M. Peralta, A Kowalski-Słodkowski theorem for 2-local $*$ -homomorphisms on von Neumann algebras, preprint 2014.
- [15] M. Burgos, F.J. Fernández-Polo, A.M. Peralta, Local triple derivations on C^* -algebras and JB^* -triples, *Bull. London Math. Soc.*, doi:10.1112/blms/bdu024. arXiv:1303.4569v2.
- [16] Ch.-H. Chu, *Jordan Structures in Geometry and Analysis*, Cambridge Tracts in Math. 190, Cambridge Univ. Press, Cambridge, 2012.
- [17] R. Crist, Local automorphisms, *Proc. Amer. Math. Soc.* **128**, 1409-1414 (1999).
- [18] R. Crist, Local derivations on operator algebras, *J. Funct. Anal.* **135**, 72-92 (1996).
- [19] C.M. Edwards, G.T. Rüttimann, On the facial structure of the unit balls in a JBW^* -triple and its predual, *J. London Math. Soc.* **38**, 317-322 (1988).
- [20] A. Fošner, 2-local Jordan automorphisms on operator algebras, *Studia Math.* **209**, no. 3, 235-246 (2012).
- [21] A. Fošner, 2-local mappings on algebras with involution, *Ann. Funct. Anal.* **5**, no. 1, 63-69 (2014).
- [22] Y. Friedman, B. Russo, Structure of the predual of a JBW^* -triple. *J. Reine Angew. Math.* **356**, 67-89 (1985).
- [23] A.M. Gleason, A characterization of maximal ideals, *J. Analyse Math.* **19**, 171-172 (1967).
- [24] H. Hanche-Olsen, E. Størmer, *Jordan operator algebras, Monographs and Studies in Mathematics 21*, Pitman, London-Boston-Melbourne 1984.
- [25] G. Horn, Characterization of the predual and ideal structure of a JBW^* -triple, *Math. Scand.* **61**, 117-133 (1987).
- [26] B.E. Johnson, Local derivations on C^* -algebras are derivations, *Trans. Amer. Math. Soc.* **353**, 313-325 (2001).
- [27] R.V. Kadison, Local derivations, *J. Algebra* **130**, 494-509 (1990).
- [28] J.P. Kahane, W. Żelazko, A characterization of maximal ideals in commutative Banach algebras, *Studia Math.* **29**, 339-343 (1968).
- [29] W. Kaup, A Riemann mapping theorem for bounded symmetric domains in complex Banach spaces, *Math. Z.* **183**, 503-529 (1983).
- [30] W. Kaup, H. Upmeyer, Jordan algebras and symmetric Siegel domains in Banach spaces, *Math. Z.* **157**, 179-200 (1977).
- [31] S.O. Kim, J.S. Kim, Local automorphisms and derivations on M_n , *Proc. Amer. Math. Soc.* **132**, no. 5, 1389-1392 (2004).

- [32] S.O. Kim, J.S. Kim, Local automorphisms and derivations on certain C^* -algebras, *Proc. Amer. Math. Soc.* **133**, no. 11, 3303-3307 (2005).
- [33] S. Kowalski, Z. Slodkowski, A characterization of multiplicative linear functionals in Banach algebras, *Studia Math.* **67**, 215-223 (1980).
- [34] D.R. Larson and A.R. Sourour, Local derivations and local automorphisms of $B(X)$, *Proc. Sympos. Pure Math.* **51**, Part 2, Providence, Rhode Island 1990, pp. 187-194.
- [35] J.-H. Liu, N.-C. Wong, 2-Local automorphisms of operator algebras, *J. Math. Anal. Appl.* **321** 741-750 (2006).
- [36] M. Mackey, Local derivations on Jordan triples, *Bull. London Math. Soc.* **45**, no. 4, 811-824 (2013). doi: 10.1112/blms/bdt007
- [37] L. Molnar, 2-local isometries of some operator algebras, *Proc. Edinburgh Math. Soc.* **45**, 349-352 (2002).
- [38] L. Molnar, Local automorphisms of operator algebras on Banach spaces, *Proc. Amer. Math. Soc.* **131**, 1867-1874 (2003).
- [39] L. Molnar, *Selected preserver problems on algebraic structures of linear operators and on function spaces*, Lecture Notes in Mathematics, 1895. Springer-Verlag, Berlin, 2007.
- [40] L. Molnar and P. Šemrl, Local automorphisms of the unitary group and the general linear group on a Hilbert space, *Expo. Math.* **18**, 231-238 (2000).
- [41] A.M. Peralta, A note on 2-local representations of C^* -algebras, preprint 2013.
- [42] F. Pop, On local representation of von Neumann algebras, *Proc. Amer. Math. Soc.* **132**, No. 12, Pages 3569-3576 (2004).
- [43] P. Šemrl, Local automorphisms and derivations on $B(H)$, *Proc. Amer. Math. Soc.* **125**, 2677-2680 (1997).
- [44] P. Šemrl, Local automorphisms of standard operator algebras, *J. Math. Anal. Appl.* **371**, no. 2, 403-406 (2010).
- [45] W. Żelazko, A characterization of multiplicative linear functionals in complex Banach algebras, *Studia Math.* **30** 83-85 (1968).
- [46] J. Zhang, F. Pan, A. Yang, Local derivations on certain CSL algebras, *Linear Algebra Appl.* **413**, 93-99 (2006).

E-mail address: maria.burgos@uca.es

DEPARTAMENTO DE MATEMATICAS, FACULTAD DE CIENCIAS SOCIALES Y DE LA EDUCACION, UNIVERSIDAD DE CADIZ, 11405, JEREZ DE LA FRONTERA, SPAIN.

E-mail address: pacopolo@ugr.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, 18071 GRANADA, SPAIN.

E-mail address: jgarces@correo.ugr.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, 18071 GRANADA, SPAIN.

E-mail address: aperalta@ugr.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, 18071 GRANADA, SPAIN.

Current address: Visiting Professor at Department of Mathematics, College of Science, King Saud University, P.O.Box 2455-5, Riyadh-11451, Kingdom of Saudi Arabia.