Dubrovin's conjecture for IG(2,6)

Sergey Galkin, Anton Mellit, Maxim Smirnov

ABSTRACT. We show that the big quantum cohomology of the symplectic isotropic Grassmanian IG(2,6) is generically semisimple, whereas its small quantum cohomology is known to be non-semisimple. This gives yet another case where Dubrovin's conjecture holds and stresses the need to consider the big quantum cohomology in its formulation.

1. Introduction

The main purpose of this paper is to give an explicit example of a smooth projective variety X such that its small quantum cohomology $qH^*(X)$ is not generically semisimple, whereas its big quantum cohomology $QH^*(X)$ is generically semisimple. Namely, we show that this pattern holds for the symplectic isotropic Grassmanian IG(2,6).

In general the small quantum cohomology of isotropic Grassmanians is studied in [BuKrTa], and an explicit presentation in terms of generators and relations is given. Based on this presentation in [ChMaPe, Sec. 7] it is shown that $qH^*(IG(2,6))$ is not generically semisimple.

The Grassmanian IG(2,6) appears to be the simplest explicit example of this sort available in the literature. More examples will appear in a forthcoming paper by Nicolas Perrin [Pe], who also independently proved the generic semisimplicity of $QH^*(IG(2,6))$ simultaneously with us.

- 1.1. **Remark.** Throughout the text we use notation $QH^*(X)$ for the big quantum cohomology of X and $qH^*(X)$ for the small quantum cohomology of X. We should mention that this notation is not standard, but it is very convenient for our purposes. To avoid any confusion, in Section 2 we will briefly recall both concepts, since they are crucial for this paper.
- 1.2. **Dubrovin's conjecture.** Dubrovin's conjecture (see [Du]) gives an intriguing relation between the quantum cohomology of a smooth projective variety X and the bounded derived category of coherent sheaves on it. Namely, it says that the generic semisimplicity of $QH^*(X)$ is equivalent to the existence of a full exceptional collection in $\mathcal{D}^b(X)$. Here $\mathcal{D}^b(X)$ denotes the bounded derived category of coherent sheaves on X.

The main motivation for studying examples like IG(2,6) comes from the conjecture above. In [Ku] it is shown that $\mathcal{D}^b(IG(2,6))$ has a full exceptional collection. Hence, it is expected that $QH^*(IG(2,6))$ is generically semisimple, whereas $qH^*(IG(2,6))$ is not.

For completeness let us just mention that one expects to find many more such examples among homogeneous spaces G/P, where G is a semi-simple algebraic group and P is a parabolic subgroup. The reason is that $\mathcal{D}^b(G/P)$ is conjectured to have a full exceptional collection (see [KuPo, Conj. 1.1]). In particular, for arbitrary symplectic isotropic Grassmanians IG(k,2n) the big quantum cohomology is expected to be generically semisimple, and one can try to find a pattern when $qH^*(IG(k,2n))$ is not. For results in this direction related to semisimplicity we refer to [ChPe, Sec. 6].

We should remark that the above formulation of Dubrovin's conjecture is not complete. Here we consider its first part only. The remaining two parts (see [Du, Conj. 4.2.2] and [GaGoIr]) are not considered in this paper. Let us just mention that for Grassmanians G(k, n) all three parts are known to hold (see [GaGoIr, Ue] and references therein). For a general introduction to this topic we refer to [Du, Ba, GaGoIr, HeMaTe].

1.2.1. Some known examples. Here we list some instances where Dubrovin's conjecture is known to hold. The simplest example is provided by the projective spaces \mathbb{P}^n . Indeed, it is well-known that $qH^*(\mathbb{P}^n)$ is generically semisimple (for example, see [KoMa]), and $\mathcal{D}^b(\mathbb{P}^n)$ has a full exceptional collection (see [Be]).

One way to generalize the example of \mathbb{P}^n is to look at Grassmanians G(k, n). By [Ab, Prop. 6.5] (or [GaGo, Cor. 8]) it is known that $qH^*(G(k, n))$ is generically semisimple, and by [K] we know that $\mathcal{D}^b(G(k, n))$ has a full exceptional collection.

Another way to generalize the example of \mathbb{P}^n is to consider arbitrary toric varieties. For a smooth projective toric variety X (no Fano assumption!) it is known that $QH^*(X)$ is generically semisimple (see

1

[Ir]). Note that from loc.cit. it is not known whether $qH^*(X)$ is generically semisimple for an arbitrary toric variety X, but only when X is a weak Fano variety. A full exceptional collection in $\mathcal{D}^b(X)$ exists for any toric X by [Ka].

In the above examples already the small quantum cohomology is generically semisimple. As far as we know IG(2,6) is the first explicit example in the literature where one really needs the big quantum cohomology in order to formulate Dubrovin's conjecture.

Acknowledgements. We are very grateful to Tarig Abdelgadir, Boris Dubrovin, Vasily Golyshev, Vassily Gorbounov, Hiroshi Iritani, Alexander Kuznetsov, Yuri Manin, Nicolas Perrin, and Evgeny Shinder for valuable discussions, comments, and attention to this work.

2. Reminder on quantum cohomology

Here we briefly recall the definition of the quantum cohomology ring for a smooth projective variety X. It is convenient for us to define it in a fixed basis of $H^*(X,\mathbb{Q})$ but, of course, the resulting ring structure is independent of it. We assume some familiarity with the subject and do not go into much detail. For a thorough introduction we refer to [Ma].

2.1. **Definition.** Let X be a smooth projective variety and assume for simplicity that $H^{odd}(X,\mathbb{Q}) = 0$. Fix a graded basis $\Delta_0, \ldots, \Delta_n$ in $H^*(X,\mathbb{Q})$ and dual linear coordinates t_0, \ldots, t_n . It is customary to choose $\Delta_0 = 1$. Let N_X be the Novikov ring of X, i.e. the ring of formal power series in q^{β} , where β runs over the Mori cone of curves on X.

The genus zero Gromov-Witten potential of X is an element $\Phi \in N_X[[t_0, \ldots, t_n]]$, i.e. a formal power series in variables t_0, \ldots, t_n with coefficients in the Novikov ring N_X , defined by the formula

$$\Phi = \sum_{(i_0, \dots, i_n)} \langle \Delta_0^{\otimes i_0}, \dots, \Delta_n^{\otimes i_n} \rangle \frac{t_0^{i_0} \dots t_n^{i_n}}{i_0! \dots i_n!}, \tag{2.1}$$

where $\langle \Delta_0^{\otimes i_0}, \dots, \Delta_n^{\otimes i_n} \rangle := \sum_{\beta} \langle \Delta_0^{\otimes i_0}, \dots, \Delta_n^{\otimes i_n} \rangle_{\beta} q^{\beta}$, and β runs over the cone of effective curve classes. The rational numbers $\langle \Delta_0^{\otimes i_0}, \dots, \Delta_n^{\otimes i_n} \rangle_{\beta}$ are Gromov-Witten invariants of X of the curve class β .

Using (2.1) one defines the quantum cohomology ring of X. Namely, on the basis elements we put

$$\Delta_a \star \Delta_b = \sum_c \Phi_{abc} \Delta^c, \tag{2.2}$$

where $\Phi_{abc} = \frac{\partial^3 \Phi}{\partial t_a \partial t_b \partial t_c}$, and $\Delta^0, \dots, \Delta^n$ is the basis dual to $\Delta_0, \dots, \Delta_n$ with respect to the Poincaré pairing. Expression (2.2) is naturally interpreted as an element of $H^*(X, \mathbb{Q}) \otimes_{\mathbb{Q}} N_X[[t_0, \dots, t_n]]$.

It is well known that (2.2) makes $H^*(X,\mathbb{Q}) \otimes_{\mathbb{Q}} N_X[[t_0,\ldots,t_n]]$ into a commutative, associative, graded $N_X[[t_0,\ldots,t_n]]$ -algebra with the identity element Δ_0 . We will denote this algebra $QH^*(X)$. For convenience we recall the definition of the grading:

$$\deg(\Delta_i) = \frac{|\Delta_i|}{2}, \quad \deg(q^\beta) = (-K_X, \beta), \quad \deg(t_i) = 1 - \frac{|\Delta_i|}{2},$$

where $|\Delta_i|$ is the cohomological degree of Δ_i .

Sometimes the algebra $QH^*(X)$ is called the *big quantum cohomology algebra* of X to distinguish it from a simpler object called the *small quantum cohomology algebra*. It is the quotient of $QH^*(X)$ with respect to the ideal (t_0, \ldots, t_n) . We will denote the latter $qH^*(X)$ and use \circ instead of \star for the product in this algebra.

It is equivalent to say that $qH^*(X) = H^*(X, \mathbb{Q}) \otimes_{\mathbb{Q}} N_X$ as a vector space, and the N_X -algebra structure is defined by putting $\Delta_a \circ \Delta_b = \sum_c \langle \Delta_a, \Delta_b, \Delta_c \rangle \Delta^c$.

- 2.1.1. **Remark.** (i) Often in the literature one uses $QH^*(X)$ both for big and small quantum cohomology. To avoid any confusion we have decided to use the non-standard notation described above.
- (ii) At first sight our definition of the small quantum cohomology, i.e. the ring structure defined by 3-pointed GW invariants, differs slightly from [Ma]. It is not hard to see that both structures are equivalent.

- (iii) In Section 4 we will use \star to denote the big quantum product in a more restricted sense. Namely, instead of a full deformation we will consider only one-parameter deformation.
- (iv) Throughout the text we work only with the even cohomology. Therefore, all degrees appearing below are Chow-ring degrees.
- 2.2. **Lemma.** Let $\operatorname{Pic} X = \mathbb{Z}, h \in H^2(X, \mathbb{Q})$ and $\gamma \in H^*(X, \mathbb{Q})$. Then

$$\frac{\partial}{\partial t_i}(h \star \gamma) = (h, \beta) \, q \frac{d}{dq}(\Delta_i \star \gamma),$$

where $q = q^{\beta}$ with β being the generator of the cone of effective curves.

Proof. We have the chain of equalities

$$\frac{\partial}{\partial t_i}(h\star\gamma) = \nabla_{\Delta_i}(h\star\gamma) = \nabla_h(\Delta_i\star\gamma) = (h,\beta)\,q\,\frac{d}{dq}(\Delta_i\star\gamma),$$

where the first one holds by definition, the second follows from potentiality of the quantum product (see the proof of Proposition 1.6 in [Ma]), and the last one holds due to the divisor axiom for GW invariants.

2.3. Corollary. For an element α in $H^*(X,\mathbb{Q})$ let M_{α} and \widetilde{M}_{α} be the matrices of the small and the big quantum multiplication by α in the basis $\Delta_0, \ldots, \Delta_n$. Moreover, to simplify notation, put $M_i = M_{\Delta_i}$ and $\widetilde{M}_i = \widetilde{M}_{\Delta_i}$.

From Lemma 2.2 it follows that for $h \in H^2(X, \mathbb{Q})$ we have

$$\widetilde{M}_h = M_h + (h, \beta) q \frac{d}{dq} \sum_i \int_0^{t_i} \widetilde{M}_i dt_i, \qquad (2.3)$$

and restricting to the first infinitesimal neighbourhood we get

$$\widetilde{M}_h = M_h + (h, \beta) q \frac{d}{dq} \sum_i t_i M_i + O(t_a t_b).$$
(2.4)

3. Small quantum cohomology of IG(2,6)

The aim of this paragraph is to fix notation and to describe the structure of the small quantum cohomology ring of IG(2,6) following [ChMaPe].

3.1. Geometry of IG(2,6). Let V be a six-dimensional complex vector space and $\omega \in \Lambda^2 V^*$ a symplectic form. The Grassmanian IG(2,V) parametrizes two-dimensional isotropic subspaces of V. It is a smooth projective variety of dimension dim X=7.

One can give an explicit presentation of IG(2,V) as a hyperplane section of G(2,V). Indeed, consider the Grassmanian G(2,V), and its Plücker embedding into $\mathbb{P}(\Lambda^2 V) \simeq \mathbb{P}^{14}$. The form ω defines a hyperplane in $\mathbb{P}(\Lambda^2 V)$. The section of G(2,V) by this hyperplane is X = IG(2,6).

Similarly to the case of usual Grassmanians, there exists a basis of cohomology indexed by the so called 1-strict partitions λ (see [ChMaPe] and references therein). The cohomology class corresponding to such a partition λ will be denoted Δ_{λ} . Explicitly, there are 12 such classes

$$\Delta_0; \quad \Delta_1; \quad \Delta_2, \Delta_{1,1}; \quad \Delta_3, \Delta_{2,1}; \quad \Delta_4, \Delta_{3,1}; \quad \Delta_{4,1}, \Delta_{3,2}; \quad \Delta_{4,2}; \quad \Delta_{4,3},$$
 (3.1)

and the Chow-ring degree of Δ_{λ} is $|\lambda| = \lambda_1 + \lambda_2$.

3.2. **Gromov-Witten invariants.** Recall that Pic $X = \mathbb{Z}$ and let l be the generator of the cone of effective curves on X. Since $-K_X = 5\Delta_1$, the dimension axiom for an n-pointed GW invariant $\langle \gamma_1, \ldots, \gamma_n \rangle$ becomes

$$n+4+5d = \sum_{i} |\gamma_i|,$$

and implies that $\langle \Delta_2, \Delta_2, \Delta_2 \rangle = \langle \Delta_2, \Delta_2, \Delta_2, \Delta_2 \rangle = 0$.

3.3. Multiplication table. As in [ChMaPe, Sec.7] we represent the small quantum cohomology of X specialized at q = 1 by the table

$$\Delta_{1} \quad \Delta_{2} \quad \Delta_{1,1} \quad \Delta_{2,1} \quad \Delta_{3} \quad \Delta_{3,1} \quad \Delta_{4} \quad \Delta_{4,1} \quad \Delta_{3,2} \quad \Delta_{4,2} \quad \Delta_{4,3}$$

$$Z_{0} \quad 0 \quad \varepsilon \quad -\varepsilon \quad 0 \quad 0 \quad 0 \quad 0 \quad -1 \quad 1 \quad 0 \quad -\varepsilon$$

$$Z_{1} \quad s \quad 0 \quad s^{2} \quad s^{3} \quad -s^{3} \quad s^{4} \quad -s^{4} \quad 0 \quad -1 \quad -s \quad -s^{2}$$

$$Z_{2} \quad t \quad \frac{2}{3}t^{2} \quad \frac{1}{3}t^{2} \quad \frac{1}{3}t^{3} \quad \frac{1}{3}t^{3} \quad \frac{1}{9}t^{4} \quad \frac{1}{9}t^{4} \quad 2 \quad 1 \quad t \quad \frac{1}{3}t^{2}$$

$$(3.2)$$

Some explanations are in order. The small quantum cohomology of X specialized at q=1 is a finite dimensional algebra A. Explicitly it is given as a direct product

$$A = A_0 \times A_1 \times A_2,$$
 (3.3)
 $A_0 = \mathbb{C}[\varepsilon]/\varepsilon^2, \quad A_1 = \mathbb{C}[s]/(s^5 + 1), \quad A_2 = \mathbb{C}[t]/(t^5 - 27).$

Therefore, its spectrum Spec (A) is a disjoint union of three components $Z_i = \text{Spec }(A_i)$. Each element Δ_{λ} is a function on Z = Spec (A), and we can look at its values at different points. Table (3.2) collects these values grouped according to components Z_i .

From (3.2) one can reconstruct all small quantum products. Moreover, using the fact that the quantum multiplication is graded, one can also put the parameter q back. For example, the small quantum product with Δ_1 is

$$\Delta_{1} \circ \Delta_{1} = \Delta_{2} + \Delta_{1,1}
\Delta_{1} \circ \Delta_{2} = \Delta_{3} + \Delta_{2,1}
\Delta_{1} \circ \Delta_{1,1} = \Delta_{2,1}
\Delta_{1} \circ \Delta_{3} = 2\Delta_{4} + \Delta_{3,1}
\Delta_{1} \circ \Delta_{2,1} = \Delta_{4} + 2\Delta_{3,1}
\Delta_{1} \circ \Delta_{4} = \Delta_{4,1} + q\Delta_{0}
\Delta_{1} \circ \Delta_{3,1} = \Delta_{4,1} + \Delta_{3,2}
\Delta_{1} \circ \Delta_{4,1} = \Delta_{4,2} + q\Delta_{1}
\Delta_{1} \circ \Delta_{3,2} = \Delta_{4,2}
\Delta_{1} \circ \Delta_{4,2} = \Delta_{4,3} + q\Delta_{2}
\Delta_{1} \circ \Delta_{4,3} = q\Delta_{3}.$$
(3.4)

Analogously one obtains multiplication tables with Δ_2 and Δ_3

Formulas (3.4) and (3.5) suffice to reconstruct the whole multiplication table. Indeed, if we let M_{λ} be the matrix of the small quantum multiplication with Δ_{λ} , then

$$\begin{split} M_{1,1} &= M_1^2 - M_2 \\ M_{2,1} &= M_1 M_2 - M_3 \\ M_{3,1} &= -\frac{1}{3} M_1 (M_3 - 2 M_{2,1}) \\ M_4 &= M_1 M_{2,1} - 2 M_{3,1} \\ M_{4,1} &= M_1 M_4 - q M_0 \\ M_{3,2} &= M_1 M_{3,1} - M_{4,1} \\ M_{4,2} &= M_1 M_{4,1} - q M_1 \\ M_{4,3} &= M_1 M_{4,2} - q M_2. \end{split}$$

3.4. **Nilpotent.** It is clear from (3.3) that $qH^*(X)$ specialized at q=1 is not semisimple. Indeed, Z has 10 reduced points and one fat point. The corresponding nilpotent (see [ChMaPe]), which is unique up to a constant factor, is

$$\Delta_{4,3} - \Delta_2 + \Delta_{1,1}. \tag{3.6}$$

Moreover, it is not hard to see that (3.6) is a specialization of a nilpotent defined globally over q. Indeed, it is given by $c_0 = \Delta_{4,3} - q\Delta_2 + q\Delta_{1,1}$.

4. Deformation along Δ_2 and semisimplicity

- 4.1. **Theorem.** The big quantum cohomology of IG(2,6) is generically semisimple.
- 4.2. **Proof.** As we will see, to exhibit the generic semisimplicity it is enough to consider the deformation of the small quantum cohomology along Δ_2 instead of the full big quantum cohomology $QH^*(X)$. Let \mathcal{A} be this deformation, i.e. \mathcal{A} is the quotient of $QH^*(X)$ with respect to the ideal generated by coordinates t_{λ} for $\lambda \neq 2$. To simplify the notation we will write t instead of t_2 . The multiplication in \mathcal{A} will be denoted \star .

To show the generic semisimplicity of \mathcal{A} (and, hence, of $QH^*(X)$) we exhibit an element that has distinct eigenvalues. Namely, we show that the element $\gamma = \Delta_2 + \Delta_1$ satisfies this property. The strategy can be described as follows. First, using Lemma 4.2.1 below, we compute the matrix \widetilde{M}_{γ} of multiplication by $\Delta_2 + \Delta_1$ in \mathcal{A} modulo terms quadratic in t. Secondly, we compute the characteristic polynomial P(x) of \widetilde{M}_{γ} . Finally, by looking at Newton polygons of P(x) and $P'_x(x)$ one deduces that the roots of P(x) are all distinct.

The computations outlined above are done using PARI/GP and the source code is available at [Source]. In the final step we use a well-known relation between roots of a polynomial over a discrete valuation ring (in our case it is the ring of formal power series in t) and its Newton polygon (for example, see [Ko, IV.3]). Namely, in PARI/GP we obtain


$$P(x) = P_0(x) + tP_1(x) + O(t^2), (4.1)$$

where

$$\begin{split} P_0(x) &= x^{12} - 60qx^9 - 90qx^8 - (96q^2 + 26q)x^7 - 60q^2x^4 - 90q^2x^3 - (96q^3 + 27q^2)x^2 \\ P_1(x) &= -30qx^{10} - 96qx^9 - 36qx^8 + 152q^2x^7 + 120q^2x^6 + (-32q^3 + 186q^2)x^5 + \\ &\quad + (240q^3 - 26q^2)x^4 - 36q^2x^3 + 152q^3x^2 - 30q^3x - 32q^4 - 9q^3. \end{split}$$

The characteristic polynomial of $\Delta_2 + \Delta_1$ in the small quantum cohomology is equal to the reduction of (4.1) modulo t, i.e. to $P_0(x)$. One can check that $P_0(x)$ has ten simple roots and one root of multiplicity two

The Newton polygon of (4.1) is



where $P(x) = a_0 x^{12} + a_1 x^{11} + \cdots + a_{11} x + a_{12}$ and v is the valuation with respect to t. Using Lemma 4 of [Ko, IV.3] we immediately obtain that P(x) has ten roots of valuation zero, i.e. their expansions start from a constant term, and two roots of valuation $\frac{1}{2}$. Since $P_0(x)$ has ten simple roots, ten roots of P(x) of valuation zero are distinct. Considering the Newton polygon for $P'_x(x)$ we see that it has ten roots of valuation zero and one root of valuation one. This implies that P(x) and $P'_x(x)$ have no common roots. Therefore, all roots of P(x) are distinct.

4.2.1. **Lemma.** Multiplication in \mathcal{A}/t^2 can be reconstructed from $qH^*(X)$, i.e. from 3-pointed Gromov-Witten invariants.

Proof. Consider elements

$$f_i = h^{\star i} \quad \text{for } 0 \le i \le 10$$
 (4.2)
 $f_{11} = \Delta_2$.

These elements form a basis of \mathcal{A} because their reductions modulo t are equal to $1, h, h^{\circ 2}, \ldots, h^{\circ 10}, \Delta_2$, which form a basis in $\mathcal{A}/t = qH^*(X)$.

In Basis (4.2) the multiplication table looks very simple. Clearly, if $i + j \le 10$, we have $f_i \star f_j = f_{i+j}$. Therefore, the only non-trivial part are the products $f_i \star f_{11}$.

For $i \neq 11$ we can compute the products $f_i \star f_{11}$ in the first neighbourhood due to Lemma 2.2. So the only possibly non-trivial product is $f_{11} \star f_{11} = \Delta_2 \star \Delta_2$.

To compute $\Delta_2 \star \Delta_2$ we do the following. First we write it as

$$\Delta_2 \star \Delta_2 = \sum_{i=0}^{10} (\Delta_2 \star \Delta_2, f_i) f^i + (\Delta_2 \star \Delta_2, f_{11}) f^{11},$$

where f^i is the dual of f_i with respect to the Poincaré pairing. Further, note that

$$(\Delta_2 \star \Delta_2, f_i) = (\Delta_2, \Delta_2 \star f_i) = (\Delta_2 \star f_i, \Delta_2),$$

so $(\Delta_2 \star \Delta_2, f_i)$ can be extracted from the products $f_i \star \Delta_2$, which we already know for $i \neq 11$. Finally, for the remaining $(\Delta_2 \star \Delta_2, f_{11})$ we have

$$(\Delta_2 \star \Delta_2, f_{11}) = (\Delta_2 \star \Delta_2, \Delta_2) = \langle \Delta_2, \Delta_2, \Delta_2 \rangle + \langle \Delta_2, \Delta_2, \Delta_2, \Delta_2 \rangle t + O(t^2).$$

By the dimension axiom for Gromov-Witten invariants we have $\langle \Delta_2, \Delta_2, \Delta_2 \rangle = \langle \Delta_2, \Delta_2, \Delta_2, \Delta_2 \rangle = 0$. Therefore, we do not need to compute any 4-pointed Gromov-Witten invariants to get the multiplication table in the first neighbourhood along t.

4.3. Spectrum of the Euler field. Here we will show that the operator of multiplication with the standard Euler vector field E of $QH^*(X)$ has simple spectrum (see [Ma, Proposition 5.3.4] for a definition). It is enough to show this for the restriction of E to the Δ_2 -direction, i.e. to consider the element

$$E = 5\Delta_1 - t\Delta_2,\tag{4.3}$$

where used E to denote this restriction as well.

Combining (2.3) and Lemma 4.2.1 one can compute the matrix \widetilde{M}_E in the second neighbourhood along t. Indeed, from Lemma 4.2.1 we know \widetilde{M}_2 modulo t^2 . Therefore, we know $t\widetilde{M}_2$ modulo t^3 . Moreover, by (2.3)

$$\widetilde{M}_1 = M_1 + q \frac{d}{dq} \int_0^t \widetilde{M}_2 dt.$$

Since we know \widetilde{M}_2 modulo t^2 , this formula gives us \widetilde{M}_1 modulo t^3 . Hence, we obtain \widetilde{M}_E modulo t^3 .

Implementing the above algorithm in PARI/GP and computing the characteristic polynomial P(x) of M_E we obtain¹

$$P(x) = P_0(x) + tP_1(x) + t^2P_2(x) - 39062500q^3t^3 + xO(t^3) + O(t^4),$$
(4.4)

where

$$P_0(x) = x^{12} - 81250qx^7 - 263671875q^2x^2$$

$$P_1(x) = -11250qx^8 - 35156250q^2x^3$$

$$P_2(x) = -900qx^9 - 78125q^2x^4.$$

Arguing the same way as after (4.1) gives that (4.4) has distinct roots. Hence, the standard Euler vector field has simple spectrum.

¹Note that here the precision of the characteristic polynomial is higher than that of \widetilde{M}_E .

References

- [Ab] L. Abrams. The quantum Euler class and the quantum cohomology of the Grassmannians. Israel J. Math. 117 (2000), 335–352.
- [Ba] A. Bayer. Semisimple quantum cohomology and blow-ups. Int. Math. Res. Not. 2004, no. 40, 2069–2083.
- [Be] A. Beilinson. Coherent sheaves on \mathbb{P}^n and problems in linear algebra. (Russian) Funktsional. Anal. i Prilozhen. 12 (1978), no. 3, 68–69.
- [BuKrTa] A. Buch, A. Kresch, H. Tamvakis. Quantum Pieri rules for isotropic Grassmannians. Invent. Math. 178 (2009), no. 2, 345–405.
- [ChMaPe] P.-E. Chaput, L. Manivel, N. Perrin. Quantum cohomology of minuscule homogeneous spaces III: semisimplicity and consequences. Canad. J. Math. 62 (2010), no. 6, 1246–1263.
- [ChPe] P.-E. Chaput, N. Perrin. On the quantum cohomology of adjoint varieties. Proc. Lond. Math. Soc. (3) 103 (2011), no. 2, 294–330.
- [Du] B. Dubrovin. Geometry and analytic theory of Frobenius manifolds. Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998). Doc. Math. 1998, Extra Vol. II, 315–326.
- [GaGo] S. S. Galkin, V. V. Golyshev. Quantum cohomology of Grassmannians, and cyclotomic fields. (Russian) Uspekhi Mat. Nauk 61 (2006), no. 1(367), 175–176; translation in Russian Math. Surveys 61 (2006), no. 1, 171–173.
- [GaGoIr] S. Galkin, V. Golyshev, H. Iritani. Gamma classes and quantum cohomology of Fano manifolds: Gamma conjectures. arXiv:1404.6407
- [HeMaTe] C. Hertling, Yu. Manin, C. Teleman. An update on semisimple quantum cohomology and F-manifolds. Proc. Steklov Inst. Math. 264 (2009), no. 1, 62–69.
- [Ir] H. Iritani. Convergence of quantum cohomology by quantum Lefschetz. J. Reine Angew. Math. 610 (2007), 29-69.
- [K] M. Kapranov. On the derived categories of coherent sheaves on some homogeneous spaces. Invent. Math. 92 (1988), no. 3, 479–508.
- [Ka] Yu. Kawamata. Derived categories of toric varieties. Michigan Math. J. 54 (2006), no. 3, 517–535.
- [Ko] N. Koblitz. p-adic numbers, p-adic analysis, and zeta-functions. Springer, 1984.
- [KoMa] M. Kontsevich, Yu. I. Manin. Gromov-Witten classes, quantum cohomology, and enumerative geometry. Comm. Math. Phys. Volume 164, Number 3 (1994), 525–562.
- [Ku] A. Kuznetsov. Exceptional collections for Grassmannians of isotropic lines. Proc. Lond. Math. Soc. (3) 97 (2008), no. 1, 155–182
- [KuPo] A. Kuznetsov, A. Polishchuk. Exceptional collections on isotropic Grassmannians. arXiv:1110.5607
- [Ma] Yu. I. Manin. Frobenius manifolds, quantum cohomology, and moduli spaces. AMS Colloquium Publ. 47, Providence RI, 1999, 303 pp.
- [Pe] N. Perrin. Semisimple quantum cohomology of some Fano varieties. to appear
- [Ue] K. Ueda. Stokes matrices for the quantum cohomologies of Grassmannians. Int. Math. Res. Not. 2005, no. 34, 2075–2086. [Source] http://users.ictp.it/~msmirnov/