

SHARP BOUNDS FOR NEUMAN MEANS IN TERMS OF GEOMETRIC, ARITHMETIC AND QUADRATIC MEANS

ZHI-JUN GUO, YAN ZHANG, YU-MING CHU, AND YING-QING SONG

ABSTRACT. In this paper, we find the greatest values $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7, \alpha_8$ and the least values $\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6, \beta_7, \beta_8$ such that the double inequalities

$$\begin{aligned} A^{\alpha_1}(a, b)G^{1-\alpha_1}(a, b) &< N_{GA}(a, b) < A^{\beta_1}(a, b)G^{1-\beta_1}(a, b), \\ \frac{\alpha_2}{G(a, b)} + \frac{1-\alpha_2}{A(a, b)} &< \frac{1}{N_{GA}(a, b)} < \frac{\beta_2}{G(a, b)} + \frac{1-\beta_2}{A(a, b)}, \\ A^{\alpha_3}(a, b)G^{1-\alpha_3}(a, b) &< N_{AG}(a, b) < A^{\beta_3}(a, b)G^{1-\beta_3}(a, b), \\ \frac{\alpha_4}{G(a, b)} + \frac{1-\alpha_4}{A(a, b)} &< \frac{1}{N_{AG}(a, b)} < \frac{\beta_4}{G(a, b)} + \frac{1-\beta_4}{A(a, b)}, \\ Q^{\alpha_5}(a, b)A^{1-\alpha_5}(a, b) &< N_{AQ}(a, b) < Q^{\beta_5}(a, b)A^{1-\beta_5}(a, b), \\ \frac{\alpha_6}{A(a, b)} + \frac{1-\alpha_6}{Q(a, b)} &< \frac{1}{N_{AQ}(a, b)} < \frac{\beta_6}{A(a, b)} + \frac{1-\beta_6}{Q(a, b)}, \\ Q^{\alpha_7}(a, b)A^{1-\alpha_7}(a, b) &< N_{QA}(a, b) < Q^{\beta_7}(a, b)A^{1-\beta_7}(a, b), \\ \frac{\alpha_8}{A(a, b)} + \frac{1-\alpha_8}{Q(a, b)} &< \frac{1}{N_{QA}(a, b)} < \frac{\beta_8}{A(a, b)} + \frac{1-\beta_8}{Q(a, b)} \end{aligned}$$

hold for all $a, b > 0$ with $a \neq b$, where G , A and Q are respectively the geometric, arithmetic and quadratic means, and N_{GA} , N_{AG} , N_{AQ} and N_{QA} are the Neuman means derived from the Schwab-Borchardt mean.

1. INTRODUCTION

For $a, b > 0$ with $a \neq b$, the Schwab-Borchardt mean $SB(a, b)$ [1-3] of a and b is given by

$$SB(a, b) = \begin{cases} \frac{\sqrt{b^2-a^2}}{\cos^{-1}(a/b)}, & a < b, \\ \frac{\sqrt{a^2-b^2}}{\cosh^{-1}(a/b)}, & a > b, \end{cases}$$

where $\cos^{-1}(x)$ and $\cosh^{-1}(x) = \log(x + \sqrt{x^2 - 1})$ are the inverse cosine and inverse hyperbolic cosine functions, respectively. Recently, the Schwab-Borchardt mean has been the subject of intensive research. In particular, many remarkable inequalities for Schwab-Borchardt mean can be found in the literature [1-7]. Very recently, the Neuman mean $N(a, b) = (a + b^2/SB(a, b))/2$ derived from the Schwab-Borchardt was introduced and researched by Neuman in [8].

2010 *Mathematics Subject Classification.* 26E60.

Key words and phrases. Schwab-Borchardt mean, Neuman mean, geometric mean, arithmetic mean, quadratic mean.

This research was supported by the Natural Science Foundation of China under Grants 11371125, 11171307 and 61374086, the Natural Science Foundation of Zhejiang Province under Grant LY13A010004, and the Natural Science Foundation of the Department of Education of Hunan Province under Grants 12C0577 and 13A013.

Let $N_{AG}(a, b) = N(A(a, b), G(a, b))$, $N_{GA}(a, b) = N(G(a, b), A(a, b))$, $N_{QA}(a, b) = N(Q(a, b), A(a, b))$ and $N_{AQ}(a, b) = N(A(a, b), Q(a, b))$ be the Neuman means, where $G(a, b) = \sqrt{ab}$, $A(a, b) = (a + b)/2$ and $Q(a, b) = \sqrt{(a^2 + b^2)/2}$ are the classical geometric, arithmetic and quadratic means of a and b , respectively. Then Neuman [8] proved that the inequalities

$$G(a, b) < N_{AG}(a, b) < N_{GA}(a, b) < A(a, b) < N_{QA}(a, b) < N_{AQ}(a, b) < Q(a, b)$$

hold for all $a, b > 0$ with $a \neq b$.

Let $a > b > 0$ and $v = (a - b)/(a + b) \in (0, 1)$. Then we clearly see that

$$(1.1) \quad G(a, b) = A(a, b)\sqrt{1 - v^2}, \quad Q(a, b) = A(a, b)\sqrt{1 + v^2},$$

and the following explicit formulas for $N_{AG}(a, b)$, $N_{GA}(a, b)$, $N_{QA}(a, b)$ and $N_{AQ}(a, b)$ are given in [8]

$$(1.2) \quad N_{AG}(a, b) = \frac{1}{2}A(a, b) \left[1 + (1 - v^2)\frac{\tanh^{-1} v}{v} \right],$$

$$(1.3) \quad N_{GA}(a, b) = \frac{1}{2}A(a, b) \left[\sqrt{1 - v^2} + \frac{\sin^{-1} v}{v} \right],$$

$$(1.4) \quad N_{QA}(a, b) = \frac{1}{2}A(a, b) \left[\sqrt{1 + v^2} + \frac{\sinh^{-1} v}{v} \right],$$

$$(1.5) \quad N_{AQ}(a, b) = \frac{1}{2}A(a, b) \left[1 + (1 + v^2)\frac{\tan^{-1} v}{v} \right],$$

where $\tanh^{-1}(x) = \log[(1 + x)/(1 - x)]/2$, $\sin^{-1}(x)$, $\sinh^{-1}(x) = \log(x + \sqrt{1 + x^2})$ and $\tan^{-1}(x)$ are the inverse hyperbolic tangent, inverse sine, inverse hyperbolic sine and inverse tangent functions, respectively.

In [8], Neuman also proved that the double inequalities

$$\alpha_1 A(a, b) + (1 - \alpha_1)G(a, b) < N_{GA}(a, b) < \beta_1 A(a, b) + (1 - \beta_1)G(a, b),$$

$$\alpha_2 Q(a, b) + (1 - \alpha_2)A(a, b) < N_{AQ}(a, b) < \beta_2 Q(a, b) + (1 - \beta_2)A(a, b),$$

$$\alpha_3 A(a, b) + (1 - \alpha_3)G(a, b) < N_{AG}(a, b) < \beta_3 A(a, b) + (1 - \beta_3)G(a, b),$$

$$\alpha_4 Q(a, b) + (1 - \alpha_4)A(a, b) < N_{QA}(a, b) < \beta_4 Q(a, b) + (1 - \beta_4)A(a, b)$$

hold for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_1 \leq 2/3$, $\beta_1 \geq \pi/4$, $\alpha_2 \leq 2/3$, $\beta_2 \geq (\pi - 2)/[4(\sqrt{2} - 1)] = 0.689\dots$, $\alpha_3 \leq 1/3$, $\beta_3 \geq 1/2$, $\alpha_4 \leq 1/3$ and $\beta_4 \geq [\log(1 + \sqrt{2}) + \sqrt{2} - 2]/[2(\sqrt{2} - 1)] = 0.356\dots$

In [9], the authors presented the best possible parameters $\alpha_1, \alpha_2, \beta_1, \beta_2 \in [0, 1/2]$ and $\alpha_3, \alpha_4, \beta_3, \beta_4 \in [1/2, 1]$ such that the double inequalities

$$G(\alpha_1 a + (1 - \alpha_1)b, \alpha_1 b + (1 - \alpha_1)a) < N_{AG}(a, b) < G(\beta_1 a + (1 - \beta_1)b, \beta_1 b + (1 - \beta_1)a),$$

$$G(\alpha_2 a + (1 - \alpha_2)b, \alpha_2 b + (1 - \alpha_2)a) < N_{GA}(a, b) < G(\beta_2 a + (1 - \beta_2)b, \beta_2 b + (1 - \beta_2)a),$$

$$Q(\alpha_3 a + (1 - \alpha_3)b, \alpha_3 b + (1 - \alpha_3)a) < N_{QA}(a, b) < Q(\beta_3 a + (1 - \beta_3)b, \beta_3 b + (1 - \beta_3)a),$$

$$Q(\alpha_4 a + (1 - \alpha_4)b, \alpha_4 b + (1 - \alpha_4)a) < N_{AQ}(a, b) < Q(\beta_4 a + (1 - \beta_4)b, \beta_4 b + (1 - \beta_4)a)$$

hold for all $a, b > 0$ with $a \neq b$.

The main purpose of this paper is to find the greatest values $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7, \alpha_8$ and the least values $\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6, \beta_7, \beta_8$ such that the double inequalities

$$A^{\alpha_1}(a, b)G^{1-\alpha_1}(a, b) < N_{GA}(a, b) < A^{\beta_1}(a, b)G^{1-\beta_1}(a, b),$$

$$\frac{\alpha_2}{G(a, b)} + \frac{1 - \alpha_2}{A(a, b)} < \frac{1}{N_{GA}(a, b)} < \frac{\beta_2}{G(a, b)} + \frac{1 - \beta_2}{A(a, b)},$$

$$A^{\alpha_3}(a, b)G^{1-\alpha_3}(a, b) < N_{AG}(a, b) < A^{\beta_3}(a, b)G^{1-\beta_3}(a, b),$$

$$\frac{\alpha_4}{G(a, b)} + \frac{1 - \alpha_4}{A(a, b)} < \frac{1}{N_{AG}(a, b)} < \frac{\beta_4}{G(a, b)} + \frac{1 - \beta_4}{A(a, b)},$$

$$Q^{\alpha_5}(a, b)A^{1-\alpha_5}(a, b) < N_{AQ}(a, b) < Q^{\beta_5}(a, b)A^{1-\beta_5}(a, b),$$

$$\frac{\alpha_6}{A(a, b)} + \frac{1 - \alpha_6}{Q(a, b)} < \frac{1}{N_{AQ}(a, b)} < \frac{\beta_6}{A(a, b)} + \frac{1 - \beta_6}{Q(a, b)},$$

$$Q^{\alpha_7}(a, b)A^{1-\alpha_7}(a, b) < N_{QA}(a, b) < Q^{\beta_7}(a, b)A^{1-\beta_7}(a, b),$$

$$\frac{\alpha_8}{A(a, b)} + \frac{1 - \alpha_8}{Q(a, b)} < \frac{1}{N_{QA}(a, b)} < \frac{\beta_8}{A(a, b)} + \frac{1 - \beta_8}{Q(a, b)}$$

hold for all $a, b > 0$ with $a \neq b$.

2. LEMMAS

In order to prove our main results we need several lemmas, which we present in this section.

Lemma 2.1. (See [10, Theorem 1.25]) Let $-\infty < a < b < \infty$, $f, g : [a, b] \rightarrow (-\infty, \infty)$ be continuous on $[a, b]$ and differentiable on (a, b) , and $g'(x) \neq 0$ on (a, b) . If $f'(x)/g'(x)$ is increasing (decreasing) on (a, b) , then so are

$$\frac{f(x) - f(a)}{g(x) - g(a)}, \quad \frac{f(x) - f(b)}{g(x) - g(b)}.$$

If $f'(x)/g'(x)$ is strictly monotone, the the monotonicity in the conclusion is also strict.

Lemma 2.2. (See [11, Lemma 1.1]) Suppose that the power series $f(x) = \sum_{n=0}^{\infty} a_n x^n$ and $g(x) = \sum_{n=0}^{\infty} b_n x^n$ have the radius of convergence $r > 0$ and $b_n > 0$ for all $n \geq 0$. If the sequence $\{a_n/b_n\}$ is (strictly) increasing (decreasing) for all $n \geq 0$, then the function $f(x)/g(x)$ is also (strictly) increasing (decreasing) on $(0, r)$.

Lemma 2.3. The function

$$(2.1) \quad f_1(x) = \frac{\log[\sin(2x)] - \log[2x + \sin(2x)] + \log 2}{\log(\cos x)}$$

is strictly increasing from $(0, \pi/2)$ onto $(2/3, 1)$.

Proof. It follows from (3.1) that

$$(2.2) \quad f_1(0) = \frac{2}{3},$$

$$(2.3) \quad f_1\left(\frac{\pi}{2}\right) = 1.$$

Let $g_1(x) = \log[\sin(2x)] - \log[2x + \sin(2x)] + \log 2$, $h_1(x) = \log(\cos x)$, $g_2(x) = \sin(2x) - 2x \cos(2x)$ and $h_2(x) = [2x + \sin(2x)] \sin^2 x$. Then simple computations lead to

$$(2.4) \quad g_1(0^+) = h_1(0) = g_2(0) = h_2(0) = 0,$$

$$(2.5) \quad f_1(x) = \frac{g_1(x)}{h_1(x)}, \quad \frac{g'_1(x)}{h'_1(x)} = \frac{g_2(x)}{h_2(x)},$$

$$(2.6) \quad \frac{g'_2(x)}{h'_2(x)} = \frac{1}{\frac{1}{2} + \frac{\sin(2x)}{2x}}.$$

It is well known that the function $\sin x/x$ is strictly decreasing on $(0, \pi)$, hence equation (2.6) leads to the conclusion that the function $g'_2(x)/h'_2(x)$ is strictly increasing on $(0, \pi/2)$. Therefore, Lemma 2.3 follows from Lemma 2.1 and (2.2)-(2.5) together with the monotonicity of $g'_2(x)/h'_2(x)$.

Lemma 2.4. *The function*

$$(2.7) \quad f_2(x) = \frac{\log[2x + \sinh(2x)] - \log[\sinh(x)] - 2 \log 2}{\log[\cosh(x)]}$$

is strictly increasing from $(0, \infty)$ onto $(1/3, 1)$.

Proof. It follows from (2.7) that

$$(2.8) \quad f_2(0^+) = \frac{1}{3},$$

$$(2.9) \quad \lim_{x \rightarrow \infty} f_2(x) = 1.$$

Let $g_3(x) = \log[2x + \sinh(2x)] - \log[\sinh(x)] - 2 \log 2$ and $h_3(x) = \log[\cosh(x)]$. Then simple computations lead to

$$(2.10) \quad f_2(x) = \frac{g_3(x)}{h_3(x)}, \quad g_3(0^+) = h_3(0) = 0,$$

$$(2.11) \quad \begin{aligned} \frac{g'_3(x)}{h'_3(x)} &= \frac{\sinh(4x) - 4x \cosh(2x) + 2 \sinh(2x) - 4x}{\sinh(4x) + 4x \cosh(2x) - 2 \sinh(2x) - 4x} \\ &= \frac{\sum_{n=1}^{\infty} \frac{(2^{2n}-2n)2^{2n+2}}{(2n+1)!} x^{2n+1}}{\sum_{n=1}^{\infty} \frac{(2^{2n}+2n)2^{2n+2}}{(2n+1)!} x^{2n+1}} \\ &= \frac{\sum_{n=0}^{\infty} \frac{(2^{2n+2}-2n-2)2^{2n+4}}{(2n+3)!} x^{2n}}{\sum_{n=0}^{\infty} \frac{(2^{2n+2}+2n+2)2^{2n+4}}{(2n+3)!} x^{2n}}. \end{aligned}$$

Let

$$(2.12) \quad a_n = \frac{(2^{2n+2} - 2n - 2) 2^{2n+4}}{(2n + 3)!}, \quad b_n = \frac{(2^{2n+2} + 2n + 2) 2^{2n+4}}{(2n + 3)!}.$$

Then

$$(2.13) \quad b_n > 0, \quad \frac{a_{n+1}}{b_{n+1}} - \frac{a_n}{b_n} = \frac{(3n + 2) 2^{2n+2}}{(2^{2n+3} + n + 2)(2^{2n+1} + n + 1)} > 0$$

for all $n \geq 0$.

It follows from Lemma 2.2 and (2.11)-(2.13) that the function $g'_3(x)/h'_3(x)$ is strictly increasing on $(0, \infty)$. Therefore, Lemma 2.4 follows from Lemma 2.1 and (2.8)-(2.10) together with the monotonicity of $g'_3(x)/h'_3(x)$.

Lemma 2.5. *The function*

$$(2.14) \quad f_3(x) = \frac{2x - \sin(2x)}{(1 - \cos x)[2x + \sin(2x)]}$$

is strictly increasing from $(0, \pi/2)$ onto $(2/3, 1)$.

Proof. It follows from (2.14) that

$$(2.15) \quad f_3(0^+) = \frac{2}{3},$$

$$(2.16) \quad f_3(\pi/2) = 1.$$

Let $g_4(x) = 2x - \sin(2x)$ and $h_4(x) = (1 - \cos x)[2x + \sin(2x)]$. Then simple computations lead to

$$(2.17) \quad f_3(x) = \frac{g_4(x)}{h_4(x)}, \quad g_4(0) = h_4(0) = 0,$$

$$g'_4(x) = 4 \sin^2 x,$$

$$h'_4(x) = 2 \sin^2 x \cos x - 4 \cos^3 x + 4 \cos^2 x + 2x \sin x,$$

$$(2.18) \quad g'_4(0) = h'_4(0) = 0,$$

$$(2.19) \quad \frac{g''_4(x)}{h''_4(x)} = \frac{4}{9 \cos x + \frac{x}{\sin x} - 4},$$

$$(2.20) \quad \left(9 \cos x + \frac{x}{\sin x}\right)' = -8 \sin x - \frac{[2x - \sin(2x)] \cos x}{2 \sin^2 x} < 0$$

for $x \in (0, \pi/2)$.

Therefore, Lemma 2.5 follows easily from Lemma 2.1 and (2.15)-(2.20).

Lemma 2.6. *The function*

$$(2.21) \quad f_4(x) = \frac{\sinh(x) \cosh^2(x) - 2 \sinh(x) \cosh(x) + x \cosh(x)}{\sinh(x) \cosh^2(x) - \sinh(x) \cosh(x) + x \cosh(x) - x}$$

is strictly increasing from $(0, \infty)$ onto $(1/3, 1)$.

Proof. It follows from (2.21) that

$$\begin{aligned}
 (2.22) \quad f_4(x) &= \frac{\frac{1}{4} \sinh(3x) + \frac{1}{4} \sinh(x) - \sinh(2x) + x \cosh(x)}{\frac{1}{4} \sinh(3x) + \frac{1}{4} \sinh(x) - \frac{1}{2} \sinh(2x) + x \cosh(x) - x} \\
 &= \frac{\sum_{n=1}^{\infty} \frac{3^{2n+1} - 2^{2n+3} + 8n+5}{4[(2n+1)!]} x^{2n+1}}{\sum_{n=1}^{\infty} \frac{3^{2n+1} - 2^{2n+2} + 8n+5}{4[(2n+1)!]} x^{2n+1}} \\
 &= \frac{\sum_{n=0}^{\infty} \frac{3^{2n+3} - 2^{2n+5} + 8n+13}{4[(2n+3)!]} x^{2n}}{\sum_{n=0}^{\infty} \frac{3^{2n+3} - 2^{2n+4} + 8n+13}{4[(2n+3)!]} x^{2n}}.
 \end{aligned}$$

Let

$$(2.23) \quad a_n = \frac{3^{2n+3} - 2^{2n+5} + 8n+13}{4[(2n+3)!]}, \quad b_n = \frac{3^{2n+3} - 2^{2n+4} + 8n+13}{4[(2n+3)!]}.$$

Then simple computations lead to

$$(2.24) \quad b_n > \frac{3^{2n+3} - 2^{2n+4}}{4[(2n+3)!]} = \frac{2^{2n+3} \left[\left(\frac{3}{2}\right)^{2n+3} - 2 \right]}{4[(2n+3)!]} > 0$$

$$(2.25) \quad \frac{a_{n+1}}{b_{n+1}} - \frac{a_n}{b_n} = \frac{(135 \times 3^{2n} - 24n - 31) 2^{2n+4}}{(3^{2n+3} - 2^{2n+4} + 8n+13)(3^{2n+5} - 2^{2n+6} + 8n+21)} > 0$$

for all $n \geq 0$.

Note that

$$(2.26) \quad f_4(0^+) = \frac{1}{3}, \quad \lim_{x \rightarrow \infty} f_4(x) = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1.$$

Therefore, Lemma 2.6 follows easily from Lemma 2.2 and (2.22)-(2.26).

3. MAIN RESULTS

Theorem 3.1. *The double inequalities*

$$(3.1) \quad A^{\alpha_1}(a, b) G^{1-\alpha_1}(a, b) < N_{GA}(a, b) < A^{\beta_1}(a, b) G^{1-\beta_1}(a, b),$$

$$(3.2) \quad \frac{\alpha_2}{G(a, b)} + \frac{1 - \alpha_2}{A(a, b)} < \frac{1}{N_{GA}(a, b)} < \frac{\beta_2}{G(a, b)} + \frac{1 - \beta_2}{A(a, b)}$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_1 \leq 2/3$, $\beta_1 \geq 1$, $\alpha_2 \leq 0$ and $\beta_2 \geq 1/3$.

Proof. We clearly see that inequalities (3.1) and (3.2) can be rewritten as

$$(3.3) \quad \left(\frac{A(a, b)}{G(a, b)} \right)^{\alpha_1} < \frac{N_{GA}(a, b)}{G(a, b)} < \left(\frac{A(a, b)}{G(a, b)} \right)^{\beta_1}$$

and

$$(3.4) \quad 1 - \beta_2 < \frac{\frac{1}{G(a, b)} - \frac{1}{N_{GA}(a, b)}}{\frac{1}{G(a, b)} - \frac{1}{A(a, b)}} < 1 - \alpha_2,$$

respectively.

Since both the geometric mean $G(a, b)$ and arithmetic mean $A(a, b)$ are symmetric and homogeneous of degree 1, without loss of generality, we assume that $a > b$.

Let $v = (a-b)/(a+b) \in (0, 1)$. Then from (1.1) and (1.3) we know that inequalities (3.3) and (3.4) are equivalent to

$$(3.5) \quad \alpha_1 < \frac{\log \left[\frac{1}{2} \left(1 + \frac{\sin^{-1}(v)}{v\sqrt{1-v^2}} \right) \right]}{\log \frac{1}{\sqrt{1-v^2}}} < \beta_1$$

and

$$(3.6) \quad 1 - \beta_2 < \frac{\sin^{-1} v - v\sqrt{1-v^2}}{(1-\sqrt{1-v^2})(v\sqrt{1-v^2} + \sin^{-1} v)} < 1 - \alpha_2,$$

respectively.

Let $x = \sin^{-1}(v)$. Then $x \in (0, \pi/2)$,

$$(3.7) \quad \frac{\log \left[\frac{1}{2} \left(1 + \frac{\sin^{-1}(v)}{v\sqrt{1-v^2}} \right) \right]}{\log \frac{1}{\sqrt{1-v^2}}} = \frac{\log[\sin(2x)] - \log[2x + \sin(2x)] + \log 2}{\log(\cos x)},$$

$$(3.8) \quad \frac{\sin^{-1} v - v\sqrt{1-v^2}}{(1-\sqrt{1-v^2})(v\sqrt{1-v^2} + \sin^{-1} v)} = \frac{2x - \sin(2x)}{(1-\cos x)[2x + \sin(2x)]}.$$

Therefore, inequality (3.1) holds for all $a, b > 0$ with $a \neq b$ follows from (3.5) and (3.7) together with Lemma 2.3, and inequality (3.2) holds for all $a, b > 0$ with $a \neq b$ follows from (3.6) and (3.8) together with Lemma 2.5.

Theorem 3.2. The double inequalities

$$(3.9) \quad A^{\alpha_3}(a, b)G^{1-\alpha_3}(a, b) < N_{AG}(a, b) < A^{\beta_3}(a, b)G^{1-\beta_3}(a, b),$$

$$(3.10) \quad \frac{\alpha_4}{G(a, b)} + \frac{1-\alpha_4}{A(a, b)} < \frac{1}{N_{AG}(a, b)} < \frac{\beta_4}{G(a, b)} + \frac{1-\beta_4}{A(a, b)}$$

hold for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_3 \leq 1/3$, $\beta_3 \geq 1$, $\alpha_4 \leq 0$ and $\beta_4 \geq 2/3$.

Proof. We clearly see that inequalities (3.9) and (3.10) can be rewritten as

$$(3.11) \quad \left(\frac{A(a, b)}{G(a, b)} \right)^{\alpha_3} < \frac{N_{AG}(a, b)}{G(a, b)} < \left(\frac{A(a, b)}{G(a, b)} \right)^{\beta_3}$$

and

$$(3.12) \quad 1 - \beta_4 < \frac{\frac{1}{G(a, b)} - \frac{1}{N_{AG}(a, b)}}{\frac{1}{G(a, b)} - \frac{1}{A(a, b)}} < 1 - \alpha_4,$$

respectively.

Without loss of generality, we assume that $a > b$. Let $v = (a-b)/(a+b) \in (0, 1)$. Then it follows from (1.1) and (1.2) that inequalities (3.11) and (3.12) are equivalent to

$$(3.13) \quad \alpha_3 < \frac{\log \left[\frac{1}{\sqrt{1-v^2}} + \frac{\sqrt{1-v^2}}{v} \tanh^{-1}(v) \right] - \log 2}{\log \frac{1}{\sqrt{1-v^2}}} < \beta_3$$

and

$$(3.14) \quad 1 - \beta_4 < \frac{v + (1-v^2) \tanh^{-1}(v) - 2v\sqrt{1-v^2}}{(1-\sqrt{1-v^2})[v + (1-v^2) \tanh^{-1}(v)]} < 1 - \beta_4,$$

respectively. Let $x = \tanh^{-1}(v) \in (0, \infty)$. Then simple computations lead to

$$(3.15) \quad \begin{aligned} & \frac{\log \left[\frac{1}{\sqrt{1-v^2}} + \frac{\sqrt{1-v^2}}{v} \tanh^{-1}(v) \right] - \log 2}{\log \frac{1}{\sqrt{1-v^2}}} \\ &= \frac{\log[2x + \sinh(2x)] - \log[\sinh(x)] - 2 \log 2}{\log[\cosh(x)]} \end{aligned}$$

and

$$(3.16) \quad \begin{aligned} & \frac{v + (1 - v^2) \tanh^{-1}(v) - 2v\sqrt{1 - v^2}}{(1 - \sqrt{1 - v^2})[v + (1 - v^2) \tanh^{-1}(v)]} \\ &= \frac{\sinh(x) \cosh^2(x) - 2 \sinh(x) \cosh(x) + x \cosh(x)}{\sinh(x) \cosh^2(x) - \sinh(x) \cosh(x) + x \cosh(x) - x}. \end{aligned}$$

Therefore, inequality (3.9) holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_3 \leq 1/3$ and $\beta_3 \geq 1$ follows from (3.13) and (3.15) together with Lemma 2.4, and inequality (3.10) holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_4 \leq 0$ and $\beta_4 \geq 2/3$ follows from (3.14) and (3.16) together with Lemma 2.6.

Theorem 3.3. *The double inequalities*

$$(3.17) \quad Q^{\alpha_5}(a, b) A^{1-\alpha_5}(a, b) < N_{AQ}(a, b) < Q^{\beta_5}(a, b) A^{1-\beta_5}(a, b),$$

$$(3.18) \quad \frac{\alpha_6}{A(a, b)} + \frac{1 - \alpha_6}{Q(a, b)} < \frac{1}{N_{AQ}(a, b)} < \frac{\beta_6}{A(a, b)} + \frac{1 - \beta_6}{Q(a, b)}$$

hold for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_5 \leq 2/3$, $\beta_5 \geq 2 \log(\pi+2)/\log 2 - 4 = 0.7244\dots$, $\alpha_6 \leq [6 + 2\sqrt{2} - (1 + \sqrt{2})\pi]/(\pi + 2) = 0.2419\dots$ and $\beta_6 \geq 1/3$.

Proof. We clearly see that inequalities (3.17) and (3.18) can be rewritten as

$$(3.19) \quad \left(\frac{Q(a, b)}{A(a, b)} \right)^{\alpha_5} < \frac{N_{AQ}(a, b)}{A(a, b)} < \left(\frac{Q(a, b)}{A(a, b)} \right)^{\beta_5}$$

and

$$(3.20) \quad 1 - \beta_6 < \frac{\frac{1}{A(a, b)} - \frac{1}{N_{AQ}(a, b)}}{\frac{1}{A(a, b)} - \frac{1}{Q(a, b)}} < 1 - \alpha_6,$$

respectively.

Without loss of generality, we assume that $a > b$. Let $v = (a - b)/(a + b) \in (0, 1)$. Then from (1.1) and (1.5) we clearly see that inequalities (3.19) and (3.20) are equivalent to

$$(3.21) \quad \alpha_5 < \frac{2 \log(1 + \frac{1+v^2}{v} \tan^{-1}(v)) - 2 \log 2}{\log(1 + v^2)} < \beta_5$$

and

$$(3.22) \quad 1 - \beta_6 < \frac{[(1 + v^2) \tan^{-1}(v) - v] \sqrt{1 + v^2}}{[(1 + v^2) \tan^{-1}(v) + v] (\sqrt{1 + v^2} - 1)} < 1 - \alpha_6,$$

respectively.

Let $x = \tan^{-1}(v)$. Then $x \in (0, \pi/4)$,

$$(3.23) \quad \frac{2 \log(1 + \frac{1+v^2}{v} \tan^{-1}(v)) - 2 \log 2}{\log(1 + v^2)}$$

$$= \frac{\log[\sin(2x)] - \log[2x + \sin(2x)] + \log 2}{\log(\cos x)} = f_1(x)$$

and

$$(3.24) \quad \begin{aligned} & \frac{[(1+v^2)\tan^{-1}(v)-v]\sqrt{1+v^2}}{[(1+v^2)\tan^{-1}(v)+v](\sqrt{1+v^2}-1)} \\ & \frac{2x-\sin(2x)}{(1-\cos x)[2x+\sin(2x)]} = f_3(x). \end{aligned}$$

Note that

$$(3.25) \quad f_1\left(\frac{\pi}{4}\right) = \frac{2\log(\pi+2)}{\log 2} - 4,$$

$$(3.26) \quad f_3\left(\frac{\pi}{4}\right) = \frac{(2+\sqrt{2})(\pi-2)}{\pi+2} = 1 - \frac{6+2\sqrt{2}-(1+\sqrt{2})\pi}{\pi+2}.$$

Therefore, inequality (3.17) holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_5 \leq 2/3$ and $\beta_5 \geq 2\log(\pi+2)/\log 2 - 4$ follows from (3.21), (3.23), (3.25) and Lemma 2.3, and inequality (3.18) holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_6 \leq [6+2\sqrt{2}-(1+\sqrt{2})\pi]/(\pi+2)$ and $\beta_6 \geq 1/3$ follows from (3.22), (3.24), (3.26) and Lemma 2.5.

Theorem 3.4. The double inequalities

$$(3.27) \quad Q^{\alpha_7}(a, b)A^{1-\alpha_7}(a, b) < N_{QA}(a, b) < Q^{\beta_7}(a, b)A^{1-\beta_7}(a, b),$$

$$(3.28) \quad \frac{\alpha_8}{A(a, b)} + \frac{1-\alpha_8}{Q(a, b)} < \frac{1}{N_{QA}(a, b)} < \frac{\beta_8}{A(a, b)} + \frac{1-\beta_8}{Q(a, b)}$$

hold for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_7 \leq 1/3$, $\beta_7 \geq 2\log[\sqrt{2} + \log(1+\sqrt{2})]/\log 2 - 2 = 0.3977\dots$, $\alpha_8 \leq [2+\sqrt{2}-(1+\sqrt{2})\log(1+\sqrt{2})]/[\sqrt{2}+\log(1+\sqrt{2})] = 0.5603\dots$ and $\beta_8 \geq 2/3$.

Proof. We clearly see that inequalities (3.27) and (3.28) can be rewritten as

$$(3.29) \quad \left(\frac{Q(a, b)}{A(a, b)}\right)^{\alpha_7} < \frac{N_{QA}(a, b)}{A(a, b)} < \left(\frac{Q(a, b)}{A(a, b)}\right)^{\beta_7}$$

and

$$(3.30) \quad 1 - \beta_8 < \frac{\frac{1}{A(a, b)} - \frac{1}{N_{QA}(a, b)}}{\frac{1}{A(a, b)} - \frac{1}{Q(a, b)}} < 1 - \alpha_8,$$

respectively.

Without loss of generality, we assume that $a > b$. Let $v = (a-b)/(a+b) \in (0, 1)$. Then from (1.1) and (1.4) we clearly see that inequalities (3.29) and (3.30) are equivalent to

$$(3.31) \quad \alpha_7 < \frac{2\log\left[\sqrt{1+v^2} + \frac{\sinh^{-1}(v)}{v}\right] - 2\log 2}{\log(1+v^2)} < \beta_7$$

and

$$(3.32) \quad 1 - \beta_8 < \frac{[v(1+v^2) + \sqrt{1+v^2}\sinh^{-1}(v)] - 2v\sqrt{1+v^2}}{(\sqrt{1+v^2}-1)[v\sqrt{1+v^2} + \sinh^{-1}(v)]} < 1 - \alpha_8,$$

respectively.

Let $x = \sinh^{-1}(v)$. Then $x \in (0, \log(1 + \sqrt{2}))$,

$$(3.33) \quad \begin{aligned} & \frac{2 \log \left[\sqrt{1+v^2} + \frac{\sinh^{-1}(v)}{v} \right] - 2 \log 2}{\log(1+v^2)} \\ &= \frac{\log[2x + \sinh(2x)] - \log[\sinh(x)] - 2 \log 2}{\log[\cosh(x)]} = f_2(x), \\ (3.34) \quad & \frac{[v(1+v^2) + \sqrt{1+v^2} \sinh^{-1}(v)] - 2v\sqrt{1+v^2}}{(\sqrt{1+v^2}-1)[v\sqrt{1+v^2} + \sinh^{-1}(v)]} \\ &= \frac{\sinh(x)\cosh^2(x) - 2\sinh(x)\cosh(x) + x\cosh(x)}{\sinh(x)\cosh^2(x) - \sinh(x)\cosh(x) + x\cosh(x) - x} = f_4(x). \end{aligned}$$

Note that

$$(3.35) \quad f_2[\log(1 + \sqrt{2})] = \frac{2 \log[\sqrt{2} + \log(1 + \sqrt{2})]}{\log 2} - 2,$$

$$(3.36) \quad \begin{aligned} f_4[\log(1 + \sqrt{2})] &= \frac{(2 + \sqrt{2}) \log(1 + \sqrt{2}) - 2}{\sqrt{2} + \log(1 + \sqrt{2})} \\ &= 1 - \frac{2 + \sqrt{2} - (1 + \sqrt{2}) \log(1 + \sqrt{2})}{\sqrt{2} + \log(1 + \sqrt{2})}. \end{aligned}$$

Therefore, inequality (3.27) holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_7 \leq 1/3$ and $\beta_7 \geq 2 \log[\sqrt{2} + \log(1 + \sqrt{2})]/\log 2 - 2$ follows from (3.31), (3.33), (3.35) and Lemma 2.4, and inequality (3.28) holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_8 \leq [2 + \sqrt{2} - (1 + \sqrt{2}) \log(1 + \sqrt{2})]/[\sqrt{2} + \log(1 + \sqrt{2})]$ and $\beta_8 \geq 2/3$ follows from (3.32), (3.34), (3.36) and Lemma 2.6.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

REFERENCES

- [1] E. Neuman and J. Sándor, On the Schwab-Borchardt mean, *Math. Pannon.*, 2003, **14**(2), 253–266.
- [2] E. Neuman and J. Sándor, On the Schwab-Borchardt mean II, *Math. Pannon.*, 2005, **17**(1), 49–59.
- [3] E. Neuman, Inequalities for the Schwab-Borchardt mean and their applications, *J. Math. Inequal.*, 2011, **5**(4), 601–609.
- [4] Z.-Y. He, Y.-M. Chu and M.-K. Wang, Optimal bounds for Neuman means in terms of harmonic and contraharmonic means, *J. Appl. Math.*, **2013**, Article ID 807623, 4 pages.
- [5] Y.-M. Chu and W.-M. Qian, Refinements of Bounds for Neuman means, *Abstr. Appl. Anal.*, **2014**, Article ID 354132, 8 pages.
- [6] E. Neuman, On some means derived from the Schwab-Borchardt mean, *J. Math. Inequal.*, 2014, **8**(1), 171–183.
- [7] E. Neuman, On some means derived from the Schwab-Borchardt mean II, *J. Math. Inequal.*, 2014, **8**(2), 361–370.
- [8] E. Neuman, On a new bivariate mean, *Aequat. Math.*, DOI 10.1007/s00010-013-0224-8.
- [9] Y. Zhang, Y.-M. Chu and Y.-L. Jiang, Sharp geometric mean bounds for Neuman means, *Abstr. Appl. Anal.*, **2014**, Article ID 949815, 6 pages.
- [10] G. D. Anderson, S.-L. Qiu, M. K. Vamanamurthy and M. Vuorinen, Generalized elliptic integrals and modular equations, *Pacific J. Math.*, 2000, **192**(1), 1–37.

- [11] S. Simić and M. Vuorinen, Landen inequalities for zero-balanced hypergeometric functions, *Abstr. Appl. Anal.*, **2012**, Article ID 932061, 11 pages.

ZHI-JUN GUO, SCHOOL OF MATHEMATICS AND COMPUTATION SCIENCE, HUNAN CITY UNIVERSITY, YIYANG 413000, CHINA
E-mail address: 492480045@qq.com

YAN ZHANG, SCHOOL OF MATHEMATICS AND COMPUTATION SCIENCE, HUNAN CITY UNIVERSITY, YIYANG 413000, CHINA
E-mail address: zhangyan080711zy@126.com

YU-MING CHU (CORRESPONDING AUTHOR), SCHOOL OF MATHEMATICS AND COMPUTATION SCIENCE, HUNAN CITY UNIVERSITY, YIYANG 413000, CHINA
E-mail address: chuyuming2005@126.com

YING-QING SONG, SCHOOL OF MATHEMATICS AND COMPUTATION SCIENCE, HUNAN CITY UNIVERSITY, YIYANG 413000, CHINA
E-mail address: 1452225875@qq.com