

# NOTE ON CERTAIN INEQUALITIES FOR NEUMAN MEANS

ZAI-YIN HE, YU-MING CHU, YING-QING SONG, AND XIAO-JING TAO

**ABSTRACT.** In this paper, we give the explicit formulas for the Neuman means  $N_{AH}$ ,  $N_{HA}$ ,  $N_{AC}$  and  $N_{CA}$ , and present the best possible upper and lower bounds for these means in terms of the combinations of harmonic mean  $H$ , arithmetic mean  $A$  and contraharmonic mean  $C$ .

## 1. INTRODUCTION

Let  $a, b, c \geq 0$  with  $ab + ac + bc \neq 0$ . Then the symmetric integral  $R_F(a, b, c)$  [1] of the first kind is defined as

$$R_F(a, b, c) = \frac{1}{2} \int_0^\infty [(t+a)(t+b)(t+c)]^{-1/2} dt.$$

The degenerate case of  $R_F$ , denoted by  $R_C$  plays an important role in the theory of special functions [1, 2], which is given by

$$R_C(a, b) = R_F(a, b, b).$$

For  $a, b > 0$  with  $a \neq b$ , the Schwab-Borchardt mean  $SB(a, b)$  [3-5] of  $a$  and  $b$  is given by

$$SB(a, b) = \begin{cases} \frac{\sqrt{b^2 - a^2}}{\cos^{-1}(a/b)}, & a < b, \\ \frac{\sqrt{a^2 - b^2}}{\cosh^{-1}(a/b)}, & a > b, \end{cases}$$

where  $\cos^{-1}(x)$  and  $\cosh^{-1}(x) = \log(x + \sqrt{x^2 - 1})$  are the inverse cosine and inverse hyperbolic cosine functions, respectively.

Carson [6] (see also [7, (3.21)]) proved that

$$SB(a, b) = [R_C(a^2, b^2)]^{-1}.$$

Recently, the Schwab-Borchardt mean has been the subject of intensive research. In particular, many remarkable inequalities for the Schwab-Borchardt mean and its generated means can be found in the literature [3-5, 8-11].

Let  $a > b > 0$ ,  $v = (a - b)/(a + b) \in (0, 1)$ ,  $p \in (0, \infty)$ ,  $q \in (0, \pi/2)$ ,  $r \in (0, \log(2 + \sqrt{3}))$  and  $s \in (0, \pi/3)$  be the parameters such that  $1/\cosh(p) = \cos(q) = 1 - v^2$ ,  $\cosh(r) = \sec(s) = 1 + v^2$ ,  $H(a, b) = 2ab/(a + b)$ ,  $G(a, b) = \sqrt{ab}$ ,  $A(a, b) = (a + b)/2$ ,  $Q(a, b) = \sqrt{(a^2 + b^2)/2}$  and  $C(a, b) = (a^2 + b^2)/(a + b)$  be respectively the harmonic, geometric, arithmetic, quadratic and contraharmonic means of  $a$  and  $b$ ,  $S_{AH}(a, b) = SB[A(a, b), H(a, b)]$ ,  $S_{HA}(a, b) = SB[H(a, b), A(a, b)]$ ,  $S_{AC}(a, b) = SB[A(a, b), C(a, b)]$ ,  $S_{CA}(a, b) = SB[C(a, b), A(a, b)]$ . Then Neuman [10] gave the explicit formulas

$$(1.1) \quad S_{AH}(a, b) = A(a, b) \frac{\tanh(p)}{p}, \quad S_{HA}(a, b) = A(a, b) \frac{\sin q}{q},$$

---

2000 *Mathematics Subject Classification.* 26E60.

*Key words and phrases.* Schwab-Borchardt mean; Neuman mean; harmonic mean; arithmetic mean; contraharmonic mean.

$$(1.2) \quad S_{CA}(a, b) = A(a, b) \frac{\sinh(r)}{r}, \quad S_{AC}(a, b) = A(a, b) \frac{\tan s}{s}.$$

Very recently, Neuman [12] found a new mean  $N(a, b)$  derived from the Schwab-Borchardt mean as follows:

$$(1.3) \quad N(a, b) = \frac{1}{2} \left[ a + \frac{b^2}{SB(a, b)} \right].$$

Let  $N_{AH}(a, b) = N[A(a, b), H(a, b)]$ ,  $N_{HA}(a, b) = N[H(a, b), A(a, b)]$ ,  $N_{AG}(a, b) = N[A(a, b), G(a, b)]$ ,  $N_{GA}(a, b) = N[G(a, b), A(a, b)]$ ,  $N_{AC}(a, b) = N[A(a, b), C(a, b)]$ ,  $N_{CA}(a, b) = N[C(a, b), A(a, b)]$ ,  $N_{AQ}(a, b) = N[A(a, b), Q(a, b)]$  and  $N_{QA}(a, b) = N[Q(a, b), A(a, b)]$  be the Neuman means. Then Neuman [12] proved that

$$G(a, b) < N_{AG}(a, b) < N_{GA}(a, b) < A(a, b) < N_{QA}(a, b) < N_{AQ}(a, b) < Q(a, b)$$

for all  $a, b > 0$  with  $a \neq b$ , and the double inequalities

$$\begin{aligned} \alpha_1 A(a, b) + (1 - \alpha_1) G(a, b) &< N_{GA}(a, b) < \beta_1 A(a, b) + (1 - \beta_1) G(a, b), \\ \alpha_2 Q(a, b) + (1 - \alpha_2) A(a, b) &< N_{AQ}(a, b) < \beta_2 Q(a, b) + (1 - \beta_2) A(a, b), \\ \alpha_3 A(a, b) + (1 - \alpha_3) G(a, b) &< N_{AG}(a, b) < \beta_3 A(a, b) + (1 - \beta_3) G(a, b), \\ \alpha_4 Q(a, b) + (1 - \alpha_4) A(a, b) &< N_{QA}(a, b) < \beta_4 Q(a, b) + (1 - \beta_4) A(a, b) \end{aligned}$$

hold for all  $a, b > 0$  with  $a \neq b$  if and only if  $\alpha_1 \leq 2/3$ ,  $\beta_1 \geq \pi/4$ ,  $\alpha_2 \leq 2/3$ ,  $\beta_2 \geq (\pi - 2)/[4(\sqrt{2} - 1)] = 0.689\dots$ ,  $\alpha_3 \leq 1/3$ ,  $\beta_3 \geq 1/2$ ,  $\alpha_4 \leq 1/3$  and  $\beta_4 \geq [\log(1 + \sqrt{2}) + \sqrt{2} - 2]/[2(\sqrt{2} - 1)] = 0.356\dots$

Zhang et. al. [13] presented the best possible parameters  $\alpha_1, \alpha_2, \beta_1, \beta_2 \in [0, 1/2]$  and  $\alpha_3, \alpha_4, \beta_3, \beta_4 \in [1/2, 1]$  such that the double inequalities

$$\begin{aligned} G(\alpha_1 a + (1 - \alpha_1)b, \alpha_1 b + (1 - \alpha_1)a) &< N_{AG}(a, b) < G(\beta_1 a + (1 - \beta_1)b, \beta_1 b + (1 - \beta_1)a), \\ G(\alpha_2 a + (1 - \alpha_2)b, \alpha_2 b + (1 - \alpha_2)a) &< N_{GA}(a, b) < G(\beta_2 a + (1 - \beta_2)b, \beta_2 b + (1 - \beta_2)a), \\ Q(\alpha_3 a + (1 - \alpha_3)b, \alpha_3 b + (1 - \alpha_3)a) &< N_{QA}(a, b) < Q(\beta_3 a + (1 - \beta_3)b, \beta_3 b + (1 - \beta_3)a), \\ Q(\alpha_4 a + (1 - \alpha_4)b, \alpha_4 b + (1 - \alpha_4)a) &< N_{AQ}(a, b) < Q(\beta_4 a + (1 - \beta_4)b, \beta_4 b + (1 - \beta_4)a) \end{aligned}$$

hold for all  $a, b > 0$  with  $a \neq b$ .

In [14], the authors found the greatest values  $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7, \alpha_8$  and the least values  $\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6, \beta_7, \beta_8$  such that the double inequalities

$$\begin{aligned} A^{\alpha_1}(a, b) G^{1-\alpha_1}(a, b) &< N_{GA}(a, b) < A^{\beta_1}(a, b) G^{1-\beta_1}(a, b), \\ \frac{\alpha_2}{G(a, b)} + \frac{1 - \alpha_2}{A(a, b)} &< \frac{1}{N_{GA}(a, b)} < \frac{\beta_2}{G(a, b)} + \frac{1 - \beta_2}{A(a, b)}, \\ A^{\alpha_3}(a, b) G^{1-\alpha_3}(a, b) &< N_{AG}(a, b) < A^{\beta_3}(a, b) G^{1-\beta_3}(a, b), \\ \frac{\alpha_4}{G(a, b)} + \frac{1 - \alpha_4}{A(a, b)} &< \frac{1}{N_{AG}(a, b)} < \frac{\beta_4}{G(a, b)} + \frac{1 - \beta_4}{A(a, b)}, \\ Q^{\alpha_5}(a, b) A^{1-\alpha_5}(a, b) &< N_{AQ}(a, b) < Q^{\beta_5}(a, b) A^{1-\beta_5}(a, b), \\ \frac{\alpha_6}{A(a, b)} + \frac{1 - \alpha_6}{Q(a, b)} &< \frac{1}{N_{AQ}(a, b)} < \frac{\beta_6}{A(a, b)} + \frac{1 - \beta_6}{Q(a, b)}, \\ Q^{\alpha_7}(a, b) A^{1-\alpha_7}(a, b) &< N_{QA}(a, b) < Q^{\beta_7}(a, b) A^{1-\beta_7}(a, b), \\ \frac{\alpha_8}{A(a, b)} + \frac{1 - \alpha_8}{Q(a, b)} &< \frac{1}{N_{QA}(a, b)} < \frac{\beta_8}{A(a, b)} + \frac{1 - \beta_8}{Q(a, b)} \end{aligned}$$

hold for all  $a, b > 0$  with  $a \neq b$ .

The main purpose of this paper is to give the explicit formulas for the Neuman means  $N_{AH}$ ,  $N_{HA}$ ,  $N_{AC}$  and  $N_{CA}$ , and present the best possible upper and lower

bounds for theses means in terms of the combinations of harmonic, arithmetic and contraharmonic means. Our main results are the following Theorems 1.1-1.3.

**Theorem 1.1.** *Let  $a > b > 0$ ,  $v = (a - b)/(a + b) \in (0, 1)$ ,  $p \in (0, \infty)$ ,  $q \in (0, \pi/2)$ ,  $r \in (0, \log(2 + \sqrt{3}))$  and  $s \in (0, \pi/3)$  be the parameters such that  $1/\cosh(p) = \cos(q) = 1 - v^2$ ,  $\cosh(r) = \sec(s) = 1 + v^2$ . Then we have*

$$(1.4) \quad N_{AH}(a, b) = \frac{1}{2}A(a, b) \left[ 1 + \frac{2p}{\sinh(2p)} \right],$$

$$(1.5) \quad N_{HA}(a, b) = \frac{1}{2}A(a, b) \left[ \cos(q) + \frac{q}{\sin(q)} \right],$$

$$(1.6) \quad N_{CA}(a, b) = \frac{1}{2}A(a, b) \left[ \cosh(r) + \frac{r}{\sinh(r)} \right],$$

$$(1.7) \quad N_{AC}(a, b) = \frac{1}{2}A(a, b) \left[ 1 + \frac{2s}{\sin(2s)} \right]$$

and

$$(1.8) \quad \begin{aligned} H(a, b) &< N_{AH}(a, b) < N_{HA}(a, b) < A(a, b) \\ &< N_{CA}(a, b) < N_{AC}(a, b) < C(a, b). \end{aligned}$$

**Theorem 1.2.** *The double inequalities*

$$(1.9) \quad \alpha_1 A(a, b) + (1 - \alpha_1)H(a, b) < N_{AH}(a, b) < \beta_1 A(a, b) + (1 - \beta_1)H(a, b),$$

$$(1.10) \quad \alpha_2 A(a, b) + (1 - \alpha_2)H(a, b) < N_{HA}(a, b) < \beta_2 A(a, b) + (1 - \beta_2)H(a, b),$$

$$(1.11) \quad \alpha_3 C(a, b) + (1 - \alpha_3)A(a, b) < N_{CA}(a, b) < \beta_3 C(a, b) + (1 - \beta_3)A(a, b),$$

$$(1.12) \quad \alpha_4 C(a, b) + (1 - \alpha_4)A(a, b) < N_{AC}(a, b) < \beta_4 C(a, b) + (1 - \beta_4)A(a, b)$$

hold for all  $a, b > 0$  with  $a \neq b$  if and only if  $\alpha_1 \leq 1/3$ ,  $\beta_1 \geq 1/2$ ,  $\alpha_2 \leq 2/3$ ,  $\beta_2 \geq \pi/4 = 0.7853\dots$ ,  $\alpha_3 \leq 1/3$ ,  $\beta_3 \geq \sqrt{3}\log(2 + \sqrt{3})/6 = 0.3801\dots$ ,  $\alpha_4 \leq 2/3$  and  $\beta_4 \geq (4\sqrt{3}\pi - 9)/18 = 0.7901\dots$

**Theorem 1.3.** *The double inequalities*

$$(1.13) \quad \frac{\alpha_5}{H(a, b)} + \frac{1 - \alpha_5}{A(a, b)} < \frac{1}{N_{AH}(a, b)} < \frac{\beta_5}{H(a, b)} + \frac{1 - \beta_5}{A(a, b)},$$

$$(1.14) \quad \frac{\alpha_6}{H(a, b)} + \frac{1 - \alpha_6}{A(a, b)} < \frac{1}{N_{HA}(a, b)} < \frac{\beta_6}{H(a, b)} + \frac{1 - \beta_6}{A(a, b)},$$

$$(1.15) \quad \frac{\alpha_7}{A(a, b)} + \frac{1 - \alpha_7}{C(a, b)} < \frac{1}{N_{CA}(a, b)} < \frac{\beta_7}{A(a, b)} + \frac{1 - \beta_7}{C(a, b)},$$

$$(1.16) \quad \frac{\alpha_8}{A(a, b)} + \frac{1 - \alpha_8}{C(a, b)} < \frac{1}{N_{AC}(a, b)} < \frac{\beta_8}{A(a, b)} + \frac{1 - \beta_8}{C(a, b)},$$

hold for all  $a, b > 0$  with  $a \neq b$  if and only if  $\alpha_5 \leq 0$ ,  $\beta_5 \geq 2/3$ ,  $\alpha_6 \leq 0$ ,  $\beta_6 \geq 1/3$ ,  $\alpha_7 \leq [2\sqrt{3} - \log(2 + \sqrt{3})]/[2\sqrt{3} + \log(2 + \sqrt{3})] = 0.4490\dots$ ,  $\beta_7 \geq 2/3$ ,  $\alpha_8 \leq (9\sqrt{3} - 4\pi)/(3\sqrt{3} + 4\pi) = 0.1701\dots$  and  $\beta_8 \geq 1/3$ .

## 2. LEMMAS

In order to prove our main results we need several lemmas, which we present in this section.

**Lemma 2.1.** (See [15, Theorem 1.25]) For  $-\infty < a < b < \infty$ , let  $f, g : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ , and be differentiable on  $(a, b)$ , let  $g'(x) \neq 0$  on  $(a, b)$ . If  $f'(x)/g'(x)$  is increasing (decreasing) on  $(a, b)$ , then so are

$$\frac{f(x) - f(a)}{g(x) - g(a)} \quad \text{and} \quad \frac{f(x) - f(b)}{g(x) - g(b)}.$$

If  $f'(x)/g'(x)$  is strictly monotone, then the monotonicity in the conclusion is also strict.

**Lemma 2.2.** (See [16, Lemma 1.1]) Suppose that the power series  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  and  $g(x) = \sum_{n=0}^{\infty} b_n x^n$  have the radius of convergence  $r > 0$  and  $a_n, b_n > 0$  for all  $n \geq 0$ . If the sequence  $\{a_n/b_n\}$  is (strictly) increasing (decreasing) for all  $n \geq 0$ , then the function  $f(x)/g(x)$  is also (strictly) increasing (decreasing) on  $(0, r)$ .

**Lemma 2.3.** (See [12, Theorem 4.1]) If  $a > b$ , then

$$N(b, a) > N(a, b).$$

**Lemma 2.4.** The function

$$\varphi_1(t) = \frac{\sinh(2t) - 4 \sinh(t) + 2t}{\sinh(2t) - 2 \sinh(t)}$$

is strictly increasing from  $(0, \infty)$  onto  $(2/3, 1)$ .

*Proof.* Making use of power series expansion we get

$$(2.1) \quad \varphi_1(t) = \frac{\sum_{n=1}^{\infty} \frac{2^{2n+1}-4}{(2n+1)!} t^{2n+1}}{\sum_{n=1}^{\infty} \frac{2^{2n+1}-2}{(2n+1)!} t^{2n+1}} = \frac{\sum_{n=0}^{\infty} \frac{2^{2n+3}-4}{(2n+3)!} t^{2n}}{\sum_{n=0}^{\infty} \frac{2^{2n+3}-2}{(2n+3)!} t^{2n}}.$$

Let

$$(2.2) \quad a_n = \frac{2^{2n+3}-4}{(2n+3)!}, \quad b_n = \frac{2^{2n+3}-2}{(2n+3)!}.$$

Then

$$(2.3) \quad a_n > 0, \quad b_n > 0$$

and  $a_n/b_n = 1 - 1/(2^{2n+2} - 1)$  is strictly increasing for all  $n \geq 0$ .

Note that

$$(2.4) \quad \varphi_1(0^+) = \frac{a_0}{b_0} = \frac{2}{3}, \quad \varphi_1(\infty) = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1.$$

Therefore, Lemma 2.4 follows easily from Lemma 2.2 and (2.1)-(2.4) together with the monotonicity of the sequence  $\{a_n/b_n\}$ .  $\square$

**Lemma 2.5.** *The function*

$$\varphi_2(t) = \frac{2t - \sin(2t)}{\sin t(1 - \cos t)}$$

*is strictly increasing from  $(0, \pi/2)$  onto  $(8/3, \pi)$ .*

*Proof.* Let  $f_1(t) = 2t - \sin(2t)$  and  $g_1(t) = \sin t(1 - \cos t)$ . Then simple computations lead to

$$(2.5) \quad \varphi_2(t) = \frac{f_1(t) - f_1(0)}{g_1(t) - g_1(0)}$$

and  $f'_1(t)/g'_1(t) = 4[1 - 1/(2 + 1/\cos t)]$  is strictly increasing on  $(0, \pi/2)$ .

Note that

$$(2.6) \quad \varphi_2(0^+) = \lim_{t \rightarrow 0^+} \frac{f'_1(t)}{g'_1(t)} = \frac{8}{3}, \quad \varphi_2(\pi/2) = \pi.$$

Therefore, Lemma 2.5 follows from Lemma 2.1, (2.5), (2.6) and the monotonicity of  $f'_1(t)/g'_1(t)$ .  $\square$

**Lemma 2.6.** *The function*

$$\varphi_3(t) = \frac{\sinh(t) \cosh(t) - t}{[\sinh(t) \cosh(t) + t](\cosh(t) - 1)}$$

*is strictly decreasing from  $(0, \infty)$  onto  $(0, 2/3)$ .*

*Proof.* Simple computations lead to

$$(2.7) \quad \begin{aligned} \varphi_3(t) &= \frac{2 \sinh(2t) - 4t}{\sinh(3t) + 4t \cosh(t) + \sinh(t) - 2 \sinh(2t) - 4t} \\ &= \frac{\sum_{n=0}^{\infty} \frac{2^{2n+4}}{(2n+3)!} t^{2n}}{\sum_{n=0}^{\infty} \frac{3^{2n+3} - 2^{2n+4} + 8n + 13}{(2n+3)!} t^{2n}}. \end{aligned}$$

Let

$$(2.8) \quad a_n = \frac{2^{2n+4}}{(2n+3)!}, \quad b_n = \frac{3^{2n+3} - 2^{2n+4} + 8n + 13}{(2n+3)!}.$$

Then

$$(2.9) \quad a_n > 0, \quad b_n > 0$$

and

$$(2.10) \quad \frac{a_{n+1}}{b_{n+1}} - \frac{a_n}{b_n} = -\frac{2^{2n+4} (5 \times 3^{2n+3} - 24n - 31)}{(3^{2n+5} - 2^{2n+6} + 8n + 21)(3^{2n+3} - 2^{2n+4} + 8n + 13)} < 0$$

for all  $n \geq 0$ .

Note that

$$(2.11) \quad \varphi_3(0^+) = \frac{a_0}{b_0} = \frac{2}{3}, \quad \varphi_3(\infty) = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0.$$

Therefore, Lemma 2.6 follows easily from (2.7)-(2.11) and Lemma 2.2.  $\square$

**Lemma 2.7.** *The function*

$$f(t) = 9 \cos t + \frac{t}{\sin t}$$

*is strictly decreasing on the interval  $(0, \pi/2)$ .*

*Proof.* Let  $f_2(t) = 9 \sin t \cos t + t$  and  $g_2(t) = \sin t$ . Then simple computations lead to

$$(2.12) \quad f(t) = \frac{f_2(t) - f_2(0)}{g_2(t) - g_2(0)},$$

$$\frac{f'_2(t)}{g'_2(t)} = \frac{18 \cos^2 t - 8}{\cos t}$$

and

$$(2.13) \quad \left[ \frac{f'_2(t)}{g'_2(t)} \right]' = -\frac{2 \sin t (9 \cos^2 t + 4)}{\cos^2(t)} < 0$$

for  $t \in (0, \pi/2)$ .

Therefore, Lemma 2.7 follows easily from (2.12) and (2.13) together with Lemma 2.1.  $\square$

**Lemma 2.8.** *The function*

$$\varphi_4(t) = \frac{\sin t \cos t - t}{(t + \sin t \cos t)(1 - \cos t)}$$

is strictly decreasing from  $(0, \pi/2)$  onto  $(-1, -2/3)$ .

*Proof.* Let  $f_3(t) = \sin t \cos t - t$  and  $g_3(t) = (t + \sin t \cos t)(1 - \cos t)$ . Then simple computations lead to

$$(2.14) \quad \varphi_4(t) = \frac{f_3(t)}{g_3(t)} = \frac{f_3(t) - f_3(0)}{g_3(t) - g_3(0)},$$

$$(2.15) \quad \frac{f'_3(t)}{g'_3(t)} = \frac{f'_3(t) - f'_3(0)}{g'_3(t) - g'_3(0)}$$

and

$$(2.16) \quad \frac{f''_3(t)}{g''_3(t)} = \frac{4}{4 - (9 \cos t + \frac{t}{\sin t})}.$$

Note that

$$(2.17) \quad \varphi_4(0^+) = \lim_{t \rightarrow 0^+} \frac{f''_3(t)}{g''_3(t)} = -\frac{2}{3}, \quad \varphi_4\left(\frac{\pi}{2}\right) = -1.$$

Therefore, Lemma 2.8 follows from Lemma 2.1 and Lemma 2.7 together with (2.14)-(2.17).  $\square$

### 3. PROOFS OF THEOREMS 1.1-1.3

**Proof of Theorem 1.1.** It follows from (1.1)-(1.3) we clearly see that

$$\begin{aligned} N_{AH}(a, b) &= \frac{1}{2} \left[ A(a, b) + \frac{H^2(a, b)}{S_{AH}(a, b)} \right] = \frac{1}{2} A(a, b) \left[ 1 + (1 - v^2)^2 \frac{p}{\tanh(p)} \right] \\ &= \frac{1}{2} A(a, b) \left[ 1 + \frac{p}{\tanh(p) \cosh^2(p)} \right] = \frac{1}{2} A(a, b) \left[ 1 + \frac{2p}{\sinh(2p)} \right], \\ N_{HA}(a, b) &= \frac{1}{2} \left[ H(a, b) + \frac{A^2(a, b)}{S_{HA}(a, b)} \right] \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2}A(a, b) \left[ (1 - v^2) + \frac{q}{\sin q} \right] = \frac{1}{2}A(a, b) \left[ \cos q + \frac{q}{\sin q} \right], \\
N_{CA}(a, b) &= \frac{1}{2} \left[ C(a, b) + \frac{A^2(a, b)}{S_{CA}(a, b)} \right] \\
&= \frac{1}{2}A(a, b) \left[ (1 + v^2) + \frac{r}{\sinh(r)} \right] = \frac{1}{2}A(a, b) \left[ \cosh(r) + \frac{r}{\sinh(r)} \right], \\
N_{AC}(a, b) &= \frac{1}{2} \left[ A(a, b) + \frac{C^2(a, b)}{S_{AC}(a, b)} \right] = \frac{1}{2}A(a, b) \left[ 1 + (1 + v^2)^2 \frac{s}{\tan(s)} \right] \\
&= \frac{1}{2}A(a, b) \left[ 1 + \frac{s}{\tan(s) \cos^2 s} \right] = \frac{1}{2}A(a, b) \left[ 1 + \frac{2s}{\sin(2s)} \right].
\end{aligned}$$

Inequalities (1.8) follows easily from  $H(a, b) < A(a, b) < C(a, b)$  and Lemma 2.3 together with the fact that  $N_{KL}(a, b)$  is a mean of  $K(a, b)$  and  $L(a, b)$  for  $K(a, b), L(a, b) \in \{H(a, b), A(a, b), C(a, b)\}$ .  $\square$

**Proof of Theorem 1.2.** Without loss of generality, we assume that  $a > b$ . Let  $v = (a - b)/(a + b) \in (0, 1)$ ,  $p \in (0, \infty)$ ,  $q \in (0, \pi/2)$ ,  $r \in (0, \log(2 + \sqrt{3}))$  and  $s \in (0, \pi/3)$  be the parameters such that  $1/\cosh(p) = \cos(q) = 1 - v^2$ ,  $\cosh(r) = \sec(s) = 1 + v^2$ . Then from (1.4)- (1.7) we have

$$\begin{aligned}
(3.1) \quad \frac{N_{AH}(a, b) - H(a, b)}{A(a, b) - H(a, b)} &= \frac{\frac{1}{2} \left[ 1 + \frac{2p}{\sinh(2p)} \right] - (1 - v^2)}{v^2} \\
&= \frac{\frac{1}{2} \left[ 1 + \frac{2p}{\sinh(2p)} \right] - \frac{1}{\cosh(p)}}{1 - \frac{1}{\cosh(p)}} = \varphi_1(p),
\end{aligned}$$

$$\begin{aligned}
(3.2) \quad \frac{N_{HA}(a, b) - H(a, b)}{A(a, b) - H(a, b)} &= \frac{\frac{1}{2} \left[ \cos q + \frac{q}{\sin q} \right] - (1 - v^2)}{v^2} \\
&= \frac{\frac{1}{2} \left[ \cos q + \frac{q}{\sin q} \right] - \cos q}{1 - \cos q} = \frac{1}{4} \varphi_2(q),
\end{aligned}$$

$$\begin{aligned}
(3.3) \quad \frac{N_{CA}(a, b) - A(a, b)}{C(a, b) - A(a, b)} &= \frac{\frac{1}{2} \left[ \cosh(r) + \frac{r}{\sinh(r)} \right] - 1}{v^2} \\
&= \frac{\frac{1}{2} \left[ \cosh(r) + \frac{r}{\sinh(r)} \right] - 1}{\cosh(r) - 1} = \frac{1}{2} \varphi_1(r).
\end{aligned}$$

$$\begin{aligned}
(3.4) \quad \frac{N_{AC}(a, b) - A(a, b)}{C(a, b) - A(a, b)} &= \frac{\frac{1}{2} \left[ 1 + \frac{2s}{\sin(2s)} \right] - 1}{v^2} \\
&= \frac{\frac{1}{2} \left[ 1 + \frac{2s}{\sin(2s)} \right] - 1}{\sec(s) - 1} = \frac{1}{4} \varphi_2(s),
\end{aligned}$$

where the functions  $\varphi_1$  and  $\varphi_2$  are defined as in Lemmas 2.4 and 2.5, respectively.

Note that

$$(3.5) \quad \varphi_1[\log(2 + \sqrt{3})] = \sqrt{3} \log(2 + \sqrt{3})/6$$

and

$$(3.6) \quad \varphi_2\left(\frac{\pi}{3}\right) = \frac{8\sqrt{3}\pi - 18}{9}.$$

Therefore, inequality (1.9) holds for all  $a, b > 0$  with  $a \neq b$  if and only if  $\alpha_1 \leq 1/3$  and  $\beta_1 \geq 1/2$  follows from (3.1) and Lemma 2.4, inequality (1.10) holds for all  $a, b > 0$  with  $a \neq b$  if and only if  $\alpha_2 \leq 2/3$  and  $\beta_2 \geq \pi/4$  follows from (3.2) and Lemma 2.5, inequality (1.11) holds for all  $a, b > 0$  with  $a \neq b$  if and only if  $\alpha_3 \leq 1/3$  and  $\beta_3 \geq \sqrt{3}\log(2 + \sqrt{3})/6$  follows from (3.3) and (3.5) together with Lemma 2.4, and inequality (1.12) holds for all  $a, b > 0$  with  $a \neq b$  if and only if  $\alpha_4 \leq 2/3$  and  $\beta_4 \geq (4\sqrt{3}\pi - 9)/18$  follows from (3.4) and (3.6) together with Lemma 2.5.  $\square$

**Proof of Theorem 1.3.** Without loss of generality, we assume that  $a > b$ . Let  $v = (a - b)/(a + b) \in (0, 1)$ ,  $p \in (0, \infty)$ ,  $q \in (0, \pi/2)$ ,  $r \in (0, \log(2 + \sqrt{3}))$  and  $s \in (0, \pi/3)$  be the parameters such that  $1/\cosh(p) = \cos(q) = 1 - v^2$ ,  $\cosh(r) = \sec(s) = 1 + v^2$ . Then from (1.4)-(1.7) we have

$$(3.7) \quad \frac{\frac{1}{N_{AH}(a,b)} - \frac{1}{A(a,b)}}{\frac{1}{H(a,b)} - \frac{1}{A(a,b)}} = \frac{\frac{2}{1 + \frac{2p}{\sinh(2p)}} - 1}{\frac{1}{1-v^2} - 1} = \frac{\frac{2 \sinh(2p)}{2p + \sinh(2p)} - 1}{\cosh(p) - 1} = \varphi_3(p),$$

$$(3.8) \quad \frac{\frac{1}{N_{HA}(a,b)} - \frac{1}{A(a,b)}}{\frac{1}{H(a,b)} - \frac{1}{A(a,b)}} = \frac{\frac{2}{\cos(q) + \frac{q}{\sin(q)}} - 1}{\frac{1}{1-v^2} - 1} = \frac{\frac{2 \sin(q) - \sin(q) \cos(q) - q}{\sin(q) \cos(q) + q} - 1}{\frac{1 - \cos(q)}{\cos(q)}} = 1 + \varphi_4(q),$$

$$(3.9) \quad \frac{\frac{1}{N_{CA}(a,b)} - \frac{1}{C(a,b)}}{\frac{1}{A(a,b)} - \frac{1}{C(a,b)}} = \frac{\frac{2}{\cosh(r) + \frac{r}{\sinh(r)}} - \frac{1}{1+v^2}}{1 - \frac{1}{1+v^2}} = \varphi_3(r)$$

and

$$(3.10) \quad \frac{\frac{1}{N_{AC}(a,b)} - \frac{1}{C(a,b)}}{\frac{1}{A(a,b)} - \frac{1}{C(a,b)}} = 1 + \varphi_4(s),$$

where the functions  $\varphi_3$  and  $\varphi_4$  are defined as in Lemmas 2.6 and 2.8, respectively.

Note that

$$(3.11) \quad \varphi_3[\log(2 + \sqrt{3})] = \frac{2\sqrt{3} - \log(2 + \sqrt{3})}{2\sqrt{3} + \log(2 + \sqrt{3})}$$

and

$$(3.12) \quad \varphi_4\left(\frac{\pi}{3}\right) = -\frac{8\pi - 6\sqrt{3}}{4\pi + 3\sqrt{3}}.$$

Therefore, Theorem 1.3 follows easily from (3.7)-(3.12) together with Lemmas 2.6 and 2.8.  $\square$

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

### Acknowledgements

This research was supported by the Natural Science Foundation of China under Grants 61374086, 11371125 and 11171307, and the Natural Science Foundation of Zhejiang Province under Grant LY13A010004.

#### REFERENCES

- [1] B. C. Carlson, Special Functions of Applied Mathematics, Academic Press, New York, 1977.
- [2] B. C. Carlson and J. L. Gustafson, Asymptotic approximations for symmetric elliptic integrals, SIAM J. Math. Anal., 1994, **25**(2): 288-303.
- [3] E. Neuman and J. Sndor, On the Schwab-Borchardt mean, Math. Pannon., 2003, **14**(2): 253-266.
- [4] E. Neuman and J. Sndor, On the Schwab-Borchardt mean II, Math. Pannon., 2005, **17**(1): 49-59.
- [5] E. Neuman, Inequalities for the Schwab-Boorhhardt mean and their applications, J. Math. Inequal., 2011, **5**(4): 601-609.
- [6] B. C. Carlson, Algorithms involving arithmetic and geometric means, Amer. Math. Monthly, 1971, **78**: 496-505.
- [7] J. L. Brenner and B. C. Carlson, Homogeneous mean values: weights and asymptotics, J. Math. Anal. Appl., 1987, **123**(1): 265-280.
- [8] Z.-Y. He, Y.-M. Chu and M.-K. Wang, Optimal bounds for Neuman means in terms of harmonic and contraharmonic means, J. Appl. Math., **2013**, Article ID 807623, 4 pages.
- [9] Y.-M. Chu and W.-M. Qian, Refinements of Bounds for Neuman means, Abstr. Appl. Anal., **2014**, Article ID 354132, 8 pages.
- [10] E. Neuman, On some means derived from the Schwab-Borchardt mean, J. Math. Inequal., 2014, **8**(1): 171-183.
- [11] E. Neuman, On some means derived from the Schwab-Borchardt mean II, J. Math. Inequal., 2014, **8**(2): 361-370.
- [12] E. Neuman, On a new bivariate mean, Aequat. Math., DOI 10.1007/s00010-013-2004-8.
- [13] Y. Zhang, Y.-M. Chu and Y.-L. Jiang, Sharp geometric mean bounds for Neuman means, Abstr. Appl. Anal., **2014**, Article ID 949815, 6 pages.
- [14] Z.-J. Guo, Y. Zhang, Y.-M. Chu and Y.-Q. Song, Sharp bounds for Neuman means in terms of geometric, arithmetic and quadratic mean, Preprint.
- [15] G. D. Anderson, S.-L. Qiu, M. K. Vamanamurthy and M. Vuorinen, Generalized elliptic integrals and modular equations, Pacific J. Math., 2000, **192**(1): 1-37.
- [16] S. Simic and M. Vuorinen, Landen inequalities for zero-balanced hypergeometric functions, Abstr. Appl. Anal., **2012**, Article ID 932061, 11 pages.

ZAI-YIN HE, SCHOOL OF MATHEMATICS AND COMPUTATION SCIENCE, HUNAN CITY UNIVERSITY, YIYANG 413000, P.R. CHINA

YU-MING CHU (CORRESPONDING AUTHOR), SCHOOL OF MATHEMATICS AND COMPUTATION SCIENCE, HUNAN CITY UNIVERSITY, YIYANG 413000, P.R. CHINA

*E-mail address:* chuyuming2005@126.com

YING-QING SONG, SCHOOL OF MATHEMATICS AND COMPUTATION SCIENCE, HUNAN CITY UNIVERSITY, YIYANG 413000, P.R. CHINA

XIAO-JING TAO, SCHOOL OF MATHEMATICS AND ECONOMETRICS, HUNAN UNIVERSITY, CHANGSHA 410082, P.R. CHINA