



34	41
1 41 224	1 34 84 224
1 34 34 41 84 224	1 1 34 41 41 41 84 224
1 1 1 34 41 41 84 224	1 1 1 34 34 41 84 84 224 224
1 34 34 41 84 224	1 1 34 41 41 41 84 224
1 41 224	1 34 84 224
34	41
84	224
1 41 84	1 34 41 224
1 34 41 84 84 224	1 1 34 41 84 224 224
1 1 34 41 41 84 224	1 1 34 41 41 84 224 22
1 34 41 84 84 224	1 1 34 41 84 224 224
1 41 84	1 34 41 224
84	224

**Theorem 2.** *The Loewy structure of the principal indecomposable modules of  $B_1$  are as follows:*

9	36	90
126 279	126 279	126
9 9 36 90	9 36 36 90	9 36 90
126 279	126 279	126
9	36	90
126	279	
9 36 90	9 36	
126 126 279	126 279	
9 36 90	9 36	
126	279	

We note that the Loewy and socle series for the principal indecomposable modules (PIMs) over  $F$  are the same, which is also the case for  $A_6, A_7, A_8$  and  $A_9$ .

The Loewy structure of the PIMs for  $FA_6$  is well-known and appears in (Benson, 1984); their module diagrams can be found in (Benson and Carlson, 1987). The structures for  $FA_7$  and  $FA_8$  have been calculated in (Scopes, 1988), while that for  $FA_9$  has been done in (Siegel, 1991). The structures for  $FA_8$  and  $FA_9$  have also been done in characteristic two in (Benson, 1983a) and (Benson, 1983b) respectively. The results for  $FA_{10}$  in characteristic 3 have not been published to our knowledge.

If  $A$  is a group algebra over  $F$  and  $M$  a finitely generated  $A$ -module, write  $L_i(M)$  and  $S_i(M)$  for the  $i^{\text{th}}$  Loewy layer of  $M$  and the  $i^{\text{th}}$  socle layer of  $M$ , respectively. We shall write  $(M, N)_A$  for  $\dim_F \text{Hom}_A(M, N)$ ,  $M^*$  for  $\text{Hom}_F(M, F)$  regarded as an  $A$ -module,  $(M, N)_A^1$  for  $\dim_F \text{Ext}_A^1(M, N)$ , and  $P_M$  for the projective cover of  $M$ . We also denote a uniserial module of Loewy length 3 with head  $S$ , heart  $T$  and socle  $U$  by  $\mathfrak{U}(S; T; U)$ , and a module  $M$  with  $L_1(M) = S_1, L_2(M) = S_2 \oplus S_3$  and  $L_3(M) = S_4$  by  $\mathfrak{D}(S_1; S_2, S_3; S_4)$  (here all the  $S_i$  are simple).

Facts about modular representation theory can be found in (Landrock, 1983). We use Brauer characters and GAP (GAP, 2013) to find the composition factors of the  $A_{10}$ -modules we study. We will occasionally use a (Benson-Carlson) module diagram to describe an  $A$ -module  $M$ . This is a finite directed graph with vertices labelled by simple modules, and with an edge from a vertex  $S$  to a vertex  $T$  corresponding to a non-trivial



$$\begin{array}{cccccc}
& & 41 & & & 35 \\
& & 1 & 7 & 35 & & & 1 & 7 & 21 & 41 & 35 \\
& 1 & 7 & 21 & 41 & 41 & 35 & & 1 & 1 & 7 & 7 & 21 & 41 & 35 & 35 & 35 \\
& 1 & 1 & 7 & 7 & 21 & 35 & 35 & 1 & 1 & 7 & 7 & 21 & 21 & 41 & 41 & 35 & 35 & 35 \\
& 1 & 7 & 21 & 41 & 41 & 35 & & 1 & 1 & 7 & 7 & 21 & 41 & 35 & 35 & 35 \\
& & 1 & 7 & 35 & & & & & 1 & 7 & 21 & 41 & 35 \\
& & 41 & & & & & & & & & & & 35
\end{array}$$

ii) *Non-principal blocks:*

$$\mathfrak{U}(27; 189; 27) , \mathfrak{U}(189; 27; 189) , 162 .$$

### 3. INDUCTION AND RESTRICTION BETWEEN $A_9$ AND $A_{10}$

#### 3.1. Restriction from $A_{10}$ to $A_9$ .

**Theorem 7.**

$$\begin{aligned}
1_{A_{10}} \downarrow &= 1_{A_9} , \quad 9_{A_{10}} \downarrow = \mathfrak{U}(1; 7; 1) , \quad 34_{A_{10}} \downarrow = 7_{A_9} \oplus 27_{A_9} , \quad 36_{A_{10}} \downarrow = \mathfrak{D}(7; 1, 21; 7) , \\
41_{A_{10}} \downarrow &= 41_{A_9} , \quad 84_{A_{10}} \downarrow = \mathfrak{D}(21; 7, 35; 21) , \quad 90_{A_{10}} \downarrow = \mathfrak{D}(41; 1, 7; 41) , \\
126_{A_{10}} \downarrow &= \mathfrak{D}(35; 21, 35; 35) , \quad 224_{A_{10}} \downarrow = 35_{A_9} \oplus 189_{A_9} , \\
279_{A_{10}} \downarrow &= 13_{A_8} \uparrow^{A_9} \oplus 162_{A_9} = \mathfrak{U}(41; 35; 41) , \quad 567_{A_{10}} \downarrow = 162_{A_9} \oplus \mathfrak{U}(189; 27; 189) .
\end{aligned}$$

*Proof.* This follows easily from Frobenius Reciprocity, block theory and self-duality of all the modules involved. For example, using Brauer characters,  $36_{A_{10}} \downarrow$  has composition factors  $1_{A_9} + 2(7_{A_9}) + 21_{A_9}$ . By Frobenius Reciprocity,  $L_1(36 \downarrow) = S_1(36 \downarrow) = 7$ . Hence, self-duality of  $36 \downarrow$  gives the result.  $\square$

#### 3.2. Induction from $A_9$ to $A_{10}$ .

**Theorem 8.**

$$\begin{aligned}
1_{A_9} \uparrow &= 1_{A_{10}} \oplus 9_{A_{10}} , \quad 7_{A_9} \uparrow = 34_{A_{10}} \oplus 36_{A_{10}} , \quad 21_{A_9} \uparrow = \mathfrak{D}(84; 1, 41; 84) , \\
35_{A_9} \uparrow &= 126_{A_{10}} \oplus 224_{A_{10}} , \quad 41_{A_9} \uparrow = 41_{A_{10}} \oplus 90_{A_{10}} \oplus 279_{A_{10}} , \\
162_{A_9} \uparrow &= P_{162_{A_9}} \uparrow = P_{162_{A_{10}}} \oplus P_{567_{A_{10}}} ,
\end{aligned}$$

$$\begin{array}{cccc}
& & 34 & & & 224 \\
& & 1 & 41 & & 1 & 41 & 224 \\
27_{A_9} \uparrow = & 34 & 84 & , & 189_{A_9} \uparrow = & 567_{A_{10}} \oplus & 1 & 34 & 84 & 224 & . \\
& & 1 & 41 & & & 1 & 41 & 224 \\
& & 34 & & & & & & 224
\end{array}$$


*Proof.* With the exception of  $27 \uparrow$  and  $189 \uparrow$ , all structures follow from block theory, Frobenius Reciprocity, Brauer characters and self-duality.

For  $27 \uparrow$  and  $189 \uparrow$ , the MeatAxe gives the claimed Loewy layers. Furthermore, the submodule lattice for  $27 \uparrow$  implies that every simple module in  $L_i(27 \uparrow)$  extends every simple module in  $L_{i+1}(27 \uparrow)$ .

For  $189 \uparrow$ , we only obtain partial information regarding extensions between modules in  $L_3(189 \uparrow)$  and modules in both  $L_2(189 \uparrow)$  and  $L_4(189 \uparrow)$ , namely,

- 34 extends 1s, 41s and 224s;
- 84 extends only 1s and 41s;
- 224 extends 1s and 41s (and may or may not extend 224s),
- 1 in  $L_3(189 \uparrow)$  extends 41s in  $L_2(189 \uparrow)$  and  $L_4(189 \uparrow)$  (and possibly other simple modules).

A partial diagram for  $189 \uparrow$  is as follows:



□

#### 4. CALCULATION OF $(S, T)_{A_{10}}^1$ FOR SIMPLE $FA_{10}$ -MODULES $S$ AND $T$

##### 4.1. Non-principal block.

**Lemma 1.**

$$(9, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S \in \{126, 279\}; \\ 0, & \text{otherwise.} \end{cases}$$

*Proof.* We have

$$\begin{aligned}
 (9, S)_{A_{10}}^1 &= (1 \oplus 9, S)_{A_{10}}^1 && \text{by block theory,} \\
 &= (1 \uparrow_{A_9}^{A_{10}}, S)_{A_{10}}^1 && \text{by Theorem 8,} \\
 &= (1_{A_9}, S \downarrow_{A_9}^{A_{10}})_{A_9}^1 && \text{by Ext Reciprocity.}
 \end{aligned}$$

By Theorem 3, Theorem 5 and Theorem 7,

$$(1_{A_9}, 9 \downarrow_{A_9}^{A_{10}})_{A_9}^1 = (1_{A_9}, \mathfrak{U}(1; 7; 1))_{A_9}^1 = (1_{A_9}, 1 \uparrow_{A_8}^{A_9})_{A_9}^1 = (1_{A_8}, 1_{A_8})_{A_8}^1 = 0.$$

By Theorem 3, Theorem 5, Theorem 7 and Theorem 8,

$$(1_{A_9}, 36 \downarrow_{A_9}^{A_{10}})_{A_9}^1 = (1_{A_9}, 36 \downarrow_{A_9}^{A_{10}} \oplus 27_{A_9})_{A_9}^1 = (1_{A_9}, 7 \uparrow_{A_8}^{A_9})_{A_9}^1 = (1_{A_8}, 7_{A_8})_{A_8}^1 = 0.$$

$$(1_{A_9}, 90 \downarrow_{A_9}^{A_{10}})_{A_9}^1 = (1_{A_9}, 90 \downarrow_{A_9}^{A_{10}} \oplus 162_{A_9})_{A_9}^1 = (1_{A_9}, 28 \uparrow_{A_8}^{A_9})_{A_9}^1 = (1_{A_8}, 28_{A_8})_{A_8}^1 = 0.$$

$$(1_{A_9}, 126 \downarrow_{A_9}^{A_{10}})_{A_9}^1 = (1_{A_9}, 126 \downarrow_{A_9}^{A_{10}} \oplus 189_{A_9})_{A_9}^1 = (1_{A_9}, 35 \uparrow_{A_8}^{A_9})_{A_9}^1 = (1_{A_8}, 35_{A_8})_{A_8}^1 = 1.$$

$$(1_{A_9}, 279 \downarrow_{A_9}^{A_{10}})_{A_9}^1 = (1_{A_9}, 13 \uparrow_{A_8}^{A_9})_{A_9}^1 = (1_{A_8}, 13_{A_8})_{A_8}^1 = 1. \quad \square$$

A similar proof gives

**Lemma 2.**

$$(36, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S \in \{126, 279\}; \\ 0, & \text{otherwise.} \end{cases}$$

$$(90, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S = 126 \\ 0, & \text{otherwise.} \end{cases}$$

$$(126, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S \in \{9, 36, 90\}; \\ 0, & \text{otherwise.} \end{cases}$$

$$(279, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S \in \{9, 36\}; \\ 0, & \text{otherwise.} \end{cases}$$

#### 4.2. Principal block.

##### Lemma 3.

$$(1, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S \in \{34, 41, 84, 224\}; \\ 0, & \text{otherwise.} \end{cases}$$

$$(34, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S \in \{1, 41, 224\}; \\ 0, & \text{otherwise.} \end{cases}$$

$$(41, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S \in \{1, 34, 84, 224\}; \\ 0, & \text{otherwise.} \end{cases}$$

$$(84, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S \in \{1, 41, 84\}; \\ 0, & \text{otherwise.} \end{cases}$$

$$(224, S)_{A_{10}}^1 = \begin{cases} 1, & \text{if } S \in \{1, 34, 41, 224\}; \\ 0, & \text{otherwise.} \end{cases}$$

*Proof.* The difficulty arises only when computing  $(84, S)_{A_{10}}^1$  (noting that, since all the simple  $FA_{10}$ -modules are self-dual,  $(S, T)_{A_{10}}^1 = (T, S)_{A_{10}}^1$ ). In all other cases, a similar proof to that in the case of non-principal blocks will suffice.

By block theory, Theorem 3, Theorem 5, Theorem 7 and Theorem 8,

$$(84, 1)_{A_{10}}^1 = (84, 1 \uparrow_{A_9}^{A_{10}})_{A_{10}}^1 = (84 \downarrow_{A_9}, 1_{A_9})_{A_9}^1 = (\mathfrak{D}(21; 7, 35; 21), 1)_{A_9}^1.$$

Furthermore, Theorem 6 implies that there are only three possibilities:

$$\begin{array}{c} 21 \\ \wedge \\ 7 \quad 35 \\ \begin{array}{c} | \quad | \\ \diagdown \quad / \\ 1 \quad 21 \end{array} \end{array}, \quad \begin{array}{c} 21 \\ \wedge \\ 7 \quad 35 \\ \begin{array}{c} | \quad | \\ | \quad | \\ 1 \quad 21 \end{array} \end{array}, \quad \begin{array}{c} 21 \\ \wedge \\ 7 \quad 35 \\ \begin{array}{c} | \quad | \\ | \quad | \\ 21 \quad 1 \end{array} \end{array}$$

However, there is a unique copy of 1 in  $L_3(P_{21A_9})$  and the first structure exists by the submodule lattice of  $21 \uparrow_{A_8}^{A_9} \cdot f_0$  obtained by the MeatAxe. Therefore, we see that the other two cases cannot occur (see 6.2.2).

The same proof gives  $(84, 41)_{A_{10}}^1 = 1$ .

Now using appropriate long exact sequences of cohomology, we have

$$(84, 84)_{A_{10}}^1 \geq 1; \quad (84, 34)_{A_{10}}^1 \leq 1; \quad (84, 224)_{A_{10}}^1 \leq 1;$$

We will finish determining the dimensions of the above Ext-spaces in Section 6.2.1 and Section 6.2.2.  $\square$

5. INDUCTION OF 2-STEP  $FA_9$  - MODULES

In this section we shall induce to  $A_{10}$  each of the non-trivial extensions of a simple module by a simple module for  $A_9$ .

By Frobenius reciprocity, we have

$$\begin{pmatrix} 1 \\ 7 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \frac{1}{34} \quad , \quad \begin{pmatrix} 1 \\ 41 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \frac{1}{41} \quad , \quad \begin{pmatrix} 1 \\ 35 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \frac{1}{224} \quad ,$$

$$\begin{pmatrix} 7 \\ 35 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \frac{34}{224} \quad , \quad \begin{pmatrix} 7 \\ 41 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \frac{34}{41} \quad ,$$

$$\begin{pmatrix} 35 \\ 35 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \frac{224}{224} \quad , \quad \begin{pmatrix} 35 \\ 41 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \frac{224}{41} \quad ,$$

$$\begin{pmatrix} 21 \\ 7 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \begin{matrix} 84 & & \\ 1 & 41 & \\ 84 & 34 & \end{matrix} \quad , \quad \begin{pmatrix} 21 \\ 35 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \begin{matrix} 84 & & \\ 1 & 41 & \\ 84 & 224 & \end{matrix} .$$

We shall determine the structure of  $\begin{pmatrix} 27 \\ 189 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0$  and  $\begin{pmatrix} 189 \\ 27 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0$  in Section 6.2.

 6. STRUCTURE OF PROJECTIVE MODULES FOR  $FA_{10}$ 

**6.1. The non-principal block.** We will use Section 4.1 and self-duality to find the structure of  $P_{90}$ ,  $P_{279}$ ,  $P_9$  and  $P_{36}$ . For  $P_{126}$ , we will additionally need Landrock's lemma.

The Cartan matrix shows that  $P_{90A_{10}}$  has composition factors  $9 + 36 + 3(90) + 2(126)$ .

Also, it has both head and socle isomorphic to 90 i.e.  $P_{90A_{10}} = \begin{matrix} 90 \\ X \\ 90 \end{matrix}$  where  $X$  has com-

position factors  $9 + 36 + 90 + 2(126)$ . Section 4.1 and self-duality shows that  $L_1(X) = \begin{matrix} 126 \\ \\ \end{matrix}$

$S_1(X) = 126$ . Hence,  $X = \begin{matrix} 9 & 36 & 90 \\ & & 126 \end{matrix}$ , and  $P_{90A_{10}}$  has Loewy structure as claimed in

Theorem 2.

• Similarly,  $P_{279A_{10}} = \begin{matrix} 279 \\ Y \\ 279 \end{matrix}$  where  $Y$  has composition factors  $2(9) + 2(36) + 126 + 279$ ,

and  $L_1(Y) = S_1(Y) = 9 \oplus 36$ . Therefore, the structure of  $Y$  can only be one of the following:

$$\begin{matrix} S & & 9 & 36 \\ 126 & 279 & T \oplus T & \\ S & & 9 & 36 \end{matrix} \quad \text{or} \quad \begin{matrix} 9 & 36 \\ 126 & 279 \\ 9 & 36 \end{matrix} .$$

Here  $\{S, T\} = \{9, 36\}$ . But the first case cannot happen since  $(S, T)_{A_{10}}^1 = 0$  by Section 4.1. Thus, the structure of  $P_{279A_{10}}$  is as claimed in Theorem 2.

• We will work out in details the structure of  $P_{9A_{10}}$ . The structure of  $P_{36A_{10}}$  can be found in a similar way.

The Cartan matrix shows that  $P_{9A_{10}} = \begin{matrix} 9 \\ Z \\ 9 \end{matrix}$  where  $Z$  has composition factors  $2(9) + 36 + 90 + 2(126) + 2(279)$ . It is easy to see that  $L_1(Z) = S_1(Z) = 126 \oplus 279$ . Consequently, the structure of  $Z$  can only be one of the following:





by the structure of  $189 \uparrow$  and Section 4.2 respectively.

Now  $84_b$  is not in either  $L_5(P_{34_{A_{10}}})$  or  $L_6(P_{34_{A_{10}}})$  by Section 4.2 and the Benson-Carlson diagram of  $189 \uparrow$ , so it has to be in  $L_4(P_{34_{A_{10}}})$ . Moreover, Section 4.2 and  $189 \uparrow$  also imply  $224_b \notin L_5(P_{34_{A_{10}}})$ . Also,  $224_b \notin L_6(P_{34_{A_{10}}})$ : otherwise, there exists a submodule  $U$  of  $\begin{pmatrix} 27 \\ 189 \end{pmatrix} \uparrow_{A_9}^{A_{10}}$  of Loewy length 4 with simple head 34 and simple socle 224. Hence, by duality, there exists a subquotient  $U^*$  of  $\begin{pmatrix} 189 \\ 27 \end{pmatrix} \uparrow_{A_9}^{A_{10}}$  with simple head 224, simple socle 34 and Loewy length 4. This means that, in Figure 2,  $34_d \in L_4(P_{224_{A_{10}}})$ , since Section 4.2 forces  $224_d \in L_i(P_{224_{A_{10}}})$  for  $i \geq 3$  so the diagram of  $189 \uparrow$  implies  $34_g \notin L_4(P_{224_{A_{10}}})$ . This in turn forces  $1_e, 41_e \in L_3(P_{224_{A_{10}}})$ . Hence,  $84_d \in L_4(P_{224_{A_{10}}})$  since  $(224, 84)_{A_{10}}^1 = 0$ . Therefore, we have

$$\text{Loewy length of } \begin{pmatrix} 189 \\ 27 \end{pmatrix} \uparrow_{A_9}^{A_{10}} \cdot e_0 \leq 7 < 8 \leq \text{Loewy length of } \begin{pmatrix} 27 \\ 189 \end{pmatrix} \uparrow_{A_9}^{A_{10}} \cdot e_0,$$

a contradiction. Therefore,  $224_b \in L_4(P_{34_{A_{10}}})$ .

Next, Landrock's Lemma implies that  $34_d \in L_4(P_{224_{A_{10}}})$ . The same argument as above shows that Loewy length of  $\begin{pmatrix} 189 \\ 27 \end{pmatrix} \uparrow_{A_9}^{A_{10}} \cdot e_0$  is at most 7, hence  $1_b \notin L_6(P_{34_{A_{10}}})$ . Moreover,  $1_b \notin L_5(P_{34_{A_{10}}})$  by the Benson-Carlson diagram of  $189 \uparrow \cdot e_0$  and Section 4.2. Therefore,  $1_b \in L_4(P_{34_{A_{10}}})$ . Now (\*) gives

$$\begin{pmatrix} 27 \\ 189 \end{pmatrix} \uparrow_{A_9}^{A_{10}} \cdot e_0 = \begin{array}{ccccccc} & & & & & & 34 \\ & & & & & & 1 & 41 & 224 \\ & & & & & & 34 & 84 & 1 & 41 & 224 \\ & & & & & & 1 & 41 & 1 & 224 & 34 & 84 \\ & & & & & & & & & & & & 34 & 1 & 41 & 224 \\ & & & & & & & & & & & & & & & 224 \end{array}.$$

This has Loewy length 6, so Loewy length of  $\begin{pmatrix} 189 \\ 27 \end{pmatrix} \uparrow_{A_9}^{A_{10}} \cdot e_0 = 6$ . This implies  $34_e \in L_6(P_{224_{A_{10}}})$ . The diagram of  $27 \uparrow$  now gives

$$\begin{pmatrix} 189 \\ 27 \end{pmatrix} \uparrow_{A_9}^{A_{10}} \cdot e_0 = \begin{array}{ccccccc} & & & & & & 224 \\ & & & & & & 1 & 41 & 224 & 34 \\ & & & & & & 1 & 224 & 34 & 84 & 1 & 41 \\ & & & & & & 1 & 41 & 224 & 34 & 84 \\ & & & & & & & & & & & & 224 & 1 & 41 \\ & & & & & & & & & & & & & & & 34 \end{array}.$$

Now, in Figure 1,  $34_a \notin L_2(P_{34_{A_{10}}})$  by Section 4.2. In fact, it is in  $L_3(P_{34_{A_{10}}})$ , otherwise, it must extend (and indeed lie below) some composition factors in the submodule  $\begin{array}{c} 1 & 41 \\ 34 \end{array}$  of  $27 \uparrow$ , which is impossible.

Also, Section 4.2 implies that  $34_f, 84_f \notin L_6(P_{34_{A_{10}}})$ ; and they are not in  $L_7(P_{34_{A_{10}}})$  either, because the diagram of  $27 \uparrow$  and Section 4.2 imply  $1_f, 41_f \notin L_6(P_{34_{A_{10}}})$ . Thus,  $34_f, 84_f \in L_5(P_{34_{A_{10}}})$ . This, in turns, forces  $1_f, 41_f \in L_4(P_{34_{A_{10}}})$ . Therefore, we obtain the structure of  $P_{34_{A_{10}}}$  as claimed in Theorem 1.

Finally we compute the structure of  $P_{224_{A_{10}}}$ . By Landrock's lemma, there is one copy of 34 in  $L_5(P_{224_{A_{10}}})$ , so  $34_g \in L_5(P_{224_{A_{10}}})$ . Thus,  $224_d \in L_3(P_{224_{A_{10}}})$  by the diagram of  $189 \uparrow$ . Also, since 84 does not extend 224,  $84_h \in L_4(P_{224_{A_{10}}})$ .

We now have  $1_h \in L_5(P_{224_{A_{10}}})$  since, otherwise, the structure of  $\begin{pmatrix} 27 \\ 189 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0$  implies that it must extend  $224_f$ . This means there exists a module  $V$  with Loewy length 4 with unique head  $224$  and unique socle  $224$  and whose head  $224$  extends 1 in  $L_2(V)$ . By duality again,  $1_h$  extends  $224_e$ , which implies that  $224_g$  is extended by  $1_h$ . Thus,  $1_h$  extends both  $224_f$  and  $224_g$ , contradicting Section 4.2. A similar argument shows that  $224_h \in L_5(P_{224_{A_{10}}})$  and finally the structure of  $P_{224_{A_{10}}}$  is as claimed in Theorem 1.

6.2.2. *The Loewy structure of  $P_{84}$ .* We have

$$\begin{pmatrix} 7 \\ 1 & 21 \\ 7 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 \sim \frac{\begin{array}{c} 34 \\ 84 \\ 1 \oplus 1 \quad 41 \\ 84 \\ 34 \end{array}}{\quad} = \begin{array}{c} 34 \quad 84 \\ \swarrow \quad \searrow \quad \swarrow \quad \searrow \\ 1 \quad 41 \quad 1 \\ \swarrow \quad \searrow \\ 34 \quad 84 \end{array} .$$

by Frobenius Reciprocity, Section 4 and Section 5.

Similarly,

$$\begin{pmatrix} 41 \\ 35 \\ 41 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \begin{array}{c} 41 \\ 224 \\ 41 \end{array}, \quad \begin{pmatrix} 35 \\ 21 & 35 \\ 35 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \begin{array}{c} 224 \quad 84 \\ 224 \quad 1 \quad 41 \\ 224 \quad 84 \end{array},$$

$$\begin{pmatrix} 1 \\ 7 \\ 1 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \begin{array}{c} 41 \\ 224 \\ 41 \end{array}, \quad \begin{pmatrix} 41 \\ 1 & 7 \\ 41 \end{pmatrix} \uparrow_{A_9}^{A_{10}} .e_0 = \begin{array}{c} 41 \\ 1 \quad 34 \\ 41 \end{array}.$$

Furthermore, we have the following filtration ((3) in (Siegel, 1991)):

$$P_{7_{A_9}} = P_{28_{A_8}} \uparrow_{A_9} .f_0$$

$$\sim \begin{array}{c} 7 \\ 1 \quad 21 \\ 7 \end{array} \left| \begin{array}{c} 41 \\ 35 \oplus 21 \quad 35 \\ 41 \quad 35 \end{array} \right| \begin{array}{c} 7 \\ 1 \quad 21 \oplus 1 \quad 21 \\ 7 \quad 7 \end{array} \left| \begin{array}{c} 7 \\ 1 \quad 21 \oplus 7 \\ 1 \quad 41 \end{array} \right| \begin{array}{c} 41 \\ 1 \quad 7 \\ 41 \end{array} \left| \begin{array}{c} 41 \\ 35 \oplus 21 \quad 35 \\ 41 \quad 35 \end{array} \right| \begin{array}{c} 7 \\ 1 \quad 21 \\ 7 \end{array}$$

Finally, the structure of  $P_{34_{A_{10}}}$  together with the above data gives the following Loewy structure:

$$P_{34_{A_{10}}} \oplus P_{84_{A_{10}}} = P_{7_{A_9}} \uparrow_{A_{10}} .e_0$$

$$= \begin{array}{c} 34 \quad 84 \\ 1 \quad 41 \quad 1 \quad 41 \quad 224 \quad 84 \\ 34 \quad 84 \quad 224 \quad 224 \quad 1 \quad 41 \quad 34 \quad 84 \quad 34 \quad 84 \quad 1 \quad 41 \\ 41 \quad 224 \quad 84 \quad 1 \quad 41 \quad 1 \quad 1 \quad 41 \quad 1 \quad 34 \quad 1 \quad 34 \quad 41 \quad 224 \quad 84 \\ 34 \quad 84 \quad 34 \quad 84 \quad 1 \quad 1 \quad 224 \quad 224 \quad 1 \quad 41 \quad 34 \quad 84 \\ 41 \quad 224 \quad 84 \quad 1 \quad 41 \quad 1 \\ 34 \quad 84 \end{array} .$$

Thus, the structure of  $P_{84A_{10}}$  is as claimed in Theorem 1 and  $(84, 84)_{A_{10}}^1 = 1$  as claimed in Section 4.

6.2.3. *The Loewy structure of  $P_{41}$ .* Landrock's Lemma, Section 6.2.1, Section 6.2.2 and the filtration of  $P_{41A_9} \uparrow^{A_{10}} .e_0$  give the positions of the unlabelled simple modules and  $1_a$  in the Loewy structure for  $P_{41A_{10}}$  as follows:

$$\begin{array}{cccccccc}
 & & & & 41 & & & \\
 & & & & 1_a & 34 & 84 & 224 \\
 & & & 34 & 224 & 84 & 1_A & 41_A & 41_D & 41_E & 41_F \\
 34 & 34 & 224 & 224 & 84 & 84 & 1_B & 41_B & 1_D & 1_H \\
 & & 34 & 224 & 84 & 1_C & 41_C & 41_G & 1_G & 41_H \\
 & & & & 34 & 224 & 84 & 1_E \\
 & & & & 1_F & 41_E \\
 & & & & 41_F
 \end{array}$$

Since we have accounted for all the 84s by Landrock's Lemma and the 84s all come from  $\mathfrak{D}(84; 1; 41; 84)$ ,  $1_A, 41_A, 1_B, 41_B, 1_C, 41_C$  are in the respective Loewy layers as above.

Next, from (Siegel, 1991), we see that

$$\begin{array}{c}
 41 \\
 \left( \begin{array}{c} 13 \\ 1 \end{array} \right) \uparrow_{A_8}^{A_9} .f_0 = \begin{array}{cc} 35 & 1 \\ 41 & 7 \end{array} \\
 1
 \end{array}$$

Thus,  $M := \left( \left( \begin{array}{c} 13 \\ 1 \end{array} \right) \uparrow_{A_8}^{A_9} .f_0 \right) \uparrow^{A_{10}} = \begin{array}{cc} 41 & \\ 224 & 1 \\ 41_a & 34_a \\ & 1_B \end{array}$ , since Frobenius Reciprocity and Lemma 3

implies  $S_1(M) = 1$  and  $41_a, 34_a \notin L_1(M), L_2(M)$ . Therefore, we get a copy of 41 and a copy of 1 in  $L_3(P_{41A_{10}})$  that do not come from  $21 \uparrow_{A_9}^{A_{10}} = \mathfrak{D}(84; 1; 41; 84)$ , say  $41_D$  and  $1_D$  respectively.

Next, Frobenius Reciprocity gives

$$\begin{array}{c}
 41 \\
 \left( \begin{array}{cc} 41 & \\ 1 & 7 \\ & 41 \end{array} \right) \uparrow_{A_9}^{A_{10}} .e_0 = \begin{array}{cc} 41 & \\ 1 & 34 \\ & 41 \end{array}
 \end{array}$$

so  $1_a$  extends a  $41_E$  in  $L_3(P_{41A_{10}})$ , and  $41_E \neq 41_A, 41_D$  since  $13 \uparrow_{A_8}^{A_9} = \begin{array}{c} 41 \\ 35 \\ 41_1 \end{array}$  and there

exists an  $FA_9$ -module  $\begin{array}{c} 41 \\ 1 \quad 7 \\ 41_2 \end{array}$ . Hence,  $41_1 \neq 41_2$  because there is only one copy of 35 in  $L_2(P_{41A_9})$ . Thus,  $41_E$  comes from  $41_2 \uparrow_{A_9}^{A_{10}}$ ,  $41_D$  comes from  $41_1 \uparrow_{A_9}^{A_{10}}$  and  $41_A$  comes from  $\mathfrak{D}(84; 1; 41; 84)$ .

Note that all the factors in  $L_2(P_{41A_{10}})$  extend some factors in  $L_3(P_{41A_{10}})$  so self-duality implies that the layer immediately above the unique bottom 41 in  $P_{41A_{10}}$  is

$$\begin{array}{cccc}
 1 & 34 & 84 & 224
 \end{array}$$

Therefore, the Loewy length of  $P_{41A_{10}}$  is 7 and there is exactly one copy of 1 in  $L_6(P_{41A_{10}})$ , say  $1_E$ . We can also fill in the bottom 41, say  $41_F$ .

Now  $41_D$  extends  $1_D$  so by self-duality and the submodule lattice of  $P_{41A_{10}}$ , we must have another copy of 41 in  $L_5(P_{41A_{10}})$ , say  $41_G$ .





positions of the four  $1^*$ . By (\*) in Section 6.2.3, at least one of the  $1^*$  must appear in  $L_4(P_{1_{A_{10}}})$ . Using the filtration (2) in Siegel (1991) below, and that  $P_{1_{A_{10}}} = P_{1_{A_9}} \uparrow^{A_{10}} .e_0$ , the remaining three  $1^*$  must be in  $L_5(P_{1_{A_{10}}})$  and the structure of  $P_{1_{A_{10}}}$  is as claimed in Theorem 1.

$$P_{1_{A_9}} = P_{1_{A_8}} \uparrow^{A_9} .f_0$$

$$\sim \begin{array}{c} 1 \\ 7 \\ 1 \end{array} \left| \begin{array}{cc} 41 & 35 \\ 35 \oplus 21 & 35 \\ 41 & 35 \end{array} \right| \begin{array}{cccc} 1 & 1 & 7 & 41 \\ 7 \oplus 7 \oplus 1 & 21 \oplus 1 & 7 & 41 \\ 1 & 1 & 7 & 41 \end{array} \left| \begin{array}{cc} 41 & 35 \\ 35 \oplus 21 & 35 \\ 41 & 35 \end{array} \right| \begin{array}{c} 1 \\ 7 \\ 1 \end{array}$$

## 7. ACKNOWLEDGEMENTS

We are grateful to Richard Parker for explaining the use of the MeatAxe and thank Juergen Manfred Muller for pointing out the inaccuracies in a previous version of this paper.



$\dim_F \text{Ext}_{A_{10}}^1(S, T)$  for  $S, T$  simple

	1	34	41	84	224	9	36	90	126	279	567
1	.	1	1	1	1	.	.	.	.	.	.
34	1	.	1	.	1	.	.	.	.	.	.
41	1	1	.	1	1	.	.	.	.	.	.
84	1	.	1	1	.	.	.	.	.	.	.
224	1	1	1	.	1	.	.	.	.	.	.
9	.	.	.	.	.	.	.	.	1	1	.
36	.	.	.	.	.	.	.	.	1	1	.
90	.	.	.	.	.	.	.	.	1	.	.
126	.	.	.	.	.	1	1	1	.	.	.
279	.	.	.	.	.	1	1	.	.	.	.
567	.	.	.	.	.	.	.	.	.	.	.

#### REFERENCES

- (2013). *GAP – Groups, Algorithms, and Programming, Version 4.7.2.*. The GAP Group.
- Benson, D. (1983a). The Loewy structure of the projective indecomposable modules for  $A_8$  in characteristic 2. *Comm. Algebra*, 11:1395–1432.
- Benson, D. (1983b). The Loewy structure of the projective indecomposable modules for  $A_9$  in characteristic 2. *Comm. Algebra*, 11:1433–1453.
- Benson, D. (1984). *Modular Representation Theory: New Trends and Methods*. Lecture Notes in Mathematics. Springer-Verlag.
- Benson, D. and Carlson, J. (1987). Diagram methods for modular representations and cohomology. *Comm. Algebra*, 15:53–121.
- James, G. and Kerber, A. (1981). *The Representation Theory of the Symmetric Group*, volume 16 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley Publishing Company.
- Landrock, P. (1983). *Finite group algebras and their modules*, volume 84 of *London Mathematical Society Lecture Notes*. Cambridge University Press.
- Scopes, J. (1988). The Loewy structure of the projective indecomposable modules of  $A_6, A_7$  and  $A_8$  in characteristic three. *Dissertation (Oxford University)*.
- Siegel, S. (1991). Projective modules for  $A_9$  in characteristic three. *Comm. Algebra*, 19:3099–3117.