

# CATEGORIES OF ORBIT TYPES FOR PROPER LIE GROUPOIDS

JACK MORAVA

ABSTRACT. It is widely understood that the quotient space of a topological group action can have a complicated combinatorial structure, indexed somehow by the isotropy groups of the action [3 II §2.8]; but how best to record this structure seems unclear. This sketch defines a database category of orbit types for a proper Lie groupoid (based on recent work [13-15] with roots in the theory of geometric quantization) as an attempt to capture some of this information.

## 1. INTRODUCTION AND BACKGROUND

A topological groupoid or stack [12]

$$\mathbb{X} := s, t : X_1 \rightrightarrows X_0$$

is **proper** if the map  $s \times t : X_1 \rightarrow X_0 \times X_0$  is proper; such an object in the category of smooth manifolds and maps is a **proper Lie groupoid**. The quotient

$$\mathbb{X} \rightarrow \mathfrak{X}$$

of  $X_0$  by the equivalence relation thus defined is a Hausdorff topological space, sometimes called the coarse moduli space of  $\mathbb{X}$ .

### Examples:

- Orbifolds [5]
- A topological **transformation group**, defined by a group action

$$G \times X \rightarrow X$$

has an associated topological groupoid  $[X/G]$  with  $X_0 = X$ ,  $X_1 = G \times X$ ; I'll write  $X/G$  for its quotient space. For instance

- **Toric varieties**, eg  $G = \mathbb{T}^{n+1}/\mathbb{T} \cong \mathbb{T}^n$  acting on  $X = \mathbb{C}P^n$  by

$$(u_0, \dots, u_n) \cdot [z_0 : \dots : z_n] = [u_0 z_0 : \dots : u_n z_n],$$

form a particularly accessible class of examples. Their quotient objects are polytopes: in the case above  $X/G \cong \Delta^n$  is a simplex. The faces of the polytope define a stratification [see §4.1 below] of the quotient, with the

---

*Date:* 14 February 2014.

1991 *Mathematics Subject Classification.* 03B65, 14K10, 22A22.

interiors of the faces as strata. This defines an interesting poset, or category, associated to the groupoid: in this example it is the category of subsets of  $\{0, \dots, n\}$  under inclusion.

An earlier paper [11] attempted to capture the sort of information encoded by the face poset of a toric variety, for more general group actions. The present note uses recent work on proper Lie groupoids [[16], cf also [1]] to propose a more general construction<sup>1</sup>.

**Acknowledgement** I am indebted to the organizers of the September 2013 Barcelona conference on homotopy type theory [2] for inspiration and hospitality, and for the opportunity to pursue these questions. I hope I will not be misunderstood by suggesting that classification problems of the sort considered here have a deep and nontrivial history in philosophy [17].

## 2. SOME TECHNICAL PRELIMINARIES

**2.1 Definition** A reasonable space  $X$  has a universal map  $X \rightarrow \pi_0 X$  to a discrete set, defined by the adjoint to the inclusion of the category of sets into that of topological spaces. The diagram

$$\begin{array}{ccccccc} \mathbf{X} : & & X_1 & \xrightarrow{\quad \cong \quad} & X_0 & \cdots \cdots \cdots & \mathfrak{X} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \pi_0 \mathbf{X} : & & \pi_0 X_1 & \xrightarrow{\quad \cong \quad} & \pi_0 X_0 & \cdots \cdots \cdots & \pi_0 \mathfrak{X} \end{array}$$

extends  $\pi_0$  to a functor from topological to discrete groupoids, such that

$$\pi_0[X/G] \cong [\pi_0(X)/\pi_0(G)] ,$$

with  $\pi_0(X)/\pi_0(G) \cong \pi_0(X/G)$  for reasonable actions. The natural transformation

$$\pi_0(\mathbf{X} \times \mathbf{Y}) \rightarrow \pi_0 \mathbf{X} \times \pi_0 \mathbf{Y}$$

is an isomorphism in such cases.

**2.2** Regarding groups as categories with a single object defines a two-category ( $\mathbf{Gps}$ ) of groups. The set of homomorphisms from  $G_0$  to  $G_1$  has an action of  $G_1$  by conjugation, defining a groupoid

$$\mathrm{Hom}_{\mathbf{Gps}}(G_0, G_1) := [\mathrm{Hom}(G_0, G_1)/G_1^{\mathrm{conj}}]$$

of morphisms from  $G_0$  to  $G_1$ .

There are many variations on this theme, eg the topological two-category ( $\mathbf{Gps}_c$ ) defined by compact groups and continuous homomorphisms. I will

<sup>1</sup>See [8, 21] for approaches based on  $\pi_1$  rather than  $\pi_0$

write  $(\mathbf{Gps}^+)$  (resp.  $(\mathbf{Gps}_c^+)$ ) for the subcategories with such groups as objects, and spaces  $\mathrm{Hom}_c^+(G_0, G_1)$  of continuous **one-to-one** homomorphisms as maps.

The construction which is the identity on objects, and is the functor

$$[\mathrm{Hom}_c(G_0, G_1)/G_1^{\mathrm{conj}}] \rightarrow \mathrm{Hom}_{\pi_0 \mathbf{Gps}}(G_0, G_1) := \pi_0[\mathrm{Hom}_c(G_0, G_1)/G_1^{\mathrm{conj}}]$$

on morphism categories, defines a monoidal two-functor

$$(\mathbf{Gps}_c) \rightarrow (\pi_0 \mathbf{Gps}_c)$$

[and similarly for  $(\mathbf{Gps}_c^+)$ ].

### 3. GROUPOIDS OF FIXED-POINTS WITH LEVEL STRUCTURE

**3.1 Definition:** If  $X$  is a proper topological groupoid, and  $H$  is a compact Lie group, let

$$X(H)_0 := \{(x, \phi) \mid x \in X_0, \phi : H \rightarrow \mathrm{Iso}(x) \in \mathbf{Gps}_c^+\}$$

and let  $X(H)_1$  be the set of commutative diagrams of the form

$$\begin{array}{ccc} H & \xrightarrow{\phi'} & \mathrm{Iso}(x') \\ \downarrow \gamma & & \downarrow g\text{-conj} \\ H & \xrightarrow{\phi} & \mathrm{Iso}(x) \end{array}$$

(with  $g : x' \rightarrow x \in X_1$ ). The resulting proper topological groupoid  $X(H)$  is a model for the subgroupoid of  $X$  defined by points fixed by a group isomorphic to  $H$ . There is a forgetful morphism  $X(H) \rightarrow X$ , but it can't be expected to be the inclusion of a subgroupoid.

**Proposition:**  $H \mapsto X(H)$  defines a (two-)functor  $X(\bullet)$  from  $(\mathbf{Gps}_c^+)$  to the two-category  $(\mathbf{Gpoids}_c)$  of proper topological groupoids.

**Proof:** First of all, if  $\alpha : H_0 \longrightarrow H_1 \in (\mathbf{Gps}_c^+)$  then

$$\begin{array}{ccc} H_0 & \xrightarrow{\alpha} & H_1 \xrightarrow{\phi'_1} \mathrm{Iso}(x') \\ \downarrow \gamma^\alpha & & \downarrow g\text{-conj} \\ H_0 & \xrightarrow{\quad} & H_1 \xrightarrow{\phi_1} \mathrm{Iso}(x) \end{array}$$

defines a functor

$$\alpha_{H_1}^{H_0} : X(H_1) \rightarrow X(H_0) .$$

Moreover, if  $\alpha : H \rightarrow H$  is an inner automorphism of  $H$  (ie  $\alpha$  is conjugation by  $a \in H$ ) then there is a natural equivalence

$$\alpha_H^H \cong \mathbf{1}_{X(H)}$$

defined by the commutative diagram

$$\begin{array}{ccccc} H & \xrightarrow{\alpha} & H & \xrightarrow{\phi} & \text{Iso}(x) \\ \vdots & & & & \downarrow \phi(a^{-1}) \\ H & \xrightarrow{1_H} & H & \xrightarrow{\phi} & \text{Iso}(x) . \end{array}$$

□

**3.2 Claim:** For any  $X$  as above, there is a commutative diagram

$$\begin{array}{ccc} X & \longrightarrow & (\text{Gpoids}_{c*}) \\ \downarrow \text{Iso} & & \downarrow \\ (\text{Gps}_c) & \xrightarrow{X[\bullet]} & (\text{Gpoids}_c) \end{array}$$

(with the category of pointed proper groupoids in the upper right corner, and the forgetful map to the category of proper groupoids along the right-hand edge). The left-hand vertical map sends  $x \in X_0$  to its isotropy group, and the top horizontal map sends  $x$  to  $X(\text{Iso}(x))$ , with  $x$  as distinguished point.

**Corollary** The universal property of a fiber product defines a continuous functor

$$X \rightarrow \Phi_0(X)$$

to the category defined by the pullback

$$\begin{array}{ccccc} \Phi_0(X) & \dashrightarrow & (\text{Gpoids}_{c*}) & \xrightarrow{\pi_0} & (\text{Gpoids})_* \\ \vdots & & \downarrow & & \downarrow \\ (\text{Gps}_c) & \xrightarrow{X[\bullet]} & (\text{Gpoids}_c) & \xrightarrow{\pi_0} & (\text{Gpoids}) \end{array}$$

(where the two right vertical arrows are the obvious forgetful functors). □

It's natural to think of  $\Phi_0(X)$  as a **database** category [11, 18]. However I don't know how to characterize  $\Phi_0$  by some universal property (such as being an adjoint).

**3.3 Example:** A proper transformation group  $[X/G]$  defines a functor

$$G > H \mapsto X^H = \{x \in X \mid \text{Iso}(x) \subset H\}$$

from the topological category  $(G - \text{Orb})$  [with closed subgroups of  $G$  as objects, and

$\text{Mor}_{G-\text{Orb}}(H_0, H_1) = \text{Maps}_G(G/H_0, G/H_1) = \{g \in G \mid gH_0g^{-1} \subset H_1\}/H_1^{\text{conj}}$  as morphism objects [3 I §10], to spaces.

This extends to a functor

$$S^0[X^\bullet] : (G - \text{Spaces}) \ni X \mapsto S^0[X^H] \in \text{Func}(G - \text{Orb}, S^0 - \text{Mod})$$

which provides a model [6 V §9, 10, 20] for the  $G$ -equivariant stable category in terms of sheaves of spectra (ie  $S^0$ -modules) over  $(G - \text{Orb})$ .

The commutative diagram

$$\begin{array}{ccc} [X/G] & \dashrightarrow & (\text{Sets})_* \\ \downarrow \text{Iso} & & \downarrow \\ (G - \text{Orb}) & \xrightarrow{\pi_0 X[*]} & (\text{Sets}) \end{array}$$

defines a functor from  $[X/G]$  to a fiber product category  $\Phi_0[X/G]$  (with objects, pairs consisting of a subgroup  $H$  of  $G$ , and a component of  $X^H$ ), analogous to the construction in the previous paragraph [11 §2.2]. The sheaf  $S^\infty X^\bullet$  of spectra pulls back to a sheaf of spectra over  $\Phi_0(X)$ .

One might hope for an unstable version of this construction, applicable in the theory of  $\infty$ -categories (cf eg [7 §5.5.6.18]); but because it depends on a presentation of  $[X/G]$  as a global quotient, it does not seem to be homotopy-invariant.

#### 4. PROPER **Lie** GROUPOIDS, AFTER PFLAUM *et al*

**4.1 A stratification  $\mathcal{S}$**  of a (paracompact, second countable) topological space  $X$  assigns to each  $x \in X$ , the germ of a closed subset  $\mathcal{S}_x$  (containing  $x$ ) of  $X$ . With suitably defined morphisms [9 §1.8, 16 §1], stratified spaces form a category. A stratification defines a locally finite partition

$$X = \coprod_{S \in \Sigma(\mathcal{S})} X_S$$

of  $X$  into locally closed subsets (called its strata), such that if  $x \in X_S$  then  $\mathcal{S}_x$  is the associated set germ.

Very interesting recent work of M. Pflaum *et al* [building on earlier work of Weinstein and Zung ([22]; cf also [13]) shows that

**Theorem** [16 Theorem 5.3, Cor 5.4] The quotient space  $\mathfrak{X}$  of a proper Lie groupoid  $\mathbf{X}$  has a canonical **Whitney** stratification. The associated decomposition of  $X_0$  into locally closed submanifolds

$$X_{0(H)} = \{x \in X_0 \mid \text{Iso}(x) \cong H\}$$

is indexed [16 Theorem 5.7] by (isomorphism classes of) compact Lie groups  $H$ .

**4.2 Definition** The **orbit** groupoid  $\mathbf{O}(x) \subset \mathbf{X}$  of  $x \in X_0$

$$\mathbf{O}_0(x) = \{y \in X_0 \mid \exists g : y \rightarrow x \in X_1\}n$$

$$\mathbf{O}_1(x) = \{g \in X_1 \mid s(g), t(g) \in \mathbf{O}_0(x)\}$$

reduces, in the case of a transformation groupoid  $[X/G]$ , to the groupoid

$$[(G/\text{Iso}(x))/G] \equiv [*/\text{Iso}(x)] .$$

A **slice** at  $x \in X_0$  is (very roughly [13 §3.3-4,3.8-9]) the germ of an  $\text{Iso}(x)$ -invariant submanifold of  $X_0$  containing  $x$ , transverse to  $\mathbf{O}_0(x)$ ; for a transformation group it is something like the image of an exponential map

$$[\mathcal{N}_x/\text{Iso}(x)] \equiv [(\mathcal{N}_x \times_{\text{Iso}(x)} G)/G] \rightarrow [X/G]$$

(where  $\mathcal{N}_x \in (\text{Iso}(x) - \text{Mod})$  is the linear representation

$$0 \rightarrow T_x G \rightarrow T_x X_0 \rightarrow \mathcal{N}_x \rightarrow 0 .$$

defining the normal bundle to the orbit of  $x$ ).

**Theorem** [16 §3.11] There is an (essentially unique) slice at every object of a proper Lie groupoid  $\mathbf{X}$ ; the corresponding set germs define the canonical stratification [16 §5.4] of  $\mathbf{X}$ .

**Definition** The **normal orbit type** of  $x \in X_0$  is the equivalence class of its normal  $\text{Iso}(x)$ -representation  $\mathcal{N}_x$ . More precisely,  $x_0 \sim x_1$  if there are isomorphisms

$$\phi : \text{Iso}(x_0) \rightarrow \text{Iso}(x_1), \quad \Phi : \mathcal{N}_{x_0} \rightarrow \phi^*(\mathcal{N}_{x_1})$$

of groups and representations. The connected components  $\nu \in \pi_0(X_0)$  of the normal orbit types of  $\mathbf{X}$  are [16 §5.7] the strata of the canonical partition of  $X_0$ .

The **condition of the frontier** [16 Prop 5.15] asserts that if  $\nu' \cap \bar{\nu} \neq \emptyset$  then  $\bar{\nu} \supset \nu'$ . This implies the existence of a partial order ( $\nu > \nu'$ ) on the set  $\Sigma(\mathbf{X})$  of connected components of normal orbit types for  $\mathbf{X}$ , which can thus be regarded as the objects of a category [4 II §2.8]. Thus

$$\overline{X_{0(K)}} = \coprod_{\pi_0(X_{0(K)}) \ni \nu > \nu'} \nu' ,$$

This gives us some control of the functor  $\Phi_0$  on proper Lie groupoids:

**4.3 Proposition** For a proper Lie groupoid  $\mathbf{X}$ , we have isomorphisms

$$\bigcup_{H < K} X_{0(K)} \times \text{Hom}_c^+(H, K) \xrightarrow{\cong} X(H)_0$$

$$\bigcup_{H < K, x \in X_{0(K)}} \mathbf{O}_1(x) \times K \xrightarrow{\cong} X(H)_1$$

and consequently

$$\bigcup_{H < K} \mathfrak{X}_K \times \text{Hom}_c^+(H, K)/K^{\text{conj}} \xrightarrow{\cong} \mathfrak{X}(H)$$

(where  $\mathfrak{X}_K \subset \mathfrak{X}$  is the space of orbits with isotropy group isomorphic to  $K$ ).

#### 4.4 Closing remarks

i) When  $\mathsf{X} = [X/G]$  this all simplifies a little. In particular, since

$$X^H = \bigcup_{H < K < G} X_K,$$

$\Phi_0[X/G]$  is essentially just the quotient of  $\Phi_0[X/G]$  which collapses the morphism spaces  $\text{Hom}_c^+(K, H)/K^{\text{conj}}$ .

ii) The subspaces  $X_{0(K)}$  are disjoint unions of strata  $\nu$  indexed by slice representations

$$K \rightarrow \text{Aut}(\mathcal{N}_\nu).$$

The resulting family of vector spaces over  $\mathsf{X}(H)$  pulls back to a fibered category

$$\mathcal{N}(\mathsf{X}) \rightarrow \Phi_0(\mathsf{X}).$$

This seems to provide a natural repository for Noether's theorem (which associates conserved quantities to elements of the Lie algebra of symmetries of states of a physical system) [11 §4.1].

iii) I don't know how generally one can associate a stratification to a topological groupoid. There are many interesting examples, coming from locally compact groupoids (eg the Thom-Boardman theory of singularities of smooth maps [18]), or from infinite-dimensional examples (Ebin's category of Riemannian metrics up to diffeomorphism, Vassiliev's finite-type invariants of immersions, ...), where a more general theory would be very interesting. The existence and good behavior of slices seem to be an essential requirement for such a theory.

## REFERENCES

1. P Cartier, Groupoides de Lie et leurs algébroïdes, Sem. Bourbaki 987 (2007-8), Astérisque 326 (2009) 165 - 196
2. A Collective, **Homotopy type theory . . .**, <http://homotopytypetheory.org/book/>
3. T tom Dieck, **Transformation groups**, de Gruyter Studies 8 (1987)
4. JJ Duistermaat, JA Kolk, **Lie groups**, Universitext. Springer (2000)
5. D Gepner, A Henriques, Homotopy theory of orbispaces, [arXiv:math/07019](https://arxiv.org/abs/math/07019)
6. LG Lewis, JP May, M Steinberger, JE McClure, **Equivariant stable homotopy theory**, Springer LNM 1213 (1986)
7. J Lurie, **Higher topos theory**, Annals of Math Studies 170 (2009)
8. W Lück, **Transformation groups and algebraic K-theory**, Springer LNM 1408 (1989)
9. J Mather, Stratifications and mappings, in **Dynamical systems** 195 - 232, ed MM Peixoto, Academic Press (1973)
10. H Miller, The Burnside bicategory of groupoids, [arXiv:1208.2360](https://arxiv.org/abs/1208.2360)
11. J Morava, Theories of anything, [arXiv:1202.0684](https://arxiv.org/abs/1202.0684)
12. B Noohi, Homotopy types of topological stacks, [arXiv:0808.3799](https://arxiv.org/abs/0808.3799)
13. R Palais, On the existence of slices for actions of non- compact Lie groups, Ann. Math. 73 (1961) 295 - 323
14. M Pflaum, **Analytic and geometric study of stratified spaces**, Springer LNM 1768 (2001)
15. —, Smooth structures on stratified spaces, in **Quantization of singular symplectic quotients** 231 - 258, Prog Math (Birkhäuser) 2001
16. —, H Posthuma, X. Tang, Geometry of orbit spaces of proper Lie groupoids, [arXiv:1101.0180](https://arxiv.org/abs/1101.0180)
17. G Scholem, **On the Kabbalah and its symbolism**, Schocken (1996), re <http://en.wikipedia.org/wiki/Sephirot>
18. DI Spivak, R Wisnesky, On The relational foundations Of functorial data migration, [arXiv:1212.5303](https://arxiv.org/abs/1212.5303)
19. R Thom, Singularities of differentiable mapping (notes by H Levine), in **Proceedings of Liverpool Singularities I** 1-89, Springer LNM 1920
20. S Schwede, **Global homotopy theory**, [www.math.uni-bonn.de/~schwede/global.pdf](http://www.math.uni-bonn.de/~schwede/global.pdf)
21. J Woolf, The fundamental category of a stratified space, [arXiv:0811.2580](https://arxiv.org/abs/0811.2580)
22. NT Zung, Proper groupoids and momentum maps . . . , Ann Sci École Norm Sup 39 (2006) 841 - 869

THE JOHNS HOPKINS UNIVERSITY, BALTIMORE, MARYLAND 21218

*E-mail address:* [jack@math.jhu.edu](mailto:jack@math.jhu.edu)