

A global approach to AECs

Hugo L. Mariano ^{*†} Pedro H. Zambrano [‡] Andrés Villaveces

December 2013

Abstract

In this work we present some general categorical ideas on Abstract Elementary Classes (AECs), inspired by the totality of AECs of the form $(Mod(T), \preceq)$, for a first-order theory T : (i) we define a natural notion of (functorial) morphism between AECs; (ii) explore the following constructions of AECs: "generalized" theories, pullbacks of AECs, (Galois) types as AECs; (iii) apply categorical and topological ideas to encode model-theoretic notions on spaces of types; (iv) present the "local" axiom for AECs here called "local Robinson's property" and an application (Robinson's diagram method); (v) introduce the category AEC of Grothendieck's gluings of all AECs (with change of basis); (vi) introduce the "global" axioms of "transversal Robinson's property" (TRP) and "global Robinson's property" (GRP) and prove that TRP is equivalent to GRP and GRP entails a natural version of Craig interpolation property.

1 Introduction

The notion of Abstract Elementary Class (AEC) was introduced by Shelah in 1987 ([She]) and is an abstract model-theoretic setting (i.e., there is no syntax and proof theory associated to it) that generalizes the natural occurrences:

- (i) $T \subseteq L_{\omega, \omega} - sent \rightsquigarrow (Mod(T), \preceq)$; (ii) $\theta \in L_{\omega_1, \omega} - sent \rightsquigarrow (Mod(\theta), \preceq)$

Definition 1.1. An AEC is a pair $(\mathcal{K}, \preceq_{\mathcal{K}})$ where $\mathcal{K} \subseteq L - str$, for some (finitary) language L and $\preceq_{\mathcal{K}}$ is a binary relation on \mathcal{K} that satisfies:

- (i) order:
 - $\preceq_{\mathcal{K}}$ is a partial order relation on \mathcal{K} ; • $M \preceq_{\mathcal{K}} N \Rightarrow M \subseteq_{L-str} N$.
- (ii) isomorphism:
 - $M \in \mathcal{K}$ and $M' \cong_{L-str} M$, then $M' \in \mathcal{K}$;
 - $M \preceq_{\mathcal{K}} N$, $f \in L - iso(N, N')$ and $f \upharpoonright M \in L - iso(M, M') \Rightarrow M' \preceq_{\mathcal{K}} N'$
- (iii) coherence:
 - $M, N, P \in \mathcal{K}$, $M \subseteq_{L-str} N \subseteq_{L-str} P$ and $M \preceq_{\mathcal{K}} P$, $N \preceq_{\mathcal{K}} P \Rightarrow M \preceq_{\mathcal{K}} N$
- (iv) reunion: Let $\{M_{\alpha} : \alpha \in \theta\} \subseteq \mathcal{K}$, $M_{\alpha} \preceq_{\mathcal{K}} M_{\beta}$ if $\alpha < \beta < \theta$, then:
 - $M := \bigcup_{\alpha \in \theta} M_{\alpha} \in \mathcal{K}$ and $M_{\alpha} \preceq_{\mathcal{K}} M$, $\forall \alpha \in \theta$; • if $N \in \mathcal{K}$ and $M_{\alpha} \preceq_{\mathcal{K}} N$, $\forall \alpha \in \theta$, then $M \preceq_{\mathcal{K}} N$.
- (v) Löwenheim-Skolem: There is an infinite cardinal κ such that for each $N \in \mathcal{K}$ and $X \subseteq |N|$, there is $M \in \mathcal{K}$, $M \preceq_{\mathcal{K}} N$, $X \subseteq |M|$ and $\#|M| \leq \#X + \kappa$. $LS(\mathcal{K}) := \min\{\kappa : \kappa \text{ satisfies the property}\}$ □

1.2. Let $(\mathcal{K}, \preceq_{\mathcal{K}})$ be an AEC with $\mathcal{K} \subseteq L - str$, then there are two natural notions:

- \mathcal{K} -substructure: $C \preceq_{\mathcal{K}} D \Rightarrow C \xrightarrow{\preceq_{\mathcal{K}}} D$ • \mathcal{K} -embedding: $(A \xrightarrow{f} B) = (A \xrightarrow{\cong} A' \xrightarrow{\preceq_{\mathcal{K}}} B)$ □

1.3. An AEC as a category: Let $cat(\mathcal{K})$ be the category with \mathcal{K} as the class of objects and \mathcal{K} -embeddings as arrows.

- This is well-defined:

identities: because $\preceq_{\mathcal{K}}$ is reflexive

composition: by the closure under isos

*Research supported by FAPESP, under the Thematic Project LOGCONS: Logical consequence, reasoning and computation (number 2010/51038-0).

[†]Instituto de Matemática e Estatística, University of São Paulo, Brazil, email: hugomar@ime.usp.br

[‡]Departamento de Matemáticas, Facultad de Ciencias, National University of Colombia, Colombia, emails: phzambranor@unal.edu.co, avillavecesn@unal.edu.co

$$\begin{array}{c} A \cong A' \overset{\sim}{\hookrightarrow} B \cong B' \overset{\sim}{\hookrightarrow} C \\ A' \cong A'' \overset{\sim}{\hookrightarrow} B' \end{array}$$

- $cat(\mathcal{K})$ has all (upward) directed colimits.
- $cat(\mathcal{K})$ is an accessible category: [AR], [Lie1] [Lie2], [BR].
- We will denote $cat(\mathcal{K})_{strict}$ the category with \mathcal{K} as the class of objects and \preceq_K -inclusions as arrows. \square

1.4. Shelah's Presentation Theorem ([She], [Bal2]): Is the first occurrence of "change of languages" in AECs-theory:
 $\mathcal{K} \subseteq L - str \Rightarrow \exists L' \supseteq L \exists T' \subseteq L' - sent \exists$ a set Γ' of (finitary) L' -types such that, if $EC(T', \Gamma') = \{M' \in L' - str : M' \models T' \text{ and } M' \text{ omits } \Gamma'\}$, then:

- $\mathcal{K} = \{M' \upharpoonright_L \in L - str : M' \in EC(T', \Gamma')\}$ (i.e. $\mathcal{K} = PC_L(T', \Gamma')$);
- $M' \preceq_{L'_{\omega\omega}} N' \in EC(T', \Gamma') \Rightarrow M' \upharpoonright_L \preceq_K N' \upharpoonright_L \in \mathcal{K}$ \square

1.5. Main Question: Shelah's categoricity conjecture ([Bal1]): There is a cardinal $\mu(\kappa)$ such that for all AEC \mathcal{K} with $LS(\mathcal{K}) \leq \kappa$, if \mathcal{K} is categorical in some cardinal greater than $\mu(\kappa)$ then \mathcal{K} is categorical in all $\lambda \geq \mu(\kappa)$. \square

1.6. AECs and infinitary languages: $LS(\mathcal{K}) \leq \kappa \Rightarrow \mathcal{K}$ is closed under \equiv in L_{∞, κ^+} ([Kue]) \square

1.7. (Generalized) types in AECs: The notion of type in AECs-theory ([Lie1], [Lie2]) is given by another application of "change of languages". We consider here an expansion of this notion, called "generalized type" or simply g-type (see, for instance, section I.4 in [Van] for motivations).

g-types: $M \in \mathcal{K}$, I set, a (generalized) I -type over M :

is an equivalence class (N, f, \bar{a}) where $N \in \mathcal{K}$, $f : M \rightarrow N \in cat(\mathcal{K})$, $\bar{a} : I \rightarrow N$;

$(N, f, \bar{a}) \sim (N', f', \bar{a}')$ iff $\exists n \in \mathbb{N}$, $P_0, \dots, P_n \in \mathcal{K}$ and \exists zigzag

$$(N, f, \bar{a}) = (P_0, f_0, \bar{a}_0) \xrightarrow{h_1} (P_1, f_1, \bar{a}_1) \xleftarrow{h_2} (P_2, f_2, \bar{a}_2) \xrightarrow{h_3} \dots \xrightarrow{h_{n-1}} (P_{n-1}, f_{n-1}, \bar{a}_{n-1}) \xrightarrow{h_n} (P_n, f_n, \bar{a}_n) = (N', f', \bar{a}')$$

- LS-axiom $\Rightarrow \exists$ a set of representatives for I -types over M ;
- if $cat(\mathcal{K})$ enjoys (AP), then $(N, f, \bar{a}) \sim (N', f', \bar{a}')$ iff $\exists P \in \mathcal{K}$ and morfisms h_1, h_2 such that:

$$(N, f, \bar{a}) \xrightarrow{h_1} (P, g, \bar{b}) \xleftarrow{h_2} (N', f', \bar{a}')$$

- if $cat(\mathcal{K})$ enjoys (AP), (JEP), has arbitrary large models, then:

– \exists monster models;

– I -types over $M \leftrightarrow$ orbits of I -sequences over a \mathcal{K} -monster model over $M \cup I$ (Galois g-types).

notation: $g - T(M, I)$

- *topology on g-types:* this is analogous to the topology on usual types considered in [Lie1]

$\lambda \geq LS(\mathcal{K})$, $M \in \mathcal{K}$:

$$\mathcal{K}^\lambda(M) := \{N \in \mathcal{K} : N \preceq_K M, \#N \leq \lambda\},$$

basis of $X^\lambda(M, I)$, the λ -topology of $g - T(M, I)$: $\{U_{p,N} : p \in g - T(N, I), N \in \mathcal{K}^\lambda(M)\}$, where:

$$U_{p,N} := \{q \in g - T(M, I) : q \upharpoonright = p\}$$

(remark: if $\#M \leq \lambda$, then $X^\lambda(M, I)$ is discrete) \square

2 The category of all AECs

2.1. Motivation: change of languages in model theory of FOL (**Institution theory**)

$\alpha : L \rightarrow L'$ (morphism of languages) $\rightsquigarrow (\hat{\alpha}, \alpha^*)$, where:

- $\hat{\alpha} : Form(L) \rightarrow Form(L')$, a function defined by recursion on complexity

- $\alpha^* : L' - str \rightarrow L - str$, a functor

$$|M'^\alpha| = |M'|, R^{M'^\alpha} = \alpha(R)^{M'}, f^{M'^\alpha} = \alpha(f)^{M'}, c^{M'^\alpha} = \alpha(c)^{M'}$$

- $M'^\alpha \models_L \theta[\bar{a}] \Leftrightarrow M' \models_{L'} \hat{\alpha}(\theta)[\bar{a}]$

- $f' : M' \rightarrow N'$ is L' -homomorfism (respect. L' -embedding, L' -elementary embedding) \Rightarrow

$f' : M'^\alpha \rightarrow N'^\alpha$ is L -homomorfism (respect. L -embedding, L -elementary embedding)

- $T \subseteq L - sent$, $T' \subseteq L' - sent$ are closed theories such that $\hat{\alpha}^{-1}[T'] \supseteq T$, then $\alpha^* \upharpoonright : (Mod(T'), \preceq_{L'}) \rightarrow (Mod(T), \preceq_L)$ \square

2.2. The category of all AECs (For foundational issues \rightsquigarrow Grothendieck's universes)

objects: the "class of all" AECs

arrows: $\mathcal{K} \subseteq L - str$, $\mathcal{K}' \subseteq L' - str$ (remark that the language is determined from the (non-empty) class, set $\emptyset \subseteq \emptyset - str$)

$\alpha : L \rightarrow L'$ such that the restriction $\alpha^* \upharpoonright : (\mathcal{K}', \preceq_{K'}) \rightarrow (\mathcal{K}, \preceq_K)$ is well defined:

$$\begin{array}{ccc}
L' - str & \xrightarrow{\alpha^*} & L - str \\
i' \uparrow & & \uparrow i \\
\mathcal{K}' & \xrightarrow{\alpha^* \upharpoonright} & \mathcal{K}
\end{array}$$

composition: $(\beta \circ \alpha)^* = \alpha^* \circ \beta^*, \dots$
identities: ok

Example: inclusion : $(K, \preceq_K) \hookrightarrow (L - str, \subseteq_L)$, *inclusion* = $(id_L)^* \upharpoonright$.

Remark: α^* is always a faithful functor. If α is a "surjective" morphism of languages, then α^* is a full functor. \square

2.1 Constructions of AECs

In this subsection we provide new examples of AECs, obtained from construction, for instance: elementary class $(\text{Mod}(T))$ in AECs; pullbacks of AECs; g-types in AECs, etc.

Theories $\text{Mod}(T)$ in AEC

2.3. (FOL) The L -satisfaction relation \models_L :

- establish a Galois connection (Mod, Th) between sets of (first-order) L -sentences and classes of L -structures;
- $(\text{Mod} \upharpoonright, \text{Th} \upharpoonright)$ establish (order reversing) bijection between closed L -theories and elementary classes in language L . \square

Definition 2.4. In an AEC $\mathcal{K} \subseteq L - str$, define:

(i) the "generalized elementary relation" in \mathcal{K} : \equiv_K is "connection relation in $\text{cat}(\mathcal{K})$ ", i.e., is the equivalence relation generated by the arrows in $\text{cat}(\mathcal{K})$:

$M \equiv_K N$ iff $\exists n \in \mathbb{N}, P_0, \dots, P_n \in \mathcal{K}$ and \exists zigzag $M = P_0 \xrightarrow{f_1} P_1 \xleftarrow{f_2} P_2 \xrightarrow{f_3} \dots \xleftarrow{f_{n-1}} P_{n-1} \xrightarrow{f_n} P_n = N$

(ii) a subclass $\mathcal{M} \subseteq \mathcal{K}$ is a "generalized elementary subclass" or a "generalized (closed) theory", when \mathcal{M} is closed under \equiv_K ;

(iii) there is only a set (of representatives) of closed theories (by LS axiom);

(iv) a "maximal consistent g-theory" is an equivalence class module \equiv_K (it is a closed theory) $(\text{Max}(\mathcal{K}) = \mathcal{K} / \equiv_K)$;

(v) a "complete g-theory" is a non-empty subset of a maximal consistent g-theory. \square

Fact 2.5. $\mathcal{M} \subseteq \mathcal{K}$ is g-elementary subclass $\Rightarrow (\mathcal{M}, (\preceq_K) \upharpoonright_{\mathcal{M}})$ is an AEC, a subAEC of \mathcal{K} , and $LS(\mathcal{M}) \leq LS(\mathcal{K})$. \square

Fact 2.6. Let $\alpha : L \rightarrow L', \alpha^* \upharpoonright : \mathcal{K}' \rightarrow \mathcal{K}$:

(i) $M' \equiv_{\mathcal{K}'} N' \Rightarrow M'^{\alpha} \equiv_{\mathcal{K}} N'^{\alpha}$

(ii) *Induced Theory:* Let $\alpha^* \upharpoonright : \mathcal{K}' \rightarrow \mathcal{K}$, $\text{Mod}(T') \subseteq \mathcal{K}'$, then exists $\text{Mod}(T')^{\alpha} :=$ the least closed theory of \mathcal{K} that contains $\{M'^{\alpha} : M' \in \text{Mod}(T')\}$ (iii) a commutative diagram:

$$\begin{array}{ccc}
\mathcal{K}' & \xrightarrow{\alpha^* \upharpoonright} & \mathcal{K} \\
Q' \downarrow & & \downarrow Q \\
\text{Max}(\mathcal{K}') & \xrightarrow{\bar{\alpha}} & \text{Max}(\mathcal{K})
\end{array}$$

$\text{Mod}(T') \subseteq \mathcal{K}'$ maximal consistent theory $\Rightarrow \text{Mod}(T')^{\alpha} \subseteq \mathcal{K}$ is a complete theory \square

Limits of AECs

We will show that the category of all AECs has all finite limits because it has terminal object and pullbacks.

2.7. The terminal AEC: is obtained taking:

- $L = \emptyset, \mathcal{K} = L - str = \text{Set}$
- $M \preceq_{\mathcal{K}} N \Leftrightarrow M \subseteq N$ \square

2.8. Pullbacks of AECs:

- L, L_0, L_1
- $\alpha_i : L \rightarrow L_i, \alpha_i^* \upharpoonright : (\mathcal{K}, \preceq_{\mathcal{K}}) \rightarrow (\mathcal{K}_i, \preceq_{\mathcal{K}_i})$
- $L' = (L_0 \sqcup L_1) / \sim$

$$\pi_i : L_i \rightarrow L' (= L_i \rightarrow L_0 \sqcup L_1 \rightarrow L')$$

$$\delta = \pi_0 \circ \alpha_0 = \pi_1 \circ \alpha_1 : L \rightarrow L'$$

$$\mathcal{K}' \subseteq L' - str$$

$$- M' \in \mathcal{K}' \Leftrightarrow M'^{\pi_0} \in \mathcal{K}_0, M'^{\pi_1} \in \mathcal{K}_1, ((M'^{\pi_0})^{\alpha_0} = (M'^{\pi_1})^{\alpha_1} \in \mathcal{K})$$


$$- M' \preceq_{\mathcal{K}'} N' \Leftrightarrow M'^{\pi_0} \preceq_{\mathcal{K}_0} N'^{\pi_0}, M'^{\pi_1} \preceq_{\mathcal{K}_1} N'^{\pi_1}, (M'^{\delta} \preceq_{\mathcal{K}} N'^{\delta})$$



• *Remark:* $cat(\mathcal{K})_{strict}$ is the pullback in CAT , the category of all (large) categories.

• **Proposition:** \mathcal{K}' is an AEC:

Proof. The validity of the axioms *order*, *isomorphism*, *coherence*, *reunion* follows from the relation $L' - str \cong (L_0 - str) \times_{L-str} (L_1 - str)$. For *Löwenhein-Skolem*: $LS(\mathcal{K}), LS(\mathcal{K}_0), LS(\mathcal{K}_1) \leq \lambda \Rightarrow LS(\mathcal{K}') \leq \lambda$. Take $X \subseteq M' \in \mathcal{K}'$ and consider the diagram:



$P' \in L' - str$ (well defined):

$$(P')^0 \stackrel{h}{=} \bigcup_{n \in \mathbb{N}} P_{2n} \preceq_0 (M')^0$$

$$(P')^1 \cong \bigcup_{n \in \mathbb{N}} P_{2n+1} \preceq_1 (M')^1$$

$$(P')^1 \subseteq_{L_1} (M')^1$$

iso condition in \mathcal{K}_1 entails $(P')^1 \preceq_1 (M')^1$

thus $P' \preceq_{\mathcal{K}'} M'$, $X \subseteq P'$, $\#P' \leq \lambda$

• *Special case:*

$$L = L_0 \cap L_1, L' = L_0 \cup L_1 \text{ (pullback/pushout)}$$

$$\begin{array}{ccc}
(L_0 \cup L_1) - str & \xrightarrow{\pi_1^* \uparrow} & L_1 - str \\
\pi_0^* \downarrow & & \alpha_1^* \downarrow \\
L_0 - str & \xrightarrow{\alpha_0^* \uparrow} & (L_0 \cap L_1) - str
\end{array}$$


□

AECs for g-types

Similar to pullback (smaller class, but with pullback relation).

2.9. Given $M \in \mathcal{K}$, consider \mathcal{K}_M the class of all pairs (N, f) where $N \in \mathcal{K}$, $f : M \rightarrow N \in \text{cat}(\mathcal{K})$.

- $\mathcal{K}_M \subseteq L(M) - str$
- binary relation: $(N_0, f_0) \preceq_{\mathcal{K}_M} (N_1, f_1)$ iff $N_0 \preceq_K N_1$ and $(N_0, f_0) \subseteq_{L(M)} (N_1, f_1)$
- $(\mathcal{K}_M, \preceq_{\mathcal{K}_M})$ is AEC $(LS(\mathcal{K}) \leq \lambda \Rightarrow LS(\mathcal{K}_M) \leq \lambda + \#M)$
- $\text{cat}(\mathcal{K}_M)$: commutative triangles in $\text{cat}(\mathcal{K})$ over M




□

2.10. An AEC for g-types

Given $M \in \mathcal{K}$ and I a set, consider $\mathcal{K}_{M,I}$ the class of all triples (N, f, \bar{a}) where $N \in \mathcal{K}$, $\bar{a} : I \rightarrow N$, $f : M \rightarrow N \in \text{cat}(\mathcal{K})$.

- $\mathcal{K}_{M,I} \subseteq L(M \cup I) - str$
- binary relation: $(N_0, f_0, \bar{a}_0) \preceq_{\mathcal{K}_{M,I}} (N_1, f_1, \bar{a}_1)$ iff $N_0 \preceq_K N_1$ and $(N_0, f_0, \bar{a}_0) \subseteq_{L(M,I)} (N_1, f_1, \bar{a}_1)$
- $(\mathcal{K}_{M,I}, \preceq_{\mathcal{K}_{M,I}})$ is AEC $(LS(\mathcal{K}) \leq \lambda \Rightarrow LS(\mathcal{K}_{M,I}) \leq \lambda + \#M + \#I)$
- $\text{cat}(\mathcal{K}_{M,I})$: commutative triangles in $\text{cat}(\mathcal{K})$ over M and I
- we may assume $M \cap I = \emptyset$ (e.g., take $I := C \times \{M\}$)



Fact 2.11. The I -types over M are precisely the g -maximal consistent theories in $\mathcal{K}_{M,I}$, thus they are AECs in the language $L(M \cup I)$. □

$$g - T(M, I) = \text{Max}(\mathcal{K}_{M,I})$$

□

Fact 2.12. Induced arrows on g-types:

$\alpha : L \rightarrow L'$, $\alpha^* \uparrow : \mathcal{K}' \rightarrow \mathcal{K}$, $h' : M'_0 \rightarrow M'_1 \in \text{cat}(\mathcal{K}')$, C a set of constants

- a commutative diagram of languages and morphisms:
 - $I_j = C \times \{|M'_j|\}$, $j = 0, 1$;
 - $M_j := M'_j{}^\alpha \Rightarrow |(M_j)^\alpha| = |M'_j|$, $j = 0, 1$;
 - $h := (h')^\alpha : M_0 \rightarrow M_1$

$$\begin{array}{ccc}
L(M_0 \cup I_0) & \xrightarrow{\tilde{h}} & L(M_1 \cup I_1) \\
\tilde{\alpha}^0 \downarrow & & \downarrow \tilde{\alpha}^1 \\
L'(M'_0 \cup I_0) & \xrightarrow{\tilde{h}'} & L'(M'_1 \cup I_1)
\end{array}$$

• a commutative diagram of AECs and morphisms:

$$\begin{array}{ccc}
\mathcal{K}_{M_0, I_0} & \xleftarrow{(\tilde{h})^*} & \mathcal{K}_{M_1, I_1} \\
(\tilde{\alpha}^0)^* \uparrow & & \uparrow (\tilde{\alpha}^1)^* \\
\mathcal{K}'_{M'_0, I_0} & \xleftarrow{(\tilde{h}')^*} & \mathcal{K}'_{M'_1, I_1}
\end{array}$$

• a commutative diagram of sets and functions:

$$\begin{array}{ccc}
\text{Max}(\mathcal{K}_{M_0, I_0}) & \xleftarrow{\tilde{h}} & \text{Max}(\mathcal{K}_{M_1, I_1}) \\
\tilde{\alpha}^0 \uparrow & & \uparrow \tilde{\alpha}^1 \\
\text{Max}(\mathcal{K}'_{M'_0, I_0}) & \xleftarrow{\tilde{h}'} & \text{Max}(\mathcal{K}'_{M'_1, I_1})
\end{array}$$

equivalently

$$\begin{array}{ccc}
g - T(M_0, I_0) & \xleftarrow{\tilde{h}} & g - T(M_1, I_1) \\
\tilde{\alpha}^0 \uparrow & & \uparrow \tilde{\alpha}^1 \\
g - T(M'_0, I_0) & \xleftarrow{\tilde{h}'} & g - T(M'_1, I_1)
\end{array}$$

□

Fact 2.13. λ -Topology on g-types (II):

(i) $\lambda \geq LS(\mathcal{K}) \Rightarrow X^\lambda(M, I)$ is the least topology such that:

– $i_N : g - T(M, I) \rightarrow g - T(N, I)$ is continuous, $N \in S^\lambda(M) := \{N \in \mathcal{K} : N \preceq_K M, \#N \leq \lambda\}$, $X^\lambda(N, I) = (g - T(N, I), \text{discrete})$

equivalently:

– $\text{can}_M^\lambda : X^\lambda(M, I) \rightarrow \lim_{N \in S^\lambda(M)} X^\lambda(N, I)$ (with induced pro-topology) is continuous;

equivalently:

– if $M \cong \text{colim}_{j \in J} P_j$ (directed colimit), $\#P_j \leq \lambda$

$\text{can} : X^\lambda(M, I) \rightarrow \lim_{j \in J} X^\lambda(P_j, I) \approx \lim_{N \in S^\lambda(M)} X^\lambda(N, I)$ (with induced pro-topology) is continuous

(ii) Properties of \mathcal{K} : pro-topology?, injective?, surjective?

– λ -Tame $\Leftrightarrow \forall M, I$, can is injective (T_2)

– “ λ -compact” $\Leftrightarrow \forall M, I$, can is surjective

(iii) Induced arrows on g-types:

$\alpha : L \rightarrow L'$, $\alpha^* \vdash : \mathcal{K}' \rightarrow \mathcal{K}$, $h' : M'_0 \rightarrow M'_1 \in \text{cat}(\mathcal{K}')$, C a set of constants, $\lambda \geq LS(\mathcal{K}), LS(\mathcal{K}')$

• $M' \cong \text{colim}_{j \in J} P'_j \Rightarrow M'^\alpha \cong \text{colim}_{j \in J} P_j^{\alpha'}$

$\hat{h}' : S^\lambda(M'_0) \rightarrow S^\lambda(M'_1)$

• two diagrams of spaces and continuous functions

$$\begin{array}{ccc}
\lim_{N_0 \in S^\lambda(M_0)} X^\lambda(N_0, I_0) & \xleftarrow{\tilde{h}} & \lim_{N_1 \in S^\lambda(M_1)} X^\lambda(N_1, I_1) \\
\uparrow \tilde{\alpha}^0 & & \uparrow \tilde{\alpha}^1 \\
\lim_{N'_0 \in S^\lambda(M'_0)} X^\lambda(N'_0, I_0) & \xleftarrow{\tilde{h}'} & \lim_{N'_1 \in S^\lambda(M'_1)} X^\lambda(N'_1, I_1)
\end{array}$$

then

$$\begin{array}{ccc}
X^\lambda(M_0, I_0) & \xleftarrow{\tilde{h}} & X^\lambda(M_1, I_1) \\
\uparrow \tilde{\alpha}^0 & & \uparrow \tilde{\alpha}^1 \\
X^\lambda(M'_0, I_0) & \xleftarrow{\tilde{h}'} & X^\lambda(M'_1, I_1)
\end{array}$$

- If \mathcal{K}' has **(AP)** $\Rightarrow \tilde{h}' : X^\lambda(M'_1, I) \rightarrow X^\lambda(M'_0, I)$ and $\tilde{h} : X^\lambda(M_1, I) \rightarrow X^\lambda(M_0, I)$ are surjective. □

2.2 New (global) axioms and applications

”Local” axioms and applications:

Where ”local” means ”for each AEC fixed”

Ex.:

(JEP): Joint Embedding Property;

(AP): Amalgamation Property;

(LRP): ”Local Robinson’s Property”: $M_0 \equiv_{\mathcal{K}} M_1$ iff $\exists N \in \mathcal{K} \exists j_0, j_1 \in \text{cat}(\mathcal{K}) j_0 : M_0 \rightarrow N, j_1 : M_1 \rightarrow N$

Robinson’s diagram method:

2.14. If \mathcal{K} is an AEC in language L , then for each set of ”new constants” $L' := L \cup \{c_i : i \in I\}$ it is available the AEC \mathcal{K}' in the language L' whose objects are pairs (M, \bar{a}) , where \bar{a} is a I -sequence of elements of M and $(M, \bar{a}) \preceq_{\mathcal{K}'} (N, \bar{b})$ iff: $M \preceq_{\mathcal{K}} N$ and $\bar{a} = \bar{b}$ ($\Leftrightarrow M \preceq_{\mathcal{K}} N$ and $(M, \bar{a}) \subseteq_{L'} (N, \bar{b})$)

- \mathcal{K}' it is the pullback:

$$\begin{array}{ccc}
\mathcal{K} & & \mathcal{K}' \xrightarrow{j^* \uparrow} \mathcal{K} \\
\downarrow \text{incl} & & \downarrow \text{incl} \\
L' - \text{str} \xrightarrow{j^*} L - \text{str} & & L' - \text{str} \xrightarrow{j^*} L - \text{str}
\end{array}$$

□

Proposition 2.15. **(LRP)** $\exists j : M \rightarrow N \in \text{cat}(\mathcal{K})$ iff $\exists \bar{b}$ such that $(M, \bar{a}) \equiv_{\mathcal{K}(M)} (N, \bar{b})$ where $\mathcal{K}(M) \subseteq L(M) - \text{str}$

$$\begin{array}{ccc}
(M, \bar{a}) & & (N, \bar{b}) \\
\downarrow t & & \downarrow h \\
(P, \bar{c}) & & (P, \bar{c})
\end{array}
\quad
\begin{array}{ccc}
(M, \bar{a}) \xrightarrow{t \uparrow} (M', \bar{c}) & \xrightarrow{\text{incl}} & (P, \bar{c}) \\
\cong \downarrow & & \cong \downarrow \\
(N, \bar{b}) \xrightarrow{h \uparrow} (N', \bar{c}) & \xrightarrow{\text{incl}} & (P, \bar{c}) \\
\cong \downarrow & & \cong \downarrow
\end{array}$$

$$\begin{array}{ccc}
(M', \bar{c}) & \xrightarrow{\subseteq_{L(M)}} & (N', \bar{c}) \\
\downarrow \cong & & \downarrow \cong \\
(P, \bar{c}) & & (P, \bar{c})
\end{array}
\quad
\begin{array}{ccccc}
M, \bar{a}) & \xrightarrow[\cong]{t!} & (M', \bar{c}) & \xrightarrow[\cong]{incl} & (N', \bar{c}) & \xrightarrow[\cong]{(h!)^{-1}} & (N, \bar{b}) \\
(M, \bar{a}) & \xrightarrow{j} & (N, \bar{b}) & & & &
\end{array}$$

□

The (global) category AEC:

The Grothendieck's gluing of all AECs

objects: pairs (M, \mathcal{K}) , where $M \in \mathcal{K}$

arrows: pairs $(h, \varphi) : (M, \mathcal{K}) \rightarrow (M', \mathcal{K}')$

$\varphi : L \rightarrow L', \Phi = \varphi^* \cdot : \mathcal{K}' \rightarrow \mathcal{K}, h : M \rightarrow \Phi(M') \in \text{cat}(\mathcal{K})$

identities: $(id_M, id_L) : (M, \mathcal{K}) \rightarrow (M, \mathcal{K})$

composition: $(M, \mathcal{K}) \xrightarrow{(h, \varphi)} (M', \mathcal{K}') \xrightarrow{(h', \varphi')} (M'', \mathcal{K}'') = (M, \mathcal{K}) \xrightarrow{(\Phi(h') \circ h, \varphi' \circ \varphi)} (M'', \mathcal{K}'')$

$(M \xrightarrow{h} \Phi(M') \xrightarrow{\Phi(h')} \Phi(\Phi'(M'')))$

□

2.16. New (global) equivalences in AEC:

– the connection relation: $(N_0, \mathcal{K}_0) \equiv (N_1, \mathcal{K}_1)$

– relative equivalence over $M \in \mathcal{K}$:

$(N_0, \mathcal{K}_0) \equiv_{(M, \mathcal{K})} (N_1, \mathcal{K}_1)$, with $h_i : M \rightarrow (N_i)^{\alpha_i}$

Ex. I -types over $M : (M, \mathcal{K}) \rightarrow ((N, \bar{a}), \mathcal{K}(I))$

□

New (global) axioms in AEC

Goal: Determine interesting subcategories of AEC satisfying interesting (global) axioms.

(GAP) "Global Amalgamation Property"

entails a simpler description of the relative equivalence over $M \in \mathcal{K}$:

$(N_0, \mathcal{K}_0) \equiv_{(M, \mathcal{K})} (N_1, \mathcal{K}_1)$, with $h_i : M \rightarrow (N_i)^{\alpha_i}$

iff $\exists \mathcal{K}' \exists P' \exists j_i : N_i \rightarrow P'^{\beta_i}$ such that the diagram below commutes

$$\begin{array}{ccc}
(M, \mathcal{K}) & \xrightarrow{(h_1, \alpha_1)} & (N_1, \mathcal{K}_1) \\
\downarrow (h_0, \alpha_0) & & \downarrow (j_1, \beta_1) \\
(N_0, \mathcal{K}_0) & \xrightarrow{(j_0, \beta_0)} & (P', \mathcal{K}')
\end{array}$$

(TRP) "Transversal Robinson Property":

• $\alpha_i : L \rightarrow L_i, i = 0, 1$

$\mathcal{K}, \mathcal{K}_0, \mathcal{K}_1$

• $(M_0)^K \equiv_K (M_1)^K \Rightarrow \exists N_0 \in \mathcal{K}_0$ such that:

– $M_0 \preceq_{\mathcal{K}_0} N_0$ in \mathcal{K}_0

– $\exists j : (M_1)^K \rightarrow (N_0)^K$ in $\text{cat}(\mathcal{K})$

(GRP) "Global Robinson Property":

• $\alpha_i : L \rightarrow L_i, \beta_i : L_i \rightarrow (L_0 \cup L_1) / \sim, i = 0, 1$

$\mathcal{K}, \mathcal{K}_0, \mathcal{K}_1, \mathcal{K}' = \text{pullback}$


• $(M_0)^K \equiv_K (M_1)^K \Rightarrow \exists P' \in \mathcal{K}'$ such that:

– $M_0 \preceq_{\mathcal{K}_0} (P')^{K_0}$ in \mathcal{K}_0

– $\exists h : M_1 \rightarrow (P')^{K_1}$ in $\text{cat}(\mathcal{K}_1)$

Proposition 2.17. $(TRP) \Rightarrow (GRP)$

Proof. Just take zig-zags as in the proof of LS property in the pullback AEC □



Craig Interpolation Property:

2.18. Craig interpolation in FOL:

- $\psi_0, \psi_1 \in L' - sent$ $\psi_0 \vdash \psi_1 \Rightarrow$ exists $\theta \in L = L_0 \cap L_1$: $\psi_0 \vdash \theta, \theta \vdash \psi_1$
- $\{\psi_0, \neg\psi_1\} \vdash \perp \Rightarrow$ exists $\theta \in L = L_0 \cap L_1$: $\psi_0 \vdash \theta, \neg\psi_1 \vdash \neg\theta$
- $T_0 \cup T_1$ has no L' -model \Rightarrow exists $\theta \in L = L_0 \cap L_1$: $T_0 \models \theta, T_1 \models \neg\theta$
- $L = L_0 \cap L_1, T_i \upharpoonright_L = \{\psi \in L - sent : T_i \vdash_L \psi\}$

$T_0 \cup T_1$ has no L' -model $\Rightarrow T_0 \upharpoonright_L \cup T_1 \upharpoonright_L$ has no L -model

- $\nexists M' \in L' - str$ such that $(M')^{L_i} \in Mod(T_i), i = 0, 1 \Rightarrow \nexists M \in L - str$ with $M \in Mod(T_0 \upharpoonright_L) \cap Mod(T_1 \upharpoonright_L)$ □

(CIP) "Craig Interpolation Property":

- $\alpha_i : L \rightarrow L_i, \beta_i : L_i \rightarrow (L_0 \cup L_1) / \sim, i = 0, 1$

$\mathcal{K}, \mathcal{K}_0, \mathcal{K}_1, \mathcal{K}' = pullback$

$Mod(T_i) \subseteq \mathcal{K}_i, i = 0, 1.$

- $\nexists M' \in \mathcal{K}'$ such that $(M')^{K_i} \in Mod(T_i), i = 0, 1 \Rightarrow \nexists M \in \mathcal{K}$ with $M \in Mod(T_0 \upharpoonright_{\mathcal{K}}) \cap Mod(T_1 \upharpoonright_{\mathcal{K}})$ (the induced theories)

Proposition 2.19. $(TRP) \Rightarrow (CIP)$

Proof. (contrapositive)

Let $N \in \mathcal{K}$ with $N \in \text{Mod}(T_0 \upharpoonright_L) \cap \text{Mod}(T_1 \upharpoonright_L)$.

Then: there are $M_i \in \text{Mod}(T_i)$, $i = 0, 1$, with $(M_0)^K \equiv_K N \equiv_K (M_1)^K$.

As $(TRP) \Rightarrow (GRP)$, there is $P' \in \mathcal{K}'$ such that:

- $M_0 \preceq_{K_0} (P')^{K_0}$ in \mathcal{K}_0
- $\exists h : M_1 \rightarrow (P')^{K_1}$ in $\text{cat}(\mathcal{K}_1)$.

Then:

- $(P')^{K_0} \equiv_{K_0} M_0 \in \text{Mod}(T_0)$;
- $(P')^{K_1} \equiv_{K_1} M_1 \in \text{Mod}(T_1)$.

□

3 Final Remarks and Future Works

- Introduce and study other global axioms (inspired on model theory of FOL).
- Consider functors induced by more general language morphisms $f : L \rightarrow L'$:
 - (i) non injective;
 - (ii) "flexible morphism":
 - $f_n \in L$ (n-ary functional symbol) $\mapsto t_n \in L'$ (n-ary term)
 - $R_n \in L$ (n-ary relational symbol) $\mapsto \varphi_n \in L'$ (n-ary atomic formula)
- Consider more general functors between AECs (ex. functors that are not "over Set ").
- Study metric abstract elementary classes (MAECs) in this global setting.

References

- [AR] J. Adámek, J. Rosický, **Locally Presentable and Accessible Categories**, Lecture Notes Series of the LMS **189**, Cambridge University Press, Cambridge, Great Britain, 1994.
- [Bal1] J. Baldwin, *Abstract Elementary Classes: Some Answers, More Questions*, July 7 2006, 18 pages.
- [Bal2] J. Baldwin, **Categoricity**, University Lecture Series **50**, American Mathematical Society, 2009.
- [BR] T. Beke, J. Rosický, *Abstract elementary classes and accessible categories*, Annals of Pure and Applied Logic **163** (**12**), 2012, 2008-2017.
- [Kue] D. Kueker, *Abstract elementary classes and infinitary logics*, Annals of Pure and Applied Logic, **156**(**11**), 2008, 274-286.
- [Lie1] M.J. Lieberman, **Topological and Category-Theoretic Aspects of Abstract Elementary Classes**, Phd, University of Michigan, 99 pages, 2009.
- [Lie2] M.J. Lieberman, *Category-theoretic aspects of abstract elementary classes*, Annals of Pure and Applied Logic, **162**(**11**), 2011, 903-915.
- [She] S. Shelah, *Classification of nonelementary classes II, abstract elementary classes*, Lecture Notes in Mathematics **1292**, 1987, 419-497.
- [Van] M. VanDieren, *Categoricity in Abstract Elementary Classes with no maximal models*, available at <http://arxiv.org/pdf/math/0510579.pdf>, 2005, 57 pages.