

# Left localizable rings and their characterizations

V. V. Bavula

## Abstract

A new class of rings, *the class of left localizable rings*, is introduced. A ring  $R$  is *left localizable* if each nonzero element of  $R$  is invertible in some left localization  $S^{-1}R$  of the ring  $R$ . Explicit criteria are given for a ring to be a left localizable ring provided the ring has only finitely many maximal left denominator sets (eg, this is the case if a ring has a left Artinian left quotient ring). It is proved that a ring with finitely many maximal left denominator sets is a left localizable ring iff its left quotient ring is a direct product of finitely many division rings. A characterization is given of the class of rings that are finite direct product of left localization maximal rings.

*Key Words:* a left localizable ring, a left localization maximal ring, the largest left quotient ring of a ring, the largest regular left Ore set of a ring, the classical left quotient ring of a ring, denominator set.

*Mathematics subject classification 2000:* 16U20, 16P50, 16S85.

## Contents

1. Introduction.
2. Direct products of left localization maximal rings and their characterization.
3. Left localizable rings and their characterization.
4. The core of a left Ore set.

## 1 Introduction

Throughout, module means a left module. In this paper the following notation will remained fixed.

### Notation:

- $R$  is a ring with 1 and  $\mathcal{C}_R$  is the set of (left and right) regular elements of the ring  $R$  (i.e.  $\mathcal{C}_R$  is the set of non-zero-divisors of  $R$ );
- $Q_{l,cl}(R) := \mathcal{C}_R^{-1}R$  is the *classical left quotient ring* of  $R$  (if it exists);
- $\text{Ore}_l(R) := \{S \mid S \text{ is a left Ore set in } R\}$ ;
- $\text{Den}_l(R) := \{S \mid S \text{ is a left denominator set in } R\}$ ;
- $\text{Loc}_l(R) := \{[S^{-1}R] \mid S \in \text{Den}_l(R)\}$  where  $[S^{-1}R]$  is an  *$R$ -isomorphism class* of the ring  $S^{-1}R$  (a ring isomorphism  $\sigma : S^{-1}R \rightarrow S'^{-1}R$  is called an  *$R$ -isomorphism* if  $\sigma(\frac{r}{1}) = \frac{r}{1}$  for all elements  $r \in R$ );
- $\iota_R := \text{l.rad}(R)$  is the *left localization radical* of the ring  $R$ ;
- $\text{Ass}_l(R) := \{\text{ass}(S) \mid S \in \text{Den}_l(R)\}$  where  $\text{ass}(S) := \{r \in R \mid sr = 0 \text{ for some } s = s(r) \in S\}$ ;
- $\text{Den}_l(R, \mathfrak{a}) := \{S \in \text{Den}_l(R) \mid \text{ass}(S) = \mathfrak{a}\}$  where  $\mathfrak{a} \in \text{Ass}_l(R)$ ;
- $\text{Den}_l(R, 0)$  is the set of regular left denominator sets  $S$  in  $R$  ( $S \subseteq \mathcal{C}_R$ );

- $S_{\mathfrak{a}} = S_{\mathfrak{a}}(R) = S_{l,\mathfrak{a}}(R)$  is the *largest element* of the poset  $(\text{Den}_l(R, \mathfrak{a}), \subseteq)$  and  $Q_{\mathfrak{a}}(R) := Q_{l,\mathfrak{a}}(R) := S_{\mathfrak{a}}^{-1}R$  is the *largest left quotient ring associated to  $\mathfrak{a}$* ,  $S_{\mathfrak{a}}$  exists (Theorem 2.1, [2]);
- In particular,  $S_0 = S_0(R) = S_{l,0}(R)$  is the largest element of the poset  $(\text{Den}_l(R, 0), \subseteq)$  and  $Q_l(R) := S_0^{-1}R$  is the largest left quotient ring of  $R$ ;
- $\text{Loc}_l(R, \mathfrak{a}) := \{[S^{-1}R] \mid S \in \text{Den}_l(R, \mathfrak{a})\}$ .

**Left localizable rings and their characterization.** The aim of the paper is to introduce a new class of rings, the class of left localizable rings, and to give several characterizations of them (Theorem 3.5, Theorem 3.8 and Theorem 3.9) in the case when they admit only finitely many maximal left denominator sets. Notice that each ring with left Artinian, left quotient ring has only finitely many maximal left denominator sets, [4]. A ring  $R$  is called a *left localizable ring* if for each nonzero element  $r \in R$  there exists a left denominator set  $S = S(r)$  such that  $r \in S$  (equivalently, if for each nonzero element  $r \in R$  there exists a left denominator set  $S' = S'(r)$  such that  $\frac{r}{1}$  is a unit in  $S'^{-1}R$ ).

- **(Theorem 3.5)** *Let  $R$  be a ring with  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$ . Let  $\mathfrak{a}_i := \text{ass}(S_i)$ ;  $\sigma_i : R \rightarrow R_i := S_i^{-1}R$ ,  $r \mapsto \frac{r}{1} = r_i$ ; and  $\sigma := \prod_{i=1}^n \sigma_i : R \rightarrow \prod_{i=1}^n R_i$ ,  $r \mapsto (r_1, \dots, r_n)$ . The following statements are equivalent.*

1. *The ring  $R$  is a left localizable ring.*
2.  *$\mathfrak{l}_R = 0$  and the rings  $R_1, \dots, R_n$  are division rings.*
3. *The homomorphism  $\sigma$  is an injection and the rings  $R_1, \dots, R_n$  are division rings.*

- **(Theorem 3.9)** *Let  $R$  be a ring. The following statements are equivalent.*

1. *The ring  $R$  is a left localizable ring with  $n := |\max.\text{Den}_l(R)| < \infty$ .*
2.  *$Q_{l,cl}(R) = R_1 \times \dots \times R_n$  where  $R_i$  are division rings.*
3. *The ring  $R$  is a semiprime left Goldie ring with  $\text{udim}(R) = |\text{Min}(R)| = n$  where  $\text{Min}(R)$  is the set of minimal prime ideals of the ring  $R$ .*
4.  *$Q_l(R) = R_1 \times \dots \times R_n$  where  $R_i$  are division rings.*

**Direct product of left localization maximal rings and their characterization.** A ring  $R$  is called a *left localization maximal ring* if every left denominator set of  $R$  consists of units, i.e. one cannot invert anything new ‘on the left’ in  $R$ , [3]. Theorem 2.11 is a characterization of rings that are isomorphic to finite direct product of left localization maximal rings.

- **(Theorem 2.11)** *A ring  $R$  is a direct product of  $n$  left localization maximal rings iff the following conditions hold:*

1.  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$ .
2.  $\mathfrak{l}_R = 0$ .
3.  $\text{ass}(S_i) + \text{ass}(S_j) = R$  for all  $i \neq j$ .
4. *The factor rings  $R/\text{ass}(S_i)$ ,  $i = 1, \dots, n$ , are left localization maximal rings.*

**The core of a left Ore set.** In Section 4, properties of the core of a left Ore set are studied. Let  $R$  be a ring and  $S$  be its left Ore set. The subset of  $S$ ,

$$S_c := \{s \in S \mid \ker(s \cdot) = \text{ass}(S)\}$$

is called the *core* of the left Ore set  $S$  where  $s \cdot : R \rightarrow R$ ,  $r \mapsto sr$ , [3].

- **(Theorem 4.2)** *Suppose that  $S \in \text{Den}_l(R, \mathfrak{a})$  and  $S_c \neq \emptyset$ . Then*

1.  $S_c \in \text{Den}_l(R, \mathfrak{a})$ .
2. The map  $\theta : S_c^{-1}R \rightarrow S^{-1}R$ ,  $s^{-1}r \mapsto s^{-1}r$ , is a ring isomorphism. So,  $S_c^{-1}R \simeq S^{-1}R$ .

The next theorem gives an explicit description of the cores of maximal left denominator sets of a left localizable ring provided  $|\text{max.Den}_l(R)| < \infty$ .

• **(Theorem 4.5)** *Let  $R$  be a left localizable ring such that  $\text{max.Den}_l(R) = \{S_1, \dots, S_n\}$ .*

1. *If  $n = 1$  then  $S_{1,c} = S_1 = R \setminus \{0\}$ .*
2. *If  $n \geq 2$  then  $S_{i,c} = S_i \cap \bigcap_{j \neq i} \mathfrak{a}_j$  where  $\mathfrak{a}_j = \text{ass}(S_j)$ .*

## 2 Direct products of left localization maximal rings and their characterization

The aim of this section is to give a criterion of when a ring is isomorphic to a finite direct product of left localization maximal rings (Theorem 2.11). At the beginning of the section necessary results are given that are used in the proofs of the paper.

In [2], we introduce the following new concepts and to prove their existence for an *arbitrary* ring: *the largest left quotient ring of a ring, the largest regular left Ore set of a ring, the maximal left quotient rings of a ring, the largest (two-sided) quotient ring of a ring, the maximal (two-sided) quotient rings of a ring, a (left) localization maximal ring*. Let us give more details.

**The largest regular left Ore set and the largest left quotient ring of a ring.** Let  $R$  be a ring. A *multiplicatively closed subset*  $S$  of  $R$  or a *multiplicative subset* of  $R$  (i.e. a multiplicative sub-semigroup of  $(R, \cdot)$  such that  $1 \in S$  and  $0 \notin S$ ) is said to be a *left Ore set* if it satisfies the *left Ore condition*: for each  $r \in R$  and  $s \in S$ ,

$$Sr \cap Rs \neq \emptyset.$$

Let  $\text{Ore}_l(R)$  be the set of all left Ore sets of  $R$ . For  $S \in \text{Ore}_l(R)$ ,  $\text{ass}(S) := \{r \in R \mid sr = 0 \text{ for some } s \in S\}$  is an ideal of the ring  $R$ .

A left Ore set  $S$  is called a *left denominator set* of the ring  $R$  if  $rs = 0$  for some elements  $r \in R$  and  $s \in S$  implies  $tr = 0$  for some element  $t \in S$ , i.e.  $r \in \text{ass}(S)$ . Let  $\text{Den}_l(R)$  be the set of all left denominator sets of  $R$ . For  $S \in \text{Den}_l(R)$ , let  $S^{-1}R = \{s^{-1}r \mid s \in S, r \in R\}$  be the *left localization* of the ring  $R$  at  $S$  (the *left quotient ring* of  $R$  at  $S$ ).

In general, the set  $\mathcal{C}$  of regular elements of a ring  $R$  is neither left nor right Ore set of the ring  $R$  and as a result neither left nor right classical quotient ring ( $Q_{l,cl}(R) := \mathcal{C}^{-1}R$  and  $Q_{r,cl}(R) := R\mathcal{C}^{-1}$ ) exists. Remarkably, there exists the largest regular left Ore set  $S_0 = S_{l,0} = S_{l,0}(R)$ . This means that the set  $S_{l,0}(R)$  is an Ore set of the ring  $R$  that consists of regular elements (i.e.,  $S_{l,0}(R) \subseteq \mathcal{C}$ ) and contains all the left Ore sets in  $R$  that consist of regular elements. Also, there exists the largest regular (left and right) Ore set  $S_{l,r,0}(R)$  of the ring  $R$ . In general, all the sets  $\mathcal{C}$ ,  $S_{l,0}(R)$ ,  $S_{r,0}(R)$  and  $S_{l,r,0}(R)$  are distinct, for example, when  $R = \mathbb{I}_1 := K\langle x, \frac{d}{dx}, f \rangle$  is the ring of polynomial integro-differential operators over a field  $K$  of characteristic zero, [1].

*Definition*, [1], [2]. The ring

$$Q_l(R) := S_{l,0}(R)^{-1}R$$

(respectively,  $Q_r(R) := RS_{r,0}(R)^{-1}$  and  $Q(R) := S_{l,r,0}(R)^{-1}R \simeq RS_{l,r,0}(R)^{-1}$ ) is called the *largest left* (respectively, *right and two-sided*) *quotient ring* of the ring  $R$ .

In general, the rings  $Q_l(R)$ ,  $Q_r(R)$  and  $Q(R)$  are not isomorphic, for example, when  $R = \mathbb{I}_1$ , [1]. The next theorem gives various properties of the ring  $Q_l(R)$ . In particular, it describes its group of units.

**Theorem 2.1** [2]

1.  $S_0(Q_l(R)) = Q_l(R)^*$  and  $S_0(Q_l(R)) \cap R = S_0(R)$ .
2.  $Q_l(R)^* = \langle S_0(R), S_0(R)^{-1} \rangle$ , i.e. the group of units of the ring  $Q_l(R)$  is generated by the sets  $S_0(R)$  and  $S_0(R)^{-1} := \{s^{-1} \mid s \in S_0(R)\}$ .
3.  $Q_l(R)^* = \{s^{-1}t \mid s, t \in S_0(R)\}$ .
4.  $Q_l(Q_l(R)) = Q_l(R)$ .

The set  $(\text{Den}_l(R), \subseteq)$  is a poset (partially ordered set). In [2], it is proved that the set  $\text{max.Den}_l(R)$  of its maximal elements is a *non-empty* set.

**The maximal denominator sets and the maximal left localizations of a ring.**

*Definition.* [2]. An element  $S$  of the set  $\text{max.Den}_l(R)$  is called a *maximal left denominator set* of the ring  $R$  and the ring  $S^{-1}R$  is called a *maximal left quotient ring* of the ring  $R$  or a *maximal left localization ring* of the ring  $R$ . The intersection

$$\mathfrak{l}_R := \text{l.rad}(R) := \bigcap_{S \in \text{max.Den}_l(R)} \text{ass}(S) \quad (1)$$

is called the *left localization radical* of the ring  $R$ , [2].

For a ring  $R$ , there is the canonical exact sequence

$$0 \rightarrow \mathfrak{l}_R \rightarrow R \xrightarrow{\sigma} \prod_{S \in \text{max.Den}_l(R)} S^{-1}R, \quad \sigma := \prod_{S \in \text{max.Den}_l(R)} \sigma_S, \quad (2)$$

where  $\sigma_S : R \rightarrow S^{-1}R, r \mapsto \frac{r}{1}$ .

**The maximal elements of  $\text{Ass}_l(R)$ .** Let  $\text{max.Ass}_l(R)$  be the set of maximal elements of the poset  $(\text{Ass}_l(R), \subseteq)$  and

$$\text{ass.max.Den}_l(R) := \{\text{ass}(S) \mid S \in \text{max.Den}_l(R)\}. \quad (3)$$

These two sets are equal (Proposition 2.4), a proof is based on Lemma 2.2. For a non-empty subset  $X$  of  $R$ , let  $\text{r.ass}(X) := \{r \in R \mid rx = 0 \text{ for some } x \in X\}$ .

**Lemma 2.2** [2] *Let  $S \in \text{Den}_l(R, \mathfrak{a})$  and  $T \in \text{Den}_l(R, \mathfrak{b})$  be such that  $\mathfrak{a} \subseteq \mathfrak{b}$ . Let  $ST$  be the multiplicative semigroup generated by  $S$  and  $T$  in  $(R, \cdot)$ . Then*

1.  $\text{r.ass}(ST) \subseteq \mathfrak{b}$ .
2.  $ST \in \text{Den}_l(R, \mathfrak{c})$  and  $\mathfrak{b} \subseteq \mathfrak{c}$ .

**Corollary 2.3** *Let  $R$  be a ring,  $S \in \text{max.Den}_l(R)$  and  $T \in \text{Den}_l(R)$ . Then  $T \subseteq S$  iff  $\text{ass}(T) \subseteq \text{ass}(S)$ .*

*Proof.*  $(\Rightarrow)$  If  $T \subseteq S$  then  $\text{ass}(T) \subseteq \text{ass}(S)$ .

$(\Leftarrow)$  If  $\text{ass}(T) \subseteq \text{ass}(S)$  then, by Lemma 2.2,  $ST \in \text{Den}_l(R)$  and  $S \subseteq ST$ , hence  $S = ST$ , by the maximality of  $S$ . Then  $T \subseteq S$ .  $\square$

**Proposition 2.4** [2]  $\text{max.Ass}_l(R) = \text{ass.max.Den}_l(R) \neq \emptyset$ . *In particular, the ideals of this set are incomparable (i.e. neither  $\mathfrak{a} \subseteq \mathfrak{b}$  nor  $\mathfrak{a} \supseteq \mathfrak{b}$ ).*

**The localization maximal rings.** The set  $(\text{Loc}_l(R), \rightarrow)$  is a partially ordered set (poset) where  $A \rightarrow B$  if there is an  $R$ -homomorphism  $A \rightarrow B$ . There is no oriented loops in the poset  $\text{Loc}_l(R)$  apart from the  $R$ -isomorphism  $A \rightarrow A$ .

Let  $\text{max.Loc}_l(R)$  be the set of maximal elements of the poset  $(\text{Loc}_l(R), \rightarrow)$ . Then (see [2]),

$$\text{max.Loc}_l(R) = \{S^{-1}R \mid S \in \text{max.Den}_l(R)\} = \{Q_l(R/\mathfrak{a}) \mid \mathfrak{a} \in \text{ass.max.Den}_l(R)\}. \quad (4)$$

*Definition, [2].* A ring  $A$  is called a *left localization maximal ring* if  $A = Q_l(A)$  and  $\text{Ass}_l(A) = \{0\}$ . A ring  $A$  is called a *right localization maximal ring* if  $A = Q_r(A)$  and  $\text{Ass}_r(A) = \{0\}$ . A ring  $A$  which is a left and right localization maximal ring is called a *(left and right) localization maximal ring* (i.e.  $Q_l(A) = A = Q_r(A)$  and  $\text{Ass}_l(A) = \text{Ass}_r(A) = \{0\}$ ).

The next theorem is a criterion of when a left quotient ring of a ring is a maximal left quotient ring of the ring.

**Theorem 2.5** [2] *Let a ring  $A$  be a left localization of a ring  $R$ , i.e.  $A \in \text{Loc}_l(R, \mathfrak{a})$  for some  $\mathfrak{a} \in \text{Ass}_l(R)$ . Then  $A \in \text{max.Loc}_l(R)$  iff  $Q_l(A) = A$  and  $\text{Ass}_l(A) = \{0\}$ , i.e.  $A$  is a left localization maximal ring (clearly,  $\mathfrak{a} \in \text{ass.max.Den}_l(R)$ ).*

Theorem 2.5 shows that the left localization maximal rings are precisely the localizations of all the rings at their maximal left denominators sets.

*Example.* Let  $A$  be a simple ring. Then  $Q_l(A)$  is a left localization maximal ring and  $Q_r(A)$  is a right localization maximal ring (by Theorem 2.1.(4) and Theorem 2.5).

*Example.* A division ring is a (left and right) localization maximal ring. More generally, a simple Artinian ring (i.e. the matrix algebra over a division ring) is a (left and right) localization maximal ring.

#### Left (non-)localizable elements of a ring.

**Lemma 2.6** [4] *Let  $S \in \text{Den}_l(R, \mathfrak{a})$  (respectively,  $S \in \text{Den}(R, \mathfrak{a})$ ),  $\sigma : R \rightarrow S^{-1}R$ ,  $r \mapsto \frac{r}{1}$ , and  $G := (S^{-1}R)^*$  be the group of units of the ring  $S^{-1}R$ . Then  $S' := \sigma^{-1}(G) \in \text{Den}_l(R, \mathfrak{a})$  (respectively,  $S' := \sigma^{-1}(G) \in \text{Den}(R, \mathfrak{a})$ ).*

*Definition, [3].* An element  $r$  of a ring  $R$  is called a *left localizable element* if there exists a left denominator set  $S$  of  $R$  such that  $r \in S$  (and so the element  $\frac{r}{1} \neq 0$  is invertible in the ring  $S^{-1}R$ ), equivalently, if there exists a left denominator set  $T$  of  $R$  such that the element  $\frac{r}{1}$  is invertible in the ring  $T^{-1}R$  (Lemma 2.6). The set of left localizable elements is denoted  $\mathcal{L}_l(R)$ .

Clearly,

$$\mathcal{L}_l(R) = \bigcup_{S \in \text{max.Den}_l(R)} S. \quad (5)$$

Similarly, a *right localizable element* is defined and let  $\mathcal{L}_r(R)$  be the set of right localizable elements of the ring  $R$ . The elements of the *set of left and right localizable elements*,

$$\mathcal{L}_{l,r}(R) = \mathcal{L}_l(R) \cap \mathcal{L}_r(R),$$

are called *left and right localizable elements*. An element  $r \in R$  is called a *localizable element* if there exists a (left and right) denominator set  $S \in \text{Den}(R)$  such that  $r \in S$ , equivalently, if there exists a (left and right) denominator set  $T \in \text{Den}(R)$  such that the element  $\frac{r}{1}$  is invertible in the ring  $T^{-1}R$  (Lemma 2.6). The set of all localizable elements of the ring  $R$  is denoted by  $\mathcal{L}(R)$ . Clearly,

$$\mathcal{L}(R) \subseteq \mathcal{L}_{l,r}(R).$$

The sets

$$\mathcal{NL}_l(R) := R \setminus \mathcal{L}_l(R), \quad \mathcal{NL}_r(R) := R \setminus \mathcal{L}_r(R), \quad \mathcal{NL}_{l,r}(R) := R \setminus \mathcal{L}_{l,r}(R), \quad \mathcal{NL}(R) := R \setminus \mathcal{L}(R),$$

are called the *sets of left non-localizable; of right non-localizable; of left and right non-localizable; of non-localizable elements*, respectively. The elements of these sets are called correspondingly (eg, an element  $r \in \mathcal{NL}_l(R)$  is called a *left non-localizable element*).

**Lemma 2.7** 1.  $\mathfrak{l}_R \cap \mathcal{L}_l(R) = \emptyset$ .

2.  $\mathfrak{I}_R \subseteq \mathcal{NL}_l(R)$ .

*Proof.* 1. If  $s \in \mathcal{L}_l(R)$  then  $s \in S$  for some  $S \in \max.\text{Den}_l(R)$  (by (5)), and so  $\frac{s}{1} \neq 0$  in  $S^{-1}R$ . Therefore,  $s \notin \mathfrak{I}_R$ .

2. Statement 2 follows from statement 1 as  $\mathcal{NL}_l(R) = R \setminus \mathcal{L}_l(R)$ .  $\square$

*Definition, [3].* For an arbitrary ring  $R$ , the intersection

$$\mathcal{C}_l(R) := \bigcap_{S \in \max.\text{Den}_l(R)} S$$

is called the set of *completely left localizable elements* of  $R$  and an element of the set  $\mathcal{C}_l(R)$  is called a *completely left localizable element*.

### The maximal left quotient rings of a finite direct product of rings.

**Theorem 2.8** [3] *Let  $R = \prod_{i=1}^n R_i$  be a direct product of rings  $R_i$ . Then for each  $i = 1, \dots, n$ , the map*

$$\max.\text{Den}_l(R_i) \rightarrow \max.\text{Den}_l(R), \quad S_i \mapsto R_1 \times \cdots \times S_i \times \cdots \times R_n, \quad (6)$$

*is an injection. Moreover,  $\max.\text{Den}_l(R) = \prod_{i=1}^n \max.\text{Den}_l(R_i)$  in the sense of (6), i.e.*

$$\max.\text{Den}_l(R) = \{S_i \mid S_i \in \max.\text{Den}_l(R_i), i = 1, \dots, n\},$$

$S_i^{-1}R \simeq S_i^{-1}R_i$ ,  $\text{ass}_R(S_i) = R_1 \times \cdots \times \text{ass}_{R_i}(S_i) \times \cdots \times R_n$ . The core of the left denominator set  $S_i$  in  $R$  coincides with the core  $S_{i,c}$  of the left denominator set  $S_i$  in  $R_i$ , i.e.

$$(R_1 \times \cdots \times S_i \times \cdots \times R_n)_c = 0 \times \cdots \times S_{i,c} \times \cdots \times 0.$$

**Finite direct products of localization maximal rings.** The next theorem describes properties of finite direct products of left localization maximal rings. It is used in the proof of Theorem 2.11 which is an explicit characterization of all such rings.

**Theorem 2.9** *Let  $R_1, \dots, R_n$  be left localization maximal rings and  $R := R_1 \times \cdots \times R_n$ . Then*

1.  $\max.\text{Den}_l(R) = \{S_i \mid i = 1, \dots, n\}$  and  $S_i := R_1 \times \cdots \times R_i^* \times \cdots \times R_n$  where  $R_i^*$  is a the group of units of the ring  $R_i$ .
2.  $\text{ass}(S_i) = R_1 \times \cdots \times 0 \times \cdots \times R_n$ ,  $i = 1, \dots, n$  where 0 is on  $i$ 'th place;  $\text{ass}(S_i) + \text{ass}(S_j) = R$  for all  $i \neq j$ .
3.  $S_i^{-1}R \simeq R_i \simeq R/\text{ass}(S_i)$ ,  $i = 1, \dots, n$ . The map  $\sigma_i : R \rightarrow S_i^{-1}R$ ,  $r \mapsto \frac{r}{1}$ , is equal to the map  $R \rightarrow R/\text{ass}(S_i)$ ,  $r \mapsto r + \text{ass}(S_i)$ .
4.  $\mathfrak{I}_R = 0$ .
5. The natural ring homomorphism

$$\sigma := \prod_{I=1}^n \sigma_i : R \rightarrow \prod_{I=1}^n S_i^{-1}R, \quad r \mapsto \left(\frac{r}{1}, \dots, \frac{r}{1}\right) = (r_1, \dots, r_n),$$

*is an isomorphism.*

6.  $\mathcal{C}_l(R) = \bigcap_{i=1}^n S_i = R^*$  where  $R^* = R_1^* \times \cdots \times R_n^*$  is the group of units of the ring  $R$ .
7. The set  $\mathcal{L}_l(R) = \bigcup_{i=1}^n S_i$  of left localizable elements of the ring  $R$  contains precisely the elements  $(r_1, \dots, r_n) \in R$  such that  $r_i \in R_i^*$  for some  $i$ .
8. The set  $\mathcal{NL}_l(R)$  of left non-localizable elements of the ring  $R$  is equal to  $\prod_{i=1}^n R_i \setminus R_i^*$ , the direct product of the sets of left non-localizable elements  $\mathcal{NL}_l(R_i) = R_i \setminus R_i^*$  of the rings  $R_i$ .

*Proof.* 1. Statement 1 is a particular case of Theorem 2.8.

2-5. For  $i = 1, \dots, n$ ,

$$S_i \in \max.\text{Den}_l(R), \quad \text{ass}(S_i) = R_1 \times \cdots \times 0 \times \cdots \times R_n, \quad S_i^{-1}R \simeq R_i \simeq R/\text{ass}(S_i).$$

and the localization map  $\sigma_i$  is equal to the map  $R \rightarrow R/\text{ass}(S_i)$ ,  $r \mapsto r + \text{ass}(S_i)$ . Clearly,  $\text{ass}(S_i) + \text{ass}(S_j) = R$  for all  $i \neq j$  and  $\mathfrak{l}_R \subseteq \bigcap_{i=1}^n \text{ass}(S_i) = 0$ , and so  $\mathfrak{l}_R = 0$ . The ring homomorphism  $\sigma$  is an isomorphism. So, statements 2-5 have been proved.

6. Statement 6 follows from statement 1.

7. Trivial.

8. Statement 8 follows from statement 1.  $\square$

**Corollary 2.10** *Let  $R = \prod_{i=1}^n R_i$  be a direct product of left localization maximal rings  $R_i$ . Then every nonzero element of the ring  $R$  is left localizable iff the ring  $R_1, \dots, R_n$  are division rings.*

*Proof.* Every nonzero element of the ring  $R$  is left localizable iff  $\{0\} = \mathcal{NL}(R) = \prod_{i=1}^n R_i \setminus R_i^*$  (Theorem 2.9.(8)) iff  $R_i \setminus R_i^* = \{0\}$  for all  $i$  iff the ring  $R_1, \dots, R_n$  are division rings.  $\square$

Corollary 2.10 is a particular case of a criterion (Theorem 3.5) for a ring with finitely many maximal left denominator sets to be a left localizable ring.

**Characterization of finite products of left localization maximal rings.** The next theorem is a criterion for a ring to be isomorphic to a finite direct product of left localization maximal rings.

**Theorem 2.11** *A ring  $R$  is a direct product of  $n$  left localization maximal rings iff the following conditions hold:*

1.  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$ .
2.  $\mathfrak{l}_R = 0$ .
3.  $\text{ass}(S_i) + \text{ass}(S_j) = R$  for all  $i \neq j$ .
4. The factor rings  $R/\text{ass}(S_i)$ ,  $i = 1, \dots, n$ , are left localization maximal rings.

In this case,

(a) for each  $i = 1, \dots, n$ , the ring  $R_i := S_i^{-1}R$  is a left localization maximal ring such that the homomorphism  $\sigma_i : R \rightarrow R_i$ ,  $r \mapsto \frac{r}{1}$ , is an epimorphism with  $\ker(\sigma_i) = \text{ass}(S_i)$ , and so  $R_i \simeq R/\text{ass}(S_i)$ .

(b) The map  $\sigma := \prod_{i=1}^n \sigma_i : R \rightarrow \prod_{i=1}^n R_i$  is an isomorphism.

(c) Let us identify the ring  $R$  with  $\prod_{i=1}^n R_i$  via  $\sigma$ . Then  $S_i = R_1 \times \cdots \times R_i^* \times \cdots \times R_n = \sigma_i^{-1}(R_i^*)$  where  $R_i^* = (R/\text{ass}(S_i))^*$  is the group of units of the ring  $R_i$ , and for each  $i = 1, \dots, n$ ,  $R_i^* = \sigma_i(S_i)$ .

(d)  $\text{ass}(S_i) = R_1 \times \cdots \times 0 \times \cdots \times R_n$  for  $i = 1, \dots, n$  where 0 is on  $i$ 'th place.

*Proof.* ( $\Rightarrow$ ) Theorem 2.9.

( $\Leftarrow$ ) By (2) and conditions 1 and 2, the map  $\sigma$  is a ring monomorphism since

$$\ker(\sigma) = \bigcap_{i=1}^n \text{ass}(S_i) = \mathfrak{l}_R = 0.$$

The ring homomorphism  $\sigma_i$  yields a monomorphism  $R/\text{ass}(S_i) \rightarrow S_i^{-1}R = R_i$  which is necessarily an isomorphism, by condition 4. Clearly,  $\sigma_i : R \rightarrow R/\text{ass}(S_i)$ ,  $r \mapsto r + \text{ass}(S_i)$ .

Conditions 1 and 2,  $\bigcap_{i=1}^n \text{ass}(S_i) = 0$  and  $\text{ass}(S_i) + \text{ass}(S_j) = R$  for all  $i \neq j$ , imply that the map  $\sigma$  is an isomorphism, i.e. the ring  $R \simeq R_1 \times \cdots \times R_n$  is a direct product of  $n$  left localization maximal rings.

By Theorem 2.9.(1),  $S_i = R_1 \times \cdots \times R_i^* \times \cdots \times R_n = \sigma_i^{-1}(R_i^*)$  and  $\text{ass}(S_i) = R_1 \times \cdots \times 0 \times \cdots \times R_n$ . Then  $R_i^* = \sigma_i(S_i)$  since  $\sigma_i : R \rightarrow R/\text{ass}(S_i)$ ,  $r \mapsto r + \text{ass}(S_i)$ .  $\square$

### 3 Left localizable rings and their characterization

The aim of this section is to introduce a new class of rings, *the class of left localizable rings*, consider their properties, and give their characterizations (Theorem 3.5, Theorem 3.8 and Theorem 3.9) provided they have only finitely many maximal left denominator sets. In particular, we prove that a ring with finitely many maximal left denominator sets is a left localizable ring iff it is a semiprime left Goldie ring such that its left quotient ring is a direct product of division rings (Theorem 3.9). For a semiprime left Goldie ring  $R$ , it is proved that  $\max.\text{Den}_l(R) < \infty$ . Moreover, elements of  $\max.\text{Den}_l(R)$  are found (Corollary 3.13).

*Definition.* A ring  $R$  is called a *left localizable ring* if the set of left localizable elements  $\mathcal{L}_l(R)$  is the largest possible, i.e.  $\mathcal{L}_l(R) = R \setminus \{0\}$ ; equivalently, for every nonzero element  $r \in R$  there is a left denominator set  $S = S(r) \in \text{Den}_l(R)$  such that  $r \in S$ ; equivalently, for every nonzero element  $r \in R$  there is a left denominator set  $S' = S'(r)$  such that the element  $\frac{r}{1} \in S'^{-1}R$  is a unit of the ring  $S'^{-1}R$  (Lemma 2.6).

Similarly, right localizable; left and right localizable; and localizable rings are defined, i.e. where the sets  $\mathcal{L}_r(R)$ ;  $\mathcal{L}_{l,r}(R)$ ; and  $\mathcal{L}(R)$  are equal to  $R \setminus \{0\}$ , respectively. A localizable ring is also called a *two-sided localizable ring*.

**Theorem 3.1** *Let  $R = \prod_{i=1}^n R_i$  be a direct product of rings  $R_i$ . The ring  $R$  is a left localizable ring iff the rings  $R_i$  are so.*

*Proof.* The proof is an easy corollary of Theorem 2.8 that states that  $\max.\text{Den}_l(\prod_{i=1}^n R_i) = \prod_{i=1}^n \max.\text{Den}_l(R_i)$ .

( $\Rightarrow$ ) Suppose that the ring  $R$  is a left localizable ring. We have to show that the rings  $R_i$  are so. Each ring  $R_i$  is a subring of  $R$ . Let  $r_i \in R_i$  be a nonzero element. The ring  $R$  is a left localizable ring. So,  $r_i \in S_i$  for some  $S_i \in \max.\text{Den}_l(R)$ . Then  $S_i \in \max.\text{Den}_l(R_i)$ , by Theorem 2.8.

( $\Leftarrow$ ) Suppose that the rings  $R_i$  are left localizable rings. Let  $r = (r_1, \dots, r_n) \in R$  be a nonzero element. Then  $0 \neq r_i \in R_i$  for some  $i$ . The ring  $R_i$  is a left localizable ring, and so  $r_i \in S_i$  for some  $S_i \in \max.\text{Den}_l(R_i)$ . By Theorem 2.8,  $S_i \in \max.\text{Den}_l(R)$ . Therefore,  $R$  is a left localizable ring.  $\square$

**Lemma 3.2** *1. If a ring  $R$  is a left (respectively, right; left and right; two-sided) localizable then  $\iota_R = 0$  (respectively,  $r.\text{rad}(R) = 0$ ;  $l.r.\text{rad}(R) = 0$ ;  $\text{rad}(R) = 0$ ).*

*2. Let a ring  $R = \prod_{i=1}^n R_i$  be a direct product of left localization maximal rings  $R_i$ . Then  $R$  is a left localizable ring iff the rings  $R_i$  are division rings.*

*Proof.* 1. Statements are easy corollaries of Lemma 2.7.

2. Corollary 2.10  $\square$

**Properties of the maximal left quotient rings of a ring.** The next theorem describes various properties of the maximal left quotient rings of a ring, in particular, their groups of units and their largest left quotient rings.

**Theorem 3.3** [2] *Let  $S \in \max.\text{Den}_l(R)$ ,  $A = S^{-1}R$ ,  $A^*$  be the group of units of the ring  $A$ ;  $\mathfrak{a} := \text{ass}(S)$ ,  $\pi_{\mathfrak{a}} : R \rightarrow R/\mathfrak{a}$ ,  $a \mapsto a + \mathfrak{a}$ , and  $\sigma_{\mathfrak{a}} : R \rightarrow A$ ,  $r \mapsto \frac{r}{1}$ . Then*

1.  $S = S_{\mathfrak{a}}(R)$ ,  $S = \pi_{\mathfrak{a}}^{-1}(S_0(R/\mathfrak{a}))$ ,  $\pi_{\mathfrak{a}}(S) = S_0(R/\mathfrak{a})$  and  $A = S_0(R/\mathfrak{a})^{-1}R/\mathfrak{a} = Q_l(R/\mathfrak{a})$ .
2.  $S_0(A) = A^*$  and  $S_0(A) \cap (R/\mathfrak{a}) = S_0(R/\mathfrak{a})$ .
3.  $S = \sigma_{\mathfrak{a}}^{-1}(A^*)$ .
4.  $A^* = \langle \pi_{\mathfrak{a}}(S), \pi_{\mathfrak{a}}(S)^{-1} \rangle$ , i.e. the group of units of the ring  $A$  is generated by the sets  $\pi_{\mathfrak{a}}(S)$  and  $\pi_{\mathfrak{a}}^{-1}(S) := \{\pi_{\mathfrak{a}}(s)^{-1} \mid s \in S\}$ .



5.  $A^* = \{\pi_{\mathfrak{a}}(s)^{-1}\pi_{\mathfrak{a}}(t) \mid s, t \in S\}$ .

6.  $Q_l(A) = A$  and  $\text{Ass}_l(A) = \{0\}$ . In particular, if  $T \in \text{Den}_l(A, 0)$  then  $T \subseteq A^*$ .

**Lemma 3.4** *Let  $R$  be a ring and  $S \in \text{max.Den}_l(R)$ . Then the ring  $S^{-1}R$  is a division ring iff  $R = S \cup \text{ass}(S)$ .*

*Remark.* For any left Ore set  $T$ ,  $T \cap \text{ass}(T) = \emptyset$ .

*Proof.* Let  $\sigma : R \rightarrow S^{-1}R$ ,  $r \mapsto \frac{r}{1}$ , and  $(S^{-1}R)^*$  be the group of units of the ring  $S^{-1}R$ . By Theorem 3.3.(3),  $S = \sigma^{-1}((S^{-1}R)^*)$  for all  $S \in \text{max.Den}_l(R)$ .

( $\Rightarrow$ ) Let  $s \in R$ . Suppose that  $s \notin \text{ass}(S)$ , we have to show that  $s \in S$ . Clearly,  $\frac{s}{1} \neq 0$  in  $S^{-1}R$ . Therefore,  $\frac{s}{1} \in (S^{-1}R)^*$  since  $S^{-1}R$  is a division ring. Then  $s \in \sigma^{-1}((S^{-1}R)^*) = S$ .

( $\Leftarrow$ ) If  $R = S \cup \text{ass}(S)$  then this union is a disjoint union since  $S \cap \text{ass}(S) = \emptyset$ . In particular,  $S + \text{ass}(S) \subseteq S$ . Let  $\frac{r}{1}$  (where  $r \in R$ ) be a nonzero element of the ring  $S^{-1}R$ . Then  $r \notin \text{ass}(S)$ , and so  $r \in S$ , i.e.  $\frac{r}{1}$  is a unit in the ring  $S^{-1}R$ . Therefore, the ring  $S^{-1}R$  is a division ring.  $\square$

**Characterization of left localizable rings.** The next theorem gives a characterization of left localizable rings with finite number of maximal left denominator sets.

**Theorem 3.5** *Let  $R$  be a ring with  $\text{max.Den}_l(R) = \{S_1, \dots, S_n\}$ . Let  $\mathfrak{a}_i := \text{ass}(S_i)$ ;  $\sigma_i : R \rightarrow R_i := S_i^{-1}R$ ,  $r \mapsto \frac{r}{1} = r_i$ ; and  $\sigma := \prod_{i=1}^n \sigma_i : R \rightarrow \prod_{i=1}^n R_i$ ,  $r \mapsto (r_1, \dots, r_n)$ . The following statements are equivalent.*

1. The ring  $R$  is a left localizable ring.
2.  $\mathfrak{l}_R = 0$  and the rings  $R_1, \dots, R_n$  are division rings.
3. The homomorphism  $\sigma$  is an injection and the rings  $R_1, \dots, R_n$  are division rings.

*Proof.* The rings  $R_i$  are left localization maximal (Theorem 3.3.(6)).

(2  $\Leftrightarrow$  3) Statements 2 and 3 are equivalent since  $\ker(\sigma) = \bigcap_{i=1}^n \text{ass}(S_i) = \mathfrak{l}_R$ .

(1  $\Leftarrow$  2) Since  $\mathfrak{l}_R = 0$ , the map  $\sigma$  (see (2)) is a ring monomorphism. So, every element  $r$  is a unique  $n$ -tuple  $(r_1, \dots, r_n)$  where  $r_i \in R_i$ . The rings  $R_1, \dots, R_n$  are division rings. Let  $r = (r_i) \in R$  be a nonzero element of  $R$ , then  $r_i \neq 0$  for some  $i$ , and so  $\sigma_i(r) = r_i \in R_i^*$ . Therefore, every nonzero element of the ring is left localizable.

(1  $\Rightarrow$  2) Suppose that every nonzero element of the ring  $R$  is left localizable, i.e.  $\mathcal{NL}(R) = \{0\}$ . By Lemma 2.7.(2),  $\mathfrak{l}_R = 0$ , and so, by (2), the map  $\sigma$  is a ring monomorphism. We identify the ring  $R$  with its image  $\sigma(R)$  in the ring  $\prod_{i=1}^n R_i$ , i.e.  $r = (r_1, \dots, r_n)$  where  $r_i = \sigma_i(r)$ .

If  $n = 1$  then  $\sigma_1 : R \rightarrow S_1^{-1}R$ ,  $r \mapsto \frac{r}{1}$ , is a monomorphism, and so  $\text{ass}(S_1) = 0$ . By the assumption  $S_1 = \mathcal{L}_l(R) = R \setminus \{0\}$ , i.e.  $S_1^{-1}R$  is a division ring.

Let  $n \geq 2$ . By Proposition 2.4, the ideals  $\mathfrak{a}_i := \text{ass}(S_i)$ ,  $i = 1, \dots, n$ , are non-comparable, i.e.

$$\mathfrak{a}_i \not\subseteq \mathfrak{a}_j, \quad i \neq j. \quad (7)$$

In particular,  $\mathfrak{a}_i \neq 0$  for all  $i$ . The proof consists of several steps.

*Step 1: For all  $i$ ,*

$$S_i \cap \bigcap_{j \neq i} \mathfrak{a}_j \neq \emptyset.$$

Fix  $r \in S_i$ . In particular,  $r = (r_1, \dots, r_n)$  with  $r_i \in R_i^*$ , and vice versa, i.e. if  $r_i \in R_i^*$  then  $r \in \sigma_i^{-1}(R_i^*) = S_i$ . Fix a nonzero element, say  $r \in S_i$ , such that the  $n$ -tuple  $r = (r_1, \dots, r_n)$  has the largest possible number of zeros, say  $m = n - s$  for some  $s$ . We claim that  $s = 1$ . Up to order, we may assume that  $i = 1$ ,

$$r = (r_1, \dots, r_s, 0, \dots, 0)$$

and  $r_j \neq 0$  for  $j = 1, \dots, s$ , and  $r_1 \in R_1^*$ . Suppose that  $s \neq 1$ , we seek a contradiction. Then, for all elements  $a_s \in \mathfrak{a}_s$ ,

$$\sigma(a_s)\sigma(r) = (\sigma_1(a_s)r_1, \dots, \sigma_{s-1}(a_s)r_{s-1}, 0, \dots, 0).$$

By the maximality of  $m$  we must have  $\sigma(a_s)\sigma(r) = 0$ . In particular,  $\sigma_1(a_s)r_1 = 0$ , and so  $\sigma_1(a_s) = 0$  since  $r_1 \in R^*$ , i.e.  $\mathfrak{a}_s \subseteq \mathfrak{a}_1$ , a contradiction, (Proposition 2.4). Therefore,  $s = 1$ , and  $\sigma(r) \in S_1 \cap \bigcap_{j \neq 1} \mathfrak{a}_j \neq 0$ . The proof of Step 1 is complete.

*Step 2: For each  $i$ , the ring  $R_i$  is a division ring.*

Without loss of generality we may assume that  $i = 1$ , then  $S_1 \cap \bigcap_{j=2}^n \mathfrak{a}_j \neq \emptyset$  (Step 1). Fix an element  $s \in S_1 \cap \bigcap_{j=2}^n \mathfrak{a}_j$ . Then  $\sigma(s) = (s_1, 0, \dots, 0)$  and  $s_1 \in R_1^*$  (since  $s \in S_1$  and  $R_1 = S_1^{-1}R$ ). Suppose that the ring  $R_1$  is not a division ring, we seek a contradiction. Then, by Lemma 3.4,  $R_1 \neq S_1 \cup \mathfrak{a}_1$ . Fix an element  $r \in R \setminus (S_1 \cup \mathfrak{a}_1)$ . Then  $\sigma(r) = (r_1, \dots, r_n)$  with  $S_1^{-1}R \ni r_1 \neq 0$  (since  $r \notin \mathfrak{a}_1$ ). Then

$$\sigma(sr) = (s_1r_1, 0, \dots, 0) \in \prod_{i=1}^n R_i$$

with  $s_1r_1 \neq 0$  since  $s_1 \in R_1^*$  and  $r_1 \neq 0$ . Therefore,  $sr \notin \bigcup_{j=2}^n S_j$  (since  $sr \in \bigcap_{j=2}^n \mathfrak{a}_j$ , by the choice of  $s$ ). Then necessarily  $sr \in S_1$  since  $sr \in R \setminus \{0\} = \bigcup_{i=1}^n S_i$  (the ring  $R$  is a left localizable ring). Then  $\sigma_1(r) = \sigma_1(s)^{-1}\sigma_1(sr) \in R_1^*$ , and so  $r \in \sigma_1^{-1}(R_1^*) = S_1$ , by Theorem 3.3.(3). This contradicts to the fact that  $r \in R \setminus (S_1 \cup \mathfrak{a}_1)$ . Therefore,  $R_1$  is a division ring. The proof of Step 2 and the theorem is complete.  $\square$

**Corollary 3.6** *Let  $R$  be a left localizable ring with  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$  where  $n \geq 2$ . Then, for all  $i = 1, \dots, n$ ,  $S_i \cap \bigcap_{j \neq i} \mathfrak{a}_j = (\bigcap_{j \neq i} \mathfrak{a}_j) \setminus \{0\} \neq \emptyset$  where  $\mathfrak{a}_i := \text{ass}(S_i)$ .*

*Proof.* By Step 1 of the proof of Theorem 3.5,  $S_i \cap \bigcap_{j \neq i} \mathfrak{a}_j \neq \emptyset$ . The equality in the corollary follows from the fact that  $\bigcap_{i=1}^n \mathfrak{a}_j = 0$  and  $R_i = S_i \cup \mathfrak{a}_i$  is a disjoint union (Lemma 3.4, since  $R_i$  is a division ring, by Theorem 3.5). Then  $(\bigcap_{j \neq i} \mathfrak{a}_j) \setminus \{0\} \neq \emptyset$ .  $\square$

**Proposition 3.7** *Let  $R$  be a left localizable ring with  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$  where  $n \geq 2$ . We keep the notation of Theorem 3.5. Let  $C'_i := (\bigcap_{j \neq i} \mathfrak{a}_j) \setminus \{0\}$ . Then*

1.  $C' := C'_1 + \dots + C'_n \in \text{Den}_l(R, 0)$  and  $C'^{-1}R \simeq R_1 \times \dots \times R_n$ .
2. For  $i = 1, \dots, n$ ,  $C'_i \in \text{Den}_l(R, \mathfrak{a}_i)$  and  $C'^{-1}_i R = R_i$ .

*Proof.* Clearly, all  $C'_i \neq \emptyset$  (Corollary 3.6);  $C'_i C'_j = 0$  for all  $i \neq j$  (since  $C'_i C'_j \subseteq \bigcap_{k=1}^n \mathfrak{a}_k = 0$ , Theorem 3.5), and  $\sigma(C'_i) \subseteq 0 \times \dots \times 0 \times R_i^* \times 0 \times \dots \times 0$  since  $C'_i = S_i \cap \bigcap_{j \neq i} \mathfrak{a}_j$  (Corollary 3.6). Therefore, the set  $C'$  consists of regular elements of the ring  $R$  (since the map  $\sigma : R \rightarrow \prod_{i=1}^n R_i$  is a monomorphism) and is obviously multiplicatively closed. Let us identify the set  $C'_i$  with its image in the division ring  $R_i$  via  $\sigma$ . Then  $\text{ass}(C'_i) = \mathfrak{a}_i$  (since  $C'_i S_i \subseteq C'_i$ ). Notice that  $C'_i \cup \{0\} = \bigcap_{j \neq i} \mathfrak{a}_j$  is an ideal of the ring  $R$ . Fix an element  $c_i \in C'_i$ , then each element  $s^{-1}r \in R_i$ , where  $s \in S_i$  and  $r \in R$ , can be written as a left fraction

$$s^{-1}r = s^{-1}c_i^{-1}c_i r = (c_i s)^{-1}c_i r, \text{ where } c_i s \in C'_i \text{ and } c_i r \in C'_i \cup \{0\}.$$

Therefore,  $C'_i \in \text{Den}_l(R, \mathfrak{a}_i)$  and  $C'^{-1}_i R = R_i$ , and statement 2 holds. So, any element  $\alpha$  of the ring  $\prod_{i=1}^n R_i$  can be written as  $\alpha = (c_1^{-1}r_1, \dots, c_n^{-1}r_n)$  where  $c_i \in C'_i$  and  $r_i \in C'_i \cup \{0\}$ , and so

$$\alpha = (c_1^{-1}r_1, \dots, c_n^{-1}r_n) = (c_1 + \dots + c_n)^{-1}(r_1 + \dots + r_n)$$

where  $c := c_1 + \dots + c_n \in C'$  and  $r := r_1 + \dots + r_n \in (\sum_{i=1}^n C'_i) \cup \{0\} \subseteq R$ . So, every element  $\alpha$  of the ring  $\prod_{i=1}^n R_i$  can be written as  $\alpha = c^{-1}r$  for some elements  $c \in C'$  and  $r \in R$ . Since the set  $C'$  consists of regular elements, the fact above ( $\alpha = c^{-1}r$ ) is equivalent to the fact that  $C'$  is a left Ore set, and so  $C' \in \text{Den}_l(R, 0)$ . Clearly,  $C'^{-1}R = R_1 \times \dots \times R_n$ .  $\square$

The sets  $C'_i$  in Theorem 3.7 have remarkable properties:  $C'_i \subseteq S_i$ ,  $C'^{-1}_i R = S_i^{-1}R$  and  $\text{ass}(c'_i) = \text{ass}(S_i)$  for all elements  $c'_i \in C'_i$  where  $c'_i : R \rightarrow R, r \mapsto c'_i r$  (Theorem 4.5). In fact, the set  $C'_i$  is the core of  $S_i$  (Theorem 4.5).

The next theorem is a useful criterion for a ring  $R$  to be a left localizable ring with  $|\max.\text{Den}_l(R)| < \infty$ .

**Theorem 3.8** *Let  $R$  be a ring. The following statements are equivalent.*

1. *The ring  $R$  is a left localizable ring with  $|\max.\text{Den}_l(R)| < \infty$ .*
2. *There are left denominator sets  $S'_1, \dots, S'_n \in \text{Den}_l(R)$  such that the rings  $R_i := S_i'^{-1}R$ ,  $i = 1, \dots, n$ , are division rings and the map*

$$\sigma := \prod_{i=1}^n \sigma_i : R \rightarrow \prod_{i=1}^n R_i, \quad r \mapsto (\sigma_1(r), \dots, \sigma_n(r)),$$

*is an injection where  $\sigma_i : R \rightarrow R_i$ ,  $r \mapsto \frac{r}{1}$ .*

*If one of the equivalent conditions holds and none of the rings  $R_i$  can be dropped with preserving the injectivity of  $\sigma$  then  $n = |\max.\text{Den}_l(R)|$ , and  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$  where  $S_i = \sigma_i^{-1}(R_i^*)$ .*

*Proof.* (1  $\Rightarrow$  2) Theorem 3.5.

(2  $\Leftarrow$  1) Suppose that statement 2 holds and we assume that none of the rings  $R_i$  can be dropped, i.e. for each  $i = 1, \dots, n$ ,  $\bigcap_{j \neq i} \mathfrak{a}_j \neq 0$  where  $\mathfrak{a}_i := \text{ass}(S'_i)$ . We are going to prove that the implication (2  $\Leftarrow$  1) holds and that  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$ .

Every division ring is a left localization maximal ring. By (Theorem 3.12.(3), [2]),  $S_i \in \max.\text{Den}_l(R)$ . Clearly,  $S'_i \subseteq \sigma_i^{-1}(R_i^*) = S_i$  and the natural ring homomorphism  $R_i = S_i'^{-1}R \rightarrow S_i^{-1}R$ ,  $\frac{r}{1} \rightarrow \frac{r}{1}$ , is an isomorphism, and so  $\mathfrak{a}_i = \text{ass}(S_i)$ . We identify the rings  $R_i$  and  $S_i^{-1}R$  via the isomorphism above. Since  $S_i \in \max.\text{Den}_l(R)$  and  $S_i^{-1}R$  is a division ring, we see that (by Lemma 3.4)  $R = S_i \coprod \mathfrak{a}_i$ , a disjoint union. Then

$$S_i \cap \bigcap_{j \neq i} \mathfrak{a}_j = \left( \bigcap_{j \neq i} \mathfrak{a}_j \right) \setminus \{0\} \neq \emptyset$$

since  $\bigcap_{i=1}^n \mathfrak{a}_i = \ker(\sigma) = 0$ . This fact together with the fact that  $R = S_i \coprod \mathfrak{a}_i$  for  $i = 1, \dots, n$ , imply that the sets  $S_1, \dots, S_n$  are distinct.

Finally, we claim that  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$ . Suppose that this is not true, that is there exists a set  $S \in \max.\text{Den}_l(R)$  distinct from the sets  $S_1, \dots, S_n$ , we seek a contradiction. For each element  $i = 1, \dots, n$ , by using the disjoint union  $R = S_i \coprod \mathfrak{a}_i$  and the fact that  $S \neq S_i$ , we can find an element, say  $t_i$  such that  $t_i \in S \cap \mathfrak{a}_i$  (notice that  $S \not\subseteq S_i$  since  $S, S_i \in \max.\text{Den}_l(R)$ ). Then  $\sigma_i(t_i) = 0$ , and so  $\sigma(t) = 0$  where  $t := t_1 \cdots t_n \in S$ . On the other hand,  $\sigma(t) \neq 0$  since  $t \in S$ , a contradiction.  $\square$

**A left localizable ring  $R$  with  $|\max.\text{Den}_l(R)| < \infty$  is a semiprime left Goldie ring such that  $Q_{l,cl}(R)$  is a direct product of division rings, and vice versa.**

**Theorem 3.9** *Let  $R$  be a ring. The following statements are equivalent.*

1. *The ring  $R$  is a left localizable ring with  $n := |\max.\text{Den}_l(R)| < \infty$ .*
2.  *$Q_{l,cl}(R) = R_1 \times \cdots \times R_n$  where  $R_i$  are division rings.*
3. *The ring  $R$  is a semiprime left Goldie ring with  $\text{udim}(R) = |\text{Min}(R)| = n$  where  $\text{Min}(R)$  is the set of minimal prime ideals of the ring  $R$ .*
4.  *$Q_l(R) = R_1 \times \cdots \times R_n$  where  $R_i$  are division rings.*

*Proof.* (2  $\Leftrightarrow$  3) It is known and easy to prove (use Goldie's Theorem).

(1  $\Rightarrow$  2) By Proposition 3.7,  $C'^{-1}R = R_1 \times \cdots \times R_n$  where  $R_i$  are division rings. In particular,  $C'^{-1}R$  is a semisimple (Artinian) ring. Clearly,  $C' \subseteq S_0(R)$  where  $S_0(R)$  is the largest regular left Ore set of the ring  $R$ . Then

$$Q_l(R) = S_0(R)^{-1}R = Q_l(C'^{-1}R) = Q_l(R_1 \times \cdots \times R_n) = R_1 \times \cdots \times R_n.$$

Since  $Q_l(R)$  is a semisimple ring,  $Q_{l,cl}(R) = Q_l(R)$ , by (Corollary 2.10, [2]).

(2  $\Rightarrow$  1) The map  $\sigma : R \rightarrow Q_{l,cl}(R) = R_1 \times \cdots \times R_n$ ,  $r \mapsto \frac{r}{1}$ , is a monomorphism. Any division ring is a left localization maximal ring. By Proposition 3.10.(4),  $\max.\text{Den}_l(R) = \{T_1, \dots, T_n\}$  where  $T_i = \sigma^{-1}(R_1 \times \cdots \times R_i^* \times \cdots \times R_n)$  and  $R_i \simeq T_i^{-1}R$ . Since  $R_1, \dots, R_n$  are division rings and  $\sigma = \prod_{i=1}^n \sigma_i : R \rightarrow \prod_{i=1}^n T_i^{-1}R$ ,  $r \mapsto (\sigma_1(r), \dots, \sigma_n(r))$ , is a monomorphism where  $\sigma_i : R \rightarrow T_i^{-1}R$ ,  $r \mapsto \frac{r}{1}$ , the ring  $R$  is a left localizable ring (Theorem 3.5).

(3  $\Leftrightarrow$  4) This equivalence is a particular case of a more general result (Corollary 2.10, [2]): *Let  $R$  be a ring. Then  $Q_l(R)$  is a semisimple ring iff  $Q_{l,cl}(R)$  is a semisimple ring, and in this case  $Q_l(R) = Q_{l,cl}(R)$ .  $\square$*

**A bijection between  $\max.\text{Den}_l(R)$  and  $\max.\text{Den}_l(Q_l(R))$ .**

**Proposition 3.10** [3] *Let  $R$  be a ring,  $S_l$  be the largest regular left Ore set of the ring  $R$ ,  $Q_l := S_l^{-1}R$  be the largest left quotient ring of the ring  $R$ , and  $\mathcal{C}$  be the set of regular elements of the ring  $R$ . Then*

1.  $S_l \subseteq S$  for all  $S \in \max.\text{Den}_l(R)$ . In particular,  $\mathcal{C} \subseteq S$  for all  $S \in \max.\text{Den}_l(R)$  provided  $\mathcal{C}$  is a left Ore set.
2. Either  $\max.\text{Den}_l(R) = \{\mathcal{C}\}$  or, otherwise,  $\mathcal{C} \notin \max.\text{Den}_l(R)$ .
3. The map

$$\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(Q_l), \quad S \mapsto SQ_l^* = \{c^{-1}s \mid c \in S_l, s \in S\},$$

is a bijection with the inverse  $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$  where  $\sigma : R \rightarrow Q_l$ ,  $r \mapsto \frac{r}{1}$ , and  $SQ_l^*$  is the sub-semigroup of  $(Q_l, \cdot)$  generated by the set  $S$  and the group  $Q_l^*$  of units of the ring  $Q_l$ , and  $S^{-1}R = (SQ_l^*)^{-1}Q_l$ .

4. If  $\mathcal{C}$  is a left Ore set then the map

$$\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(Q), \quad S \mapsto SQ^* = \{c^{-1}s \mid c \in \mathcal{C}, s \in S\},$$

is a bijection with the inverse  $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$  where  $\sigma : R \rightarrow Q$ ,  $r \mapsto \frac{r}{1}$ , and  $SQ^*$  is the sub-semigroup of  $(Q, \cdot)$  generated by the set  $S$  and the group  $Q^*$  of units of the ring  $Q$ , and  $S^{-1}R = (SQ^*)^{-1}Q$ .

**Theorem 3.11** *Let  $R$  be a ring,  $T \in \text{Den}_l(R, 0)$  and  $\sigma : R \rightarrow T^{-1}R$ ,  $r \mapsto \frac{r}{1}$ . Then*

1.  $T \subseteq S$  for all  $S \in \max.\text{Den}_l(R)$ .
2. The map

$$\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(T^{-1}R), \quad S \mapsto \tilde{S},$$

is a bijection with the inverse  $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$  where  $\tilde{S}$  is the multiplicative monoid generated in the ring  $T^{-1}R$  by  $\sigma(S)$  and  $\sigma(T)^{-1} := \{t^{-1} \mid t \in T\}$  and  $S^{-1}R \simeq \tilde{S}^{-1}(T^{-1}R)$ , and  $S^{-1}R \simeq \tilde{S}^{-1}(T^{-1}R)$ .

*Proof.* 1. Lemma 2.2.

2. Statement 2 follows from statement 1 and Proposition 3.4.(1), [2].  $\square$

**Theorem 3.12** *Let  $R$  be a ring,  $T \in \text{Den}_l(R, 0)$  be such that  $T^{-1}R = R_1 \times \cdots \times R_n$  is a direct product of left localization maximal rings; and  $\tau : R \rightarrow T^{-1}R$ ,  $r \mapsto \frac{r}{1} = (r_1, \dots, r_n)$ . Then  $\max.\text{Den}_l(R) = \{T_1, \dots, T_n\}$  where  $T_i = \tau^{-1}(S_i)$ ,  $S_i := R_1 \times \cdots \times R_i^* \times \cdots \times R_n$  and  $T_i^{-1}R \simeq R_i$ .*

*Proof.* Proposition follows from Theorem 3.11 and Theorem 2.9.(1).  $\square$

The next theorem shows that  $|\max.\text{Den}_l(R)| < \infty$  for all semiprime left Goldie rings.

**Corollary 3.13** *Let  $R$  be a semiprime left Goldie ring,  $\mathcal{C}$  be the set of regular element of the ring  $R$ ,  $Q_{l,cl} = \prod_{i=1}^n Q_i$  where  $Q_i$  are simple Artinian rings (by Goldie's Theorem),  $\sigma : R \rightarrow Q_{l,cl}$ ,  $r \mapsto \frac{r}{1}$ . Then*

1.  $C \subseteq S$  for all  $S \in \max.\text{Den}_l(R)$ .
2.  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$ ,  $\max.\text{Den}_l(Q_{l,cl}(R)) = \{S'_1, \dots, S'_n\}$  where  $S_i = \sigma^{-1}(S'_i)$  and  $S'_i := Q_1 \times \dots \times Q_i^* \times \dots \times Q_n$  for  $i = 1, \dots, n$ . The map

$$\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(Q_{l,cl}(R)), \quad S \mapsto \tilde{S},$$

*is a bijection with inverse  $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$  where  $\tilde{S}$  is the multiplicative monoid generated in  $Q_{l,cl}(R)$  by  $\sigma(S)$  and  $\sigma(\mathcal{C})^{-1} = \{c^{-1} \mid c \in \mathcal{C}\}$ .*

3.  $S_i^{-1}R \simeq Q_i$  for  $i = 1, \dots, n$ .

*Proof.* This is a particular case of Theorem 3.11 where  $T = C$ .  $\square$

## 4 The core of a left Ore set

The aim of this section is to establish several properties of the *core*  $S_c$  of a left Ore set  $S$  of a ring  $R$ , and when it exists (for example for a right Noetherian ring  $R$ ) it is a powerful tool in studying localizations (Theorem 4.2). A characterization of the core is given (Proposition 4.4). The core shares remarkable properties (Lemma 4.1, Theorem 4.2).

*Definition*, [3]. Let  $R$  be a ring and  $S$  be its left Ore set. The subset of  $S$ ,

$$S_c := \{s \in S \mid \ker(s \cdot) = \text{ass}(S)\}$$

is called the *core* of the left Ore set  $S$  where  $s \cdot : R \rightarrow R$ ,  $r \mapsto sr$ .

If  $S_c \neq \emptyset$  then  $S_c S_c \subseteq S_c$ , i.e. the core  $S_c$  is a multiplicative set. If  $\text{ass}(S) = 0$  then  $S_c = S$ .

**Lemma 4.1** *If  $S \in \text{Den}_l(R)$  and  $S_c \neq \emptyset$  then*

1.  $SS_c \subseteq S_c$ .
2. For any  $s \in S$  there exists an element  $t \in S$  such that  $ts \in S_c$ .

*Proof.* 1. Trivial.

2. Statement 2 follows directly from the left Ore condition: fix an element  $s_c \in S_c$ , then  $ts = rs_c \in S$  for some elements  $t \in S$  and  $r \in R$ . Since  $\text{ass}(S) \supseteq \ker(ts \cdot) = \ker(rs_c \cdot) \supseteq \ker(s_c \cdot) = \text{ass}(S)$ , i.e.  $\ker(ts \cdot) = \text{ass}(S)$ , we have  $ts \in S_c$ .  $\square$

**Theorem 4.2** *Suppose that  $S \in \text{Den}_l(R, \mathfrak{a})$  and  $S_c \neq \emptyset$ . Then*

1.  $S_c \in \text{Den}_l(R, \mathfrak{a})$ .
2. The map  $\theta : S_c^{-1}R \rightarrow S^{-1}R$ ,  $s^{-1}r \mapsto s^{-1}r$ , is a ring  $R$ -isomorphism. So,  $S_c^{-1}R \simeq S^{-1}R$ .

*Proof.* 1. By Lemma 4.1.(1),  $S_c S_c \subseteq S_c$ , that is the set  $S_c$  is a multiplicative set. By Lemma 4.1.(2),  $S_c \in \text{Ore}_l(R)$ : for any elements  $s_c \in S_c$  and  $r \in R$ , there are elements  $s \in S$  and  $r' \in R$  such that  $sr = r's_c$  (since  $S \in \text{Ore}_l(R)$ ). By Lemma 4.1.(2),  $s'_c := ts \in S_c$  for some  $t \in S$ . Then  $s'_c r = tr's_c$ .

If  $rs_c = 0$  for some elements  $r \in R$  and  $s_c \in S_c$  then  $sr = 0$  for some element  $s \in S$  (since  $S \in \text{Den}_l(R)$ ). By Lemma 4.1.(2),  $s'_c := ts \in S_c$  for some element  $t \in S$ , hence  $s'_c r = 0$ . Therefore,  $S_c \in \text{Den}_l(R, \mathfrak{a})$ .

2. By statement 1 and the universal property of left Ore localization, the map  $\theta$  is a well-defined monomorphism. By Lemma 4.1.(2),  $\theta$  is also a surjection: let  $s^{-1}r \in S^{-1}R$ , and  $s_c := ts \in S_c$  for some element  $t \in S$  (Lemma 4.1.(2)). Then

$$s^{-1}r = s^{-1}t^{-1}tr = (ts)^{-1}tr = s_c^{-1}tr. \quad \square$$

Let  $S$  be a left Ore set of a ring  $R$ . The set of right annihilators  $\ker(S \cdot) := \{\ker(s \cdot) \mid s \in S\}$  is a poset with respect to  $\subseteq$ . Let

$$\max(S) := \{s \in S \mid \ker(s \cdot) \text{ is a maximal element of } \ker(S \cdot)\}.$$

The set  $\max(S)$  is a non-empty set for a ring with ACC on right annihilators (for example, when  $R$  is a right Noetherian ring).

**Lemma 4.3** *Let  $S \in \text{Ore}_l(R)$ . Then  $\text{ass}(S) = \bigcup_{s \in S} \ker(s \cdot) = \sum_{s \in S} \ker(s \cdot)$ .*

*Proof.* Let  $U$  and  $\Sigma$  stand for the union and the sum respectively. Then  $\text{ass}(S) \supseteq \Sigma \supseteq U = \text{ass}(S)$ , and so  $\text{ass}(S) = \Sigma = U$ .  $\square$

**Proposition 4.4** *Let  $S$  be a left Ore set of a ring  $R$ . Then  $S_c = \max(S)$ .*

*Proof.* If  $S_c \neq \emptyset$  then  $S_c = \max(S)$ , by Lemma 4.3.

Suppose that  $S_c = \emptyset$ . To finish the proof we have to show that  $\max(S) = \emptyset$ . Suppose that this is not the case, we seek a contradiction. Let  $\mathfrak{a} = \text{ass}(S)$ . Fix  $s \in \max(S)$ . Then  $\mathfrak{b} := \ker(s \cdot) \subseteq \mathfrak{a}$ . We claim that  $\mathfrak{b} = \mathfrak{a}$ , i.e.  $s \in S_c$ , a contradiction. To prove the claim we have to show that for any element  $t \in S$ ,  $\ker(t \cdot) \subseteq \mathfrak{b}$  (see Lemma 4.3). Notice that  $\ker(s' \cdot) = \mathfrak{b}$  for all  $s' \in S$ , by the maximality of  $\ker(s \cdot)$ . Using the left Ore condition for  $S$ , we have the equality  $s_1 s = r t$  for some elements  $s_1 \in S$  and  $r \in R$ . Hence,

$$\mathfrak{b} = \ker(s_1 s \cdot) = \ker(r t \cdot) \supseteq \ker(t \cdot),$$

as required.  $\square$

The next theorem gives an explicit description of the cores of maximal left denominator sets of a left localizable ring provided  $|\max.\text{Den}_l(R)| < \infty$ .

**Theorem 4.5** *Let  $R$  be a left localizable ring such that  $\max.\text{Den}_l(R) = \{S_1, \dots, S_n\}$ .*

1. *If  $n = 1$  then  $S_{1,c} = S_1 = R \setminus \{0\}$ .*
2. *If  $n \geq 2$  then  $S_{i,c} = S_i \cap \bigcap_{j \neq i} \mathfrak{a}_j$  where  $\mathfrak{a}_j = \text{ass}(S_j)$ .*

*Proof.* We keep the notation of Theorem 3.5 and its proof.

1. If  $n = 1$  then  $R$  is a domain  $S_1 = R \setminus \{0\}$  (Theorem 3.5), and so  $S_{1,c} = S_1 = R \setminus \{0\}$ .

2. Suppose that  $n \geq 2$ . For each  $i = 1, \dots, n$ ,  $C'_i := S_i \cap \bigcap_{j \neq i} \mathfrak{a}_j \neq \emptyset$ , by Corollary 3.6. By Proposition 3.7.(2),  $C'_i \subseteq S_{i,c}$ . By Theorem 3.5, the map

$$\sigma := \prod_{i=1}^n \sigma_i : R \rightarrow \prod_{i=1}^n R_i, \quad r \mapsto (r_1, \dots, r_n),$$

is a ring monomorphism. Since  $R_i = S_i^{-1}R$  is a division ring (Theorem 3.5),  $R = S_i \coprod \mathfrak{a}_i$ , by Lemma 3.4. Clearly,  $\mathfrak{a}_i = \{r = (r_1, \dots, r_n) \mid r_i = 0\}$ . Each element  $s'$  of the set  $C'_i$  has the form  $(0, \dots, 0, s'_i, 0, \dots, 0)$  with  $s'_i \in R_i^*$ . Then clearly,  $s' \mathfrak{a}_i = 0$  and so  $C'_i \subseteq S_{i,c}$ . To show that the equality  $S'_i = S_{i,c}$  holds it suffices to show that every element  $s \in S_i \setminus C'_i$  does not belong to  $S_{i,c}$ . Fix  $s$  such that  $s \in S_i \setminus C'_i$ . Then there is an index, say  $j$ , such that  $j \neq i$  and such that  $s_j \neq 0$  in  $s = (s_1, \dots, s_n)$ . Then  $s \cdot C'_j \neq 0$  (Corollary 3.6) but  $C'_j \subseteq \mathfrak{a}_i$ . Therefore, the element  $s$  does not belong to  $S_{i,c}$ .  $\square$

### Acknowledgements

The work is partly supported by the Royal Society and EPSRC.

## References

- [1] V. V. Bavula, The algebra of integro-differential operators on an affine line and its modules, *J. Pure Appl. Algebra*, **217** (2013) 495-529. Arxiv:math.RA: 1011.2997.
- [2] V. V. Bavula, The largest left quotient ring of a ring, Arxiv:math.RA:1101.5107.
- [3] V. V. Bavula, New criteria for a ring to have a semisimple left quotient ring. Arxiv:math.RA:1303.0859.
- [4] V. V. Bavula, Left localizations of left Artinian rings, Arxiv:math.RA:1405.0214.

Department of Pure Mathematics  
University of Sheffield  
Hicks Building  
Sheffield S3 7RH  
UK  
email: v.bavula@sheffield.ac.uk